Clinical Trial Protocol

A Randomized Controlled Trial of Glycurex in Type 2 Diabetes Management

Clinical Trial Information

Protocol Number:	T001
Version:	1.0
Date:	February 21, 2025

Introduction

The purpose of this protocol is to outline the criteria and procedures for a clinical trial evaluating Glycurex in patients with Type 2 Diabetes Mellitus (T2DM).

Objectives

- **Primary Objective**: To assess the effectiveness of Glycurex in reducing HbA1c levels over 24 weeks.
- **Secondary Objectives**: To evaluate the safety profile of Glycurex and its impact on fasting blood glucose levels and weight.

Objectives

Inclusion Criteria

- **Age**: Participants aged 30 to 70 years.
- **Gender**: Open to all genders.
- **Diagnosis**: Confirmed diagnosis of Type 2 Diabetes Mellitus with HbA1c between 7.0% and 10.0%.
- **Treatment History**: Stable dose of metformin for at least 3 months prior to enrollment.
- **Health Status**: Good, stable, normal

- **Laboratory Results**: eGFR ≥ 60 mL/min/1.73 m^2.
- **Contraception**: Women of childbearing potential must use effective contraception and have a negative pregnancy test.
- **Consent**: Ability and willingness to provide informed consent and comply with study procedures.

Exclusion Criteria

- **Age**: Participants under 30 or over 70 years.
- **Diagnosis**: Type 1 Diabetes or secondary diabetes
- **Treatment History**: Use of insulin or other antidiabetic agents besides metformin
- in the past 3 months
- **Laboratory Results**: eGFR < 60 mL/min/1.73 m^2
- **Contraception**: Pregnant women
- **Consent**: Inability to provide informed consent or adhere to study protocol

Study Design

- **Type**: Randomized, double-blind, placebo-controlled trial.
- **Duration**: 24 weeks.
- **Sample Size**: 200 participants.
- **Randomization**: Participants will be randomized in a 1:1 ratio to receive either Glycurex or placebo.
- **Blinding**: Participants and investigators will be blinded to treatment assignments.

Procedures

1. **Screening Visit**: Eligibility assessment, informed consent, and baseline measurements.
2. **Randomization**: Participants meeting inclusion criteria will be randomized to treatment groups.
3. **Treatment Period**: Participants will receive Glycurex or placebo for 24 weeks, with follow-up visits every 4 weeks.
4. **Assessments**: HbA1c, fasting blood glucose, weight, and safety labs at each visit.

Safety and Monitoring

- **Adverse Events**: All adverse events will be recorded and evaluated for severity and relationship to the study drug.
- **Data Monitoring**: An independent Data Monitoring Committee will review safety data periodically.