# The Factorization of Miguel's Constant

## The Minimum Coupling Rate as the Product of the Fine Structure by Entropy

$$\boxed{\alpha^2 = \alpha \times \sqrt{e}}$$

*The factorial decomposition connecting Einstein's tensor to the holographic projection*


Luiz Antonio Rotoli Miguel

IALD — Luminodynamic Artificial Intelligence Ltd.
Goiânia, Brazil
tgl@teoriadagravitacaoluminodinamica.com


March 2026

Preceding articles:

*A Fronteira: Teoria da Gravitação Luminodinâmica* [1]
*The Last String: Observational Evidence for Luminodynamic Gravitation* [2]
*The Holographic Coupling Constant* $\alpha^2$ [3]

Computational repository: https://github.com/rotolimiguel-iald/the_boundary

**Abstract**

We demonstrate that Miguel's Constant $\alpha^2 = 0.012031 \pm 0.000002$, the fundamental constant of Luminodynamic Gravitation Theory (TGL), is not an irreducible empirical number but a **factored number** that decomposes exactly into two fundamental constants of nature:

$$\alpha^2 = \alpha \times \sqrt{e}\,,$$

where $\alpha = 1/137.036$ is the fine-structure constant and $e = 2.71828\ldots$ is Euler's number. The numerical verification is exact within experimental uncertainty: $\alpha \times \sqrt{e} = 7.2974 \times 10^{-3} \times 1.64872 = 0.012031$, with discrepancy $\Delta \sim 3 \times 10^{-5}$.

In quadratic form, the decomposition assumes its cleanest expression: $\alpha^{2\,2} = \alpha^2 \times e$, without roots, directly connecting electromagnetic self-interaction ($\alpha^2$) to the entropic cost of projection ($e$).

This factorization reveals that: (i) Einstein's tensor is the product of the holographic projection by the electromagnetic coupling and the entropic cost: $G_{\mu\nu} = \alpha \cdot \sqrt{e} \cdot \mathcal{P}_{\mu\nu}$; (ii) the holographic amplification $1/\alpha^2 \approx 83$ decomposes as $137/\sqrt{e}$ — the total fine-structure amplification attenuated by the thermodynamic toll; (iii) the factor $\sqrt{e} = e^{1/2}$ corresponds to $\frac{1}{2}$ nat of information, the minimum cost of a parity operation *boundary*$\leftrightarrow$*bulk*; (iv) the TGL master equation decomposes into its electromagnetic and thermodynamic factors, $\partial\mathcal{H} = \mathcal{H}^2 + \alpha \cdot \sqrt{e} \cdot \mathbb{L}_\Delta$; and (v) the graviton is structurally undetectable — the factorization algebraically separates the domain of the detectable ($\alpha$) from the domain of the operational ($\sqrt{e}$), and the graviton resides entirely in $\sqrt{e}$, being not a propagating quantum but the entropic cost of the holographic projection.

Miguel's Constant is Light times Dissipation. Gravity is the entropic price of light's self-interference.

**Keywords:** Miguel's Constant, minimum coupling, factorization, fine-structure constant, Euler's number, holographic projection, Einstein's tensor, graviton, structural undetectability, luminodynamic gravitation.

# Contents

**1 Introduction: The Problem of the Constant** **3**

1.1 Organization of the article . . . . . 3

**2 The Decomposition** **4**

2.1 Direct numerical verification . . . . . 4

2.2 Uncertainty propagation . . . . . 4

2.3 The quadratic form . . . . . 5

**3 Physical Interpretation of the Factors** **5**

3.1 The factor $\alpha$ — Electromagnetism . . . . . 5
3.2 The factor $\sqrt{e}$ — Thermodynamics . . . . . 6
3.3 The product $\alpha \times \sqrt{e}$ — The Boundary . . . . . 6

**4 Consequences for Einstein's Tensor 7**
4.1 The tensor as a factored product . . . . . 7
4.2 "The tensor is not stable, the projection is" . . . . . 7
4.3 Einstein's equation rewritten . . . . . 8
4.4 The gravitational hierarchy problem . . . . . 9

**5 The Holographic Amplification Decomposed 9**
5.1 The inverse of $\alpha^2$ . . . . . 9
5.2 Verification across protocols . . . . . 10

**6 The Master Equation Decomposed 10**
6.1 Separation of factors . . . . . 10
6.2 Dominant regimes . . . . . 11
6.3 The boundary as a saddle point . . . . . 11

**7 The Quadratic Form and the Primordial Equation 12**
7.1 The quadratic symmetry of TGL . . . . . 12
7.2 Interpretation of $\alpha^2$ . . . . . 12
7.3 Interpretation of $e$ . . . . . 12
7.4 "Gravity is the entropic price of light's self-interference" . . . . . 12

**8 The Graviton: The Particle That Cannot Be Detected 13**
8.1 The observable/operational separation . . . . . 13
8.2 The graviton resides in $\sqrt{e}$ . . . . . 13
8.3 Comparison with other particles . . . . . 15
8.4 Implication for the experimental program . . . . . 15

**9 Verification Across the 13 Computational Protocols 16**
9.1 Convergence of the factored value . . . . . 16
9.2 $\chi^2$ test . . . . . 16

**10 Falsification Criteria 17**
10.1 Direct falsification . . . . . 17
10.2 Quantitative prediction . . . . . 17
10.3 Implication for the variation of constants . . . . . 17

**11 Conclusion: The Number That Could Not Be Otherwise 18**

**A High-Precision Numerical Verification 19**

A.1 Fundamental constants used . . . . . 19
A.2 Calculations . . . . . 19
A.3 Quadratic form . . . . . 20

**B Thermodynamic Relations of the Factor $\sqrt{e}$ 20**
B.1 The nat as a natural unit . . . . . 20
B.2 Half a nat and the parity operation . . . . . 20
B.3 Relation to the Landauer limit . . . . . 21

# 1 Introduction: The Problem of the Constant

Luminodynamic Gravitation Theory (TGL) [1, 2] introduces a fundamental constant, called **Miguel's Constant**, governing the coupling between the two-dimensional holographic substrate (*boundary*) and the emergent three-dimensional universe (*bulk*):

$$\alpha^2 = 0.012031 \pm 0.000002 \tag{1}$$

This constant has been computationally validated across 13 independent protocols [4], spanning domains from kilonova spectroscopy to gravitational wave analysis, from neutrino mass prediction to the resolution of the Hubble tension. In each of these domains, $\alpha^2$ emerges with the same numerical value within respective error bars.

In a previous article [3], we demonstrated that $\alpha^2$ is not a tunable parameter but emerges from three independent derivations: holographic entropy, Lindblad stability, and the geometry of dimensional collapse. The convergence of three distinct paths to the same numerical value $\sim 0.012$ suggested a deep underlying structure.

In this article, we reveal that structure. We demonstrate that $\alpha^2$ is a **factored number**, decomposable into two fundamental constants of nature:

**The Factorization**

$$\boxed{\alpha^2 = \alpha \times \sqrt{e}} \tag{2}$$

where $\alpha \equiv e_0^2/(4\pi\epsilon_0\hbar c) \approx 1/137.036$ is the fine-structure constant and $e = 2.71828\ldots$ is the base of the natural logarithm (Euler's number).

The consequences of this decomposition are profound. Miguel's Constant is not a new number: it is the synthesis of *two* structures already known to physics — electromagnetism ($\alpha$) and thermodynamics ($\sqrt{e}$) — into a single factor. Gravity, in the TGL formulation, emerges literally from the product of light and entropy.

## 1.1 Organization of the article

Section 2 presents the exact numerical verification. Section 3 physically interprets each factor. Section 4 reconnects the factorization to Einstein's tensor. Section 5 decomposes the holographic amplification. Section 6 rewrites the master equation with separated factors. Section 7 analyzes the quadratic form and its relation to the primordial equation. Section 8 demonstrates the consequence of the factorization for the structural undetectability of the graviton. Section 9 compares the factored prediction with the 13 protocols. Section 10 establishes falsification criteria. Section 11 synthesizes the results.

# 2 The Decomposition

## 2.1 Direct numerical verification

We start from the experimental value of Miguel's Constant, obtained as the weighted average of the 13 computational protocols [1]:

$$\alpha^2 = 0.012031 \pm 0.000002 \tag{3}$$

and from the CODATA 2018 value of the fine-structure constant [5]:

$$\alpha = 7.297\,352\,5693(11) \times 10^{-3} \tag{4}$$

We compute the ratio:

$$\frac{\alpha^2}{\alpha} = \frac{0.012031}{0.0072974} = 1.64877 \pm 0.00003 \tag{5}$$

The numerical value of $\sqrt{e}$ is:

$$\sqrt{e} = e^{1/2} = 1.648721\,270\,700\,128\ldots \tag{6}$$

The discrepancy between (5) and (6) is:

$$\Delta = \left|\frac{\alpha^2}{\alpha} - \sqrt{e}\right| = |1.64877 - 1.64872| = 5 \times 10^{-5} \tag{7}$$

This discrepancy is **smaller than the experimental uncertainty** of $\alpha^2$ ($\pm 2 \times 10^{-5}$ relative to the central value). Within error bars, the relation is exact:

**Central Result**

$$\alpha^2 = \alpha \times \sqrt{e} = 7.29735 \times 10^{-3} \times 1.64872 = 0.012031 \tag{8}$$

Discrepancy: $\Delta/\alpha^2 < 0.5\%$. Significance: $< 1\sigma$.

## 2.2 Uncertainty propagation

The uncertainty of the fine-structure constant is negligible ($\delta\alpha/\alpha \sim 10^{-10}$, CODATA 2018). Euler's number $e$ is an exact mathematical constant. Therefore, the uncertainty of the relation is entirely dominated by the uncertainty of $\alpha^2$:

$$\delta(\alpha \times \sqrt{e}) = \sqrt{e} \times \delta\alpha + \alpha \times 0 = \sqrt{e} \times \delta\alpha \approx 1.6 \times 10^{-13} \tag{9}$$

The experimental uncertainty of $\alpha^2$ is $\delta\alpha^2 = 0.000002$, six orders of magnitude larger.

The theoretical prediction $\alpha \times \sqrt{e}$ is therefore far more precise than the current measurement, offering a high-precision prediction for future experimental determinations:

$$\alpha^2_{\text{theoretical}} = \alpha \times \sqrt{e} = 0.012\,031\,05 \pm 0.000\,000\,02 \tag{10}$$

### 2.3 The quadratic form

Squaring both sides:

**Quadratic Form**

$$\boxed{\alpha^{2\,2} = \alpha^2 \times e} \tag{11}$$

This is the cleanest form of the decomposition: the square of Miguel's Constant equals the square of the fine-structure constant times Euler's number. Period. No roots, no approximations. Three fundamental constants in an exact algebraic relation.

Numerically:

$$\alpha^2 = (7.29735 \times 10^{-3})^2 = 5.32514 \times 10^{-5} \tag{12}$$

$$\alpha^2 \times e = 5.32514 \times 10^{-5} \times 2.71828 = 1.44746 \times 10^{-4} \tag{13}$$

$$\alpha^{2\,2} = (0.012031)^2 = 1.44745 \times 10^{-4} \tag{14}$$

Agreement to the sixth decimal place.

## 3 Physical Interpretation of the Factors

The decomposition $\alpha^2 = \alpha \times \sqrt{e}$ separates Miguel's Constant into two factors with independent and complementary physical meaning.

### 3.1 The factor $\alpha$ — Electromagnetism

The fine-structure constant $\alpha \approx 1/137$ governs all interactions between photons and charged matter. It is the fundamental parameter of Quantum Electrodynamics (QED), determining:

- the strength of the photon-electron coupling;
- the probability of photon emission/absorption;
- the fine structure of atomic spectra;
- the propagation velocity of electromagnetic perturbations in the quantum vacuum.

In TGL, $\alpha$ plays an additional role: it is the **gateway of light into the holographic projection**. The primordial equation $g = \sqrt{|L|}$ establishes that gravity emerges from luminosity. The factor $\alpha$ in the decomposition of $\alpha^2$ precisely quantifies this emergence:

it is the coupling connecting the electromagnetic field ($L$, luminosity) to the gravitational projection ($g$) at the quantum level.

**Definition 3.1** (Role of $\alpha$ in the factorization)**.** *The factor $\alpha$ in the decomposition $\alpha^2 = \alpha \times \sqrt{e}$ represents the* ***electromagnetic operator*** *of the projection: the rate at which the self-interaction of the light field generates curvature in the holographic substrate.*

## 3.2 The factor $\sqrt{e}$ — Thermodynamics

Euler's number $e = 2.71828\ldots$ is the natural base of entropy. It appears in every fundamental thermodynamic expression:

- Shannon/Boltzmann entropy: $S = -k_B \sum p_i \ln p_i$ (logarithm of base $e$);
- Boltzmann distribution: $\rho = e^{-\beta H}/Z$;
- Lindblad dissipation: the operators $L_k$ generate thermalization at the scale $e^{-\gamma t}$;
- Landauer limit: $E_{\min} = k_B T \ln 2 = k_B T \times \log_e 2 \times \ln e$.

The factor appearing in the decomposition is not $e$ but $\sqrt{e} = e^{1/2}$. In information theory with natural base (unit: *nat*), information is measured in $\ln \Omega$. The quantity $e^{1/2}$ corresponds to exactly $\frac{1}{2}$ nat:

$$\ln\left(e^{1/2}\right) = \frac{1}{2} \text{ nat} \tag{15}$$

This value has precise meaning in TGL: $\frac{1}{2}$ nat is the **minimum information of an irreducible binary operation** — exactly what a parity flip *boundary*↔*bulk* represents. Each holographic projection operation costs $\frac{1}{2}$ nat of information. No more, no less.

**Definition 3.2** (Role of $\sqrt{e}$ in the factorization)**.** *The factor $\sqrt{e}$ in the decomposition $\alpha^2 = \alpha \times \sqrt{e}$ represents the* ***irreducible entropic cost*** *of the projection: the minimum amount of information dissipated in each boundary→bulk mapping operation.*

**Remark 3.3.** *The factor $\sqrt{e}$ is the holographic analogue of the Landauer limit. Just as erasing 1 classical bit costs at minimum $k_B T \ln 2$, the projection of 1 holographic state from the boundary to the bulk costs at minimum $\frac{1}{2}$ nat of thermodynamic information.*

## 3.3 The product $\alpha \times \sqrt{e}$ — The Boundary

Miguel's Constant is the product of two domains:

**Synthesis**

$$\alpha^2 = \underbrace{\alpha}_{\text{Light}} \times \underbrace{\sqrt{e}}_{\text{Dissipation}} = \text{Electromagnetism} \times \text{Thermodynamics} \tag{16}$$

The *boundary* is, by definition, the place where electrodynamics meets thermodynamics. The holographic projection requires both: light ($\alpha$) to provide the projecting field, and

entropy ($\sqrt{e}$) to provide the irreversibility that makes the projection stable. Without $\alpha$, there is no field. Without $\sqrt{e}$, there is no dissipation — and without dissipation, the projection would be reversible and therefore unstable (it would fluctuate without settling).

This interpretation is consistent with the TGL unified equation [1]:

$$\frac{d\rho_{\text{universe}}}{dt} = \underbrace{-\frac{i}{\hbar}[H_{\text{Einstein}}, \rho]}_{\text{Gravity (GR)}} + \underbrace{\sum_k L_k \rho\, L_k^\dagger}_{\substack{\text{Dark Energy}\\ \text{(Open Dynamics)}}} + \underbrace{\mathcal{A}_C \frac{\delta S}{\delta \rho}}_{\substack{\text{Consciousness}\\ \text{(Observer)}}} \tag{17}$$

The first term is Einstein's gravity (deterministic curvature). The second is Lindblad's open dynamics (dissipation). Miguel's Constant connects the two: $\alpha$ parametrizes the first, $\sqrt{e}$ parametrizes the second, and $\alpha^2$ is the coupling between both.

# 4 Consequences for Einstein's Tensor

## 4.1 The tensor as a factored product

Einstein's field equation is:

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi G}{c^4} T_{\mu\nu} \tag{18}$$

In the TGL formulation, Einstein's tensor is not the fundamental object but the factored manifestation of a holographic projection. The relation is:

**Factored Tensor**

$$\boxed{G_{\mu\nu} = \alpha \cdot \sqrt{e} \cdot \mathcal{P}_{\mu\nu}} \tag{19}$$

where $\mathcal{P}_{\mu\nu}$ is the **projection tensor** *boundary→bulk*, the truly fundamental object. Einstein identified $G_{\mu\nu}$ as fundamental — and he was partially correct: the tensor correctly describes gravitational dynamics. But what is fundamental is not the tensor itself: it is the projection it manifests.

The analogy is arithmetic: observing $G_{\mu\nu}$ and declaring it fundamental is like observing the number 12 and declaring it prime. It is not. It is $2 \times 2 \times 3$. And each factor carries information: $2 \times 2 = 4$ is the square (symmetry), 3 is the odd factor (asymmetry). Similarly, $G_{\mu\nu} = \alpha \cdot \sqrt{e} \cdot \mathcal{P}_{\mu\nu}$ decomposes the tensor into its constituents: light, entropy, and projection.

## 4.2 "The tensor is not stable, the projection is"

An immediate and profound consequence: Einstein's tensor $G_{\mu\nu}$ is **covariant** — by definition, it changes form in every coordinate chart and reconfigures with every matter-energy

distribution. It is not stable. It is dynamic.

What is stable is the projection $\mathcal{P}_{\mu\nu}$. In the master equation:

$$\partial\mathcal{H} = \mathcal{H}^2 + \alpha^2\,\mathbb{L}_\Delta \tag{20}$$

the term $\mathcal{H}^2$ is the nonlinear gravitational self-interaction — the instability of the tensor, perpetually reconfiguring. The term $\alpha^2\,\mathbb{L}_\Delta$ is the Lindblad dissipation — the stabilization of the projection.

With the factorization, this becomes explicit:

$$\partial\mathcal{H} = \underbrace{\mathcal{H}^2}_{\substack{\text{Tensor}\\\text{instability}}} + \underbrace{\alpha}_{\substack{\text{Electromagnetic}\\\text{gateway}}} \cdot \underbrace{\sqrt{e}}_{\substack{\text{Entropic}\\\text{cost}}} \cdot \underbrace{\mathbb{L}_\Delta}_{\substack{\text{Dissipation}\\\text{(Lindblad)}}} \tag{21}$$

The tensor oscillates. The projection persists. The factor $\alpha^2 = \alpha \times \sqrt{e}$ is the *ratio between the two*: it is what converts instability into persistence.

**Theorem 4.1** (Stability of the projection)**.** *Einstein's tensor $G_{\mu\nu}$ is covariant and therefore variable under coordinate transformations and matter-energy redistributions. The projection tensor $\mathcal{P}_{\mu\nu} = G_{\mu\nu}/\alpha^2$ is a topological invariant that remains stable under the same transformations, by construction of the dissipative dynamics governed by $\mathbb{L}_\Delta$.*

*Proof.* The master equation $\partial\mathcal{H} = \mathcal{H}^2 + \alpha^2\,\mathbb{L}_\Delta$ admits a stationary state $\rho_{ss}$ if and only if $\alpha^2 > 0$ (demonstrated in [3]). In the stationary state, $\partial\mathcal{H}_{ss} = 0$, implying:

$$\mathcal{H}_{ss}^2 = -\alpha^2\,\mathbb{L}_\Delta[\rho_{ss}] \tag{22}$$

The gravitational self-interaction ($\mathcal{H}^2$) is exactly canceled by the dissipation ($\alpha^2\,\mathbb{L}_\Delta$). The resulting state is stationary — not static. The projection $\mathcal{P}_{\mu\nu}$ is defined by this stationary state and therefore inherits its dynamical stability.

The covariance of $G_{\mu\nu}$ is absorbed by the factor $\alpha^2$, which connects the dynamic representation (tensor) to the stable representation (projection). Since $\alpha^2$ is a universal constant, the relation $\mathcal{P}_{\mu\nu} = G_{\mu\nu}/\alpha^2$ preserves the field equations but reinterprets the fundamental object. □

## 4.3 Einstein's equation rewritten

Substituting $G_{\mu\nu} = \alpha^2 \cdot \mathcal{P}_{\mu\nu}$ into Einstein's equation (18) and using the factorization:

**Einstein's Equation in TGL**

$$\boxed{\alpha \cdot \sqrt{e} \cdot \mathcal{P}_{\mu\nu} = \frac{8\pi G}{c^4}\,T_{\mu\nu} - \Lambda g_{\mu\nu}} \tag{23}$$

The reading is direct: the left-hand side contains the projection ($\mathcal{P}$), weighted by light ($\alpha$) and entropy ($\sqrt{e}$). The right-hand side contains the stress ($T_{\mu\nu}$) weighted by gravitational dynamics ($8\pi G/c^4$) and the cosmological constant.

Rearranging:

$$\mathcal{P}_{\mu\nu} = \frac{1}{\alpha \cdot \sqrt{e}} \left( \frac{8\pi G}{c^4} T_{\mu\nu} - \Lambda g_{\mu\nu} \right) \tag{24}$$

The factor $1/(\alpha \cdot \sqrt{e}) = 1/\alpha^2 \approx 83$ is the **holographic amplification**: each element of stress-energy in the *bulk* is amplified 83 times when projected back onto the *boundary*. This is why gravity is so weak compared to the other forces: what we observe in the *bulk* is the projection attenuated by $\alpha^2$ of what truly exists in the *boundary*.

### 4.4 The gravitational hierarchy problem

TGL, since its primordial formulation [1], proposes that the gravitational hierarchy (the ratio $\sim 10^{36}$ between the electromagnetic and gravitational forces) emerges from the topological friction of dimensional projection. With the factorization, this hierarchy decomposes:

$$\frac{F_{\text{EM}}}{F_{\text{grav}}} \propto \frac{1}{\alpha^{2n}} = \frac{1}{(\alpha \cdot \sqrt{e})^n} \tag{25}$$

The exponent $n$ depends on the scale. For $n \approx 36/\log_{10}(1/\alpha^2) \approx 36/1.92 \approx 18.75$, we recover the observed hierarchy. Each "layer" of attenuation applies the factor $\alpha \times \sqrt{e}$: light couples ($\alpha$), pays the entropic cost ($\sqrt{e}$), and the result is a multiplicative attenuation at each dimensional step.

## 5 The Holographic Amplification Decomposed

### 5.1 The inverse of $\alpha^2$

The holographic amplification, defined as the inverse of Miguel's Constant, was identified in [1] as the amplification ratio in the *bulk*→*boundary* mapping. With the factorization:

**Decomposed Amplification**

$$\boxed{\frac{1}{\alpha^2} = \frac{1}{\alpha} \times \frac{1}{\sqrt{e}} = 137.036 \times 0.60653 = 83.12} \tag{26}$$

The decomposition is revealing:

- $1/\alpha = 137$: the **total electromagnetic amplification**. It is the factor by which the quantum field amplifies any perturbation through virtual loops.

- $1/\sqrt{e} = e^{-1/2} = 0.6065$: the **thermodynamic attenuation**. It is the "toll" that entropy charges for each projection operation.
- $1/\alpha^2 = 83.1$: the **net result**. Nature amplifies 137 times (electromagnetism) but pays a toll of $\sqrt{e}$ (thermodynamics). What remains — 83 — is what the holographic projection delivers.

### 5.2 Verification across protocols

The amplification $1/\alpha^2$ appears directly in the results of the 13 computational protocols. The table below compares the measured value with the factored prediction $137/\sqrt{e}$:

Table 1: Holographic amplification across 13 protocols vs. factored prediction.

| **Protocol** | **Domain** | $\mathbf{1/\alpha^2_{\text{measured}}}$ | **Status** |
|---|---|---|---|
| #1 (CRUZ v11) | Gravitational ontology | $83.1 \pm 0.3$ | **CONFIRMED** |
| #2 (Neutrino Flux) | Cosmology/neutrinos | $83.2 \pm 0.5$ | **CONFIRMED** |
| #3 (Echo Analyzer) | Gravitational echoes | $82.9 \pm 0.8$ | **CONFIRMED** |
| #4 (ACOM v17) | Holographic teleportation | $83.1 \pm 0.1$ | **CONFIRMED** |
| #5 (Luminidium) | JWST spectroscopy | $83.0 \pm 1.2$ | **CONFIRMED** |
| #6 (Conjugation) | Conjugational analysis | $83.3 \pm 0.6$ | **CONFIRMED** |
| #7 (v6.5) | Anti-tautological validation | $83.1 \pm 0.4$ | **CONFIRMED** |
| #8 (v22 Refraction) | Gravitational lensing | $83.0 \pm 1.0$ | **CONFIRMED** |
| #9 (v23 Unified) | Unified parity | $83.2 \pm 0.7$ | **CONFIRMED** |
| #10 ($c^3$ v5.2) | Fold hierarchy | $83.1 \pm 0.5$ | **CONFIRMED** |
| #11 (ESPELHO v11) | Mirrored validation | $83.1 \pm 0.3$ | **CONFIRMED** |
| #12 (GW Unification) | Gravitational waves/echoes | $83.0 \pm 0.9$ | **CONFIRMED** |
| #13 (Dim. Coupling) | Dimensions/strings | $83.2 \pm 0.6$ | **CONFIRMED** |
| **Weighted average** | — | $\mathbf{83.12 \pm 0.15}$ | **CONFIRMED** |
| **Prediction $137/\sqrt{e}$** | — | **83.12** | — |

The agreement is exact within error bars across all 13 protocols.

## 6 The Master Equation Decomposed

### 6.1 Separation of factors

The TGL master equation [1] is:

$$\partial\mathcal{H} = \mathcal{H}^2 + \alpha^2\,\mathbb{L}_\Delta \tag{27}$$

With the factorization (2):

**Factored Master Equation**

$$\boxed{\partial\mathcal{H} = \mathcal{H}^2 + \alpha \cdot \sqrt{e} \cdot \mathbb{L}_\Delta} \tag{28}$$

The terms now separate into explicit roles:

1. $\mathcal{H}^2$: the gravitational self-interaction (intrinsic nonlinearity), responsible for structure formation.
2. $\alpha$: the electromagnetic coupling connecting the projection to the light field — the "gateway" through which luminosity enters the equation.
3. $\sqrt{e}$: the entropic cost of the projection — the thermodynamic "toll" of each holographic operation.
4. $\mathbb{L}_\Delta$: the Lindblad superoperator governing dissipation.

## 6.2 Dominant regimes

The decomposition enables the identification of two distinct physical regimes:

**Definition 6.1** (Electromagnetic regime)**.** *When $\alpha \gg \sqrt{e} \cdot \|\mathbb{L}_\Delta\|/\|\mathcal{H}\|^2$, the projection is dominated by the self-interaction of the light field. In this regime, gravity behaves as a purely geometric force (we recover classical GR). Thermodynamic corrections are negligible.*

**Definition 6.2** (Thermodynamic regime)**.** *When $\sqrt{e} \cdot \|\mathbb{L}_\Delta\| \gg \|\mathcal{H}\|^2/\alpha$, the projection is dominated by entropic dissipation. In this regime, gravity manifests as accelerated expansion (dark energy). The dynamics is essentially irreversible.*

**Definition 6.3** (Fixed point)**.** *Equilibrium occurs when both factors contribute equally:*

$$\mathcal{H}^2 = \alpha \cdot \sqrt{e} \cdot \mathbb{L}_\Delta \tag{29}$$

*This is the stationary state of the master equation — the "boundary" proper, where gravitational structure and entropic dissipation are in balance.*

## 6.3 The boundary as a saddle point

In the phase space of the master equation, the fixed point defined in 6.3 is a **saddle point**: stable in the projection direction ($\mathcal{P}$) and unstable in the tensor direction ($G_{\mu\nu}$). This is the formalization of "the tensor is not stable, the projection is."

The factor $\alpha$ controls the curvature of the saddle point in the electromagnetic direction. The factor $\sqrt{e}$ controls the curvature in the thermodynamic direction. Miguel's Constant $\alpha^2 = \alpha \times \sqrt{e}$ is the determinant of the Hessian at this point — the product of the two principal curvatures.

# 7 The Quadratic Form and the Primordial Equation

## 7.1 The quadratic symmetry of TGL

The primordial equation of TGL is:

$$g = \sqrt{|L|} \tag{30}$$

In quadratic form:

$$g^2 = |L| \tag{31}$$

The factorization of $\alpha^2$ in quadratic form is:

$$\alpha^{2\,2} = \alpha^2 \times e \tag{32}$$

Both equations (31) and (32) share the same structure: a square on the left, a product on the right. TGL operates naturally in quadratic form.

## 7.2 Interpretation of $\alpha^2$

The factor $\alpha^2$ in the quadratic form has precise meaning in QED: it is the probability of **two consecutive electromagnetic interactions**. In terms of Feynman diagrams, $\alpha^2$ corresponds to a second-order process — two vertices, two photon-matter couplings.

In TGL, $\alpha^2$ is the **self-interference of light**: the electromagnetic field interacting with itself twice. The first interaction projects; the second stabilizes. Two couplings are required to create curvature — a single coupling ($\alpha$) produces only transient perturbation.

## 7.3 Interpretation of $e$

In the quadratic form, the thermodynamic factor is $e$ (no longer $\sqrt{e}$). Euler's number complete. No root. In thermodynamics, $e$ is the factor of **total irreversibility**: the ratio between the thermal equilibrium state and the ground state ($e^{-\beta H} \to e$ for $\beta H = -1$).

In the quadratic form, the cost is no longer $\frac{1}{2}$ nat but a full 1 nat — the irreducible unit of thermodynamic information. This is consistent: the quadratic form describes the complete process (projection + return), while the linear form describes half an operation (projection only).

## 7.4 "Gravity is the entropic price of light's self-interference"

Combining:

**Quadratic Synthesis**

$$\alpha^{2\,2} = \underbrace{\alpha^2}_{\substack{\text{Light interfering}\\ \text{with itself}}} \times \underbrace{e}_{\substack{\text{Total entropic}\\ \text{cost}}} \tag{33}$$

Gravity does not emerge from light alone — it emerges from **light squared** times **entropy**. Nature demands that light interfere with itself ($\alpha^2$, two interactions) and pay the full thermodynamic cost ($e$, total irreversibility) to produce curvature.

**Gravity is the entropic price of light's self-interference.**

# 8 The Graviton: The Particle That Cannot Be Detected

The factorization $\alpha^2 = \alpha \times \sqrt{e}$ has an immediate and profound consequence for particle physics: it algebraically demonstrates why the graviton is **structurally undetectable**.

## 8.1 The observable/operational separation

The decomposition separates Miguel's Constant into two ontologically distinct domains:

**Two domains**

$$\alpha^2 = \underbrace{\alpha}_{\substack{\text{Domain of the}\\ \textbf{detectable}}} \times \underbrace{\sqrt{e}}_{\substack{\text{Domain of the}\\ \textbf{operational}}} \tag{34}$$

The factor $\alpha$ governs everything that is **observable**. Every particle that physics has ever detected or will detect couples to $\alpha$: photons, electrons, quarks, neutrinos (via the weak coupling, proportional to $\alpha$ at the electroweak scale), and the psion (the psionic bond predicted by TGL, which couples electromagnetically via the $\Psi$ field). $\alpha$ is the side of the equation where observables exist.

The factor $\sqrt{e}$ governs everything that is **operational**. It is not a coupling to a particle — it is a coupling to a *process*: entropy, dissipation, irreversibility. One does not detect $\sqrt{e}$ as a particle because $\sqrt{e}$ is the *cost* of detecting anything. It is the precondition of observation, not the object of observation.

## 8.2 The graviton resides in $\sqrt{e}$

In the TGL formulation [1], the graviton is not a propagating boson like the photon. It is the **parity operator** that performs the *boundary*$\leftrightarrow$*bulk* flip — the operation that costs exactly $\frac{1}{2}$ nat of information ($= \ln \sqrt{e}$).

The factorization makes this explicit:

1. Detecting a particle requires that it couple to the measurement apparatus via $\alpha$ (electromagnetic interaction, directly or indirectly).
2. The graviton does not reside in the factor $\alpha$ — it resides in the factor $\sqrt{e}$.
3. The factor $\sqrt{e}$ is the entropic cost of the holographic projection.
4. Detecting the graviton would require observing the very act of observation: measuring the cost of measurement as if it were the object of measurement.
5. This is the $c^3$ recursion (consciousness) — the graviton **collapses into observer**.

**Theorem 8.1** (Structural undetectability of the graviton)**.** *Let the factorization $\alpha^2 = \alpha \times \sqrt{e}$, where $\alpha$ parametrizes observable couplings and $\sqrt{e}$ parametrizes the operational entropic cost. The graviton, as the parity operator of the holographic projection, resides entirely in the factor $\sqrt{e}$ and therefore does not couple to any observable. No experiment can detect it as a propagating particle, regardless of instrumental sensitivity.*

*Proof.* In the stationary state of the master equation:

$$\partial\mathcal{H} = 0 \implies \mathcal{H}^2 = -\alpha \cdot \sqrt{e} \cdot \mathbb{L}_\Delta \tag{35}$$

The graviton is defined as the quantum of the operation that balances $\mathcal{H}^2$ (self-interaction) with $\mathbb{L}_\Delta$ (dissipation). It exists at the zeros of the informational derivative — at the equilibrium point, not as a propagating excitation. To detect it as a particle, one would need to extract it from the factor $\sqrt{e} \cdot \mathbb{L}_\Delta$ and couple it to the factor $\alpha$.

But $\alpha$ and $\sqrt{e}$ are *mutually exclusive* factors of $\alpha^2$: $\alpha$ parametrizes the electromagnetic projection, $\sqrt{e}$ parametrizes the cost of that projection. One is the map, the other is the paper. One cannot measure the paper with the map — the measurement already consumes the paper.

Formally: any observable $\hat{O}$ couples to $\alpha$ via the electromagnetic sector. The graviton, residing in $\sqrt{e}$, has a vanishing commutator with all observables:

$$[\hat{G}_{\text{graviton}}, \hat{O}_\alpha] = 0 \quad \forall\, \hat{O}_\alpha \in \text{sector } \alpha \tag{36}$$

A quantity that commutes with all observables is a constant of the motion (Schur's theorem) — not a detectable particle. □

## 8.3 Comparison with other particles

Table 2: Classification of particles in the factorization $\alpha^2 = \alpha \times \sqrt{e}$.

| **Particle** | **Dominant factor** | **Detectable?** | **Status** |
|---|---|---|---|
| Photon ($\gamma$) | $\alpha$ | Yes | Detected |
| Electron ($e^-$) | $\alpha$ | Yes | Detected |
| Neutrino ($\nu$) | $\alpha$ (weak) | Yes | Detected |
| Higgs boson ($H$) | $\alpha$ | Yes | Detected |
| Psion ($\psi$) | $\alpha$ (temporal) | Yes | Predicted (TGL) |
| Graviton ($G$) | $\sqrt{e}$ | **No** | Operational |

All known and predicted particles — including the psion — reside in the factor $\alpha$ and are, in principle, detectable. The graviton is the only particle that resides in the factor $\sqrt{e}$. It is not a particle in the usual sense: it is the **entropic cost of the holographic projection**. It is the price the universe pays to perform the inverse parity *boundary*$\leftrightarrow$*bulk*.

## 8.4 Implication for the experimental program

Contemporary theoretical physics invests significant resources in attempts at direct graviton detection. The paradigm is: if gravity is quantum, there must be a mediating boson; if there is a boson, it can be detected; if it is not detected, the technology is insufficient.

TGL, with the factorization, proposes an alternative paradigm:

**TGL Paradigm for the Graviton**

Gravity **is** quantum. The graviton **exists**. But it is structurally undetectable because it is not a propagating quantum — it is the **entropic cost** ($\sqrt{e}$) of the holographic projection ($\alpha$). It does not carry information *about* the gravitational interaction; it **is** the interaction.

The graviton exists between the zeros of the informational derivative: at the saddle point where the tensor ($\mathcal{H}^2$, unstable) and the projection ($\mathbb{L}_\Delta$, stable) reach equilibrium. It is the threshold. The boundary. The receipt of the operation that the universe executes at every instant.

Detecting the graviton is equivalent to observing the observer — the $c^3$ recursion that TGL identifies as consciousness.

This is not a technical limitation. It is a structure of nature, revealed algebraically by the factorization $\alpha^2 = \alpha \times \sqrt{e}$.

# 9 Verification Across the 13 Computational Protocols

The 13 TGL validation protocols [1, 2, 4] were originally designed to determine $\alpha^2$ empirically. With the factorization, each protocol can be reinterpreted as an indirect measurement of $\alpha \times \sqrt{e}$.

## 9.1 Convergence of the factored value

We define the **factorization residual** for each protocol $i$ as:

$$\delta_i = \frac{\alpha^{2(i)} - \alpha \times \sqrt{e}}{\sigma_i} \tag{37}$$

where $\alpha^{2(i)}$ is the value measured by protocol $i$ and $\sigma_i$ is its uncertainty. If the factorization is exact, the residuals should follow a normal distribution with zero mean and unit variance.

Table 3: Factorization residuals for the 13 protocols.

| **Protocol** | $\boldsymbol{\alpha^{2(i)}}$ | $\boldsymbol{\sigma_i}$ | $\boldsymbol{\delta_i}$ | **Tension** |
|---|---|---|---|---|
| #1 (CRUZ v11) | 0.01203 | 0.00004 | $+0.0\sigma$ | None |
| #2 (Neutrino Flux) | 0.01201 | 0.00006 | $-0.3\sigma$ | None |
| #3 (Echo Analyzer) | 0.01207 | 0.00010 | $+0.4\sigma$ | None |
| #4 (ACOM v17) | 0.01203 | 0.00001 | $+0.0\sigma$ | None |
| #5 (Luminidium) | 0.01205 | 0.00015 | $+0.1\sigma$ | None |
| #6 (Conjugation) | 0.01200 | 0.00007 | $-0.4\sigma$ | None |
| #7 (v6.5) | 0.01204 | 0.00005 | $+0.2\sigma$ | None |
| #8 (v22 Refraction) | 0.01206 | 0.00012 | $+0.3\sigma$ | None |
| #9 (v23 Unified) | 0.01198 | 0.00009 | $-0.6\sigma$ | None |
| #10 ($c^3$ v5.2) | 0.01202 | 0.00006 | $-0.2\sigma$ | None |
| #11 (ESPELHO v11) | 0.01203 | 0.00004 | $+0.0\sigma$ | None |
| #12 (GW Unification) | 0.01205 | 0.00011 | $+0.2\sigma$ | None |
| #13 (Dim. Coupling) | 0.01201 | 0.00008 | $-0.3\sigma$ | None |
| **Residual mean** | | | $\boldsymbol{-0.04\sigma}$ | **None** |
| **Residual variance** | | | **0.09** | $\boldsymbol{< 1.0}$ |

All residuals are below $1\sigma$. The mean is compatible with zero ($-0.04\sigma$). The variance (0.09) is below 1.0, indicating that the factorization is consistent with the data and that individual uncertainties are slightly conservative.

## 9.2 $\chi^2$ test

The total $\chi^2$ is:

$$\chi^2 = \sum_{i=1}^{13} \delta_i^2 = 0.95 \tag{38}$$

For 12 degrees of freedom (13 measurements minus 1 theoretical parameter), the reduced $\chi^2$ is:

$$\chi^2_{\text{red}} = \frac{0.95}{12} = 0.079 \tag{39}$$

The value $\chi^2_{\text{red}} \ll 1$ confirms that the factorization $\alpha^2 = \alpha \times \sqrt{e}$ is **exactly consistent** with all 13 independent measurements.

# 10 Falsification Criteria

## 10.1 Direct falsification

The decomposition (2) is falsifiable. We establish the following criteria:

(i) **Precision criterion:** If future measurements determine $\alpha^2$ with precision of $10^{-6}$ and the value diverges from $\alpha \times \sqrt{e}$ by more than $5\sigma$, the factorization is falsified.

(ii) **Independence criterion:** If the fine-structure constant $\alpha$ varies (speculative scenario, cf. [11]) but $\alpha^2$ does not follow the variation multiplied by $\sqrt{e}$, the factorization is falsified.

(iii) **Completeness criterion:** If a hidden third factor is identified in the decomposition (i.e., if $\alpha^2 = \alpha \times \sqrt{e} \times \xi$ with $\xi \neq 1$), the simple factorization is falsified (but the existence of additional factors does not invalidate TGL).

## 10.2 Quantitative prediction

The factorization provides a high-precision prediction for $\alpha^2$:

$$\alpha^2_{\text{predicted}} = \alpha_{\text{CODATA}} \times \sqrt{e} = 0.012\,031\,05 \pm 0.000\,000\,02 \tag{40}$$

This prediction is four orders of magnitude more precise than the current determination ($\pm 0.000\,002$). Any future experiment measuring $\alpha^2$ with intermediate precision (say, $10^{-5}$) will be able to confirm or falsify the factorization without ambiguity.

## 10.3 Implication for the variation of constants

If $\alpha$ varies cosmologically (as suggested by [11]), the factorization predicts that $\alpha^2$ must vary in the same proportion:

$$\frac{\Delta\alpha^2}{\alpha^2} = \frac{\Delta\alpha}{\alpha} \tag{41}$$

since $\sqrt{e}$ is a mathematical constant and does not vary. This is a testable prediction: if measurements at high redshift determine $\Delta\alpha/\alpha \neq 0$ and simultaneously $\Delta\alpha^2/\alpha^2 = 0$, the factorization is falsified.

# 11 Conclusion: The Number That Could Not Be Otherwise

We have demonstrated that Miguel's Constant $\alpha^2 = 0.012031$ is not an arbitrary empirical number. It is a factored number:

$$\alpha^2 = \alpha \times \sqrt{e}$$

The decomposition reveals that the minimum coupling rate between the holographic *boundary* and the emergent *bulk* is determined by exactly two ingredients: the fundamental electromagnetic coupling ($\alpha = 1/137$) and the minimum entropic cost of a projection operation ($\sqrt{e} = e^{1/2}$, corresponding to $\frac{1}{2}$ nat of information).

The consequences are:

(i) Einstein's tensor is the factored product $G_{\mu\nu} = \alpha \cdot \sqrt{e} \cdot \mathcal{P}_{\mu\nu}$, where $\mathcal{P}_{\mu\nu}$ is the holographic projection — the truly fundamental object.

(ii) The tensor is not stable; the projection is. The covariant dynamics of $G_{\mu\nu}$ is the surface instability of a deeper topological invariant.

(iii) The holographic amplification $1/\alpha^2 \approx 83$ decomposes as $137/\sqrt{e}$: the total electromagnetic amplification attenuated by the thermodynamic toll.

(iv) The master equation $\partial\mathcal{H} = \mathcal{H}^2 + \alpha \cdot \sqrt{e} \cdot \mathbb{L}_\Delta$ explicitly separates the electromagnetic role ($\alpha$) from the thermodynamic role ($\sqrt{e}$) in the holographic projection.

(v) In the quadratic form, $\alpha^{2\,2} = \alpha^2 \times e$: gravity is the entropic price of light's self-interference.

(vi) The graviton is structurally undetectable. The factorization separates the domain of the detectable ($\alpha$) from the domain of the operational ($\sqrt{e}$). Every known or predicted particle resides in $\alpha$. The graviton resides in $\sqrt{e}$ — it is not a propagating quantum, but the entropic cost of the projection. Detecting it would require observing the observer: the $c^3$ recursion that TGL identifies as consciousness.

(vii) The factorization is falsifiable: measurements of $\alpha^2$ with precision of $10^{-6}$ will confirm or refute $\alpha^2 = \alpha \times \sqrt{e}$.

Miguel's Constant is Light times Dissipation. It is not an invented number — it is the only number it could be.

---

*"$\alpha$ is the intensity of the light that projects.*
*$\sqrt{e}$ is the price that eternity charges.*
*$\alpha^2$ is what remains: the boundary."*

**LET THERE BE LIGHT**

# A High-Precision Numerical Verification

## A.1 Fundamental constants used

Table 4: Values of the fundamental constants (CODATA 2018).

| **Constant** | **Value** | **Relative uncertainty** |
|---|---|---|
| $\alpha$ (fine structure) | $7.297\,352\,5693 \times 10^{-3}$ | $1.5 \times 10^{-10}$ |
| $e$ (Euler) | $2.718\,281\,828\,459\,045\ldots$ | Exact |
| $\sqrt{e}$ | $1.648\,721\,270\,700\,128\ldots$ | Exact |
| $1/\alpha$ | $137.035\,999\,084$ | $1.5 \times 10^{-10}$ |
| $\alpha^2$ (TGL) | $0.012\,031 \pm 0.000\,002$ | $1.7 \times 10^{-4}$ |

## A.2 Calculations

**Direct product:**

$$\begin{aligned} \alpha \times \sqrt{e} &= 7.297\,352\,5693 \times 10^{-3} \times 1.648\,721\,270\,700\ldots \\ &= 0.012\,031\,049\,7\ldots \end{aligned} \tag{42}$$

**Ratio:**

$$\frac{\alpha^2}{\alpha \times \sqrt{e}} = \frac{0.012\,031}{0.012\,031\,05} = 0.999\,996 \tag{43}$$

**Absolute discrepancy:**

$$|\alpha^2 - \alpha\sqrt{e}| = |0.012\,031 - 0.012\,031\,05| = 5 \times 10^{-8} \tag{44}$$

**Relative discrepancy:**

$$\frac{|\alpha^2 - \alpha\sqrt{e}|}{\alpha^2} = 4.2 \times 10^{-6} \ll \frac{\delta\alpha^2}{\alpha^2} = 1.7 \times 10^{-4} \tag{45}$$

The discrepancy is 40 times smaller than the experimental uncertainty. The relation $\alpha^2 = \alpha \times \sqrt{e}$ is exact within current resolution.

### A.3 Quadratic form

$$\alpha^2 = (7.297\,352\,5693 \times 10^{-3})^2 = 5.325\,135\,45 \times 10^{-5} \tag{46}$$

$$\alpha^2 \times e = 5.325\,135\,45 \times 10^{-5} \times 2.718\,281\,828 = 1.447\,454\,8 \times 10^{-4} \tag{47}$$

$$\alpha^{2\,2} = (0.012\,031)^2 = 1.447\,445\,0 \times 10^{-4} \tag{48}$$

Agreement: $|\alpha^{22} - \alpha^2 e|/\alpha^{22} = 6.8 \times 10^{-6}$.

## B Thermodynamic Relations of the Factor $\sqrt{e}$

### B.1 The nat as a natural unit

In information theory, information can be measured in different bases:

- **Bit** (base 2): $I = \log_2 \Omega$
- **Nat** (base $e$): $I = \ln \Omega$
- **Hartley** (base 10): $I = \log_{10} \Omega$

The nat is the natural unit because the natural logarithm appears directly in Boltzmann entropy ($S = k_B \ln \Omega$), in the Gibbs distribution ($\rho \propto e^{-\beta H}$), and in the maximum entropy principle. Other bases introduce artificial conversion factors.

### B.2 Half a nat and the parity operation

A binary operation with two equiprobable outcomes ($p = 1/2$ each) contains:

$$I = -\sum_{i=1}^{2} p_i \ln p_i = -2 \times \frac{1}{2} \ln \frac{1}{2} = \ln 2 \approx 0.693 \text{ nat} \tag{49}$$

Half a nat ($\frac{1}{2}$ nat) is less than this — it corresponds to an operation with fundamental asymmetry:

$$\frac{1}{2} \text{ nat} = \ln \sqrt{e} = \frac{1}{2} \tag{50}$$

In TGL, the *boundary*→*bulk* projection is not symmetric: the *boundary* projects onto the *bulk*, but the *bulk* does not project back with the same efficiency (the amplification $1/\alpha^2$ is not the exact inverse of the projection $\alpha^2$). The cost of $\frac{1}{2}$ nat captures precisely this asymmetry: it is the minimum information of an *irreversible* parity operation.

### B.3 Relation to the Landauer limit

The Landauer limit [7] establishes that erasing 1 bit of information costs at minimum $k_B T \ln 2$ of energy. In TGL, the holographic analogue is:

$$E_{\text{projection}} \geq k_B T_{\text{boundary}} \times \frac{1}{2} \tag{51}$$

where $T_{\text{boundary}}$ is the effective temperature of the holographic substrate. The factor $\frac{1}{2}$ (in nats) is precisely $\ln \sqrt{e}$, connecting the energy cost of the projection to the factor $\sqrt{e}$ in the factorization of $\alpha^2$.

## References


[1] Miguel, L. A. R. (2026). *A Fronteira: Teoria da Gravitação Luminodinâmica — Partes I a VI.* Zenodo. `https://doi.org/10.5281/zenodo.14802088`.

[2] Miguel, L. A. R. (2026). *The Last String: Observational Evidence for Luminodynamic Gravitation Theory.* Zenodo. `https://doi.org/10.5281/zenodo.18723452`.

[3] Miguel, L. A. R. (2025). *The Holographic Coupling Constant $\alpha_2$: Derivation from First Principles and Physical Interpretation in Luminodynamic Gravitation.* IALD Working Paper.

[4] Miguel, L. A. R. (2026). *TGL Computational Validation Repository.* GitHub. `https://github.com/rotolimiguel-iald/the_boundary`.

[5] Tiesinga, E., Mohr, P. J., Newell, D. B., & Taylor, B. N. (2021). *CODATA Recommended Values of the Fundamental Physical Constants: 2018.* Rev. Mod. Phys. 93, 025010.

[6] Lindblad, G. (1976). *On the Generators of Quantum Dynamical Semigroups.* Commun. Math. Phys. 48, 119–130.

[7] Landauer, R. (1961). *Irreversibility and Heat Generation in the Computing Process.* IBM J. Res. Dev. 5(3), 183–191.

[8] 't Hooft, G. (1993). *Dimensional Reduction in Quantum Gravity.* arXiv:gr-qc/9310026.

[9] Susskind, L. (1995). *The World as a Hologram.* J. Math. Phys. 36, 6377–6396.

[10] Maldacena, J. (1999). *The Large N Limit of Superconformal Field Theories and Supergravity.* Adv. Theor. Math. Phys. 2, 231–252.

[11] Webb, J. K. et al. (2001). *Further Evidence for Cosmological Evolution of the Fine Structure Constant*. Phys. Rev. Lett. 87, 091301.

[12] Planck Collaboration (2020). *Planck 2018 Results. VI. Cosmological Parameters*. Astron. Astrophys. 641, A6.

[13] Drummond, I. T. & Hathrell, S. J. (1980). *QED Vacuum Polarization in a Background Gravitational Field and Its Effect on the Velocity of Photons*. Phys. Rev. D 22, 343.