

Handbook on excursion sets

Emmanuel Roubin

Version: February 11, 2014

Contents

1	Priliminary calculations	2
1.1	Differentiation under the integral sign	2
1.2	Functions	2
1.2.1	Gamma function: Γ	2
1.2.2	Error function: erf	2
1.3	Lipschitz-Killing curvatures (LKC)	2
1.3.1	For a cube in 1,2 and 3 dimensions	2
1.4	Volume of the unit ball	3
1.5	Probabilistic Hermite polynomials	3
1.6	Flag coefficients	3
1.7	Second spectral moment	3
2	Gaussian Minkowsky functionals	4
2.1	Preliminary	4
2.1.1	Gaussian measure: $\gamma^k(\mathcal{H})$	4
2.1.2	Tube Taylor expansion	4
2.2	GMFs for Gaussian distribution	4
2.2.1	Application to one-dimensional tail hitting set: $k = 1$ and $\mathcal{H} = [\kappa, \infty)$	4
2.2.2	Application to one-dimensional cumulative hitting set: $k = 1$ and $\mathcal{H} = (-\infty, \kappa]$	5
2.3	GMFs for Gaussian related Random Fields	8
2.3.1	Log-normal distribution	8
3	Expectation Formula	10
3.1	Gaussian distribution	10
3.1.1	General case	10
3.1.2	One dimension	10

3.1.3	Two dimensions	10
4	Size effect	10
4.1	One dimensional case	10
4.2	Two dimensional case	10
4.2.1	Behavior of the Euler characteristic for different spec- imen length	10

1 Preliminary calculations

1.1 Differentiation under the integral sign

$$\frac{d}{dx} \left(\int_{a(x)}^{b(x)} f(x, t) dt \right) = f(x, b(x))b'(x) - f(x, a(x))a'(x) + \int_{a(x)}^{b(x)} \frac{\partial}{\partial x} f(x, t) dt \quad (1)$$

1.2 Functions

1.2.1 Gamma function: Γ

$$\Gamma : z \rightarrow \int_0^\infty t^{z-1} e^{-t} dt \quad (2)$$

1.2.2 Error function: erf

$$\text{erf}(x) = \frac{2}{\sqrt{\pi}} \int_0^x e^{-t^2} dt \quad (3)$$

1.3 Lipschitz-Killing curvatures (LKC)

1.3.1 For a cube in 1,2 and 3 dimensions

$$\text{For } \mathcal{M}_N = \prod_{i=1}^N [0, a], \quad \mathcal{L}_j(\mathcal{M}_N) = \binom{N}{j} a^j \quad (4)$$

N	LKC	Value	Meaning
1	$\mathcal{L}_0(\mathcal{M})$	1	Euler Characteristic
1	$\mathcal{L}_1(\mathcal{M})$	a	Length of the segment
2	$\mathcal{L}_0(\mathcal{M})$	1	Euler Characteristic
2	$\mathcal{L}_1(\mathcal{M})$	$2a$	Half the boundary length
2	$\mathcal{L}_2(\mathcal{M})$	a^2	Surface area
3	$\mathcal{L}_0(\mathcal{M})$	1	Euler Characteristic
3	$\mathcal{L}_1(\mathcal{M})$	$3a$	Twice the caliper diameter
3	$\mathcal{L}_2(\mathcal{M})$	$3a^2$	Half the surface area
3	$\mathcal{L}_3(\mathcal{M})$	a^3	Volume

1.4 Volume of the unit ball

$$\omega_N = \frac{\pi^{N/2}}{\Gamma(1 + N/2)} \quad (5) \quad \left| \begin{array}{l} \omega_0 = 1 \\ \omega_1 = 2 \\ \omega_2 = \pi \\ \omega_3 = 4\pi/3 \\ \dots \end{array} \right. \quad (6)$$

1.5 Probabilistic Hermite polynomials

$$\mathcal{H}_n = (-1)^n e^{x^2/2} \frac{d^n}{dx^n} e^{-x^2/2} \quad (7) \quad \left| \begin{array}{l} \mathcal{H}_0(x) = 1 \\ \mathcal{H}_1(x) = x \\ \mathcal{H}_2(x) = x^2 - 1 \\ \mathcal{H}_3(x) = x^3 - 3x \\ \dots \end{array} \right. \quad (8)$$

1.6 Flag coefficients

$$\left[\begin{array}{c} n \\ j \end{array} \right] = \binom{n}{j} \frac{\omega_n}{\omega_{n-j}\omega_j} \quad (9) \quad \begin{array}{c|c|c} n & j & \text{flag} \\ \hline 0 & 0 & 1 \\ 1 & 0 & 1 \\ 1 & 1 & 1 \\ 2 & 0 & 1 \\ 2 & 1 & \pi/2 \\ 2 & 2 & 1 \\ 3 & 0 & 1 \\ 3 & 1 & 2 \\ 3 & 2 & 2 \\ 3 & 3 & 1 \end{array} \quad (10)$$

1.7 Second spectral moment

$$\lambda_2 = \mathbb{V} \left\{ \frac{\partial g(\mathbf{x})}{\partial x_i} \right\} = \frac{d^2 \mathcal{C}(h)}{dh^2} \quad (11)$$

Covariance	Parameter	λ_2
Gaussian	—	$2\sigma^2/L_c^2$

(12)

2 Gaussian Minkowsky functionals

2.1 Preliminary

2.1.1 Gaussian measure: $\gamma^k(\mathcal{H})$

$$\gamma^k(\mathcal{H}) = P\{\mathbf{X} \in \mathcal{H}\} = \frac{1}{\sigma^k (2\pi)^{k/2}} \int_{\mathcal{H}} e^{-\|\mathbf{x}-\mu\|^2/2\sigma^2} d\mathbf{x} \quad (13)$$

2.1.2 Tube Taylor expansion

$$\gamma^k(\text{tube}(\mathcal{H}, \rho)) = \sum_{j=0}^{\infty} \frac{\rho^j}{j!} \mathcal{M}_j^k(\mathcal{H}) \quad (14)$$

2.2 GMFs for Gaussian distribution

2.2.1 Application to one-dimensional tail hitting set: $k = 1$ and $\mathcal{H} = [\kappa \infty)$

Tail probability \bar{F} :

$$\gamma^1([\kappa \infty)) = P\{X \geq \kappa\} = \frac{1}{\sigma\sqrt{2\pi}} \int_{\kappa}^{\infty} e^{-(x-\mu)^2/2\sigma^2} dx = \bar{F}(\kappa) \quad (15)$$

Error function erf:

$$\begin{aligned} \gamma^1([\kappa \infty)) &= \bar{F}(\kappa) = \frac{1}{\sigma\sqrt{2\pi}} \int_{\frac{\kappa-\mu}{\sigma\sqrt{2}}}^{\infty} e^{-t^2} \sqrt{2}\sigma dt = \frac{1}{\sqrt{\pi}} \int_{\frac{\kappa-\mu}{\sigma\sqrt{2}}}^{\infty} e^{-t^2} dt \\ &= \frac{1}{\sqrt{\pi}} \left[\int_0^{\infty} e^{-t^2} dt - \int_0^{\frac{\kappa-\mu}{\sigma\sqrt{2}}} e^{-t^2} dt \right] \\ &= \frac{1}{\sqrt{\pi}} \left[\frac{\sqrt{\pi}}{2} - \frac{\sqrt{\pi}}{2} \text{erf}\left(\frac{\kappa-\mu}{\sigma\sqrt{2}}\right) \right] \\ &= \frac{1}{2} \left(1 - \text{erf}\left(\frac{\kappa-\mu}{\sigma\sqrt{2}}\right) \right) \end{aligned}$$

Tube expansion:

$$\gamma^1(\text{tube}([\kappa \infty), \rho)) = \gamma^1([\kappa - \rho \infty)) = \bar{F}(\kappa - \rho) = \sum_{j=0}^{\infty} \frac{(-\rho)^j}{j!} \frac{d^j \bar{F}(\kappa)}{d\kappa^j} \quad (17)$$

Identification:

- Notations:

$$\text{Tail hitting set: } \bar{\mathcal{H}}^{1\text{D}} = [\kappa \infty) \quad (18a)$$

$$\text{Variable substitution: } \tilde{\kappa} \leftarrow \frac{\kappa - \mu}{\sigma} \quad (18b)$$

- For $j = 0$:

$$\mathcal{M}_0^{\gamma^1}(\bar{\mathcal{H}}^{1\text{D}}) = \bar{F}(\kappa) \quad (19)$$

- For $j > 0$:

$$\mathcal{M}_j^{\gamma^1}(\bar{\mathcal{H}}^{1\text{D}}) = (-1)^j \frac{d^j \bar{F}(\kappa)}{d\kappa^j} = \frac{e^{-(\kappa-\mu)^2/2\sigma^2}}{\sigma^j \sqrt{2\pi}} \mathcal{H}_{j-1}((\kappa - \mu)/\sigma) \quad (20)$$

- First values:

$$\left| \begin{array}{l} \mathcal{M}_0^{\gamma^1}(\bar{\mathcal{H}}^{1\text{D}}) = \frac{1}{2} \left(1 - \text{erf} \left(\frac{\tilde{\kappa}}{\sqrt{2}} \right) \right) \\ \mathcal{M}_1^{\gamma^1}(\bar{\mathcal{H}}^{1\text{D}}) = \frac{1}{\sigma \sqrt{2\pi}} e^{-\tilde{\kappa}^2/2} \\ \mathcal{M}_2^{\gamma^1}(\bar{\mathcal{H}}^{1\text{D}}) = \frac{1}{\sigma^2 \sqrt{2\pi}} \tilde{\kappa} e^{-\tilde{\kappa}^2/2} \\ \mathcal{M}_3^{\gamma^1}(\bar{\mathcal{H}}^{1\text{D}}) = \frac{1}{\sigma^3 \sqrt{2\pi}} (\tilde{\kappa}^2 - 1) e^{-\tilde{\kappa}^2/2} \\ \mathcal{M}_4^{\gamma^1}(\bar{\mathcal{H}}^{1\text{D}}) = \frac{1}{\sigma^4 \sqrt{2\pi}} (\tilde{\kappa}^3 - 3\tilde{\kappa}) e^{-\tilde{\kappa}^2/2} \\ \dots \end{array} \right. \quad (21)$$

- Derivatives:

$$\forall j, \quad \frac{d\mathcal{M}_j^{\gamma^1}}{d\kappa}(\bar{\mathcal{H}}^{1\text{D}}) = (-1)^j \frac{d^{j+1} \bar{F}(\kappa)}{d\kappa^{j+1}} = -\mathcal{M}_{j+1}^{\gamma^1}(\bar{\mathcal{H}}^{1\text{D}}) \quad (22)$$

2.2.2 Application to one-dimensional cumulative hitting set: $k = 1$ and $\mathcal{H} = (-\infty \kappa]$

Cumulative probability F :

$$\gamma^1((-\infty \kappa]) = P\{X \leq \kappa\} = \frac{1}{\sigma \sqrt{2\pi}} \int_{-\infty}^{\kappa} e^{-(x-\mu)^2/2\sigma^2} dx = F(\kappa) \quad (23)$$




Figure 1: First 4 GMF for $\sigma^2 = 2$ and $\mathcal{H} = \bar{\mathcal{H}}^{1D} = [\kappa \infty)$

Error function erf:

$$\gamma^1((-\infty \kappa]) = F(\kappa) = \frac{1}{2} \left(1 + \operatorname{erf} \left(\frac{\kappa - \mu}{\sigma\sqrt{2}} \right) \right) \quad (24)$$

Tube expansion:

$$\gamma^1(\text{tube}((-\infty \kappa], \rho)) = \gamma^1((-\infty \kappa + \rho]) = F(\kappa + \rho) = \sum_{j=0}^{\infty} \frac{\rho^j}{j!} \frac{d^j F(\kappa)}{d\kappa^j} \quad (25)$$

Identification:

- Notations:

$$\text{Tail hitting set: } \mathcal{H}^{1D} = (-\infty \kappa] \quad (26a)$$

$$\text{Variable substitution: } \tilde{\kappa} \leftarrow \frac{\kappa - \mu}{\sigma} \quad (26b)$$

- For $j = 0$:

$$\mathcal{M}_0^{\gamma^1}(\mathcal{H}^{1D}) = F(\kappa) \quad (27)$$

- For $j > 0$:

$$\mathcal{M}_j^{\gamma^1}(\mathcal{H}^{1D}) = \frac{d^j F(\kappa)}{d\kappa^j} = (-1)^{j+1} \frac{e^{-(\kappa-\mu)^2/2\sigma^2}}{\sigma^j \sqrt{2\pi}} \mathcal{H}_{j-1}((\kappa - \mu)/\sigma) \quad (28)$$

- First values:

$$\begin{aligned}
 \mathcal{M}_0^{\gamma^1}(\mathcal{H}^{1\text{D}}) &= \frac{1}{2} \left(1 - \operatorname{erf} \left(\frac{\tilde{\kappa}}{\sqrt{2}} \right) \right) \\
 \mathcal{M}_1^{\gamma^1}(\mathcal{H}^{1\text{D}}) &= \frac{1}{\sigma\sqrt{2\pi}} e^{-\tilde{\kappa}^2/2} \\
 \mathcal{M}_2^{\gamma^1}(\mathcal{H}^{1\text{D}}) &= \frac{-1}{\sigma^2\sqrt{2\pi}} \tilde{\kappa} e^{-\tilde{\kappa}^2/2} \\
 \mathcal{M}_3^{\gamma^1}(\mathcal{H}^{1\text{D}}) &= \frac{1}{\sigma^3\sqrt{2\pi}} (\tilde{\kappa}^2 - 1) e^{-\tilde{\kappa}^2/2} \\
 \mathcal{M}_4^{\gamma^1}(\mathcal{H}^{1\text{D}}) &= \frac{-1}{\sigma^4\sqrt{2\pi}} (\tilde{\kappa}^3 - 3\tilde{\kappa}) e^{-\tilde{\kappa}^2/2} \\
 &\dots
 \end{aligned} \tag{29}$$




Figure 2: First 4 GMF for $\sigma^2 = 2$ and $\mathcal{H} = \mathcal{H}^{1\text{D}} = (-\infty, \kappa]$

- Derivatives:

$$\forall j, \quad \frac{d\mathcal{M}_j^{\gamma^1}}{d\kappa}(\mathcal{H}^{1\text{D}}) = \frac{d^{j+1}F(\kappa)}{d\kappa^{j+1}} = \mathcal{M}_{j+1}^{\gamma^1}(\mathcal{H}^{1\text{D}}) \tag{30}$$

2.3 GMFs for Gaussian related Random Fields

$$g_r : \Omega \times \mathbf{R}^N \xrightarrow{g} \mathbf{R}^k \xrightarrow{S} \mathbf{R} \quad (31)$$

$$P \{g_r \in \mathcal{H}\} = P \{S(g) \in \mathcal{H}\} = P \{g \in S^{-1}(\mathcal{H})\} = \gamma^k(S^{-1}(\mathcal{H})) \quad (32)$$

2.3.1 Log-normal distribution

$$k = 1 \quad , \quad S = \exp \quad \text{and} \quad S^{-1} = \ln \quad (33)$$

$$\left| \begin{array}{l} \mu = \ln(\mu_{\ln}) - \frac{1}{2} \ln(1 + \sigma_{\ln}^2 / \mu_{\ln}^2) \\ \sigma^2 = \ln(1 + \sigma_{\ln}^2 / \mu_{\ln}^2) \end{array} \right. \quad (34)$$

- Notations:

$$\text{Log tail hitting set: } \bar{\mathcal{H}}_{\ln}^{1D} = [\ln(\kappa) \infty) \quad (35a)$$

$$\text{Log cumulative hitting set: } \mathcal{H}_{\ln}^{1D} = (-\infty \ln(\kappa)] \quad (35b)$$

$$\text{Variable substitution: } \tilde{\kappa}_{\ln} \leftarrow \frac{\ln(\kappa) - \mu}{\sigma} \quad (35c)$$

- For $\mathcal{H} = \bar{\mathcal{H}}_{\ln}^{1D} = [\kappa \infty) \rightarrow S^{-1}(\mathcal{H}) = \bar{\mathcal{H}}_{\ln}^{1D} = [\ln(\kappa) \infty)$

$$\left| \begin{array}{l} \mathcal{M}_0^{\gamma^1}(\bar{\mathcal{H}}_{\ln}^{1D}) = \frac{1}{2} \left(1 - \operatorname{erf} \left(\frac{\tilde{\kappa}_{\ln}}{\sqrt{2}} \right) \right) \\ \mathcal{M}_1^{\gamma^1}(\bar{\mathcal{H}}_{\ln}^{1D}) = \frac{1}{\sigma \sqrt{2\pi}} e^{-\tilde{\kappa}_{\ln}^2/2} \\ \mathcal{M}_2^{\gamma^1}(\bar{\mathcal{H}}_{\ln}^{1D}) = \frac{1}{\sigma^2 \sqrt{2\pi}} \tilde{\kappa}_{\ln} e^{-\tilde{\kappa}_{\ln}^2/2} \\ \mathcal{M}_3^{\gamma^1}(\bar{\mathcal{H}}_{\ln}^{1D}) = \frac{1}{\sigma^3 \sqrt{2\pi}} (\tilde{\kappa}_{\ln}^2 - 1) e^{-\tilde{\kappa}_{\ln}^2/2} \\ \mathcal{M}_4^{\gamma^1}(\bar{\mathcal{H}}_{\ln}^{1D}) = \frac{1}{\sigma^4 \sqrt{2\pi}} (\tilde{\kappa}_{\ln}^3 - 3\tilde{\kappa}_{\ln}) e^{-\tilde{\kappa}_{\ln}^2/2} \\ \dots \end{array} \right. \quad (36)$$

Derivatives:

$$\begin{aligned}
& \text{Help: } \left(e^{-(\ln(x)-\mu)^2/2\sigma^2} \right)' = -\frac{1}{\sigma^2} \frac{\ln(x)-\mu}{x} e^{-(\ln(x)-\mu)^2/2\sigma^2} \\
& \frac{d\mathcal{M}_0^{\gamma^1}(\tilde{\mathcal{H}}_{\ln}^{\text{1D}})}{d\kappa} = \frac{-1}{\sigma\sqrt{2\pi}} \frac{1}{\kappa} e^{-(\ln(\kappa)-\mu)^2/2\sigma^2} \\
& \frac{d\mathcal{M}_1^{\gamma^1}(\tilde{\mathcal{H}}_{\ln}^{\text{1D}})}{d\kappa} = \frac{-1}{\sigma^2\sqrt{2\pi}} \frac{1}{\kappa} \frac{\ln(\kappa)-\mu}{\sigma} e^{-(\ln(\kappa)-\mu)^2/2\sigma^2} \\
& \frac{d\mathcal{M}_2^{\gamma^1}(\tilde{\mathcal{H}}_{\ln}^{\text{1D}})}{d\kappa} = \frac{-1}{\sigma^3\sqrt{2\pi}} \frac{1}{\kappa} \left[\frac{(\ln(\kappa)-\mu)^2}{\sigma^2} - 1 \right] e^{-(\ln(\kappa)-\mu)^2/2\sigma^2} \\
& \frac{d\mathcal{M}_3^{\gamma^1}(\tilde{\mathcal{H}}_{\ln}^{\text{1D}})}{d\kappa} = \frac{-1}{\sigma^4\sqrt{2\pi}} \frac{1}{\kappa} \left[\frac{(\ln(\kappa)-\mu)^3}{\sigma^3} - 3\frac{(\ln(\kappa)-\mu)}{\sigma} \right] e^{-(\ln(\kappa)-\mu)^2/2\sigma^2} \\
& \dots \\
& \text{Guess would be: } \forall j \geq 0, \\
& \frac{d\mathcal{M}_j^{\gamma^1}(\mathcal{H}_{\ln}^{\text{1D}})}{d\kappa} = \frac{-1}{\sigma^{j+1}\sqrt{2\pi}} \frac{1}{\kappa} \mathcal{H}_j(\tilde{\kappa}_{\ln}) e^{-\tilde{\kappa}_{\ln}^2/2}
\end{aligned} \tag{37}$$

- For $\mathcal{H} = [0 \ \kappa] \rightarrow S^{-1}(\mathcal{H}) = \mathcal{H}_{\ln}^{\text{1D}} = (-\infty \ \ln(\kappa)]$

$$\begin{aligned}
& \mathcal{M}_0^{\gamma^1}(\mathcal{H}_{\ln}^{\text{1D}}) = \frac{1}{2} \left(1 + \text{erf} \left(\frac{\tilde{\kappa}_{\ln}}{\sqrt{2}} \right) \right) \\
& \mathcal{M}_1^{\gamma^1}(\mathcal{H}_{\ln}^{\text{1D}}) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\tilde{\kappa}_{\ln}^2/2} \\
& \mathcal{M}_2^{\gamma^1}(\mathcal{H}_{\ln}^{\text{1D}}) = \frac{-1}{\sigma^2\sqrt{2\pi}} \tilde{\kappa}_{\ln} e^{-\tilde{\kappa}_{\ln}^2/2} \\
& \mathcal{M}_3^{\gamma^1}(\mathcal{H}_{\ln}^{\text{1D}}) = \frac{1}{\sigma^3\sqrt{2\pi}} (\tilde{\kappa}_{\ln}^2 - 1) e^{-\tilde{\kappa}_{\ln}^2/2} \\
& \mathcal{M}_4^{\gamma^1}(\mathcal{H}_{\ln}^{\text{1D}}) = \frac{-1}{\sigma^4\sqrt{2\pi}} (\tilde{\kappa}_{\ln}^3 - 3\tilde{\kappa}_{\ln}) e^{-\tilde{\kappa}_{\ln}^2/2} \\
& \dots
\end{aligned} \tag{38}$$

Derivatives:

$$\begin{aligned}
& \text{Guess would be: } \forall j \geq 0, \\
& \frac{d\mathcal{M}_j^{\gamma^1}(\mathcal{H}_{\ln}^{\text{1D}})}{d\kappa} = \frac{(-1)^j}{\sigma^{j+1}\sqrt{2\pi}} \frac{1}{\kappa} \mathcal{H}_j(\tilde{\kappa}_{\ln}) e^{-\tilde{\kappa}_{\ln}^2/2}
\end{aligned} \tag{39}$$

3 Expectation Formula

3.1 Gaussian distribution

3.1.1 General case

$$\mathbb{E} \{ \mathcal{L}_j(\mathcal{E}) \} = \sum_{i=0}^{N-j} \begin{bmatrix} i+j \\ i \end{bmatrix} \left(\frac{\lambda_2}{2\pi} \right)^{i/2} \mathcal{L}_{i+j}(\mathcal{M}) \mathcal{M}_i^{\gamma^k}(S^{-1}(\mathcal{H})) \quad (40)$$

Notations:

$$\text{LKC of } \mathcal{M}: \mathcal{L}_j(\mathcal{M}) = \mathcal{L}_j^{\mathcal{M}} \quad (41a)$$

$$\text{LKC of } \mathcal{E}: \mathcal{L}_j(\mathcal{E}) = \mathcal{L}_j^{\mathcal{E}} \quad (41b)$$

$$\text{GMF of } S^{-1}(\mathcal{H}): \mathcal{M}_j^{\gamma^k}(S^{-1}(\mathcal{H})) = \mathcal{M}_j^{\gamma^k} \quad (41c)$$

3.1.2 One dimension

$$\mathbb{E} \{ \mathcal{L}_0^{\mathcal{E}} \} = \left(\frac{\lambda_2}{2\pi} \right)^{1/2} \mathcal{L}_1^{\mathcal{M}} \mathcal{M}_1^{\gamma^k} + \mathcal{L}_0^{\mathcal{M}} \mathcal{M}_0^{\gamma^k} \quad (42a)$$

$$\mathbb{E} \{ \mathcal{L}_1^{\mathcal{E}} \} = \mathcal{L}_1^{\mathcal{M}} \mathcal{M}_0^{\gamma^k} \quad (42b)$$

3.1.3 Two dimensions

$$\mathbb{E} \{ \mathcal{L}_0^{\mathcal{E}} \} = \frac{\lambda_2}{2\pi} \mathcal{L}_2^{\mathcal{M}} \mathcal{M}_2^{\gamma^k} + \sqrt{\frac{\lambda_2}{2\pi}} \mathcal{L}_1^{\mathcal{M}} \mathcal{M}_1^{\gamma^k} + \mathcal{L}_0^{\mathcal{M}} \mathcal{M}_0^{\gamma^k} \quad (43a)$$

$$\mathbb{E} \{ \mathcal{L}_1^{\mathcal{E}} \} = \sqrt{\frac{\lambda_2 \pi}{8}} \mathcal{L}_2^{\mathcal{M}} \mathcal{M}_1^{\gamma^k} + \mathcal{L}_1^{\mathcal{M}} \mathcal{M}_0^{\gamma^k} \quad (43b)$$

$$\mathbb{E} \{ \mathcal{L}_2^{\mathcal{E}} \} = \mathcal{L}_2^{\mathcal{M}} \mathcal{M}_0^{\gamma^k} \quad (43c)$$

4 Size effect

4.1 One dimensional case

4.2 Two dimensional case

4.2.1 Behavior of the Euler characteristic for different specimen length

It is first important to understand well the different behaviors (regimes) of the Euler characteristic in terms of the specimen length a .




Figure 3: Expectation of LKC for $a = 100$, $\sigma^2 = 2$ and $\mathcal{H} = [\kappa, \infty)$ in 1D.




Figure 4: Expectation of LKC for $a = 100$, $\mu = 0.5$, $\sigma^2 = 2$, $L_c = 0.5$, and $\mathcal{H} = [0, \kappa]$ (log-normal distribution with cumulative hitting set) in 1D. Experimental results are calculated over 1 000 realizations.




Figure 5: Expectation of LKC for $a = 100$, $\mu = 0.0$, $\sigma^2 = 2$, $L_c = 5$, and $\mathcal{H} = [\kappa, \infty)$ (gaussian distribution with tail hitting set) in 2D. Experimental results are calculated over 1 000 realizations.







Figure 6: Expectation of LKC for $a = 100$, $\mu = 0.0$, $\sigma^2 = 2$, $L_c = 5$, and $\mathcal{H} = [0, \kappa]$ (lognormal distribution with cumulative hitting set) in 2D. Experimental results are calculated over 1 000 realizations.



(a) Variation of the quantile



(b) Variation of the variance



(c) Variation of the mean

Figure 7: Size Effect: variations around $\mu_{\ln} = 10$, $\sigma_{\ln}^2 = 10$ and $q = 1\%$ in 1D.




Figure 8: Size Effect: variations around $\mu_{\text{ln}} = 10$, $\sigma_{\text{ln}}^2 = 10$ and $q = 1\%$ in 1D.