


Introduction To Sampling

Jeff Rouder

May, 2022

Estimate the parameters in a normal

- I mean, how hard is this
- $\hat{\mu} = \frac{\sum y_i}{n}$
- $\hat{\sigma}^2 = \frac{\sum (y_i - \bar{y})^2}{n-1}$
- $CI = \pm t(n-1, .975) \times \sqrt{n} \times \frac{\hat{\mu}}{\hat{\sigma}}$



Who Wakes Up First?

Let X denote when Becky wakes up. It is minutes after 7a.


$$X \sim \text{Gamma}(2, 10)$$

Let Y denote when Jeff wakes up. It is minutes after 6:30a.

$$Y \sim \text{Gamma}(1, 30)$$

What is the probability Becky wakes up before Jeff, $Pr(X < Y)$?

Setup




Simulation Code

```
M=100000
j=rgamma(M,shape=1,scale=30)
b=rgamma(M,shape=2,scale=10)+30
hist(b,breaks=seq(0,500,2),col=rgb(1,0,0,.5),
      xlim=c(0,180),prob=T)
hist(j,add=T,col=rgb(0,1,0,.5),breaks=seq(0,500,2),prob=T)
print(mean(b<j))
```

Simulations

Histogram of b



[1] 0.2064

- Samples from posterior
- e.g. $(\mu, \sigma^2 | \mathbf{Y})$
- Samples are enough to learn

How To?

- Program Your Own Sampler To Learn About MCMC
 - fast to execute
 - you feel smart
 - insightful
- JAGS
 - fast but autocorrelation
 - great manual
 - stable
- stan
 - best samplers,
 - changing, info scattered
 - very popular (cool kids)

$$Y_i \sim N(\mu, \sigma^2)$$

$$\mu \sim N(a, b)$$

$$\sigma^2 \sim \text{Inverse Gamma}(q, r)$$

$$Y = 92, 115, 100, 98$$

$$\mu \mid \sigma^2, \mathbf{Y} \sim \text{Normal}(vc, v), \quad v = \left(\frac{N}{\sigma^2} + \frac{1}{b} \right)^{-1}, \quad c = \frac{N\bar{Y}}{\sigma^2} + \frac{a}{b}$$

$$\sigma^2 \mid \mu, \mathbf{Y} \sim \text{IG} \left(q + \frac{N}{2}, r + \frac{SSE}{2} \right)$$

]



Demo, Roll My Own

```
Y=c(92,115,100,98)
ybar=mean(Y)
N=length(Y)
a=100
b=25
q=.5
r=.5*15^2
M=20000 # number of samples
mu=1:M
s2=1:M
mu[1]=100 #initial value
s2[1]=100 #initial value
```

Demo, Roll My Own


```
for (m in 2:M){  
  v=1/(N/s2[m-1]+1/b)  
  c=N*ybar/s2[m-1]+a/b  
  mu[m]=rnorm(1,c*v,sqrt(v))  
  SSE=sum((Y-mu[m])^2)  
  s2[m]=rinvgamma(1,N/2+q,SSE/2+r)  
}
```

Demo Roll My Own




Demo Roll My Own

Histogram of μ



Histogram of $\sqrt{s_2}$



- Examples in their directories
- It is too hard to typeset here

Two-sample analysis

- Half the kids get Smarties (treatment)
- Half don't (control)
- All get an IQ test an hour later
- What was the effect of Smarties?
- You make up the data to explore the following questions
- Let Y_{ij} denote the IQ for the j th kiddo, $j = 1, \dots, J$ in the i th group, $i = 1, 2$. ($i = 1$ for control, $i = 2$ for treatment)
- $Y_1 = 101, 82, 136, 77, 102, 99$
- $Y_2 = 104, 101, 115, 88, 83, 122$

Two Models, Which Is Better?

Cell Means Parameterization:

$$Y_{1j} \sim N(\mu_1, \sigma^2)$$

$$Y_{2j} \sim N(\mu_2, \sigma^2)$$

Same, but more compact:

$$Y_{ij} \sim N(\mu_i, \sigma^2)$$

Effects Parameterization:

$$Y_{1j} \sim N(\alpha - \theta/2, \sigma^2)$$

$$Y_{2j} \sim N(\alpha + \theta/2, \sigma^2)$$

Same but more compact:

$$Y_{ij} \sim N(\alpha + x_i\theta, \sigma^2), \quad x_i = c(-1/2, 1/2)$$

Two Models, Which Is Better?

Cell Means:

$$Y_{ij} \sim N(\mu_i, \sigma^2)$$

$$\mu_i \sim N(a, b)$$

$$\sigma^2 \sim \text{IG}(1/2, r/2)$$

Effects:

$$Y_{ij} \sim N(\alpha + x_i\theta, \sigma^2)$$

$$\alpha \sim N(a_1, b_1)$$

$$\theta \sim N(a_2, b_2)$$

$$\sigma^2 \sim \text{IG}(1/2, r/2)$$

- $a = a_1 = 100$, $b = b_1 = 5^2$, $r=15^2$