

Quiz 9

Friday, November 20 — inverses, rank, and determinants (Week 11)

Last updated: October 8, 2026.

Covers the prior instructional week only: [Inverses](#), [Rank](#), [Determinants](#) (Week 11). Midterm 2 week is skipped when choosing the prior week. Same-day [Lab 5](#) / eigenvectors are **not** on this quiz.

Textbooks: [Boyd & Vandenberghe \(VMLS\)](#) · [Strang §3.5](#) · [Lachniet Octave PDF](#) · also on [Resources](#).

[Schedule](#)

Core

Fluency (complete prompts — practice these by hand)

C9.1. Let

$$A = \begin{bmatrix} 2 & 1 \\ 0 & 3 \end{bmatrix}.$$

(a) Compute A^{-1} using the 2×2 formula (or by solving $AB = I$). (b) Check $A^{-1}A = I$. (c) Solve $Ax = (4, 3)$ using $x = A^{-1}b$.

C9.2. Decide whether each matrix is invertible. Give a one-line reason (dependent columns, $Ax = 0$, or $\det$).

(a) $\begin{bmatrix} 1 & 2 \\ 3 & 4 \end{bmatrix}$

(b) $\begin{bmatrix} 1 & 2 \\ 2 & 4 \end{bmatrix}$

(c) $\begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}$

C9.3. True or false, with a one-line reason: for a single right-hand side b , forming A^{-1} explicitly and then computing $A^{-1}b$ is generally preferable to solving $Ax = b$ with a solver / elimination.

C9.4. Let

$$A = \begin{bmatrix} 1 & 2 & 0 \\ 2 & 4 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

The RREF of A is

$$\begin{bmatrix} 1 & 2 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}$$

(after row swaps / scaling as needed — use this RREF). Find $\text{rank}(A)$, a basis for $C(A)$ using original columns, and a basis for $N(A)$.

C9.5. Let

$$A = \begin{bmatrix} 1 & 0 \\ 2 & 0 \\ 0 & 1 \end{bmatrix}.$$

Describe $C(A)$ in words. Is $b = (1, 2, 3)$ in $C(A)$? Find a basis for $N(A)$ (or state that $N(A) = \{0\}$).

C9.6. True or false, with a one-line reason: if $\text{rank}(A) = n$ for an $m \times n$ matrix with $m \geq n$, then $Ax = 0$ has only the zero solution.

C9.7. Compute each determinant.

(a) $\det \begin{bmatrix} 3 & 1 \\ 0 & 2 \end{bmatrix}$

(b) $\det \begin{bmatrix} 1 & 4 & 0 \\ 0 & -2 & 5 \\ 0 & 0 & 3 \end{bmatrix}$

(c) $\det \begin{bmatrix} 2 & 1 \\ 4 & 2 \end{bmatrix}$

C9.8. Let $A = \begin{bmatrix} 0 & -2 \\ 1 & 0 \end{bmatrix}$. Compute $\det A$. Where do e_1 and e_2 go under $x \mapsto Ax$? What area does the unit square map to?

C9.9. True or false, with a one-line reason: if $\det A = 0$ for square A , then A is not invertible and $Ax = b$ fails to have a unique solution for every b .

C9.10. Connect the ideas. For square A , match each statement to the equivalent view (write the letter):

- (i) A is invertible
- (ii) $\text{rank}(A) = n$ (full)
- (iii) $\det A \neq 0$
- (iv) $N(A) = \{0\}$

Which of (ii)–(iv) are equivalent to (i)?

C9.11 (cumulative, light). Let $A = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}$. What geometric map is $x \mapsto Ax$? Is A invertible? What is $\det A$?

Textbook Core

- B 11.1, 11.2, 11.4, 11.10
- S 3.5 #1 and #2 (rank / subspace portions used in class)
- L Ch. 1 Ex. 3(a), 3(b), and 3(d): determinant, rank, and inverse portions

Stretch

B 11.15, 11.25; area scaling from $|\det A|$ for a new 2×2 map (C-M2 style).

What a quiz item might change

A new 2×2 matrix, an RREF with a free variable, whether $\det A = 0$ in context, or asking for $C(A)$ vs $N(A)$ from the same matrix.