

Module 3 — Synthesis Quiz

[Question bank](#) · [After exercise lab](#) · [open notes](#) · [two attempts](#)

Module 3, Synthesis quiz

Readings: [Read 0](#), [Read 1](#), [Read 2](#)

Labs: [Demo](#), [Exercise](#)

Canvas: Synthesis quiz (Module 3) · **two attempts**, highest score counts

Full integration for [Project 3](#): five-number summary, boxplot comparisons, scatterplot description, two-way tables and proportions, ggplot2 calls, and **five demo-lab EDA situations**.

Image items: Section F includes histograms, boxplots, scatterplots, and bar/stacked-bar figures (also collected in the [Graph Interpretation Pack](#)).

Instructions

- Open notes ([methods map](#), [R reference](#)).
 - Use the graph-selection table from Read 0 as a guide.
 - For R items, match the workflow in the [demo lab](#) and your [Project 3 notebook](#).
-

Section A — Five-number summary and boxplots (Read 1)

A1. Compute five-number summary

Data (sorted): 2, 5, 7, 8, 10, 12, 15, 18, 22, 30. Compute the **five-number summary** (min, Q_1 , median, Q_3 , max).

A2. IQR and outlier fence

Using the five-number summary above, compute IQR and the upper outlier **fence** ($Q_3 + 1.5 \times \text{IQR}$).

A3. Compare groups from boxplots

Two groups have boxplots: Group A median=25, $Q_1 = 18$, $Q_3 = 32$; Group B median=40, $Q_1 = 35$, $Q_3 = 45$. Which group has higher typical values? Which has more spread?

A4. R code for grouped boxplot

Write the ggplot2 call for boxplots of `body_mass_g` grouped by `species` in the `penguins` dataset.

A5. `group_by` + `summarize` in R

Write the pipeline that computes **median** and **IQR** of `body_mass_g` for each `species`.

Answer key — Section A — Five-number summary and boxplots (Read 1)

1. $\text{Min}=2$, $Q_1 = (7 + 8)/2 = 7.5$, $\text{Median}=(10 + 12)/2 = 11$, $Q_3 = (15 + 18)/2 = 16.5$, $\text{Max}=30$.
 2. $\text{IQR} = 16.5 - 7.5 = 9$. Upper fence $= 16.5 + 1.5(9) = 30$. The value 30 is on the fence — borderline outlier.
 3. Group B has **higher typical values** (median 40 vs 25). Group A has **more spread** ($\text{IQR}=14$ vs $\text{IQR}=10$).
 4. `ggplot(penguins, aes(x = species, y = body_mass_g)) + geom_boxplot()` — optionally add `fill = species` for color.
 5. `penguins %>% group_by(species) %>% summarize(med = median(body_mass_g, na.rm = TRUE), iqr = IQR(body_mass_g, na.rm = TRUE))`
-

Section B — Scatterplot description (Read 2 §2)

B1. Describe a scatterplot

A scatterplot of `bill_length_mm` vs `flipper_length_mm` shows points trending upward in a roughly straight band. Describe the association.

B2. Negative association

A scatterplot shows higher values of x paired with lower values of y , with points scattered loosely. Describe the association.

B3. No association

A scatterplot shows no discernible trend — points form a cloud with no slope. Describe the association.

B4. R code for scatterplot

Write the `ggplot2` call for a scatterplot of `bill_length_mm` (x) vs `flipper_length_mm` (y).

B5. Color points by group

How would you modify the scatterplot to color points by `species`?

Answer key — Section B — Scatterplot description (Read 2 §2)

1. **Form:** linear. **Direction:** positive. **Strength:** moderate to strong.
 2. **Form:** linear (loosely). **Direction:** negative. **Strength:** weak.
 3. **Form:** no clear pattern. **Direction:** none. **Strength:** none (weak/no association).
 4. `ggplot(penguins, aes(x = bill_length_mm, y = flipper_length_mm)) + geom_point()`
 5. Add `color = species` to `aes()`: `aes(x = bill_length_mm, y = flipper_length_mm, color = species)`.
-

Section C — Two-way tables and proportions (Read 2 §3)

C1. Build a two-way table in R

Write the tidyverse pipeline that counts the number of penguins of each `species` on each `island`.

C2. Compute row proportions

Two-way count table (`species × island`) has row totals: Adelie=152, Chinstrap=68, Gentoo=124. Adelie on Biscoe=44. What is the row proportion of Adelie on Biscoe?

C3. Compute column proportions

Two-way count table (`species × island`): Biscoe column total=168 penguins, and Gentoo on Biscoe=124. What is the **column proportion** of Gentoo among Biscoe penguins?

C4. Stacked bar chart in R

Write the `ggplot2` call for a **100% stacked bar** of `island` (x) with `species` fill.

C5. `pivot_wider` usage

After `penguins %>% count(species, island)`, you want a wide table with `species` as rows and `islands` as columns. Which function converts it?

Answer key — Section C — Two-way tables and proportions (Read 2 §3)

1. `penguins %>% count(species, island)` — or `penguins %>% group_by(species, island) %>% summarize(n = n())`.
 2. $44/152 \approx 0.289$ — about 29% of Adelie penguins are on Biscoe Island.
 3. $124/168 \approx 0.738$ — about 74% of Biscoe penguins are Gentoo.
 4. `ggplot(penguins, aes(x = island, fill = species)) + geom_bar(position = "fill") + labs(y = "Proportion")`
 5. `pivot_wider(names_from = island, values_from = n)` — pivots island names to column headers and fills cells with counts.
-

Section D — Choosing the right graph/summary (integration)

D1. Match to variable types

Match each scenario to the correct graph:

- a) Penguin mass by species
- b) Bill length vs flipper length
- c) Species distribution on each island
- d) Distribution of flipper lengths alone

D2. When `group_by` before `summarize`

You have `penguins` and want the **mean** flipper length for each **sex**. Write the pipeline.

D3. `filter` before `plot`

You want a scatterplot of only Adelie penguins. What step comes before `ggplot`?

D4. `na.rm` in summaries

Why is `na.rm = TRUE` often needed inside `mean()` or `median()`?

Answer key — Section D — Choosing the right graph/summary (integration)

1. a) **Side-by-side boxplots**. b) **Scatterplot**. c) **Stacked bar chart**. d) **Histogram** or dot plot.
 2. `penguins %>% group_by(sex) %>% summarize(mean_flip = mean(flipper_length_mm, na.rm = TRUE))`
 3. `filter(species == 'Adelie')` — use `penguins %>% filter(species == 'Adelie') %>% ggplot(...)`.
 4. If any values are NA (missing), R returns NA for the whole mean by default. `na.rm = TRUE` removes missing values before computing the mean.
-

Section E — Project 3 workflow (demo lab R patterns)

E1. Load data from URL

You want to load a CSV file from a URL in R. Which function do you use, and where does the URL go?

E2. Recode a variable

Column `sex` has values 1 and 2 (numeric). You want 'male' and 'female'. Which tidyverse function and how?

E3. Handling missing values before plot


Before making a scatterplot of `bill_length_mm` vs `flipper_length_mm`, how do you drop rows with NA in either variable?

Answer key — Section E — Project 3 workflow (demo lab R patterns)

1. `read_csv('https://...')` from the **readr** package (loaded with `library(tidyverse)`).
 2. `mutate(sex = ifelse(sex == 1, 'male', 'female'))` — or `case_when()` for more than two levels.
 3. `filter(!is.na(bill_length_mm), !is.na(flipper_length_mm))` — or `drop_na(bill_length_mm, flipper_length_mm)`.
-


Section F — Graph interpretation (images)

IMG4. Histogram interpretation




What does this graph primarily display?

IMG5. Paired histograms




Which group appears to have the higher typical value?

IMG6. Single boxplot feature




In a boxplot, the horizontal line inside the box marks the:

IMG7. Side-by-side boxplots




Which group has larger spread (IQR)?

IMG8. Scatterplot form-direction-strength




Best description of this scatterplot?

IMG9. Bar plot variable type




The x-axis categories in a bar plot represent what variable type?

IMG10. Stacked bar counts



This is a stacked bar chart with raw counts. What is easiest to compare?

IMG11. 100% stacked bar



This is a 100% stacked bar chart. What is easiest to compare?

Answer key — Section F — Graph interpretation (images)

1. The distribution of **one numerical variable**.
 2. **Group A** (its distribution is shifted higher).
 3. The **median**.
 4. **Group B** (wider box / longer whiskers).
 5. Roughly **linear**, **positive** direction, **moderate-to-strong** association.
 6. **Categorical**.
 7. **Total counts** by group.
 8. **Within-group proportions** across groups.
-