

Math 340 / 640 Fall 2024

Final Examination

Professor Adam Kapelner

December 19, 2024

Full Name _____

Code of Academic Integrity

Since the college is an academic community, its fundamental purpose is the pursuit of knowledge. Essential to the success of this educational mission is a commitment to the principles of academic integrity. Every member of the college community is responsible for upholding the highest standards of honesty at all times. Students, as members of the community, are also responsible for adhering to the principles and spirit of the following Code of Academic Integrity.

Activities that have the effect or intention of interfering with education, pursuit of knowledge, or fair evaluation of a student's performance are prohibited. Examples of such activities include but are not limited to the following definitions:

Cheating Using or attempting to use unauthorized assistance, material, or study aids in examinations or other academic work or preventing, or attempting to prevent, another from using authorized assistance, material, or study aids. Example: using an unauthorized cheat sheet in a quiz or exam, altering a graded exam and resubmitting it for a better grade, etc.

I acknowledge and agree to uphold this Code of Academic Integrity.

signature

date

Instructions

This exam is 120 minutes (variable time per question) and closed-book. You are allowed **three** pages (front and back) of a “cheat sheet”, blank scrap paper (provided by the proctor) and a graphing calculator (which is not your smartphone). Please read the questions carefully. Within each problem, I recommend considering the questions that are easy first and then circling back to evaluate the harder ones. No food is allowed, only drinks.

Problem 1 Let $X_n \sim \text{Beta}(\alpha, n\beta)$ where $n \in \mathbb{N}, \alpha > 0, \beta > 0$ and $Y_n := nX_n$.

(a) [5 pt / 5 pts] What is $f_{X_n}(x)$? Your answer must be an explicit function of x without using a brand name. Don't forget the indicator function.

(b) [5 pt / 10 pts] What is $f_{Y_n}(y)$? Your answer must be an explicit function of y without using a brand name. Simplify as much as possible as it will help for the next question.

(c) [7 pt / 17 pts] $Y_n \xrightarrow{d} Y$. Find $f_Y(y)$ by taking the limit of $f_{Y_n}(y)$. If the answer is a brand name rv, mark it and find the value(s) of its parameter(s). You will need to use the following limit to solve that we didn't prove in class:

$$\lim_{n \rightarrow \infty} \frac{\Gamma(\alpha + n\beta)}{n^\alpha \Gamma(n\beta)} = \beta^\alpha$$

Problem 2 Some theoretical problems.

- (a) [7 pt / 24 pts] Let $X_1, \dots, X_n \stackrel{iid}{\sim} \text{Gamma}(\alpha, \beta)$ and $\mathbf{X} = [X_1 \ X_2 \ \dots \ X_n]^\top$, the column vector of all the X_i 's. Find the JDF of $\mathbf{X}$. Simplify as much as possible. Historical note: this is a type of standard multivariate gamma distribution.

- (b) [6 pt / 30 pts] Let $\mathbf{X}$ be a n -dimensional vector rv whose chf is below where α, β are positive constants:

$$\phi_{\mathbf{X}}(\mathbf{t}) = \left(\left(1 - \frac{it_1}{\beta}\right) \left(1 - \frac{it_2}{\beta}\right) \cdot \dots \cdot \left(1 - \frac{it_n}{\beta}\right) \right)^{-\alpha}$$

Find the distribution of X_1 , the first element of the vector. If it is a brand name rv, specify its name and its parameter(s).

- (c) [7 pt / 37 pts] Let $X \sim F_{a,b}$ and $Y = \frac{1}{2} \ln(X)$. Find the PDF of Y . Simplify as much as you can. Historical note: this is known as “Fisher’s Z distribution” which he discovered in 1924. Hint: the answer is not a standard normal.

- (d) [5 pt / 42 pts] Prove $\frac{B(\alpha + 1, \beta)}{B(\alpha, \beta)} = \frac{\alpha}{\alpha + \beta}$ using the properties of the gamma function.

- (e) [7 pt / 49 pts] Let $Y|X = x \sim \text{Poisson}(x)$ and $X \sim \text{Exp}(2)$. Find $p_Y(y)$ using a step-by-step derivation from scratch. Simplify enough to demonstrate that Y is distributed as a brand name rv that we studied in MATH 340. Specify its name and its parameter(s). Hint: use the gamma-factorial identity and one of the gamma integral identities.

- (f) [4 pt / 53 pts] Let $Y|X = x \sim \text{Exp}(x^{-1})$ and $X \sim \text{Gamma}(\alpha, \beta)$. Find $\mathbb{E}[Y]$. Recall that if $U \sim \text{Exp}(\lambda)$ then $\mathbb{E}[U] = 1/\lambda$.

- (g) [7 pt / 60 pts] Let $Y|X = x \sim \text{Binomial}(n, x)$ and $X \sim \text{Beta}(\alpha, \beta)$. Find $\text{Var}[Y]$. Your answer cannot be in sum form; it can only be a function n, α, β and fundamental constants. You do not need to simplify. Hints: $\mathbb{E}[X] = \frac{\alpha}{\alpha+\beta}$ and $\text{Var}[X] = \frac{\alpha\beta}{(\alpha+\beta)^2(\alpha+\beta+1)}$.

- (h) [6 pt / 66 pts] Let $X \sim F_{4,4}$. Find the numerical value of $\text{Mode}[X]$ exactly.

- (i) [4 pt / 70 pts] If a person's wealth in a certain country is modeled by $W \sim \text{ParetoI}(1, 1.005191)$ then fill in the following sentence as best as you can. Hint: guess different values until you get something approximate.

The top _____% own _____% of the wealth in that country.

- (j) [6 pt / 76 pts] If $X \sim \text{Weibull}(0.8, 3.7)$, circle the larger quantity. If both are the same, circle both:

$$\mathbb{P}(X > 1) \quad \text{or} \quad \mathbb{P}(X > 2 \mid X > 1)$$

If $X \sim \text{Weibull}(1.8, 3.7)$, circle the larger quantity. If both are the same, circle both:

$$\mathbb{P}(X > 1) \quad \text{or} \quad \mathbb{P}(X > 2 \mid X > 1)$$

If $X \sim \text{Weibull}(1.0, 3.7)$, circle the larger quantity. If both are the same, circle both:

$$\mathbb{P}(X > 1) \quad \text{or} \quad \mathbb{P}(X > 2 \mid X > 1)$$

- (k) [7 pt / 83 pts] Let $X_1, \dots, X_n \stackrel{iid}{\sim} \text{Weibull}(k, \lambda)$. Find $F_{X_{(1)}}(x)$. Simplify until you show that $X_{(1)}$ is a brand name rv that we studied in MATH 340, then specify its name and its parameter(s).

- (l) [7 pt / 90 pts] Let $X \sim \text{Laplace}(0, \pi)$. Let $Y = X \mid X > 0$. Show that Y is a brand name rv that we studied in MATH 340, then specify its name and its parameter(s).

- (m) [6 pt / 96 pts] Let $\mathbf{X} \sim \mathcal{N}_3([1 \ 2 \ 3]^\top, \mathbf{I}_3)$. Show that $\begin{bmatrix} X_2 \\ X_3 \end{bmatrix}$ is a brand name rv that we studied in MATH 340, then specify its name and its parameter(s).

- (n) [4 pt / 100 pts] Fill in the two missing values in the identity below:

$$\int_0^\infty \frac{\lambda^k}{\Gamma(k)} x^{k-1} e^{-\lambda x} dx = \sum_{x=0}^\infty \frac{\lambda^x e^{-\lambda}}{x!}$$