



A LETTERS JOURNAL EXPLORING
THE FRONTIERS OF PHYSICS

OFFPRINT

**Generation of platicons and frequency combs
in optical microresonators with normal GVD by
modulated pump**

VALERY E. LOBANOV, GRIGORY LIHACHEV and MICHAEL L.
GORODETSKY

EPL, **112** (2015) 54008

Please visit the website
www.epljournal.org

Note that the author(s) has the following rights:

- immediately after publication, to use all or part of the article without revision or modification, **including the EPLA-formatted version**, for personal compilations and use only;
- no sooner than 12 months from the date of first publication, to include the accepted manuscript (all or part), **but not the EPLA-formatted version**, on institute repositories or third-party websites provided a link to the online EPL abstract or EPL homepage is included.

For complete copyright details see: <https://authors.epletters.net/documents/copyright.pdf>.



A LETTERS JOURNAL EXPLORING
THE FRONTIERS OF PHYSICS

AN INVITATION TO SUBMIT YOUR WORK

www.epljournal.org

The Editorial Board invites you to submit your letters to EPL

EPL is a leading international journal publishing original, innovative Letters in all areas of physics, ranging from condensed matter topics and interdisciplinary research to astrophysics, geophysics, plasma and fusion sciences, including those with application potential.

The high profile of the journal combined with the excellent scientific quality of the articles ensures that EPL is an essential resource for its worldwide audience. EPL offers authors global visibility and a great opportunity to share their work with others across the whole of the physics community.

Run by active scientists, for scientists

EPL is reviewed by scientists for scientists, to serve and support the international scientific community. The Editorial Board is a team of active research scientists with an expert understanding of the needs of both authors and researchers.



www.epljournal.org

OVER

560,000

full text downloads in 2013

24 DAYS

average accept to online
publication in 2013

10,755

citations in 2013

*"We greatly appreciate
the efficient, professional
and rapid processing of
our paper by your team."*

Cong Lin
Shanghai University

Six good reasons to publish with EPL

We want to work with you to gain recognition for your research through worldwide visibility and high citations. As an EPL author, you will benefit from:

- 1 Quality** – The 50+ Co-editors, who are experts in their field, oversee the entire peer-review process, from selection of the referees to making all final acceptance decisions.
- 2 Convenience** – Easy to access compilations of recent articles in specific narrow fields available on the website.
- 3 Speed of processing** – We aim to provide you with a quick and efficient service; the median time from submission to online publication is under 100 days.
- 4 High visibility** – Strong promotion and visibility through material available at over 300 events annually, distributed via e-mail, and targeted mailshot newsletters.
- 5 International reach** – Over 2600 institutions have access to EPL, enabling your work to be read by your peers in 90 countries.
- 6 Open access** – Articles are offered open access for a one-off author payment; green open access on all others with a 12-month embargo.

Details on preparing, submitting and tracking the progress of your manuscript from submission to acceptance are available on the EPL submission website www.epletters.net.

If you would like further information about our author service or EPL in general, please visit www.epljournal.org or e-mail us at info@epljournal.org.

EPL is published in partnership with:



European Physical Society



Società Italiana di Fisica



EDP Sciences

IOP Publishing

IOP Publishing

Generation of platicons and frequency combs in optical microresonators with normal GVD by modulated pump

VALERY E. LOBANOV¹, GRIGORY LIHACHEV^{1,2} and MICHAEL L. GORODETSKY^{1,2}

¹ *Russian Quantum Center - Skolkovo 143025, Russia*

² *Faculty of Physics, M. V. Lomonosov Moscow State University - Moscow 119991, Russia*

received 10 August 2015; accepted in final form 7 December 2015

published online 29 December 2015

PACS 42.60.Da – Resonators, cavities, amplifiers, arrays, and rings

PACS 42.65.Ky – Frequency conversion; harmonic generation, including higher-order harmonic generation

PACS 42.65.Tg – Optical solitons; nonlinear guided waves

Abstract – We demonstrate that flat-topped dissipative solitonic pulses, “platicons”, and corresponding frequency combs can be excited in optical microresonators with normal group velocity dispersion using either the amplitude modulation of the pump or the bichromatic pump. Soft excitation may occur in a particular frequency range if the modulation depth is large enough and the modulation frequency is close to the free spectral range of the microresonator.



Copyright © EPLA, 2015

Optical frequency combs generated in mode-locked lasers have revolutionized optical measurements. Last years optical microresonators have been attracting growing attention as a promising platform for microresonator-based Kerr frequency combs [1–4]. It was demonstrated both theoretically and experimentally that frequency comb consisting of equidistant optical lines in spectral domain may appear in a nonlinear microresonator pumped by a continuous wave (c.w.) laser due to the cascaded four-wave mixing processes. The frequency spacing between comb lines corresponds to the free spectral range (FSR, typically tens of GHz up to THz) of the microresonator which is the inverted roundtrip time of light in the cavity. Microresonator-based combs were shown to perform at the level required for optical-frequency metrology applications [5] and for optical communications [6]. However, such systems often suffer from significant frequency and amplitude noise [7,8] due to the formation of sub-combs [9]. In contrast to conventional mode-locked laser-based frequency combs microresonator combs do not correspond, generally, to stable ultrashort pulses in the time domain because of arbitrary phase relations between the comb lines obtained in the process of formation. Kerr solitons solve this problem with a c.w. laser beam converted into a train of pulses, corresponding to a low-noise frequency comb having smooth spectral envelope in the spectral domain. This regime was demonstrated experimentally in optical crystalline and integrated ring

microresonators with anomalous group velocity dispersion (GVD) [10,11]. The limitation of this approach is the difficulty to obtain anomalous GVD in broad band for arbitrary centered wavelength in microresonators since material GVD in the visible and near IR is mostly normal. In this way, the development of new methods that enable the implementation of materials with normal GVD for frequency comb generation is interesting.

Frequency combs in normal GVD microresonators are theoretically possible [12–14]. Besides, narrow comb-like spectra from normal GVD crystalline resonators [15–17] and integrated microrings [18] have been demonstrated experimentally. Microresonators with normal GVD can also support dark optical solitons [14,19].

A novel type of solitonic pulses, “platicons”, was predicted in microresonators with normal dispersion under the condition of the shifted pump mode frequency resonance [20]. In real microresonators such frequency shift may occur either due to the normal mode coupling between different mode families [19,21,22], or as a result of self-injection locking [16], both providing equivalent local anomalous dispersion [21–24]. It was demonstrated that one may continuously change in wide range the duration of generated platicons varying the pump detuning and that the generation of platicons is significantly more efficient than the generation of bright soliton trains in microresonators for the same absolute value of anomalous GVD in terms of conversion of the c.w. power into the power of the




Fig. 1: (Color online) (a) Averaged intracavity intensity U *vs.* normalized detuning ζ_0 at $f = 4.11$ for different relative sideband pump amplitudes ε . (b) Profiles and (c) spectra of low-contrast wave (black line) and platonic (red line) at $\varepsilon = 0.15$, $f = 4.11$. Excitation domains (d) for different primary pump amplitudes f and (e) for different relative sideband pump amplitudes ε . (f) Platonic generation critical value of relative sideband pump amplitude *vs.* primary pump amplitude. In all cases $D_2/\kappa \approx -1.036 \times 10^{-2}$, $\Delta = 0$.

comb [20]. The problem with the proposed method is that it requires an artificial perturbation of the local dispersion. In this paper we show that platonic generation is possible even in the absence of the pump mode shift when a bichromatic or amplitude-modulated pump is used. This method is efficient if the pump modulation frequency or the frequency difference between two pump waves is close to one or several FSRs. The proposed approach is experimentally feasible since comb generation from a bichromatic pump has been already studied experimentally [25] and theoretically [26] in the case of anomalous dispersion. Recently, frequency comb generation by dual-pump input signal was also discussed in normally dispersive optical fibers [27].

We consider first the case of the amplitude-modulated pump $|P| = P_0 \{1 + 2\varepsilon \cos(\Omega t)\}^2$ with modulation

frequency Ω and relative amplitude of the sideband pump ε . Our numerical model is based on the system of dimensionless coupled nonlinear mode equations [9,28] modified to take into account the complexity of the pump field. Assuming that the pump modulation frequency is close to one FSR, we have

$$\frac{\partial a_\mu}{\partial \tau} = -(1 + i\zeta_\mu) a_\mu + i \sum_{\mu' \leq \mu''} (2 - \delta_{\mu'\mu''}) a_{\mu'} a_{\mu''} a_{\mu'+\mu''-\mu}^* + \delta_{0\mu} f + \varepsilon \{ \delta_{1\mu} f \exp(i\Delta\tau) + \delta_{-1\mu} f \exp(-i\Delta\tau) \}. \quad (1)$$

Here $\delta_{\mu'\mu''}$ is the Kronecker delta, a_μ is the slowly varying amplitude of the comb modes for the mode frequency ω_μ , $\tau = \kappa t/2$ denotes the normalized time, $\kappa = \omega_p/Q$ is the cavity decay rate, Q is the loaded quality factor, ω_p is




Fig. 2: (Color online) (a) Averaged intracavity intensity U of platons *vs.* normalized detuning ζ_0 at $f = 4.11$ for $\varepsilon = 0.15$ (black line) and $\varepsilon = 0.25$ (blue line). Thin red lines indicate the results of soft excitation. (b) Profiles and (c) spectra of platons for different values of ζ_0 at $\varepsilon = 0.25$, $f = 4.11$. (d) Averaged intracavity intensity U of platons *vs.* normalized detuning ζ_0 for different values of f at $\varepsilon = 0.25$. In all cases $D_2/\kappa \approx -1.036 \times 10^{-2}$, $\Delta = 0$.

the pump frequency; f stands for the dimensionless pump amplitude, $\Delta = 2(D_1 - \Omega)/\kappa$ is the normalized modulation frequency mismatch, D_1 is FSR. All mode numbers μ are defined relative to the pumped mode. We consider the Taylor expansion of the dispersion law $\omega_\mu = \omega_0 + D_1\mu + \frac{1}{2}D_2\mu^2 + \dots$ and neglecting third-order dispersion we get the following expressions for the normalized detunings: $\zeta_0 = 2(\omega_0 - \omega_p)/\kappa$, $\zeta_\mu = \zeta_0 + (D_2/\kappa)\mu^2$. Note that $D_2 < 0$ for normal GVD. For the numerical analysis we consider the range of parameters corresponding to experimentally feasible (realistic) microresonators: $f \in [1.5; 6.5]$ (the value $f = 1$ corresponds to bistability and comb threshold [10,14]), $D_2/\kappa \in [-5 \times 10^{-2}; -10^{-3}]$, $\zeta_0 \in [0; 15]$. For example, for MgF_2 microresonator [16] ($\lambda = 780 \text{ nm}$, $Q = 2.5 \times 10^9$, $D_2/2\pi = -2.96 \text{ kHz}$) $D_2/\kappa \approx -0.019$ and the pump power of 1 mW corresponds to $f \approx 3.83$; for Si_3N_4 microrings [18] ($\lambda = 1550 \text{ nm}$, $Q = 1.1 \times 10^6$, $D_2/2\pi = -300 \text{ kHz}$) $D_2/\kappa \approx -0.0017$ and the pump of 1 W corresponds to $f \approx 2.76$.

In our simulations we used initial weak noise-like inputs for seeding. Coupled-mode equations for 501 modes were numerically propagated in time using the Runge-Kutta integrator. Nonlinear terms were calculated using a fast method proposed in [29]. We also checked that results do not change with the increase of the number of modes. For the analysis we calculated the average

intracavity intensity $U = \sum_\mu |a_\mu|^2$ for different values of the normalized detuning ζ_0 (see fig. 1(a)) and corresponding waveforms $\psi(\phi) = \sum_\mu a_\mu \exp(i\mu\phi)$.

For the resonant modulation ($\Delta = 0$) it was found that at $\varepsilon \neq 0$ due to the presence of the sideband pump conventional c.w. solutions turn into low-contrast waves with narrow spectrum (see figs. 1(b) and (c)). For small sideband pump amplitudes ε internal intensity *vs.* detuning dependence has a conventional triangular shape and only low-amplitude sideband modes may be observed. However, if the sideband pump amplitude is large enough, this dependence deviates from the expected triangular shape, the characteristic step-like dependence of intensity *vs.* detuning appears and at a particular frequency range, in the vicinity of this step, one may observe the generation of high-contrast optical pulses with a pronounced dip located at the minima of the low-contrast wave (see fig. 1(b)) analogously to platons [20]. Platicon generation is accompanied by a dramatic spectrum widening (fig. 1(c)). Similarly to experimental spectra observed in [16] these are characterized by two pronounced wings (figs. 1(c) and 2(c)). Thus, the amplitude modulation of the pump power may provide soft excitation (from noise-like inputs) of platons without a tricky laser scanning necessary for bright solitons in case of anomalous GVD [10]. Such dramatic profile and spectrum

transformation can be explained by the phenomenon of optical wave breaking [30] due to the double shock formation described in ref. [27]. Shock formation is accompanied by the emergence of multiple frequency components that mix nonlinearly to produce new frequency components by four-wave mixing resulting in a significant spectrum broadening.

Generated platons were found to be stable against perturbations. To elucidate their stability we propagated them up to large periods of time (up to $\tau \sim 500$) in the presence of added white-noise-like perturbations ($U_{noise} \sim 0.1$) by solving the governing equations (eq. (1)). Platons rapidly (at time periods ~ 10) clean up the noise and propagate in a stable fashion over indefinitely large periods of time.

Platons generated in optical microresonators look similar to “flatons” demonstrated in optical fibers with normal GVD [31] or dispersive shock waves in an externally driven passive Kerr resonator with weak normal dispersion [32]. However, in contrast to quickly evolving flatons and shock waves platons are stationary soliton-like waveforms in time. It should be noted that in addition to the coupled-mode approach platons may be also described by the Lugiato-Lefever equation (driven and damped nonlinear Schrödinger equation) [14,20] as it was done for dispersive shock waves [32]. However, the considered microresonators are comparatively small and pulse duration is comparable with the roundtrip time. Thus, periodic boundary conditions play an important if not determining role providing stationary solutions.

We found also that the growth of the sideband pump amplitude results in a widening of the excitation domain (fig. 1(d)). The position of the excitation domain depends on the primary pump amplitude f : for larger amplitude values the excitation domain shifts to larger values of ζ_0 and becomes narrower (fig. 1(e)). It is interesting to note, that for a larger primary pump amplitude platon generation requires larger values of the relative sideband pump amplitude ε (fig. 1(f)).

In order to study the possibility of soft excitation by a phase-modulated pump we modified eq. (1) as

$$\frac{\partial a_\mu}{\partial \tau} = -(1 + i\zeta_\mu) a_\mu + i \sum_{\mu' \leq \mu''} (2 - \delta_{\mu' \mu''}) a_{\mu'} a_{\mu''} a_{\mu' + \mu'' - \mu}^* + \delta_{0\mu} f + \varepsilon \{ \delta_{1\mu} f \exp(i\Delta\tau) - \delta_{-1\mu} f \exp(-i\Delta\tau) \}. \quad (2)$$

and found that phase modulation does not lead to the generation of platons. Contrastingly, temporal cavity bright soliton excitation via direct phase modulation of the cavity driving field was demonstrated numerically [33] and experimentally [34].

Using an initial platon as an input for our simulations we found all possible platon solutions (fig. 2(a)). Interestingly, the excitation domain, defined as detuning range where platon excitation from noise-like inputs is possible, is significantly narrower than the full existence domain and lies at its lower bound (in terms of ζ_0) (compare

widths of characteristic steps on thin red lines in fig. 2(a) corresponding to excitation domains and widths of existence domains where the thick blue and black lines lie). Moreover, it was revealed that noise-like inputs provide the excitation of wide platons only. However, having a wide platon, generated inside the excitation domain, one may further find all other possible platon solutions varying slowly the pump frequency (or the detuning value ζ_0) and, consequently, may determine full existence domain. It should be noticed that if several platon solutions may exist at the same pump frequency, soft excitation provides the generation of a low-power platon.

It was found that while the sideband pump amplitude increases, the existence domain of platons becomes wider and the energy spectrum contains a smaller number of steps. In this way, in the considered system the parameter ε behaves analogously to the parameter describing the pump mode shift in [20] and similarly the growth of ε results in the transformation of the discrete energy spectrum of dark solitons into the quasi-continuous spectrum of platons.

We revealed that similarly to platons described in [20] one can effectively control the duration and, consequently, the power of generated pulses by slow pump frequency tuning. Figure 2(b) shows that while the pump frequency decreases (the corresponding detuning ζ_0 increases), the pulse duration also decreases. Consequently, the width of the corresponding frequency comb may also be tuned (fig. 2(c)).

It was found that while the pump power decreases, the existence domain of platons becomes narrower and shifts to smaller values of ζ_0 (fig. 2(d)). In this way, one can tune the platon duration varying the primary pump power: decreasing the pump one may generate shorter pulses.

To prove that the proposed approach is applicable for a wide range of materials used for the fabrication of microresonators with normal GVD we make calculations for a wide range of GVD values from $D_2/\kappa \approx -1.036 \times 10^{-3}$ up to $D_2/\kappa \approx -4.142 \times 10^{-2}$. It was found that the growth of the absolute value of D_2 results in a weak increase of the sideband pump power necessary for platon generation (see fig. 3(a)). The decrease of the absolute value of the dispersion coefficient results in more localized pulses with sharper profiles and wider spectrum (figs. 3(b) and (c)). The positions of spectrum peaks and, consequently, the spectrum width may be estimated rather accurately by the simple phase-matching condition [35]: $\zeta_\mu = \zeta_0 + (D_2/\kappa) \mu^2 = 0$.

Also, while the existence domain weakly depends on D_2 , the increase of the absolute GVD value leads to a more pronounced discreteness of the platon energy spectrum (fig. 3(d)).

We also found that the method considered is very sensitive to the pump modulation frequency. While the mismatch between modulation frequency and FSR increases, existence (fig. 4(a)) and excitation (fig. 4(b)) domains




Fig. 3: (Color online) (a) Excitation domains for different values of GVD at $f = 4.11$. (b) Profiles and (c) spectra of platons for different values of GVD at $\varepsilon = 0.25$, $f = 4.11$, $\zeta_0 = 14$. (d) Averaged intracavity intensity U of platons *vs.* normalized detuning ζ_0 at $f = 4.11$, $\varepsilon = 0.25$ for different values of GVD. In all cases $\Delta = 0$.




Fig. 4: (Color online) (a) Averaged intracavity intensity U of platons *vs.* normalized detuning ζ_0 at $f = 4.11$, $\varepsilon = 0.25$ for different values of the modulation frequency mismatch Δ . (b) Excitation domain *vs.* mismatch value at $f = 4.11$, $\varepsilon = 0.25$ (black line) and $\varepsilon = 0.225$ (red line). $\varepsilon = 0.25$. In all cases $D_2/\kappa \approx 1.036 \times 10^{-2}$.

become significantly narrower and disappear at some critical value of mismatch. The critical for platicon excitation mismatch value increases with the sideband pump amplitude ε (see fig. 4(b)). This may be explained by the inefficiency of sideband pumping at large mismatches. In this case pump becomes effectively monochromatic and, thus, it cannot provide soft excitation of platons or dark solitons at the considered dispersion law as it was shown earlier [14,20].

It was also revealed that for frequency combs generated by the amplitude-modulated pump the spacing

between the comb lines is equal to the modulation frequency but not to the FSR of a microresonator. Interestingly, but not surprisingly, if the modulation frequency is equal to an integer number of FSR, the corresponding number of platons per roundtrip can be generated.

Rewriting eq. (1) in the following form:

$$\begin{aligned} \frac{\partial a_\mu}{\partial \tau} = & -(1 + i\zeta_\mu) a_\mu + i \sum_{\mu' \leq \mu''} (2 - \delta_{\mu'\mu''}) a_{\mu'} a_{\mu''} a_{\mu'+\mu''-\mu}^* \\ & + \delta_{0\mu} f + \tilde{\varepsilon} \delta_{1\mu} f \exp(i\Delta\tau), \end{aligned} \quad (3)$$

we also studied the usage of a biharmonic pump for platicon generation. It was found that the results obtained for a biharmonic pump practically coincide with the results for the amplitude-modulated pump if $\tilde{\varepsilon} \approx 2\varepsilon$. The main difference is that in contrast with the amplitude-modulated pump biharmonic pump is sensitive to the sign of modulation frequency mismatch (compare eqs. (1) and (3)). Interestingly, platicon generation is possible even if there is a constant phase shift between two pumping waves.

To summarize, we showed that soft excitation (from noise-like inputs) of platicons is possible for an amplitude-modulated or a biharmonic pump if the sideband pump power is large enough. For the effective platicon excitation the modulation frequency (or the frequency difference between two pump waves) should be close to the FSR of a microresonator. The excitation domain strongly depends on the sideband pump power and platicon duration may be altered significantly by tuning the pump frequency.

* * *

The work was supported by the Ministry of Education and Science of the Russian Federation (project No. 14.585.21.0005).

REFERENCES

- [1] DEL'HAYE P., SCHLIESSER A. *et al.*, *Nature*, **450** (2007) 1214.
- [2] SAVCHENKOV A. A., MATSKO A. B. *et al.*, *Phys. Rev. Lett.*, **101** (2008) 93902.
- [3] KIPPENBERG T. J., HOLZWARTH R. and DIDDAMS S. A., *Science*, **332** (2011) 555.
- [4] LI J., LEE H. *et al.*, *Phys. Rev. Lett.*, **109** (2012) 233901.
- [5] DEL'HAYE P., PAPP S. B. and DIDDAMS S. A., *Phys. Rev. Lett.*, **109** (2012) 263901.
- [6] PFEIFLE J., BRASCH V. *et al.*, *Nat. Photon.*, **8** (2014) 375.
- [7] FERDOUS F., MIAO H. X., LEAIRD D. E., SRINIVASAN K., WANG J., CHEN L., VARGHESE L. T. and WEINER A. M., *Nat. Photon.*, **5** (2011) 770.
- [8] DEL'HAYE P., HERR T. *et al.*, *Phys. Rev. Lett.*, **107** (2011) 063901.
- [9] HERR T., HARTINGER K. *et al.*, *Nat. Photon.*, **6** (2012) 48.
- [10] HERR T., BRASCH V. *et al.*, *Nat. Photon.*, **8** (2014) 145.
- [11] BRASCH V., HERR T. *et al.*, <http://arxiv.org/abs/1410.8598> (2014).
- [12] MATSKO A. B., SAVCHENKOV A. A. and MALEKI L., *Opt. Lett.*, **37** (2012) 43.
- [13] HANSSON T., MODOTTO D. and WABNITZ S., *Phys. Rev. A*, **88** (2013) 023819.
- [14] GODEY C., BALAKIREVA I. *et al.*, *Phys. Rev. A*, **89** (2014) 063814.
- [15] COILLET A., BALAKIREVA I. *et al.*, *IEEE Photon. J.*, **5** (2013) 6100409.
- [16] LIANG W., SAVCHENKOV A. A. *et al.*, *Opt. Lett.*, **39** (2014) 2920.
- [17] HENRIET R., LIN G. *et al.*, *Opt. Lett.*, **40** (2015) 1567.
- [18] HUANG S. W., ZHOU H. *et al.*, *Phys. Rev. Lett.*, **114** (2015) 053901.
- [19] XUE X., XUAN Y. *et al.*, *Nat. Photon.*, **9** (2015) 594.
- [20] LOBANOV V. E., LIHACHEV G. *et al.*, *Opt. Express*, **23** (2015) 7713.
- [21] XUE X., XUAN Y. *et al.*, *Laser Photon. Rev.*, **9** (2015) L-23.
- [22] HERR T., BRASCH V. *et al.*, *Phys. Rev. Lett.*, **113** (2014) 123901.
- [23] SAVCHENKOV A. A., MATSKO A. B. *et al.*, *Opt. Express*, **20** (2012) 27290.
- [24] LIU Y., XUAN Y. *et al.*, *Optica*, **1** (2014) 137.
- [25] STREKALOV D. V. and YU N., *Phys. Rev. A*, **79** (2009) 041805(R).
- [26] HANSSON T. and WABNITZ S., *Phys. Rev. A*, **90** (2014) 013811.
- [27] ANTIKAINEN A. and AGRAWAL G. P., *J. Opt. Soc. Am. B*, **32** (2015) 1705.
- [28] CHEMBO Y. K. K. and YU N., *Phys. Rev. A*, **82** (2010) 33801.
- [29] HANSSON T., MODOTTO D., WABNITZ S., *Opt. Commun.*, **312** (2014) 134.
- [30] FINOT C., KIBLER B. *et al.*, *J. Opt. Soc. Am. B*, **25** (2008) 1938.
- [31] VARLOT B., WABNITZ S. *et al.*, *Opt. Lett.*, **38** (2013) 3899.
- [32] MALAGUTI S., BELLANCA G. and TRILLO S., *Opt. Lett.*, **39** (2014) 2475.
- [33] TAHERI H., EFTEKHAR A. A. *et al.*, *IEEE Photon. J.*, **7** (2015) 2200309.
- [34] JANG J. K., ERKINTALO M. *et al.*, *Opt. Lett.*, **40** (2015) 4755.
- [35] COEN S. and HAELTERMAN M., *Phys. Rev. Lett.*, **79** (1997) 4139.