

# COMPSCI 590N

## Lecture 10: Probability 2

Roy J. Adams

College of Information and Computer Sciences  
University of Massachusetts Amherst

# Outline

## 1 Visualizing Distributions

## 2 Sampling

## 3 Sources of Randomness

# Histograms and Counting

- An important tool for exploring data is visualizing the distribution it came from.

# Histograms and Counting

- An important tool for exploring data is visualizing the distribution it came from.
  - Allows us to verify and inform modeling decisions.

# Histograms and Counting

- An important tool for exploring data is visualizing the distribution it came from.
  - Allows us to verify and inform modeling decisions.
  - Summary statistics (e.g. mean, standard deviation, skewness, kurtosis, etc.), while important and useful, discard information about the data.

# Histograms and Counting


- An important tool for exploring data is visualizing the distribution it came from.
  - Allows us to verify and inform modeling decisions.
  - Summary statistics (e.g. mean, standard deviation, skewness, kurtosis, etc.), while important and useful, discard information about the data.
- A flexible tool for visualizing data distributions is the histogram.

# Histograms and Counting

- An important tool for exploring data is visualizing the distribution it came from.
  - Allows us to verify and inform modeling decisions.
  - Summary statistics (e.g. mean, standard deviation, skewness, kurtosis, etc.), while important and useful, discard information about the data.
- A flexible tool for visualizing data distributions is the histogram.

# Histograms and Counting

- An important tool for exploring data is visualizing the distribution it came from.
  - Allows us to verify and inform modeling decisions.
  - Summary statistics (e.g. mean, standard deviation, skewness, kurtosis, etc.), while important and useful, discard information about the data.
- A flexible tool for visualizing data distributions is the histogram.





# Histograms and Counting

Numpy provides functions for counting and computing histograms.

```
>>> import numpy as np
>>> A = np.random.randn(1000)
>>> np.histogram(A,nbins=5)
>>> counts,bin_edges = np.histogram(A,bins=7)
>>> counts
array([ 38, 180, 381, 289,  98,  11,   3])
>>> bin_edges
array([-2.75925765, -1.77917762, -0.7990976 ,
        0.18098242,  1.16106244,
        2.14114246,  3.12122248,  4.1013025 ])
```

# Histograms in Python

- There are a variety of tools for plotting distributions:

# Histograms in Python

- There are a variety of tools for plotting distributions:
  - `matplotlib.pyplot.histogram` computes and plots a histogram from an array.

# Histograms in Python

- There are a variety of tools for plotting distributions:
  - `matplotlib.pyplot.histogram` computes and plots a histogram from an array.
  - `pandas.DataFrame.hist` plots a histogram directly from a `pandas.DataFrame`.

# Histograms in Python

- There are a variety of tools for plotting distributions:
  - `matplotlib.pyplot.histogram` computes and plots a histogram from an array.
  - `pandas.DataFrame.hist` plots a histogram directly from a `pandas.DataFrame`.
  - Seaborn is an extension to `matplotlib` design explicitly for plotting distributions.

# Outline

1 Visualizing Distributions

2 Sampling

3 Sources of Randomness

# Applications of Randomness

Randomness is a core operation for many commonly used algorithms:

# Applications of Randomness

Randomness is a core operation for many commonly used algorithms:

- Cryptography and encryption.



# Applications of Randomness

Randomness is a core operation for many commonly used algorithms:

- Cryptography and encryption.
- Randomized storage (e.g. random hash tables used in Python dictionaries).

# Applications of Randomness

Randomness is a core operation for many commonly used algorithms:

- Cryptography and encryption.
- Randomized storage (e.g. random hash tables used in Python dictionaries).
- Randomized communications (e.g. LAN).

# Applications of Randomness

Randomness is a core operation for many commonly used algorithms:

- Cryptography and encryption.
- Randomized storage (e.g. random hash tables used in Python dictionaries).
- Randomized communications (e.g. LAN).
- Approximate integration (we will look at this later).

# Applications of Randomness

Randomness is a core operation for many commonly used algorithms:

- Cryptography and encryption.
- Randomized storage (e.g. random hash tables used in Python dictionaries).
- Randomized communications (e.g. LAN).
- Approximate integration (we will look at this later).
- Randomized experiments.

# Applications of Randomness

Randomness is a core operation for many commonly used algorithms:

- Cryptography and encryption.
- Randomized storage (e.g. random hash tables used in Python dictionaries).
- Randomized communications (e.g. LAN).
- Approximate integration (we will look at this later).
- Randomized experiments.
- Statistical modeling:

# Applications of Randomness

Randomness is a core operation for many commonly used algorithms:

- Cryptography and encryption.
- Randomized storage (e.g. random hash tables used in Python dictionaries).
- Randomized communications (e.g. LAN).
- Approximate integration (we will look at this later).
- Randomized experiments.
- Statistical modeling:
  - Evaluating models.

# Applications of Randomness

Randomness is a core operation for many commonly used algorithms:

- Cryptography and encryption.
- Randomized storage (e.g. random hash tables used in Python dictionaries).
- Randomized communications (e.g. LAN).
- Approximate integration (we will look at this later).
- Randomized experiments.
- Statistical modeling:
  - Evaluating models.
  - Approximate inference.

# Applications of Randomness

Randomness is a core operation for many commonly used algorithms:

- Cryptography and encryption.
- Randomized storage (e.g. random hash tables used in Python dictionaries).
- Randomized communications (e.g. LAN).
- Approximate integration (we will look at this later).
- Randomized experiments.
- Statistical modeling:
  - Evaluating models.
  - Approximate inference.
  - Approximate estimation.



# Applications of Randomness

Randomness is a core operation for many commonly used algorithms:

- Cryptography and encryption.
- Randomized storage (e.g. random hash tables used in Python dictionaries).
- Randomized communications (e.g. LAN).
- Approximate integration (we will look at this later).
- Randomized experiments.
- Statistical modeling:
  - Evaluating models.
  - Approximate inference.
  - Approximate estimation.
- **Many** other randomized algorithms.

# Random Sampling

- Problem: Given a distribution  $p(x)$  we would like to generate samples from this distribution.

# Random Sampling

- Problem: Given a distribution  $p(x)$  we would like to generate samples from this distribution.
- Example: To generate samples from a Bernoulli distribution (binary), flip a bunch of coins. The result of each coin flip is a sample from the Bernoulli.

# Types of Sampling

As you might imagine, sampling has been the focus of intense study for a long time. Some common sampling algorithms are:

# Types of Sampling

As you might imagine, sampling has been the focus of intense study for a long time. Some common sampling algorithms are:

- Inverse CDF sampling (we will look at this algorithm).

# Types of Sampling

As you might imagine, sampling has been the focus of intense study for a long time. Some common sampling algorithms are:

- Inverse CDF sampling (we will look at this algorithm).
- Rejection sampling.

# Types of Sampling

As you might imagine, sampling has been the focus of intense study for a long time. Some common sampling algorithms are:

- Inverse CDF sampling (we will look at this algorithm).
- Rejection sampling.
- Markov Chain Monte Carlo (MCMC).

# Types of Sampling

As you might imagine, sampling has been the focus of intense study for a long time. Some common sampling algorithms are:

- Inverse CDF sampling (we will look at this algorithm).
- Rejection sampling.
- Markov Chain Monte Carlo (MCMC).
- Metropolis-Hastings.



# Types of Sampling

As you might imagine, sampling has been the focus of intense study for a long time. Some common sampling algorithms are:

- Inverse CDF sampling (we will look at this algorithm).
- Rejection sampling.
- Markov Chain Monte Carlo (MCMC).
- Metropolis-Hastings.
- Slice sampling.

# Types of Sampling

As you might imagine, sampling has been the focus of intense study for a long time. Some common sampling algorithms are:

- Inverse CDF sampling (we will look at this algorithm).
- Rejection sampling.
- Markov Chain Monte Carlo (MCMC).
- Metropolis-Hastings.
- Slice sampling.
- The list goes on.

# Cumulative Distribution Functions

Given a distribution  $p(x)$  for a random variable  $X$ , the associated Cumulative Distribution Function (CDF)  $F(x)$ , gives the probability that  $X$  is less than some value, i.e.  $F(x) = P(X \leq x)$ .

# Cumulative Distribution Functions

Given a distribution  $p(x)$  for a random variable  $X$ , the associated Cumulative Distribution Function (CDF)  $F(x)$ , gives the probability that  $X$  is less than some value, i.e.  $F(x) = P(X \leq x)$ .

## Cumulative Distribution Function

The CDF  $F(x)$  for a distribution  $p(x)$  is given by,

$$F(x) = \int_{-\infty}^x p(x') dx'$$

for continuous variables and by

$$F(x) = \sum_{x'=-\infty}^x P(x')$$

for discrete variables.

# Cumulative Distribution Functions: Multinomial

For example: Consider the CDF for  $Multinomial(1, \frac{1}{6}, \dots, \frac{1}{6})$  (i.e. a six sided dice roll).

# Cumulative Distribution Functions: Multinomial

For example: Consider the CDF for *Multinomial* $(1, \frac{1}{6}, \dots, \frac{1}{6})$  (i.e. a six sided dice roll).

$$F(x) = P(X \leq x) = \sum_{i=1}^x \frac{1}{6}$$

$$F(3) = P(X \leq 3) = \frac{3}{6} = \frac{1}{2}$$

# Cumulative Distribution Functions: Multinomial

For example: Consider the CDF for *Multinomial*(1,  $\frac{1}{6}$ , ...,  $\frac{1}{6}$ ) (i.e. a six sided dice roll).

$$F(x) = P(X \leq x) = \sum_{i=1}^x \frac{1}{6}$$

$$F(3) = P(X \leq 3) = \frac{3}{6} = \frac{1}{2}$$

# Sampling Algorithms: Inverse CDF Sampling

One of the most fundamental sampling algorithms is inverse CDF sampling. For any invertible CDF, we can use the following algorithm.



# Sampling Algorithms: Inverse CDF Sampling

One of the most fundamental sampling algorithms is inverse CDF sampling. For any invertible CDF, we can use the following algorithm.

## Inverse CDF Sampling

Let  $F(x)$  be an invertible CDF. Then we can sample from the associated distribution by:

- 1 Sample  $u^{(s)}$  from a Uniform(0,1).
- 2  $x^{(s)} = F^{-1}(U)$

# Monte Carlo Integration

- **Monte Carlo Integration** is a method for approximating the value of an integral that is difficult (or impossible) to compute.

# Monte Carlo Integration

- **Monte Carlo Integration** is a method for approximating the value of an integral that is difficult (or impossible) to compute.
- The core idea is based on approximating **expected values**.

# Expected Values

A common computation when working with probability distributions is an expected value.

# Expected Values

A common computation when working with probability distributions is an expected value.

## Expected Value

Given a distribution  $p(x)$  over a random variable  $X$  and a function of the random variable  $f : \Omega \rightarrow \mathbb{R}$ , the expected value of the function is defined as

$$\mathbb{E}[f(X)] = \int_{\Omega} p(x)f(x)dx$$

where the integral is a sum when  $X$  is discrete.

# Expected Values: Multinomial

- Let  $X$  be the result of a fair six-sided dice roll.

# Expected Values: Multinomial

- Let  $X$  be the result of a fair six-sided dice roll.
- What is  $\mathbb{E}[X]$ ?

# Expected Values: Multinomial

- Let  $X$  be the result of a fair six-sided dice roll.
- What is  $\mathbb{E}[X]$ ?



# Expected Values: Multinomial

- Let  $X$  be the result of a fair six-sided dice roll.
- What is  $\mathbb{E}[X]$ ?

$$\begin{aligned}\mathbb{E}[X] &= \sum_x P(X = x) \cdot x \\ &= \sum_{x=1}^6 \frac{1}{6} \cdot x \\ &= \frac{1}{6}(1 + 2 + 3 + 4 + 5 + 6) = 3.5\end{aligned}$$

# Expected Values: Normal

- Let  $X$  be a Normally distributed random variable with mean  $\mu$  and standard deviation  $\sigma$ .

# Expected Values: Normal

- Let  $X$  be a Normally distributed random variable with mean  $\mu$  and standard deviation  $\sigma$ .
- What is  $\mathbb{E}[X^2]$ ?

# Expected Values: Normal

- Let  $X$  be a Normally distributed random variable with mean  $\mu$  and standard deviation  $\sigma$ .
- What is  $\mathbb{E}[X^2]$ ?

# Expected Values: Normal

- Let  $X$  be a Normally distributed random variable with mean  $\mu$  and standard deviation  $\sigma$ .
- What is  $\mathbb{E}[X^2]$ ?

$$\begin{aligned}\mathbb{E}[X^2] &= \int_{-\infty}^{\infty} \frac{1}{\sigma\sqrt{2\pi}} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right) x^2 dx \\ &= \frac{1}{\sigma\sqrt{2\pi}} \int_{-\infty}^{\infty} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right) x^2 dx \\ &= \frac{1}{\sigma\sqrt{2\pi}} \int_{-\infty}^{\infty} \exp\left(-\frac{(x-\mu)^2}{2\sigma^2}\right) x^2 dx\end{aligned}$$

# The Importance of Expected Values

# Difficult Expected Values

- In many cases, the expected value is too complex to evaluate.

# Difficult Expected Values

- In many cases, the expected value is too complex to evaluate.
- This may occur because:



# Difficult Expected Values

- In many cases, the expected value is too complex to evaluate.
- This may occur because:
  - The normalization constant of the distribution is difficult to evaluate.

# Difficult Expected Values

- In many cases, the expected value is too complex to evaluate.
- This may occur because:
  - The normalization constant of the distribution is difficult to evaluate.
  - There is not analytical solution to the integral.

# Difficult Expected Values

- In many cases, the expected value is too complex to evaluate.
- This may occur because:
  - The normalization constant of the distribution is difficult to evaluate.
  - There is not analytical solution to the integral.
- What do we do? Approximate!

# Difficult Expected Values

- In many cases, the expected value is too complex to evaluate.
- This may occur because:
  - The normalization constant of the distribution is difficult to evaluate.
  - There is not analytical solution to the integral.
- What do we do? Approximate!
- Types of approximations:

# Difficult Expected Values

- In many cases, the expected value is too complex to evaluate.
- This may occur because:
  - The normalization constant of the distribution is difficult to evaluate.
  - There is not analytical solution to the integral.
- What do we do? Approximate!
- Types of approximations:
  - 1 Approximate the distribution with a simpler one that allows us to take the expectation (e.g. Variational Bayes).

# Difficult Expected Values

- In many cases, the expected value is too complex to evaluate.
- This may occur because:
  - The normalization constant of the distribution is difficult to evaluate.
  - There is not analytical solution to the integral.
- What do we do? Approximate!
- Types of approximations:
  - 1 Approximate the distribution with a simpler one that allows us to take the expectation (e.g. Variational Bayes).
  - 2 Upper and lower-bound based approximations (e.g. Variational Approximations).

# Difficult Expected Values

- In many cases, the expected value is too complex to evaluate.
- This may occur because:
  - The normalization constant of the distribution is difficult to evaluate.
  - There is not analytical solution to the integral.
- What do we do? Approximate!
- Types of approximations:
  - 1 Approximate the distribution with a simpler one that allows us to take the expectation (e.g. Variational Bayes).
  - 2 Upper and lower-bound based approximations (e.g. Variational Approximations).
  - 3 Sampling based approximations (Monte Carlo Integration).

# Monte Carlo Integration

**Monte Carlo integration** approximates an expected value by sampling from the associated distribution and then taking an average.



# Monte Carlo Integration

**Monte Carlo integration** approximates an expected value by sampling from the associated distribution and then taking an average.

## Monte Carlo Integration

Let  $X$  be a random variable drawn from  $p(x)$ , let  $f(x)$  be a function of  $X$ , and let  $x^{(1)}, \dots, x^{(S)}$  be samples from  $p(x)$ , then

$$\mathbb{E}(f(x)) \approx \frac{1}{S} \sum_s x^{(s)}$$

# Monte Carlo Integration

How can we use Monte Carlo integration to calculate general integrals?

# Monte Carlo Integration

How can we use Monte Carlo integration to calculate general integrals?

$$\begin{aligned}\int_a^b f(x)dx &= \int_a^b \frac{b-a}{b-a} f(x)dx \\ &= (b-a) \int_a^b \frac{1}{b-a} f(x)dx \\ &= (b-a) \mathbb{E}[f(X)]\end{aligned}$$

where  $X$  is draw from  $Uniform(a, b)$ .

# Monte Carlo Integration

How can we use Monte Carlo integration to calculate general integrals?

$$\begin{aligned}\int_a^b f(x)dx &= \int_a^b \frac{b-a}{b-a} f(x)dx \\ &= (b-a) \int_a^b \frac{1}{b-a} f(x)dx \\ &= (b-a) \mathbb{E}[f(X)]\end{aligned}$$

where  $X$  is draw from  $Uniform(a, b)$ .

- Now we can approximate this expected value using sampling.

# Monte Carlo Integration

How can we use Monte Carlo integration to calculate general integrals?

$$\begin{aligned}\int_a^b f(x)dx &= \int_a^b \frac{b-a}{b-a} f(x)dx \\ &= (b-a) \int_a^b \frac{1}{b-a} f(x)dx \\ &= (b-a) \mathbb{E}[f(X)]\end{aligned}$$

where  $X$  is draw from  $Uniform(a, b)$ .

- Now we can approximate this expected value using sampling.
- This works for multidimensional integrals as well.

# Monte Carlo Integration: Area of a Circle

- How can we use Monte Carlo integration to estimate the area of a circle with radius  $r$ ?

# Monte Carlo Integration: Area of a Circle

- How can we use Monte Carlo integration to estimate the area of a circle with radius  $r$ ?
- Step 1: Formulate the problem as a bounded integral.

# Monte Carlo Integration: Area of a Circle

- How can we use Monte Carlo integration to estimate the area of a circle with radius  $r$ ?
- Step 1: Formulate the problem as a bounded integral.



# Monte Carlo Integration: Area of a Circle

- How can we use Monte Carlo integration to estimate the area of a circle with radius  $r$ ?
- Step 1: Formulate the problem as a bounded integral.

A circle is defined by  $x^2 + y^2 = r^2$  so we can define a piecewise function that tells us whether we are inside or outside of the circle.

$$f(x, y) = \begin{cases} 1 & : x^2 + y^2 \leq r^2 \\ 0 & : x^2 + y^2 > r^2 \end{cases}$$

# Monte Carlo Integration: Area of a Circle

- How can we use Monte Carlo integration to estimate the area of a circle with radius  $r$ ?
- Step 1: Formulate the problem as a bounded integral.

A circle is defined by  $x^2 + y^2 = r^2$  so we can define a piecewise function that tells us whether we are inside or outside of the circle.

$$f(x, y) = \begin{cases} 1 & : x^2 + y^2 \leq r^2 \\ 0 & : x^2 + y^2 > r^2 \end{cases}$$

The area of a circle is given by

$$A_r = \int_{-r}^r \int_{-r}^r f(x, y) dx dy$$

# Monte Carlo Integration: Area of a Circle

- Now that the area is formulated as an integral, how do we approximate it with sampling?

# Monte Carlo Integration: Area of a Circle

- Now that the area is formulated as an integral, how do we approximate it with sampling?
- Step 1: Identify the sampling distribution.

# Monte Carlo Integration: Area of a Circle

- Now that the area is formulated as an integral, how do we approximate it with sampling?
- Step 1: Identify the sampling distribution.

# Monte Carlo Integration: Area of a Circle

- Now that the area is formulated as an integral, how do we approximate it with sampling?
- Step 1: Identify the sampling distribution.
- In this case we are integrating over the  $2r \times 2r$  square.

# Monte Carlo Integration: Area of a Circle

- Now that the area is formulated as an integral, how do we approximate it with sampling?
- Step 1: Identify the sampling distribution.
- In this case we are integrating over the  $2r \times 2r$  square.
- How do we sample uniformly from a square?

# Monte Carlo Integration: Area of a Circle

- Now that the area is formulated as an integral, how do we approximate it with sampling?
- Step 1: Identify the sampling distribution.
- In this case we are integrating over the  $2r \times 2r$  square.
- How do we sample uniformly from a square?
- Sample  $x$  and  $y$  independently from  $Uniform(-r, r)$



# Monte Carlo Integration: Area of a Circle

Now we can estimate the area of a circle as:

$$A_r = \int_{-r}^r \int_{-r}^r f(x, y) dx dy$$

$$\approx 2r^2 \frac{1}{S} \sum_s f(x^{(s)}, y^{(s)})$$

where

$$f(x, y) == \begin{cases} 1 & : x^2 + y^2 \leq r^2 \\ 0 & : x^2 + y^2 > r^2 \end{cases}$$

# Monte Carlo Integration: Area of a Circle

# Outline

1 Visualizing Distributions

2 Sampling

3 Sources of Randomness

# Sources of Randomness

- Most sampling methods are based on the assumption that we can generate a uniform random number.

# Sources of Randomness

- Most sampling methods are based on the assumption that we can generate a uniform random number.
- How do we generate a uniform random number?

# Sources of Randomness

- Most sampling methods are based on the assumption that we can generate a uniform random number.
- How do we generate a uniform random number?
- There are two main methods:

# Sources of Randomness

- Most sampling methods are based on the assumption that we can generate a uniform random number.
- How do we generate a uniform random number?
- There are two main methods:
  - 1 Harvested natural randomness

# Sources of Randomness

- Most sampling methods are based on the assumption that we can generate a uniform random number.
- How do we generate a uniform random number?
- There are two main methods:
  - 1 Harvested natural randomness
  - 2 Pseudo-random number generation



# Harvesting Natural Randomness

- The idea behind harvesting natural randomness is that there are natural phenomena that are sufficiently random, due to physical entropy, that we can treat them as random.

# Harvesting Natural Randomness

- The idea behind harvesting natural randomness is that there are natural phenomena that are sufficiently random, due to physical entropy, that we can treat them as random.
- Examples of harvested sources of natural randomness include:

# Harvesting Natural Randomness

- The idea behind harvesting natural randomness is that there are natural phenomena that are sufficiently random, due to physical entropy, that we can treat them as random.
- Examples of harvested sources of natural randomness include:
  - The amount of current flowing through an electrical circuit is actually a discrete number as it is caused by individual electrons moving through a conductor. **Shot noise** is the random fluctuations in this discrete number.

# Harvesting Natural Randomness

- The idea behind harvesting natural randomness is that there are natural phenomena that are sufficiently random, due to physical entropy, that we can treat them as random.
- Examples of harvested sources of natural randomness include:
  - The amount of current flowing through an electrical circuit is actually a discrete number as it is caused by individual electrons moving through a conductor. **Shot noise** is the random fluctuations in this discrete number.
  - Source of nuclear radiation decays randomly and the amount of decay can be used as a source of randomness.

# Harvesting Natural Randomness

- The idea behind harvesting natural randomness is that there are natural phenomena that are sufficiently random, due to physical entropy, that we can treat them as random.
- Examples of harvested sources of natural randomness include:
  - The amount of current flowing through an electrical circuit is actually a discrete number as it is caused by individual electrons moving through a conductor. **Shot noise** is the random fluctuations in this discrete number.
  - Source of nuclear radiation decays randomly and the amount of decay can be used as a source of randomness.
  - When a beam of photons is directed at a semi-transparent mirror, the photons will randomly be reflected or transmitted.

# Harvesting Natural Randomness

- The idea behind harvesting natural randomness is that there are natural phenomena that are sufficiently random, due to physical entropy, that we can treat them as random.
- Examples of harvested sources of natural randomness include:
  - The amount of current flowing through an electrical circuit is actually a discrete number as it is caused by individual electrons moving through a conductor. **Shot noise** is the random fluctuations in this discrete number.
  - Source of nuclear radiation decays randomly and the amount of decay can be used as a source of randomness.
  - When a beam of photons is directed at a semi-transparent mirror, the photons will randomly be reflected or transmitted.
  - The temperature of a resistor in a circuit.

# Harvesting Natural Randomness

- The idea behind harvesting natural randomness is that there are natural phenomena that are sufficiently random, due to physical entropy, that we can treat them as random.
- Examples of harvested sources of natural randomness include:
  - The amount of current flowing through an electrical circuit is actually a discrete number as it is caused by individual electrons moving through a conductor. **Shot noise** is the random fluctuations in this discrete number.
  - Source of nuclear radiation decays randomly and the amount of decay can be used as a source of randomness.
  - When a beam of photons is directed at a semi-transparent mirror, the photons will randomly be reflected or transmitted.
  - The temperature of a resistor in a circuit.
  - Atmospheric noise detected by a radio receiver.

# Harvesting Natural Randomness

- The idea behind harvesting natural randomness is that there are natural phenomena that are sufficiently random, due to physical entropy, that we can treat them as random.
- Examples of harvested sources of natural randomness include:
  - The amount of current flowing through an electrical circuit is actually a discrete number as it is caused by individual electrons moving through a conductor. **Shot noise** is the random fluctuations in this discrete number.
  - Source of nuclear radiation decays randomly and the amount of decay can be used as a source of randomness.
  - When a beam of photons is directed at a semi-transparent mirror, the photons will randomly be reflected or transmitted.
  - The temperature of a resistor in a circuit.
  - Atmospheric noise detected by a radio receiver.
- The rate at which these can be harvested may be limited.



# Pseudo-Random Numbers

- A **Pseudo-random number generator** returns a sequence of numbers that has the statistical properties of a uniform random sequence, but is actually deterministic given an initial value.

# Pseudo-Random Numbers

- A **Pseudo-random number generator** returns a sequence of numbers that has the statistical properties of a uniform random sequence, but is actually deterministic given an initial value.
- This initial value is called a **seed**.

# Pseudo-Random Numbers

- A **Pseudo-random number generator** returns a sequence of numbers that has the statistical properties of a uniform random sequence, but is actually deterministic given an initial value.
- This initial value is called a **seed**.
- Without knowledge of the seed, the sequence should look random.

# Pseudo-Random Numbers

- A **Pseudo-random number generator** returns a sequence of numbers that has the statistical properties of a uniform random sequence, but is actually deterministic given an initial value.
- This initial value is called a **seed**.
- Without knowledge of the seed, the sequence should look random.
- Many such sequences exist.

# Pseudo-Random Numbers

- A **Pseudo-random number generator** returns a sequence of numbers that has the statistical properties of a uniform random sequence, but is actually deterministic given an initial value.
- This initial value is called a **seed**.
- Without knowledge of the seed, the sequence should look random.
- Many such sequences exist.
- Given the seed, computing pseudo-random number is extremely fast.

# Randomness in Numpy

`numpy.random` provides functions for generating pseudo-random samples from many common distributions.

# Randomness in Numpy

`numpy.random` provides functions for generating pseudo-random samples from many common distributions.

```
>>> np.random.rand(5)
array([ 0.05321291,  0.7837193 ,  0.21213337,
        0.26499685,  0.4338131 ])
```

```
>>> np.random.randn(5)
array([-1.23805748, -0.18105506,  0.96736693,
        1.25355742, -0.08038946])
```

```
>>> np.random.randint(1,100,5)
array([28, 35, 32, 19, 94])
```

# Sources of Randomness

## Demo