

Table 1 (Continued). HIV/AIDS diagnoses and deaths from January 1, 2021, to December 31, 2021; and people diagnosed with HIV, reported in NYC and presumed to be living as of December 31, 2021

| | HIV Diagnoses ¹ | | | | | | | AIDS Diagnoses ³ | | PWH as of Dec. 31, 2021 | | Deaths ⁴ | |
|---------------------------------------|----------------------------|------|--------------|------|---|------|-------|-----------------------------|------|-------------------------|------|---------------------|------|
| | Total | | Without AIDS | | Concurrent with AIDS Diagnosis ² | | | | | | | | |
| | | | | | | | | | | | | | |
| | N | % | N | % | N | % | Row % | N | % | N | % | N | % |
| Borough of Residence ⁸ | | | | | | | | | | | | | |
| Bronx | 371 | 23.3 | 301 | 23.4 | 70 | 22.8 | 18.9 | 296 | 28.1 | 31,035 | 23.9 | 623 | 28.6 |
| Brooklyn | 416 | 26.1 | 330 | 25.6 | 86 | 28.0 | 20.7 | 246 | 23.3 | 30,810 | 23.8 | 445 | 20.4 |
| Manhattan | 308 | 19.3 | 251 | 19.5 | 57 | 18.6 | 18.5 | 192 | 18.2 | 32,749 | 25.3 | 381 | 17.5 |
| Queens | 295 | 18.5 | 231 | 17.9 | 64 | 20.8 | 21.7 | 181 | 17.2 | 19,140 | 14.8 | 156 | 7.2 |
| Staten Island | 40 | 2.5 | 30 | 2.3 | 10 | 3.3 | 25.0 | 29 | 2.8 | 2,538 | 2.0 | 40 | 1.8 |
| Outside NYC | 135 | 8.5 | 115 | 8.9 | 20 | 6.5 | 14.8 | 104 | 9.9 | 13,201 | 10.2 | 79 | 3.6 |
| Unknown | 29 | 1.8 | 29 | 2.3 | 0 | 0 | 0 | 6 | 0.6 | 187 | 0.1 | 454 | 20.8 |
| Area-based Poverty Level ⁹ | | | | | | | | | | | | | |
| Low poverty | 195 | 12.2 | 144 | 11.2 | 51 | 16.6 | 26.2 | 122 | 11.6 | 18,722 | 14.4 | 144 | 6.6 |
| Medium poverty | 553 | 34.7 | 455 | 35.4 | 98 | 31.9 | 17.7 | 329 | 31.2 | 42,718 | 32.9 | 516 | 23.7 |
| High poverty | 377 | 23.7 | 291 | 22.6 | 86 | 28.0 | 22.8 | 253 | 24.0 | 27,593 | 21.3 | 424 | 19.5 |
| Very high poverty | 304 | 19.1 | 253 | 19.7 | 51 | 16.6 | 16.8 | 237 | 22.5 | 25,981 | 20.0 | 559 | 25.7 |
| Unavailable | 165 | 10.4 | 144 | 11.2 | 21 | 6.8 | 12.7 | 113 | 10.7 | 14,646 | 11.3 | 535 | 24.6 |

PWH=People with HIV.

⁸For HIV and AIDS diagnoses, residence at diagnosis. For PWH and deaths, residence based on most recent record available.⁹Area-based poverty level is determined by the proportion of residents living below the federal poverty level (FPL) in the NYC ZIP code of residence at diagnosis or most recent residence (see footnote 8). Low poverty=<10% below FPL; Medium poverty=10 to <20% below FPL; High poverty=20 to <30% below FPL; Very high poverty=≥30% below FPL.

Table 1 (Continued). HIV/AIDS diagnoses and deaths from January 1, 2021, to December 31, 2021; and people diagnosed with HIV, reported in NYC and presumed to be living as of December 31, 2021

| | HIV Diagnoses ¹ | | | | | | | AIDS Diagnoses ³ | | PWH as of Dec. 31, 2021 | | Deaths ⁴ | |
|---|----------------------------|------|--------------|------|---|------|-------|-----------------------------|------|-------------------------|------|---------------------|------|
| | Total | | Without AIDS | | Concurrent with AIDS Diagnosis ² | | | N | % | N | % | N | % |
| | N | % | N | % | N | % | Row % | | | | | | |
| Transmission Category¹⁰ | | | | | | | | | | | | | |
| MSM | 748 | 46.9 | 631 | 49.0 | 117 | 38.1 | 15.6 | 398 | 37.8 | 56,540 | 43.6 | 573 | 26.3 |
| IDU | 15 | 0.9 | 14 | 1.1 | 1 | 0.3 | 6.7 | 54 | 5.1 | 13,330 | 10.3 | 505 | 23.2 |
| MSM-IDU | 21 | 1.3 | 20 | 1.6 | 1 | 0.3 | 4.8 | 26 | 2.5 | 3,194 | 2.5 | 100 | 4.6 |
| Heterosexual contact | 236 | 14.8 | 177 | 13.8 | 59 | 19.2 | 25.0 | 219 | 20.8 | 24,993 | 19.3 | 416 | 19.1 |
| TG-SC | 57 | 3.6 | 54 | 4.2 | 3 | 1.0 | 5.3 | 31 | 2.9 | 2,021 | 1.6 | 25 | 1.1 |
| Perinatal | 2 | 0.1 | 1 | 0.1 | 1 | 0.3 | 50.0 | 12 | 1.1 | 2,537 | 2.0 | 20 | 0.9 |
| Other | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 191 | 0.1 | 4 | 0.2 |
| Unknown | 515 | 32.3 | 390 | 30.3 | 125 | 40.7 | 24.3 | 314 | 29.8 | 26,854 | 20.7 | 535 | 24.6 |

PWH=People with HIV; MSM=Men who have sex with men; IDU=Injection drug use history; MSM-IDU=Men who have sex with men and inject drugs;

TG-SC=Transgender people with sexual contact.

¹⁰“Heterosexual contact” includes people who had heterosexual sex with a person they know to have HIV, a person who has injected drugs or a person who has received blood products. For women only, it also includes history of sex work, multiple sex partners, sexually transmitted infection, crack or cocaine use, sex with a bisexual man, probable heterosexual transmission as noted in a medical chart or sex with a man and negative history of injection drug use. “Transgender people with sexual contact” includes people identified as transgender at any time by self-report, a medical provider or chart review, or ongoing data collection, who have reported sexual contact and have a negative history of injection drug use. “Other” includes people who received treatment for hemophilia, people who received a transfusion or transplant, people with other health care-associated transmission and children with nonperinatal transmission category.