

A Four-Term Counterexample to the Special Image Conjecture in Three Pairs

Roy van Rijn
Independent researcher

28 July 2026

Abstract

We give an explicit counterexample to the Special Image Conjecture in three contraction pairs. In $\mathbb{C}[\tau, w, v, t, z, y]$, let

$$f = \tau(t - y)(wz + vt), \quad g = y.$$

For the standard contraction map $\mathcal{E}$, we prove for every $m \geq 1$ that

$$\mathcal{E}(f^m) = 0, \quad [t]\mathcal{E}(gf^m) = (-1)^{m-1}(m+1)!m!.$$

Thus every positive power of f lies in the image of the operators $\partial_t - \tau, \partial_z - w, \partial_y - v$, whereas $gf^m \notin \mathcal{M}_3$ for every $m \geq 1$. The polynomial f has four terms and the multiplier is linear. Its two-pair dehomogenization is exactly Christopher D. Long's counterexample to the Mathieu conjecture for $SU(2)$; the present construction is its one-pair bihomogeneous lift. Consequently the minimum number of pairs in which the Special Image Conjecture can fail is either two or three.

1 The counterexample

For $r \geq 1$, write

$$R_r = \mathbb{C}[\zeta_1, \dots, \zeta_r, z_1, \dots, z_r], \quad \mathcal{M}_r = \sum_{i=1}^r (\partial_{z_i} - \zeta_i) R_r.$$

The Special Image Conjecture asserts that $\mathcal{M}_r$ is a Mathieu subspace: if $h^m \in \mathcal{M}_r$ for every $m \geq 1$, then $qh^m \in \mathcal{M}_r$ for every $q \in R_r$ and all sufficiently large m ; see Zhao [1] and van den Essen, Wright, and Zhao [2].

Define the polynomial-valued contraction map

$$\mathcal{E}_r : R_r \longrightarrow \mathbb{C}[z_1, \dots, z_r], \quad \mathcal{E}_r(\zeta^\alpha z^\beta) = \partial_z^\alpha z^\beta.$$

Lemma 1. *For every $r \geq 1$, one has $\mathcal{M}_r = \ker \mathcal{E}_r$.*

Proof. For $h \in R_r$,

$$\mathcal{E}_r(\partial_{z_i} h) = \partial_{z_i} \mathcal{E}_r(h) = \mathcal{E}_r(\zeta_i h).$$

Thus $\mathcal{E}_r((\partial_{z_i} - \zeta_i)h) = 0$, so $\mathcal{M}_r \subseteq \ker \mathcal{E}_r$. Conversely,

$$\zeta_i h \equiv \partial_{z_i} h \pmod{\mathcal{M}_r}.$$

Repeatedly applying this congruence gives, for every monomial,

$$\zeta^\alpha z^\beta \equiv \partial_z^\alpha z^\beta = \mathcal{E}_r(\zeta^\alpha z^\beta) \pmod{\mathcal{M}_r}.$$

By linearity, $p \equiv \mathcal{E}_r(p) \pmod{\mathcal{M}_r}$. Since $\mathcal{E}_r$ is the identity on $\mathbb{C}[z_1, \dots, z_r]$, $\mathcal{E}_r(p) = 0$ implies $p \in \mathcal{M}_r$. $\square$

Use the pairs

$$(\tau, t), \quad (w, z), \quad (v, y)$$

and put

$$\mathcal{M}_3 = (\partial_t - \tau)R_3 + (\partial_z - w)R_3 + (\partial_y - v)R_3.$$

Theorem 2. *Let*

$$f = \tau(t - y)(wz + vt), \quad g = y.$$

Then, for every $m \geq 1$,

$$\mathcal{E}_3(f^m) = 0, \quad [t]\mathcal{E}_3(gf^m) = (-1)^{m-1}(m+1)!m! \neq 0.$$

In particular, $\mathcal{E}_3(gf^m) \neq 0$. Consequently $f^m \in \mathcal{M}_3$ and $gf^m \notin \mathcal{M}_3$ for every $m \geq 1$.

In expanded form,

$$f = \tau vt^2 - \tau vty + \tau wtz - \tau wyz.$$

This displays the advertised four terms directly.

2 Proof

First omit the pair (τ, t) . On $\mathbb{C}[W, Z, V, Y]$, define

$$\mathcal{F}(W^a Z^b V^c Y^d) = \delta_{a,b} \delta_{c,d} a! c!$$

and set

$$P = (1 - Y)(WZ + V), \quad Q = Y.$$

Expanding the second factor gives

$$P^m = \sum_{k=0}^m \binom{m}{k} W^{m-k} Z^{m-k} V^k (1 - Y)^m.$$

For a nonzero contraction, the V, Y pair selects $[Y^k](1 - Y)^m = (-1)^k \binom{m}{k}$. Hence

$$\mathcal{F}(P^m) = \sum_{k=0}^m (-1)^k \binom{m}{k}^2 (m-k)! k! = m! \sum_{k=0}^m (-1)^k \binom{m}{k} = 0. \quad (1)$$

After multiplication by $Q = Y$, the V, Y pair instead selects $[Y^k]Y(1 - Y)^m = (-1)^{k-1} \binom{m}{k-1}$ for $1 \leq k \leq m$. Therefore

$$\begin{aligned} \mathcal{F}(YP^m) &= m! \sum_{k=1}^m (-1)^{k-1} \binom{m}{k-1} \\ &= (-1)^{m-1} m!. \end{aligned} \quad (2)$$

Homogenize in the coordinate variables Z, Y , treating W, V as coefficient, or dual, variables:

$$\tilde{P}(t, w, z, v, y) = t^2 P\left(w, \frac{z}{t}, v, \frac{y}{t}\right) = (t - y)(wz + vt).$$

Each factor of P has total W, V -degree one. Every monomial of P^m that survives $\mathcal{F}$ therefore has total (Z, Y) -degree m , matching its total W, V -degree. Its coordinate homogenization contains $t^{2m-m} = t^m$, exactly matching the power τ^m . The new contraction contributes $m!$, and (1) gives

$$\mathcal{E}_3(f^m) = m! \mathcal{F}(P^m) = 0.$$

For YP^m , a surviving term of P^m has total (Z, Y) -degree $m - 1$. Its homogenization contains $t^{2m-(m-1)} = t^{m+1}$. Contracting against τ^m leaves one power of t and contributes $(m + 1)!$. By (2),

$$[t]\mathcal{E}_3(yf^m) = (m + 1)! \mathcal{F}(YP^m) = (-1)^{m-1}(m + 1)! m! \neq 0.$$

Lemma 1 completes the proof of Theorem 2.

3 Relation with the xz - and $SU(2)$ counterexamples

Write the coordinate functions on $SU(2)$ as

$$\begin{pmatrix} a & c \\ b & d \end{pmatrix}.$$

Under the substitution

$$W = a, \quad Z = d, \quad V = b, \quad Y = -c,$$

the two-pair seed and multiplier become

$$P = (1 - Y)(WZ + V) \mapsto (1 + c)(ad + b), \quad Q = Y \mapsto -c.$$

This is precisely Long's $SU(2)$ counterexample [3]. Indeed normalized Haar integration satisfies

$$\int_{SU(2)} a^r b^s c^t d^u dg = (-1)^s \delta_{r,u} \delta_{s,t} \frac{r! s!}{(r + s + 1)!}.$$

Because every monomial of P^m has total W, V -degree m , this formula gives

$$\mathcal{F}(P^m) = (m + 1)! \int_{SU(2)} ((1 + c)(ad + b))^m dg$$

and the analogous identity with multipliers Y and $-c$. Thus equations (1)–(2) recover Long's moments

$$\int_{SU(2)} ((1 + c)(ad + b))^m dg = 0, \quad \int_{SU(2)} (-c)((1 + c)(ad + b))^m dg = \frac{(-1)^{m-1}}{m + 1}.$$

Under the M\"uger–Tuset integration substitution used by Long,

$$(a, b, c, d) = ((1 - x)z_2, xz_1, -z_1^{-1}, z_2^{-1}),$$

the same seed becomes

$$(1 - z_1^{-1})((1 - x) + xz_1),$$

which is Long's three-term counterexample to $xz(1, 1)$. Accordingly, the seed and its xz - and $SU(2)$ -interpretations are due to Long. The contribution here is the degree-two one-pair bihomogenization that produces the four-term counterexample to the Special Image Conjecture in three pairs.

Corollary 3. *If r_{SIC} denotes the minimum number of contraction pairs in which the Special Image Conjecture fails, then*

$$2 \leq r_{\text{SIC}} \leq 3.$$

Proof. The upper bound is Theorem 2. The one-pair case follows from [2, Theorem 2.8]: take the $\mathbb{Q}$ -algebra $A = \mathbb{C}[\tau]$ and $a = \tau$. Then a is a non-zero-divisor and the principal ideal $Aa = (\tau)$ is radical, precisely the hypotheses of that theorem. $\square$

4 Size, scope, and verification

The coefficients of f lie in $\{1, -1\}$. With respect to (τ, w, v) -degree and (t, z, y) -degree, f has bidegree $(2, 2)$ and ordinary degree four; $g = y$ has bidegree $(0, 1)$. The polynomial f is not of the separated form $\lambda(\zeta)P(z)$; consequently the example does not give a three-variable Generalized Vanishing Conjecture counterexample.

Thus the known one-pair result and Theorem 2 give

$$2 \leq r_{\text{SIC}} \leq 3.$$

We make no claim here concerning the two-pair case or absolute literature priority.

The all-order proof above is independent of computation. An exact sparse-arithmetic checker expands f , verifies the contractions through $m = 10$, and checks the two binomial identities through $m = 99$. The checker is included in the accompanying Zenodo record and in the companion repository.¹ As a low-order checksum,

$$\mathcal{E}_3(f) = 0, \quad \mathcal{E}_3(yf) = 2t - y.$$

Acknowledgment. Computational and editorial assistance from OpenAI Codex was used in searching for, checking, and presenting the example. The author takes responsibility for the mathematical statements and proof.

References

- [1] W. Zhao, *Images of commuting differential operators of order one with constant leading coefficients*, J. Algebra **324** (2010), 231–247. doi:10.1016/j.jalgebra.2010.04.022.
- [2] A. van den Essen, D. Wright, and W. Zhao, *On the Image Conjecture*, J. Algebra **340** (2011), 211–224. doi:10.1016/j.jalgebra.2011.04.036.
- [3] C. D. Long, *Counterexamples to the xz -Conjecture and the Mathieu Conjecture for $SU(2)$* , arXiv:2607.19012v1 (2026). arXiv:2607.19012.

¹Repository path: `scripts/verify_three_pair_image_mathieu_counterexample.py`.