


**Waveform capnography - Neonatal**


*Percent of Neonatal patient transport contacts with advanced airways in whom continuous waveform capnography was used.*



group	num	den	prop
GAMUT	1174	2742	0.4281546
[redacted]	5	115	0.0434783
[redacted] Neo	0	91	0.0000000
[redacted] Neo	0	16	0.0000000

**Waveform capnography - Pediatric**


*Percent of Pediatric patient transport contacts with advanced airways in whom continuous waveform capnography was used*



group	num	den	prop
CA TX Adult Peds	83	83	1.0000000
KY Adult Peds	14	14	1.0000000
MD VA Adult Peds	47	47	1.0000000
MI Adult Peds	21	21	1.0000000
NM Adult Peds	12	12	1.0000000
PHI AZ Adult Peds	67	68	0.9852941
IN Adult Peds	41	42	0.9761905
	285	312	0.9134615
GAMUT	3892	4439	0.8767740
Neo	0	23	0.0000000

**Waveform capnography - Adult**

*Percent of Adult patient transport contacts with advanced airways  
in whom continuous waveform capnography was used*



group	num	den	prop
MI Adult Peds	71	71	1.0000000
NM Adult Peds	117	117	1.0000000
KY Adult Peds	259	260	0.9961538
MD VA Adult Peds	710	713	0.9957924
CA TX Adult Peds	899	903	0.9955703
IN Adult Peds	2856	2875	0.9933913
AZ Adult Peds	299	302	0.9900662
GAMUT	501	509	0.9842829
GAMUT	20656	21158	0.9762737