

Chapter 1

Introduction

1.1 Overview

Microseconds after the Big Bang, the Universe existed in a state known as the Quark Gluon Plasma (QGP) [1, 2]. In the QGP, quarks and gluons are not in hadronic bondage, forced to the confines of bound states such as protons and neutrons. The Large Hadron Collider (LHC) [3] produces QGP in the lab in lead-lead (PbPb) collisions. The high energies and rates of the collisions at the LHC make it possible to do detailed studies of the QGP. The LHC is producing rare experimental probes such as suppressed jets [4] and heavy quarkonia [5] at unprecedented rates in heavy-ion collisions. As a result of recent LHC studies described in this thesis, physicists now have better constraints on the QGP properties such as temperature, viscosity, and energy density.

Detailed studies of the state of the nuclei before they collide are needed to further constrain the properties of the QGP. Without more knowledge of the initial state, physicists cannot determine which experimental effects are due to the QGP and which effects are inherent to the nuclei themselves. For example, suppression of heavy quarkonia is a signature of the QGP but also appears to occur in deuterium-gold (dAu) collisions where the QGP is not expected to arise [6]. Another important example is the measurement of viscosity, which depends on the relationship between the observed azimuthal anisotropy and the initial eccentricity of the overlap of the two colliding nuclei

(see Section 1.4). A clean probe of the initial state is needed by physicists to comprehensively understand the QGP. Ultra-peripheral heavy-ion collisions (UPC) at the LHC provide such a probe. A recent review on UPC physics can be found in [7, 8].

The current understanding of heavy-ion collisions evolved over the last 30 years. Relativistic heavy-ion collisions were first studied at Belavac at the Lawrence Berkeley National Laboratory (LBNL) in 1974, then at the Alternating Gradient Synchrotron (AGS) at Brookhaven National Laboratory (BNL) in Upton, NY, followed by the Super Proton Synchrotron (SPS) at CERN near Geneva, Switzerland. From the numerous AGS and SPS experiments two main observables emerged, namely, J/ψ suppression and strangeness enhancement [9]. These results pioneered the search for the QGP.

At AGS the ion isotopes ^{16}O , ^{28}Si , and ^{197}Au beams were collided with fix targets. At SPS the same fix target configuration was used, but ~~the ion isotopes were with~~ ^{16}O , ^{32}S , and ^{208}Pb beams. The center of mass energies per nucleon pair, $\sqrt{s_{NN}}$, for these experiments ranged from just below 5 GeV to 20 GeV. Although the strangeness enhancement and J/ψ suppression signals indicated that a deconfined state of quarks and gluons was likely created, at the energies of the AGS and SPS this state perished quickly. The threshold for creating the QGP requires an energy density of $\sim 0.15 \text{ GeV}/\text{fm}^3$ fm³ and a temperature near 170 MeV [18, 19]. Because of this, the QGP signals at the AGS and SPS energies could not have lasted long enough to study their properties.

Plans for a colliding beam machine dedicated to heavy-ions, designed to reach energies of 200 GeV per nucleon pair, was first proposed in 1983. It was believed that in these collisions signs of a hot gas of quarks and gluons would emerge. In 2000, RHIC began collisions and the four experiments, STAR, PHENIX, BRAHMS, and PHOBOS started taking data. The energies at RHIC were a factor of 10 higher than was previously achieved ~~reaching when running at~~ the designed maximum energy. RHIC experiments confirmed the existence of a thermalized state of quarks and gluons. Contrary to expectations, the state at RHIC was found to be a strongly coupled fluid with nearly no viscosity [10].

The LHC heavy-ion program began collisions in 2010, colliding PbPb at a center of mass

energy of 2.76 TeV per nucleon pair. This corresponds to an increase in the colliding energy by an order of magnitude with respect to RHIC. The LHC experiments, ALICE [11], ATLAS [12], and CMS [13] have been studying the heavy-ion collisions since then. In 2013 LHCb [14] joined the LHC heavy-ion program. Thanks to the LHC and RHIC physics programs, a new era of precision heavy-ion measurements is underway. The latest results from the LHC have come from the 2013 proton-lead (pPb) run. This period of data taking was originally designed to be a control measurement. However, CMS and ALICE have both shown an elliptical flow-like signal present in the pPb data [15, 16], which was unexpected. Understanding the origin of this nuclear effect is one of the most important priorities of the heavy-ion physics program at present. The latest data from the pPb measurements [17] confirm the need to understand the nature of the initial state. UPC events fulfill this need by probing the nucleus through photon interactions. By measuring UPC J/ψ events, theoretical models of the initial state can be constrained. In addition, these constraints will provide the important benchmarks needed to design a future electron-ion collider dedicated to exploring the nuclear initial state. In this thesis, the CMS capability for measuring this process, the description of the analysis, and the comparison between the measured coherent J/ψ cross section to theoretical models are given. The correlations between the rapidity of J/ψ 's and neutrons in photo-nuclear interactions are also explored.

In this chapter, the measurement of transverse energy, direct photons, and elliptic flow will be discussed. The transverse energy and direct photons measurements are related to the energy density and temperature of the QGP. The elliptic flow measurement is related to the viscosity of the QGP. Before discussing these results, several terms used to describe heavy-ion measurements and detectors will be presented. This discussion will be divided into two parts, a generic description of the main detector elements, and an explanation of variables used in heavy-ion measurements.

1.2 Variables in heavy-ion measurements

Two types of generic detectors are in use in heavy-ion experiments, trackers and calorimeters. A tracker measures the trajectory of the particles as they pass through the detector. The particle trajectories measured by a tracker detector are called tracks. From these tracks, the particles' momenta can be deduced. The radius of curvature of the track is related to the momentum due to the curved path of the particle in a magnetic field. In CMS tracks are mapped out from measuring charge deposits in silicon cells. A tracker will typically be designed to interact minimally with the particles in order to preserve the trajectories of the particles. The second generic type of detector, a calorimeter, records the energy of the particles that hit it. A calorimeter is typically filled with dense material used to initiate a shower of particle collisions. The light from this shower of particles is collected and used to measure the energy of the initial particle or particles which hit the calorimeter. Unlike a tracker, a calorimeter typically is designed to stop the particles which hit it and collect as much of the energy of these particles as possible. The details the CMS detector will be discussed in more detail in Chapter 3.

Heavy-ion measurements will typically be comprised of combinations of the variables, p , for momentum, E , for energy, ϕ , for the azimuthal angle, p_T , the transverse momentum, y , for rapidity, and η , for pseudorapidity. Energy ~~are~~is typically measured by a calorimeter and momentum by a tracker. The ϕ angle is measured around the beam axis. The transverse momentum, p_T , is the component of the momentum that points perpendicular to the beam line. Rapidity is related to energy and momentum by $y = \frac{1}{2} \ln \left(\frac{E+p_z}{E-p_z} \right)$, where p_z is the momentum component that points in the direction of the beam axis. Here c is taken to be equal to 1. Pseudorapidity is defined by $\eta = \frac{1}{2} \ln \left(\frac{|\mathbf{p}|+p_z}{|\mathbf{p}|-p_z} \right)$, which can be shown to be equal to $-\ln(\tan(\theta/2))$. The polar angle for the beam with respect to the beam line defines θ . If $m \ll p$, where m is the mass of the particle, the relativistic equation for energy, $E^2 = p^2 + m^2 \rightarrow E = p$, and pseudorapidity is equal to the rapidity. Tracker measurements will typically provide a rapidity value with the mass assigned based on a particle identification criteria, where as calorimeter based measurements will typically be assigned a rapidity based on the angle θ with respect to the interaction point.

In addition, the variable called centrality is used in heavy-ion collisions to characterize the degree of activity. It is typically either measured using activity in either a calorimeter in the forward region at high η values or by the number of tracks. Centrality is reported as a percentage where 0% corresponds to the most energetic collisions and 100% corresponds to the least energetic. For example, events assigned a centrality between 0-10% would be the top ten percent of events in terms of number of hits in a tracker detector or energy in a forward calorimeter depending on which variable is used for centrality. The number of nucleons that participate in a collision, N_{part} , the number of collisions between these participants, N_{col} , and the closest distance between the center of the two colliding nuclei, the impact parameter, b , are all related to the measured centrality percentage by use of a simulation. The convention of labeling the most active centralities 0% and the least active 100% is due to the correlation between the impact parameter. When the nuclei collide exactly head on the impact parameter is 0, which correlates with a 0% centrality value.

1.3 Temperature and energy density of the QGP

Lattice QCD predicts that the QGP forms above a critical temperature and energy density, 170 MeV and 1.5 GeV/fm³ [18, 19]. The measurement of $\frac{dE_T}{d\eta}$ provides a means of estimating the energy density of the hot state created in heavy-ion collisions. The value of E_T for this measurement is defined as $E_T = \sum_i E_i \sin\theta_i$, where E_i is the energy measured by the i th calorimeter element and θ is the angle between the center of the detector element and the interaction point. The temperature can be estimated from the transverse momentum spectrum of the direct photons, which are photons that are produced directly from the QGP. The measurements from CMS and ALICE [11] at the LHC and STAR and PHENIX at RHIC confirm that the critical values for energy density and temperature are exceeded.

The values of $\frac{dE_T}{d\eta}$ measured by CMS [20] and PHENIX [21] were done using the experiments' calorimeter systems. For the calculated $\frac{dE_T}{d\eta}$ at $\eta = 0$, the interval $|\eta| > 0.35$ was used by both experiments and corresponds to the full coverage of the PHENIX calorimeters. From the $\frac{dE_T}{d\eta}$

at $\eta = 0$ measurements, the energy density of the created medium at the LHC and RHIC were estimated from the Bjorken energy density formula, $\varepsilon_{Bj} = \frac{1}{A\tau} \frac{dE_T}{dy}$, where A is the region of overlap between the two colliding nuclei and τ is the formation time of the medium [22]. The formation time times the energy density, $\tau \cdot \varepsilon_{Bj}$, from PHENIX was measured at RHIC to be $5.4 \pm 0.6 \text{ GeV fm}^{-2} \text{ fm}^{-2} \text{ c}^{-1}$ for the centrality interval 0-5%. At the LHC CMS measured $\tau \cdot \varepsilon_{Bj} = 14 \text{ GeV fm}^{-2} \text{ fm}^{-2} \text{ c}^{-1}$ for the same centrality interval. Both values are well above the critical energy density of about 1.5 GeV/fm^3 calculated from lattice QCD when assuming a medium formation time $\tau = 1 \text{ fm/c}$.




Figure 1.1: Comparison of $\frac{dE_T}{d\eta}$ as a function of collision energy, $\sqrt{s_{NN}}$, normalized by the number participating nucleons, N_p , to account for the difference in the ion species collided at the various different experiments.

Direct photons defined as thermal photons created in the QGP, have been measured at RHIC by PHENIX [23] and at the LHC by ALICE [24] through the measurements of electron-positron pairs. Each experiment measured the inclusive p_T spectrum from electron-positron pairs, all pairs from the sample are taken without regard to the creation mechanism. The PHENIX measurement was

taken from pp collisions and the top 20% most energetic AuAu collisions. The ALICE measurement analyzed the 0-40% centrality and 40%-80% centrality PbPb collisions, and pp collisions. In the PHENIX and ALICE measurements, the contribution to the inclusive spectrum was sorted into a direct component and a background component. The latter consists mainly of decays from hadrons such as π and η particles. In the PHENIX measurement this was done by fitting to the mass distribution of the electron-positron pair for each p_T bin. For the ALICE measurement, the double ratio between the inclusive photons to ~~pions~~ photons from pion decays over the ratio between photons coming from all decays ~~and pions~~ to photons from pion decays was measured to estimate the direct contribution. This double ratio is equal to the fraction of all photons which come from decays. All photons that do not come from decays are direct photons; ~~therefore the fraction of inclusive photons that are direct can be found from the ratio of decay photons to inclusive photons.~~ The direct photon fraction is estimated by subtracting the decay photon fraction from 1. The inclusive photon p_T spectra were then rescaled by the direct photon fractions. The slope of an exponential fit to the low p_T portion of the direct photon spectrum is used to measure the temperature.




Figure 1.2: Direct photon yield as a function of p_T from PHENIX [23] (left) and ALICE [24] (right)

The temperature obtained from direct photon measurements offers a clear way of establishing whether the critical temperature for deconfinement was achieved. The direct photon p_T spectra in Fig. 1.2 shows a clear enhancement at low p_T compared to the rescaling to the pp spectra that fits the high p_T part, indicating a clear deviation from pp collisions. The low p_T photons are then consistent with thermal activity of the QGP. The thermal spectrum of the QGP is therefore imprinted on this part of the spectrum. The temperature from PHENIX was measured to be 221 ± 21 MeV and 304 ± 51 MeV from ALICE, both well above the critical temperature estimated from lattice QCD of ~ 170 MeV. The combination of the energy density and temperature measurements create a consistent picture, that both RHIC and the LHC have created a deconfined state.

1.4 Elliptic flow and viscosity in the QGP

Prior to RHIC, the QGP was thought to be a hot gas of quarks and gluons. At RHIC, elliptic flow, v_2 , was measured [25] showing that the medium appears to obey hydrodynamic equations and flows like a fluid. This same signal was also measured by CMS [26] at the LHC.

Elliptic flow [27, 28] is the second order Fourier expansion coefficient of the azimuthal distribution of measured tracks. This expansion has the form

$$1 + \sum_{n=1}^{\infty} 2v_n \cos[n(\phi - \Psi)], \quad (1.1)$$

where v_n is the n th coefficient of the Fourier expansion, ϕ is the azimuthal angle, and Ψ is the event-plane angle. The event-plane angle is the azimuthal angle of the maximal flow of the final state particles. The event-plane angle accounts for the random orientation of the nuclei event-by-event and is correlated with the reaction-plane, Ψ_R (see Fig. 1.3). The reaction-plane is defined by the line which runs through the center of the two nuclei when they reach the distance of closest approach.

The v_2 signal is believed to arise from pressure gradients. These pressure gradients originate from the initial elliptical shape of the overlap region between the colliding nuclei. The pressure




Figure 1.3: Schematic of the initial nuclear overlap and elliptic flow.

gradient is higher along the shorter axis compared to the longer axis of the ellipsoid in Fig 1.3. This difference in pressure gradient will create a flow of the QGP medium in the direction of the shorter axis. The flow is responsible for the measured anisotropic distribution of tracks.

The extent to which the initial shape of the overlap translates to flow in the medium is controlled by viscosity. A viscous fluid will tend to smooth out anisotropies and results in a smaller flow signal for the same overlap configuration of the initial nuclei. The ratio between the measured v_2 signal and the initial eccentricity, ϵ , based on the given centrality is sensitive to the viscosity of the medium.

In Fig. 1.4, v_2/ϵ is plotted as a function of the number of charged particle, N_{ch} , per unit of pseudorapidity, η , per area of overlap between the colliding nuclei, S [29]. The model calculation was performed for the four values of the ratio of the shear viscosity over entropy density, η/s , of 0.0, 0.08, 0.16, and 0.24. The η/s value which gives the best agreement with data in Fig 1.4 depends strongly on the initial state model calculation. The KLN model in the upper right portion of Fig. 1.4 is based on the Color Glass Condensate approach, which contains gluon saturation effects. The upper left portion of Fig. 1.4 is a Glauber model approach, where the nucleus is treated as a super-position of nucleons. The observed strong dependence is one of the primary




Figure 1.4: v_2/ϵ as a function of $1/S \frac{dN_{ch}}{dy}$ for different different values of η/s for KLN initial state (upper left) and Glauber initial state (upper right) compared to STAR (overlaid) and CMS and PHOBOS data (bottom).

187 reason why more studies on the nature of the initial state are needed.

1.5 Recent results from heavy-ion control measurements

The proton-nucleus runs at the LHC and RHIC were pursued as control measurements. It was believed that these configurations could measure initial state backgrounds to heavy-ion measurements. However, a flow-like signal has been very recently measured in pPb collisions by LHC experiments. A v_2 signal, like that indicating flow of the QGP in PbPb collisions, has been measured by CMS [15].




Figure 1.5: v_2 measured using the cumulant method in pPb collisions [15].

The v_2 measurement shown in Fig 1.5 uses a cumulant method [30]. This method differs from the event plan method discussed above, in that it uses the correlations between tracks rather than the ϕ angle of the tracks with respect to the event plane angle. This method can be done using the correlations between any number of tracks. The number of tracks that are used for a given method is indicated by the number in brackets. In addition a method called the Lee-Yang Zeros (LYZ) method [31, 32] was used, which uses the correlations between all the tracks. The measurement is done over a range of total number of tracks, $N_{trk}^{offline}$. The range shown in Fig. 1.5 corresponds to the number of tracks present in PbPb collisions for centrality values between 50-100%. The v_2 measurements using more than 2 track correlations are all consistent with each other. This is an indication that v_2 in pPb collisions at the LHC is the result of collective motion. The pPb v_2 measurement underscores the need for a clean probes of the initial [state](#).

In this thesis the analysis of ultra-peripheral J/ψ photoproduction is discussed. In UPC events,

206 the interaction between the nuclei occurs through exchange of photons. This results in a clean
207 probe of the initial state of the nucleus. In the following chapter, the theory of J/ψ photoproduction
208 in UPC events is presented.

Chapter 2

Photo-nuclear interactions

In ultra-peripheral heavy-ion collision, the colliding nuclei interact only electromagnetically [7, 8]. In such events, no QGP state emerges, and the effects arising from the QGP no longer obscure the initial state effects. Other initial state probes such as peripheral nuclear collisions and proton-nucleus collisions have the potential to create the QGP obscuring effects that come from the initial state. Hadronic interactions do ~~not~~ occur in ultra-peripheral collisions (UPC), therefore final state effects coming from the QGP are not expected. Thus, UPC events provide clarity by enhancing ~~physicists'~~ the understanding of the initial state.

In particular, these interactions between the field of photons surrounding the colliding nuclei and the gluons within the nuclei can produce a J/ψ , probing the gluon density [33, 34]. The J/ψ can be produced either coherently or incoherently. In the case of coherent interactions, the photon couples to the nucleus as a whole. In the incoherent case, the photon couples to the nucleons within the nucleus. The UPC J/ψ photoproduction cross section is therefore a probe of the initial state of the nucleus.

This cross section can be calculated using three steps. First, the Weizsäcker-Williams approximation [35, 36] provides a way to calculate the density of probing photons that surrounds the nucleus. Second, the electron-proton scattering data gives a value for the proton photoproduction cross section at lower energies [37]. Last, a specific model is used to combined the previous two

steps in order to calculate the nuclear photoproduction cross section. In this thesis the Adeluyi and Bertulani (AB) [38, 39], STARlight [40, 41], and the Leading Twist (LTA) [42, 43] models are discussed. Each of these methods handle the gluon density of the nucleus differently producing a measurable difference in the value of the J/ψ photoproduction cross section. There are other UPC models available [44, 45, 46, 47], but this thesis focuses on the STARlight, the AB, and the LTA models. The analysis presented in Chapter 5 adds to the existing data in a new kinematic region. In this Chapter, the theoretical framework for photon-nuclear interactions, together with an experimental review of recent results on ultra-peripheral heavy-ion collisions are described.

2.1 Weizsäcker-Williams approximation

The density of photons surrounding the colliding nuclei can be calculated using the Weizsäcker-Williams approximation [35, 36]. This approximation relates the electric field of a stationary point charge to the photon field that arises at ultra relativistic velocities. The approximation is semi-classical. In the Weizsäcker-Williams approximation, a Fourier transform of Maxwell's equations is combined with the quantum mechanical equation for the energy of the photon. The frequency




Figure 2.1: The electromagnetic field boosted (right) and at rest (left) [48].

modes of the electrostatic field are treated as photons. The Weizsäcker-Williams approximation makes the calculation of electromagnetic interactions with the nucleus tractable.

The Weizsäcker-Williams approximation begins with the equation for the electric field of the projectile nucleus at rest. To calculate the photon flux on the target nucleus, the electromagnetic field only needs to be considered at the position of the target nucleus. From the projectile's point of view, the target is moving and contributes $-vt$ to Eq. 2.1, the equation for the electric field of the projectile nucleus at rest.

$$x' = -vt', \quad y' = b, \quad z' = 0, \quad \vec{\mathbf{E}}' = \left(\frac{eZ}{4\pi\epsilon_0 \left((-vt')^2 + b^2 \right)^{3/2}} \right) \left(-vt' \hat{\mathbf{x}}' + b \hat{\mathbf{y}}' \right), \quad (2.1)$$

where b is the impact parameter defined as the distance of separation at closest approach, v is the velocity of the projectile nucleus, Z is the number of protons in the nucleus, and e is the charge of the electron. Two simplifications occur due to the choice of coordinates in Eq. 2.1. The magnetic field is equal to zero, as the projectile is at rest, and the z coordinate can be ignored, reducing the equation to two dimensions.

The Lorentz transformation converts the field equations in the projectile's frame to equations in the target's frame. Eq. 2.2 gives the result of the transformation from the projectile's primed frame to the target's rest frame for the field components [48]:

$$\begin{aligned} E'_x &= E_x, & \gamma(E'_y/c + \beta B'_z) &= E_y/c, & \gamma(E'_z/c + \beta B'_y) &= E_z/c, \\ B'_x &= B_x, & \gamma(B'_y - \beta E'_z/c) &= B_y, \text{ and } & \gamma(B'_z + \beta E'_y/c) &= B_z. \end{aligned} \quad (2.2)$$

The transformation equations for the fields, Eq. 2.2, and the transformation of the coordinates reduce to Eq. 2.3 [48]:

$$\begin{aligned} E'_x &= E_x, & \gamma E'_y &= E_y, & \gamma \beta E'_y/c &= B_z, \\ ct' &= \gamma ct, \text{ and } & x' &= -\gamma \beta ct. \end{aligned} \quad (2.3)$$

The Lorentz transformation reduces the six components of the electromagnetic field in the target's

frame to the three equations in Eq. 2.2 by relating them to the fields in the projectile's frame.

By combining Eq. 2.1 and Eq. 2.2, equations for the electric and magnetic fields in the target's rest frame are obtained

$$\begin{aligned}\vec{\mathbf{E}} &= \left(\frac{\gamma e Z}{4\pi\epsilon_0 \left((\gamma vt)^2 + b^2 \right)^{3/2}} \right) (vt\hat{\mathbf{x}} + b\hat{\mathbf{y}}), \text{ and} \\ \vec{\mathbf{B}} &= \frac{\gamma\beta e Z b}{4\pi c\epsilon_0 \left((\gamma vt)^2 + b^2 \right)^{3/2}} \hat{\mathbf{z}} = \frac{\gamma\mu_0 v e Z b}{4\pi \left((\gamma vt)^2 + b^2 \right)^{3/2}} \hat{\mathbf{z}}.\end{aligned}\tag{2.4}$$

If the impact parameter b goes to zero, the target sits in the line of the projectile particle's motion, and the denominator carries a factor of γ squared. When vt goes to zero, the projectile particle position lays on the y -axis, and the numerator carries a factor of γ . This results in fields that are a factor of γ^3 higher in the y direction than in the x direction (see Fig. 2.1). The boost compresses the electric field of the charge in the direction of the boost and produces a magnetic field resulting in a form similar to radiation. The point charge at ultra relativistic velocities produces a strong electric field in the plane transverse to its motion resembling a plane wave.

Separating the electromagnetic field into even and odd functions of time simplifies the decomposition of the field equations into Fourier frequency modes. The even functions decompose into cosine functions, odd functions into sine functions. The y -component of the electric field and the z -component of the magnetic field are even functions in time, and the x -component of the electric field is an odd function in time. Eq. 2.5 gives the Fourier transformation integrals.

$$\begin{aligned}E_x(\omega) &= \sqrt{\frac{2}{\pi}} \frac{eZ}{4\pi\epsilon_0 b^2} \int_0^\infty \frac{(\gamma vt/b) \sin(\omega t)}{\left((\gamma vt/b)^2 + 1 \right)^{3/2}} dt, \\ E_y(\omega) &= \sqrt{\frac{2}{\pi}} \frac{\gamma e Z}{4\pi\epsilon_0 b^2} \int_0^\infty \frac{\cos(\omega t)}{\left((\gamma vt/b)^2 + 1 \right)^{3/2}} dt, \text{ and} \\ B_z(\omega) &= \frac{\beta E_y(\omega)}{c}.\end{aligned}\tag{2.5}$$

275 The solutions to the integrals of Eq. 2.5 are the following [49]:

$$u = \frac{\gamma vt}{b}, \quad du \left(\frac{b}{\gamma v} \right) = dt, \quad \omega' = \frac{\omega b}{\gamma v},$$

$$\int_0^\infty \frac{u \sin(\omega' u)}{(u^2 + 1)^{3/2}} du = \omega' K_0(\omega'), \text{ and } \int_0^\infty \frac{\cos(\omega' u)}{(u^2 + 1)^{3/2}} du = \omega' K_1(\omega'). \quad (2.6)$$

276 In Eq. 2.6, ω can be related to the energy of a photon by $E = \hbar\omega$. The components of the electric
277 field in terms of ω are:

$$E_x(\omega) = \sqrt{\frac{2}{\pi}} \frac{eZ}{4\pi\epsilon_0 b^2} \frac{b}{\gamma v} \frac{\omega b}{\gamma v} K_0 \left(\frac{\omega b}{\gamma v} \right), \text{ and } E_y(\omega) = \sqrt{\frac{2}{\pi}} \frac{\gamma eZ}{4\pi\epsilon_0 b^2} \frac{b}{\gamma v} \frac{\omega b}{\gamma v} K_1 \left(\frac{\omega b}{\gamma v} \right). \quad (2.7)$$

278 The y-component of the electric field does not have a factor of t in the numerator in Eq. 2.5,
279 therefore the factor of γ remains outside of the integral for the Bessel functions in Eq. 2.7. In
280 Eq. 2.7, E_y carries an additional factor of γ in the numerator relative to the E_x , therefore in the case
281 when $\gamma \gg 1$, E_x can be neglected.

282 When v approaches c , $\beta \approx 1$, the y-component of the electric field and the z-component of the
283 magnetic field are related by a factor of c , $E_y/c = B_z$. E_y is approximately equally to γE_x because
284 $K_0(x)$ is smaller than $K_1(x)$ for all x . The conditions imposed by the ultra-relativistic limit result in
285 the following relationship

$$\gamma \gg 1 \Rightarrow \gamma E_x \gg E_x \Rightarrow E_y \gg E_x. \quad (2.8)$$

286 The six field components are reduced to one electric component and one perpendicular magnetic
287 field component, which have a configuration identical to a plane wave.

288 As with plane waves, the energy per area per time transferred by the electromagnetic field
289 is given by the Poynting vector. The Poynting vector takes the simple form of a plane pulse
290 propagating in the x direction as given by

$$\vec{S} \equiv \vec{E} \times \vec{B}/\mu_0 = (E_y^2/c\mu_0) \hat{x} = c\epsilon_0 E_y^2 \hat{x}. \quad (2.9)$$

291 The Poynting vector relates to the fluence (energy per unit area) by the expression [50]

$$I(b) = \hat{\mathbf{x}} \cdot \int_0^\infty \vec{\mathbf{S}} d\omega = \int_0^\infty (c\epsilon_0 E_y^2) d\omega = \int_0^\infty \left(\frac{dI}{d\omega} \right) d\omega, \quad (2.10)$$

292 and the spectral fluence (energy per area per frequency) is given by

$$\frac{dI}{d\omega} = c\epsilon_0 E_y^2 = \frac{e^2 Z^2 c}{4\pi^3 b^2 v^2 \epsilon_0} \left(\frac{\omega b}{\gamma v} \right)^2 K_1^2 \left(\frac{\omega b}{\gamma v} \right) = \alpha \hbar \left(\frac{Z}{b\beta\pi} \right)^2 \left(\frac{\omega b}{\gamma v} \right)^2 K_1^2 \left(\frac{\omega b}{\gamma v} \right). \quad (2.11)$$

293 The spectral fluence given by Eq. 2.11 relates the frequency to energy. The quantum mechanical
 294 equation, $E = \hbar\omega$, gives the energy of a photon, which is related to the spectral fluence. The
 295 relationship between the photon number density and the spectral fluence is [48]

$$\frac{dI}{d\omega} d\omega = \hbar\omega N(\omega) d(\hbar\omega) \Rightarrow \frac{1}{\hbar^2 \omega} \frac{dI}{d\omega} = N(\omega). \quad (2.12)$$

296 Substituting Eq. 2.11 into Eq. 2.12 yields the semiclassical photon flux:

$$N(\omega, b) = \frac{\alpha}{\hbar\omega} \left(\frac{Z}{b\beta\pi} \right)^2 \left(\frac{\omega b}{\gamma v} \right)^2 K_1^2 \left(\frac{\omega b}{\gamma v} \right). \quad (2.13)$$

297 This replaces the classical electric field of a point charge with a semiclassical field of photons.
 298 The photon flux in Eq. 2.13 provides the electromagnetic input to the J/ψ photoproduction cross
 299 section calculation.

300 **2.2 The STARlight model**

301 The STARlight model [40, 41] for calculating the J/ψ photoproduction cross section has three
 302 main components. The STARlight approach is constructed from the Weizsäcker-Williams photon
 303 flux, uses the vector meson dominance [51] fit to the proton-electron data, and uses the Glauber
 304 model [52] for calculating the nuclear cross sections from the proton-electron cross sections. The
 305 Weizsäcker-Williams photon flux provides the probe. The proton-electron scattering data com-

306 bined with the Glauber model create a picture of the initial state of the nucleus. Each of the
 307 different approaches discussed in this thesis to calculating the UPC J/ψ photoproduction cross
 308 section use these same elements. However, the different models each use the last two elements
 309 differently to produce different pictures of the nucleus and different cross sections values.

310 The photon flux in the photoproduction cross section calculation must be finite in order for the
 311 cross section to be meaningful. The Weizsäcker-Williams approximation, Eq. 2.13, diverges at
 312 $b = 0$. The probability of the nuclei interacting would exceed one if the photon flux were infinite.
 313 Special treatment of impact parameter, b , where the colliding nuclei overlap cancels the divergency.

314 A convolution of the photon flux with the nucleon number density functions removes the di-
 315 vergency at $b = 0$. The nucleon density of a single nucleus is given by

$$\rho_A(s) = \frac{\rho_0}{1 + \exp[(s - R_{WS})/d]}, \quad (2.14)$$

316 where s is the distance from the center of the nucleus, R_{WS} is the radius of the nucleus, and d is the
 317 skin depth, which determines how quickly the nucleon density falls off beyond the nuclear radius.
 318 Here WS refers to the Wood-Saxon distribution. In Eq. 2.15 the depth of the nucleus is integrated
 319 out leaving just the transverse dimensions in $T_A(\vec{r})$. The average number of nucleons in the overlap
 320 region, T_{AA} , is given by a convolution of T_A from each of the two nuclei.

$$\begin{aligned} T_A(\vec{r}) &= \int dz \rho_A(\sqrt{|\vec{r}|^2 + z^2}), \text{ and} \\ T_{AA}(|\vec{b}|) &= \int d^2\vec{r} T_A(\vec{r}) T_A(\vec{r} - \vec{b}). \end{aligned} \quad (2.15)$$

321 For a given impact parameter b , the product of $T_{AA}(|\vec{b}|)$ and σ_{NN} gives the average number of
 322 nucleon-nucleon collisions. The Poisson distribution gives the probability that no collisions occur:

$$P_0(b) = \exp[-T_{AA}(b)\sigma_{NN}], \quad (2.16)$$

323 where σ_{NN} is the cross section for a nucleon-nucleon interaction, which gives the probability that

a collision will occur given the average number of nucleons in the overlap region. T_{AA} is the mean of the Poisson distribution in Eq. 2.16. The photon flux is averaged over impact parameter by integration of the b -dependent photon flux multiplied by the b -dependent probability of having no nucleon-nucleon interactions:

$$\frac{dN_\gamma(k)}{dk} = \int_0^\infty 2\pi b db P_0(b) \int_0^R \frac{r dr}{\pi R_A^2} \int_0^{2\pi} d\phi \frac{d^3 N_\gamma(k, b + r \cos(\phi))}{dk d^2 r}. \quad (2.17)$$

Although Eq. 2.17 goes down to $b = 0$ where the photon flux is infinite, the fact that the probability of having a nucleon-nucleon collisions is high eliminates the divergency. T_{AA} reduces Eq. 2.13 at values of b where the nuclei overlap significantly and eliminates the divergency in the photon flux. The convolution of the photon flux with the b -dependent probability that no nucleon-nucleon collisions occur removes the divergency in Eq. 2.13.

A power-law fit to the photoproduction data off of proton targets gives an analytic expression for the energy dependence of the this photoproduction cross section. The forward photoproduction cross section is parameterized by

$$\left. \frac{d\sigma(\gamma p \rightarrow V p)}{dt} \right|_{t=0} = b_v (XW^\epsilon + YW^{-\eta}), \quad (2.18)$$

where W is the center of mass energy of the proton-photon system in Eq. 2.18. The remaining variables in Eq. 2.18 are power-law fit parameters. The XW^ϵ term characterizes Pomeron[53] mediated interactions, and the $YW^{-\eta}$ term characterizes meson mediated interactions [40]. The Pomeron can be considered to be a pair of gluons in a color singlet state. J/ψ 's high mass relative to the π and ρ^0 renders the second term in Eq. 2.18 negligible as the term falls rapidly with increasing W . Eq. 2.18 allows for extrapolation and interpolation of the measured forward proton photoproduction cross section. The fit to the data provides estimates for energies that have not yet been probed experimentally. The proton photoproduction cross sections from the electron-proton scattering data is a direct input to the STARlight model. In this method, a power-law fit to the proton photoproduction data is the input for the Glauber calculation.

346 Vector meson dominance and the optical theorem allow for the calculation of the total proton-
 347 meson scattering cross section from the fit given by Eq. 2.18. The optical theorem relates the
 348 total cross section, σ , to the corresponding forward scattering cross section, $d\sigma/dt|_{t=0}$, where t is
 349 the momentum transfer squared. The most likely fluctuations of the photon are to vector mesons
 350 because of the quantum numbers of the photon. For this reason, vector meson dominance asserts
 351 that only the vector meson fluctuations of the photon need to be considered. The forward scattering
 352 cross section is given by the following

$$\begin{aligned} \frac{d\sigma(\gamma p \rightarrow Vp)}{dt} \Big|_{t=0} &= \frac{4\pi\alpha}{f_v^2(M_V, \Gamma_{l+l-})} \frac{d\sigma(Vp \rightarrow Vp)}{dt} \Big|_{t=0}, \\ \sigma(Vp)_{tot}^2 &= 16\pi \frac{d\sigma(Vp \rightarrow Vp)}{dt} \Big|_{t=0}, \end{aligned} \quad (2.19)$$

353 where M_V , is the mass of the vector meson, and Γ_{l+l-} , is leptonic decay width. The result of
 354 combining vector meson dominance and the optical theorem in Eq.2.19 provides the cross section
 355 for a meson to scatter off a proton.

356 The Glauber model allows for Eq. 2.19, the proton-meson scattering cross section, to be used to
 357 calculate a nucleus-meson scattering cross section. The Glauber model is used to calculate nuclear
 358 cross sections from nucleon interaction cross sections by use of T_{AA} . The combination of the mean
 359 number of nucleons in the overlapping region of a nucleus-nucleus collision, T_{AA} , the nucleon cross
 360 section, σ , and the Poisson distribution make-up the core of Eq. 2.20. For the total nucleus-meson
 361 scattering cross section, the equation has the following form:

$$\sigma_{tot}(VA) = \int d^2\vec{r} (1 - e^{-\sigma_{tot}(Vp)T_{AA}(\vec{r})}). \quad (2.20)$$

362 In Eq. 2.20, the term $e^{\sigma_{tot}(Vp)T_{AA}}$ gives the probability of having no meson-nucleon scatterings from
 363 the Poisson distribution. The probability of having at least one scattering is given by subtracting
 364 one from the term $e^{\sigma_{tot}(Vp)T_{AA}}$ in Eq. 2.20.

365 Reversing the process used for the proton, Eq. 2.20, the meson nucleus scattering cross section,

relates to forward nuclear photoproduction cross section through the optical theorem. Using the optical theorem, the nuclear photoproduction cross section is given by

$$\begin{aligned} \left. \frac{d\sigma(\gamma A \rightarrow VA)}{dt} \right|_{t=0} &= \frac{\alpha \sigma_{tot}^2(VA)}{4\pi f_v^2}, \\ \sigma(\gamma A \rightarrow VA) &= \left. \frac{d\sigma(\gamma A \rightarrow VA)}{dt} \right|_{t=0} \int_{t_{min}}^{\infty} dt |F(t)|^2, \end{aligned} \quad (2.21)$$

where F is the Fourier transform of the nuclear density function, ρ_A . Eq. 2.21 is joined with the photon flux incident on the nucleus resulting in the following

$$\sigma(AA \rightarrow AAV) = 2 \int dk \frac{dN_\gamma}{dk} \sigma(\gamma A \rightarrow VA). \quad (2.22)$$

The factor of 2 in Eq. 2.22 comes from the fact that both of the two colliding nuclei contribute. Vector meson production rates in UPC collisions can be calculated by Eq. 2.22. In this thesis the measured UPC J/ψ cross section is compared to the STARlight predictions.

2.3 The Adeluyi and Bertulani model

To calculate the UPC J/ψ photoproduction cross section, the Adeluyi and Bertulani model [38, 39, 54, 55] uses the nuclear gluon density to characterize the nucleus and the Weizsäcker-Williams approximation for the probing photon flux. The AB method combines these components such that the nuclear gluon density is a direct variable. The nuclear gluon density term in the AB formulation allows for the use of a variety of nuclear gluon density models [56, 57]. A range of nuclear gluon densities are present in the available models resulting in a wide range of possible cross section values. The UPC J/ψ photoproduction cross section is correlated with the gluon density of the nucleus, increasing with higher densities and decreasing with lower densities. In the AB approach, the calculation of the UPC J/ψ photoproduction cross section allows experiments to constrain many different nuclear gluon density models.

In the AB method, the photon interacts with the nucleus by fluctuating to a quark-antiquark

pair. For J/ψ [production](#), the photon fluctuates to a $c\bar{c}$ pair. The probability for the photon to fluctuate to a $c\bar{c}$ pair depends on $M_{J/\psi}$, the mass of J/ψ , Γ_{l+l^-} , the J/ψ leptonic decay width, and α , the electromagnetic coupling constant. These three variables connect the c quark to the electromagnetic force mediator, the photon. Recast as a $c\bar{c}$ pair, the photon couples to the nuclear gluon density. The AB method uses the fluctuation of the photon to a $c\bar{c}$ pair as the foundation for calculating the forward J/ψ photoproduction cross section.

The $c\bar{c}$ pair arising from the photon fluctuation scatters off the gluons of the nucleus. The density of gluons in the nucleus determines how likely and therefore how large the cross section is for the quarks to scatter and form a J/ψ . The forward scattering cross section is the portion of those scattering events which transfer the minimum amount of momentum between the photon and the nucleus. The forward cross section for J/ψ photoproduction in the nucleus has the following form [38]:

$$\left. \frac{d\sigma_{\gamma A \rightarrow J/\psi A}}{dt} \right|_{t=0} = \xi_{J/\psi} \left(\frac{16\pi^3 \alpha_s^2 \Gamma_{l+l^-}}{3\alpha M_{J/\psi}^5} \right) [xG_A(x, \mu^2)]^2, \quad (2.23)$$

where $\xi_{J/\psi}$ is an experimentally derived correction factor, α_s is the strong coupling constant, x is the momentum fraction of the nucleus the scattering gluons carry, and G_A is the gluon density of the nucleus. Both the c and \bar{c} couple to the gluon density, and the double coupling results in the squared dependence of the cross section on the gluon density in Eq. 2.23. Fitting Eq. 2.23 to proton-electron scattering data sets $\xi_{J/\psi}$ [38]. The forward scattering cross section given by Eq. 2.23 connects the photon flux to the gluon density and provides the input to calculate the total cross section by the optical theorem.

The optical theorem relates the forward cross section in Eq. 2.23 to the total photoproduction cross section. The total cross section gives the probability that a photon incident on the nucleus will produce a J/ψ regardless of the momentum transferred in the interaction. The total cross section equation is given by

$$\sigma_{\gamma A \rightarrow J/\psi A}(k) = \left. \frac{d\sigma_{\gamma A \rightarrow J/\psi A}}{dt} \right|_{t=0} \int_{t_{min}(k)}^{\infty} dt |F(t)|^2, \quad (2.24)$$

where $t_{min} = (M_{J/\psi}^2/4k\gamma)^2$, which is the minimum amount of momentum transfer required to produce a J/ψ given the photon wave number k . The k dependence of t_{min} translates to the rapidity dependence of the total cross section. The total cross section for photoproduction, Eq. 2.24, provides the input to Eq. 2.28, which gives the rapidity dependence of the UPC photoproduction cross section. Eq. 2.24 as input to Eq. 2.28 allows for experimental comparison of the AB method to measurements of UPC photoproduction cross sections. With the AB method's direct use of the nuclear gluon density in Eq. 2.23, the AB method allows for experimental exploration of any gluon density model.

2.4 The Leading Twist Approach model

The Leading Twist Approach [42, 43] is another method for calculating UPC photoproduction cross sections. Contrary to the STARlight model, the LTA model introduces additional nuclear effects originating from modification of the nuclear gluon density. The LTA method uses the Weizsäcker-Williams approximation to calculate the photon flux created by the colliding nuclei. As in the STARlight method, the probability of having no hadronic collisions modulates the flux. The photon flux for the LTA method has the following form [43]:

$$n_{\gamma/A}^i(\omega_\gamma) = \frac{2\alpha Z^2}{\pi} \int_{b_{min}}^{\infty} db \frac{x^2}{b} \left[K_1^2(x) + \frac{K_0^2(x)}{\gamma_L^2} \right] P_0(b) P_C^i(b), \quad (2.25)$$

where $x = \frac{\omega b}{\gamma_L}$, and $K_0(x)$ term contributes a photon flux in the transverse direction, and $P_C^i(b)$ is an modulation factor that requires various additional interactions. These interactions result in emission of neutrons from the receding nuclei as the nuclei relax from excited states. The term P_C^i provides additional ways to distinguish UPC events from nuclear collisions experimentally but leaves the underlying interaction mechanism the same. For example, this additional term in the LTA formulation of the photon flux produces calculations of asymmetric neutron emission, which separate UPC events from nuclear collisions. In UPC events the number of forward neutrons on either side of the interaction can differ significantly, whereas in nuclear collisions this is rare.

The LTA model derives the nucleon cross section from derivations of the nucleon gluon densities from electron-proton scattering data and leading order perturbative quantum field theory calculations. The forward photoproduction cross section of the nucleon has the following form [43]:

$$\frac{d\sigma_{\gamma N \rightarrow J/\psi N}(t=0)}{dt} = \frac{16\Gamma_{l+l-}\pi^3}{3\alpha M_{J/\psi}^5} [\alpha_s \mu^2 x G_N(x, \mu^2)]^2, \quad (2.26)$$

where G_N is the gluon density of the nucleon, x is the fraction of the nucleon's momentum the gluon carries, and μ is related to momentum at which the nucleon is being probed, which is equal to $M_{J/\psi}/2$ for J/ψ photoproduction. By connecting the gluon density to the cross section, Eq. 2.26 allows for the gluon density to be experimentally probed.

The LTA model exploits the optical theorem to relate the forward photoproduction cross section of the nucleon to the nuclear cross section. The relation is

$$\sigma_{\gamma A \rightarrow J/\psi A}(\omega) = \frac{d\sigma_{\gamma N \rightarrow J/\psi N}}{dt}(\omega, t_{min}) R_g^2 \int_{t_{min}}^{\infty} dt |F(t)|^2, \quad (2.27)$$

where R_g is the nuclear modification function, the ratio between the gluon density of the nucleon, G_N , to the gluon density of the nucleus, G_A . As with the STARlight method, the optical theorem relates the forward cross section, $\frac{d\sigma_{\gamma N \rightarrow J/\psi N}}{dt}(\omega, t_{min})$, to the total cross section, $\sigma_{\gamma A \rightarrow J/\psi A}$.

From Eq. 2.27, the LTA method can predict the angular distribution of photoproduced J/ψ with respect to the beam axis. The angular distribution is expressed in the form of the rapidity dependency of the UPC photoproduction cross section given by

$$\frac{d\sigma_{A_1 A_2 \rightarrow A_1 A_2 J/\psi}}{dy} = n_{\gamma/A_1}(y) \sigma_{\gamma A_2 \rightarrow J/\psi A_2}(y) + n_{\gamma/A_2}(-y) \sigma_{\gamma A_1 \rightarrow J/\psi A_1}(-y), \quad (2.28)$$

where $y = \ln\left(\frac{2\omega}{M_{J/\psi}}\right)$. Eq. 2.28 is comprised of two terms, one for photons from the forward going nucleus interacting with the backward going nucleus, and a second for the reverse situation. The integration of Eq. 2.28 over y produces the factor of 2 that is present in Eq. 2.22. The rapidity distribution of the photoproduction cross section given in Eq. 2.28 provides a more detailed prediction

and allows for more direct experimental comparisons.

2.5 Photon-induced nuclear break-up

In addition to the photoproduction of quark anti-quark resonances such as the J/ψ , photo-nuclear interactions can also result in emission of a neutron from the struck target nucleus [58, 59]. To calculate the cross section for neutron emission in UPC events, the photon flux calculated from the Weizsäcker-Williams approximation is combined with the nuclear photon absorption cross section. The absorption of photons can be described by two processes, namely the Giant Dipole Resonance (GDR) [60, 61], and the dissociation of deuterium. In this section, the theoretical description of photon-induced nuclear break-up is discussed.

Following the formulation in [62], the cross section for a nucleus to absorb a photon, σ_{PN} , is given by

$$\sigma_{PN} = \sigma_{GDR} + \sigma_{QD}. \quad (2.29)$$

The Giant Dipole Resonance (GDR) cross section, σ_{GDR} is the result of the collective motion of the protons relative to the neutrons and dominates at lower photon frequencies. The Quasi-Deuterium (QD) cross section, σ_{QD} , represents the nucleus as a collection of proton-neutron pairs (deuterium) and dominates at higher photon energies.

There are two models for the GDR: the Goldhaber and Teller model [63], and the Steinwedel and Jensen model [64]. Goldhaber and Teller treats the protons and neutrons as two separate and ridged density profiles that, when excited, oscillate with respect to each other [65]. Steinwedel and Jensen modeled the protons and neutrons as fluids contained in a single sphere that have shifting density profiles [65].

In the Goldhaber and Teller model, the potential that holds protons and neutrons together depends on the difference of the neutron and proton densities squared. Assuming the neutron and proton densities have the same shape, if the two fully overlap, there is no difference in the densities and the potential energy is zero. If the two density distributions are separated, the overlap in the

475 shape will not cancel. In the separated configuration, there will be a non-zero potential energy.
 476 The potential energy has the form of a harmonic oscillator with a spring constant that depends on
 477 the initial density distribution [65] given by

$$U = \frac{1}{2}Kz^2, \quad K = k \int d^3r (\nabla \rho_0)^2. \quad (2.30)$$

478 If the nucleus has a shape cut off in density at its edge, then the integral is dominated by the
 479 region at the surface, and the spring constant, K , becomes proportional to $A^{2/3}$, where A is the
 480 mass number of the nucleus. The surface area of a sphere is proportional to its volume to the $2/3$
 481 power explaining the mass dependence. Due to this dependence, the frequency of the giant dipole
 482 resonance in the Goldhaber and Teller model is given by

$$\omega = \sqrt{\frac{K}{M}} \propto \sqrt{\frac{A^{2/3}}{A}} = A^{-1/6}. \quad (2.31)$$

483 This dependence describes light nuclei well, but it does not describe heavier nuclei [65]. The
 484 Steinwedel and Jensen model can be used to describe heavier nuclei. In this model, the proton
 485 and neutron fluids are confined to a single sphere where they are allowed to slosh back and forth
 486 creating the same effect as the Goldhaber and Teller model. Here there is no global separation of
 487 the proton and neutron fluids. The dipole is created by under-densities and over-densities of the
 488 proton and neutron fluids. It can be shown that this results in a frequency of oscillation which
 489 depends on one over the radius of the nucleus [65]:

$$\omega \propto \frac{1}{R} \propto A^{-1/3}. \quad (2.32)$$

490 As before, the relationship in Eq. 2.32 arises from the geometry of a sphere. The dependence of
 491 the giant dipole resonance that is seen in the Steinwedel and Jensen model describes medium and
 492 heavier mass nuclei well. Empirically, both models are put together to give the following mass

493 number dependence of the dipole resonance [65]

$$E_{GDR} = 32.2A^{-1/3} + 20.6A^{1/6}. \quad (2.33)$$

494 In order to compute the effect of an excitation in either model, the harmonic oscillator solutions
 495 found earlier can be driven by an interacting force. The resulting differential equation can then
 496 be solved using a Fourier transform to eliminate the time derivatives. In this model the driven
 497 harmonic has the following Lorentzian form:

$$\sigma_{GDR}(E_\gamma) = \frac{\sigma_{max} E_\gamma \Gamma_{GDR}^2}{\left(E_\gamma^2 - E_{GDR}^2\right)^2 + E_\gamma^2 \Gamma^2}, \quad (2.34)$$

498 where σ_{max} is the maximum cross section reached when $E_\gamma = E_{GDR}$; E_{GDR} is the peak resonance
 499 energy, and Γ_{GDR} is the width of the resonance. The width of this distribution lies in a range from
 500 4-8 MeV and depends on the orbital arrangement to the neutrons and protons in the given nucleus
 501 [65].

502 For higher energy photons, the quasi-deuterium cross section is needed. The nucleus is treated
 503 as a collection of proton-neutron pairs, which are screened by the rest of the nucleus, in the quasi-
 504 deuterium approach. This behavior is modeled by [62]

$$\sigma_{QD}(E_\gamma) = L \frac{(L-A)Z}{A} \sigma_d(E_\gamma) F(E_\gamma), \quad (2.35)$$

505 where σ_d is the deuterium disintegration cross section for $\gamma + d \rightarrow p + n$; F is a function from Pauli
 506 blocking of fermions, and L is an empirical parameter set by data to 6.5 [62]. Certain energy levels
 507 are not available to the products of the deuterium disintegration process because of the presence of
 508 the rest of the nucleus. The result is a reduction of the cross section relative to free deuterium.

509 F can be modeled with an exponential cutoff below 20 MeV, a polynomial in the intermediate
 510 range with nearly linear dependence on E_γ , and an inverted exponential above 140 MeV, pushing
 511 F to one at higher values of E_γ [62]. Essentially, the model at low photon energies disallows deu-

terium disintegration because the products have no available state to occupy, and at high energies the rest of the nucleus becomes transparent and looks more and more like a collection of deuterium. The deuterium disintegration cross section is found empirically and is fit to the following function [62]:

$$\sigma_d(E_\gamma) = 61.2 (E_\gamma - 2.224)^{3/2} / E_\gamma. \quad (2.36)$$

In order to produce a final state for the target nucleus, a branching ratio is needed. The branching ratio gives the probability that the photo-excited nucleus will end up in a particular state. This determines the sort of emission that will result from the de-excitation process [62].

All the tools are now assembled to calculate the cross section for neutron emission. The first step in the calculation is to assume that the number of photons absorbed by either nucleus in the collision obeys the Poisson distribution [66, 62]. The average number of absorptions as a function of impact parameter, $m(b)$,

$$m(b) = \int_{\omega_{min}}^{\omega_{max}} N(\omega, b) \sigma_{PN}(\omega) d\omega. \quad (2.37)$$

where σ_{PN} is the photo-nuclear cross section, and N is the photon flux (see Eq. 2.13).

In [62], the probability of any final state i due to the absorption of a single photon is given by

$$P_i(b) = \int_{\omega_{min}}^{\omega_{max}} P_a(b, 1) q(b, \omega) f_i(\omega) d\omega = e^{-m(b)} \int_{\omega_{min}}^{\omega_{max}} N(b, \omega) \sigma_{PN}(\omega) f_i(\omega) d\omega, \quad (2.38)$$

where P_a is the probability that the target absorbs a single photon as calculated by following the Poisson distribution; q is the probability that the photon will have the frequency ω , and f_i is the branching ratio to a given final state. The following equation describes q [67]

$$q(b, \omega) = \frac{N(b, \omega) \sigma_{PN}(\omega)}{m(b)} \quad (2.39)$$

Integrating over the impact parameter to get an area that is weighted by the probability func-

tions gives [66, 62]

$$\sigma_i = 2\pi \int_{b_0}^{\infty} b P_i(b) db. \quad (2.40)$$

Three parameters arise when calculating the cross section in Eq. 2.40, the minimum impact parameter b_0 , the minimum emitted photon frequency ω_{min} , and the maximum emitted photon frequency ω_{max} . A minimum impact parameter ensures that the Bessel function in the photon flux in Eq. 2.13 is not evaluated at zero. At zero, the modified Bessel function does not converge. Physically, a minimum impact parameter is selected in order to separate the domains between electromagnetic interactions and the strong interactions that happen inside the nucleus. To serve this end, the minimum impact parameter is set to the radius of the nuclei [66, 62]. This excludes collisions where the nuclei overlap in the calculation, and ensures that only electromagnetic interactions are involved.

2.6 Experimental results

2.6.1 UPC measurements at RHIC

One of the first results from RHIC was a UPC measurement, the measurement of the neutron spectrum from photon-induced nuclear break-up [68]. Neutrons due to the nuclear photon absorption are emitted with momenta on the order of the giant dipole resonance, about 60 MeV. Momenta of 60 MeV compared to the 100 GeV beam energy at RHIC result in neutron emission at very small angles, $\sim 2\text{mrad}$, relative to the beam. These neutrons are captured by calorimeters, which are set in line with the beam called Zero Degree Calorimeters (ZDCs). These detectors are described in Section 3.5. Figure 2.2 shows the charge recorded by the ZDCs (see Section 3.5). The peak at ~ 30 in Fig. 2.2 is due to emission of a single neutron. Each successive peak is due to emission of an additional neutron. From these peaks, the cross section for nuclear break-up by neutron emission by both nuclei was measured. The fraction of nuclear interactions in all events with neutrons on both sides of the interaction point was found to be 0.661 ± 0.014 [68], which is in agreement the value 0.659 predicted by the model described in [62] and Section 2.5.




Figure 2.2: ZDC neutron spectrum at RHIC with each a Gaussian fit to the 1, 2, 3, 5, and 5 neutron peak [68].

Nuclear collisions were separated from photon-induced break up by counting hits in the central scintillating counter similar to a tracker. These scintillators covered the pseudo-rapidity region $3.05 < |\eta| < 3.85$. Events with hits on both sides of the interaction point were categorized as nuclear interactions, and those with no hits or hits on only one side were categorized as electromagnetic interactions. The difference in the charge measured on each of the two sides of the interaction point, by the two ZDCs, is divided by the total charge for both ZDCs (see Fig. 2.3). Figure 2.3 shows that events created by electromagnetic interactions result in asymmetric neutron emission, whereas nuclear interactions produce neutrons on both sides. In the analysis presented in this thesis, the energy asymmetry in the ZDCs is used to select UPC events. This will be discussed in Chapter 4.

Both the ρ^0 and J/ψ meson photoproduction cross sections in UPC events were measured at RHIC. STAR measured the photoproduction of the ρ^0 meson at collisions energies per nucleon of 62.4 GeV [69], 130 GeV [70], and 200 GeV [71]. The J/ψ was measured by PHENIX at 200 GeV [72]. The ρ^0 measurements by STAR were in good agreement with the STARlight calculation described in Section 2.2. The J/ψ measurement ~~where~~ were limited by statistic as only 10 J/ψ




Figure 2.3: The energy asymmetry in the ZDCs for electromagnetic interactions, coulomb events, and nuclear interaction, hadronic events [68].

candidates were found.

2.6.2 UPC J/ψ at the LHC

The increase in beam energy from 200 GeV at RHIC to 2.76 TeV at the LHC results in an increase in the Lorentz γ of about 10. Because the photon flux depends on γ^2 , the photon flux at the LHC increases by a factor of about 100 compared to RHIC. ALICE and CMS have studied the production of UPC events. ALICE has studied coherent J/ψ photoproduction in ultra-peripheral PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV. The cross section for this process was measured at both forward-rapidity, $y = 3$ [73], and mid-rapidity, $y = 0$ [74]. With an integrated luminosity of $55 \mu b^{-1}$, ALICE measured $78 \pm 10(\text{stat})^{+7}_{-11}(\text{syst})$ coherent J/ψ candidates at forward rapidity. The measured cross section was $1.00 \pm 0.18(\text{stat})^{+0.24}_{-0.26}(\text{syst})$ mb. For a symmetric system like such as with PbPb collisions, as opposed to pPb collisions, there is an ambiguity between which ion is the target and which is the photon emitter. Therefore, the cross section has a contribution from the low- x and high- x parts of the gluon density. At $y = 3$ for PbPb collisions at the LHC, the cross section has a contribution from both $x = 5 \times 10^{-5}$ and $x = 2 \times 10^{-2}$, where x is the fraction nucleon's

momentum carried by the struck parton. This ambiguity is not present at $y = 0$. ALICE has also measured the coherent J/ψ photoproduction cross section at $y = 0$, using an integrated luminosity of about $23 \mu b^{-1}$. 291 ± 18 (stat) ± 4 (syst) and 265 ± 40 (stat) ± 12 (syst) coherent J/ψ candidates were measured in the dimuon and dielectron channels, respectively. The combined cross section from both channels was measured to be $2.38^{+0.34}_{-0.24}$ (stat+syst) mb. At $y = 0$, $x \sim 10^{-3}$, which is a smaller x than at forward rapidity, and more sensitive to the nuclear gluon shadowing (see Fig. 2.4).




Figure 2.4: Coherent J/ψ photoproduction cross section in ultra-peripheral PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV, measured by the ALICE experiment at forward and mid-rapidity [73, 74].

587

The ALICE data points have been compared to several theoretical models. The UPC photoproduction cross section calculations depend significantly on how the nucleus is represented in the calculation. The results from the STARlight, LTA, and AB methods vary from a relatively large cross section in the STARlight model, ranging through a variety of values in the AB method, to a relatively small cross section in the LTA method. Each of these methods utilizes the same probe of the nucleus, the equivalent photon flux that is calculated using the Weizsäcker-Williams approximation (see Section 2.1). The three methods deviate in how they calculate the forward photoproduction scattering cross section. The differences in the UPC photoproduction cross sections predicted by the different models demonstrates the amount of experimental sensitivity there

Model	$\sigma_{AA \rightarrow AAJ/\psi}(mb)$
STARlight/STARlight MC	23
LTA	9
AB-MSTW08	34
AB-EPS08	7
AB-EPS09	14
AB-HKN07	23

Table 2.1: $\sigma_{AA \rightarrow AAJ/\psi}(mb)$ obtained by the LTA, STARlight, AB methods for PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV. Four different gluon density models [38, 39, 54] are used in the AB method. STARlight is a simulation software package that utilizes the STARlight model.

is to distinguishing between the models. The dependence of the cross section on rapidity is clearly visible.

The cross section value calculated by Eq. 2.22 in the STARlight, LTA, and the various gluon density models in AB method vary significantly. Table 2.1 gives the predicted values for the three main methods taken from [39], [43], and [40]. The cross sections in Table 2.1 differ by a factor of 4 from the smallest to largest and create an experimental opportunity.

The nuclear suppression factor, S , demonstrates the difference between how the models represent the nucleus. S is the ratio between the nuclear photoproduction cross section and the free nucleon photoproduction cross section. It is a measure of how the nuclear gluon densities evolve in each of the models. Figure. 2.5 from [75] shows the nuclear suppression, which is equivalent to R_g in Eq. 2.27, for the LTA and AB methods. Fig. 2.6 shows the nuclear suppression for the STARlight method [75]. S is not directly related to the gluon density because the STARlight uses a Glauber approach, which uses the $\gamma - p$ photoproduction cross section rather than the nuclear gluon density directly. Fig. 2.5 and Fig. 2.6 show that as the momentum of the probing photon goes up, increasing $W_{\gamma p}$, and momentum of the probed gluon goes down, decreasing x , the nuclear gluon density decreases relative to the free nucleon. The nuclear suppression factor, S , allows for the different models' representations of the gluon content of the nucleus to be directly compared to each other and to data. S can be measured from data by assuming a Weizsäcker-Williams photon flux and provides insight into nuclear gluon densities.




Figure 2.5: Both panels show the nuclear suppression factor, S , obtained from the ALICE data points and calculations by the AB (top) and LTA (bottom) models [75]. S is denoted as R_{pb} in this figure.

In addition, ALICE reported the measurement of the incoherent J/ψ photoproduction cross section at mid-rapidity [74]. This provided additional constraints to the models for gluon shadowing.

In Chapter 5, the coherent UPC J/ψ photoproduction cross section using CMS is described. The measurement in this thesis adds to existing ALICE results, covering an intermediate range of x values.




Figure 2.6: Nuclear suppression factor, S , in STARlight ~~method~~model [75]. The y-axis is labeled S because for the STARlight model S is not ~~neccessary~~necessarily equal to R .

Chapter 3

The Compact Muon Solenoid detector

The Compact Muon Solenoid (CMS) is housed at interaction point 5 of the LHC. The LHC is designed to pursue physics at the TeV scale. This is the scale where electroweak symmetry breaking is believed to occur [76]. While the search for the standard model Higgs boson was the central driving design consideration, the wide range of possibilities for finding new physics signals requires a general purpose detector. CMS was optimized for efficient muon triggering and identification. Muon capabilities developed for the Higgs boson studies can be used to study J/ψ [production](#). A versatile trigger is needed to accommodate the high interaction rates that accompany the high luminosities. By exploiting the versatility of the trigger and muon systems it is possible to explore processes like UPC J/ψ production, which push to the low energy edge of the experiment's capabilities.

CMS is dominated by the massive 4T superconducting solenoid at its core. The magnet is 13m long with a 6m diameter, and pushes the limits of power and compactness [13]. These two conflicting limits are achieved through the novel design of interweaving structural and conducting elements together in the coil of the solenoid.

Within the solenoid resides three different sub detectors. The innermost is the world's largest silicon tracker [13]. The tracker is surrounded by a highly effective lead tungstate crystal electromagnetic calorimeter (ECAL) [77]. ECAL is encapsulated in a brass scintillating hadronic




Figure 3.1: The Compact Muon Solenoid layout [13].

calorimeter (HCAL). Outside the magnet, muon chambers are used to aid in the measurement and triggering of muon events. Altogether CMS weighs 12,500 metric tons, has a diameter of 14.6m, and a length of 21.6m [13].

3.1 Tracker

The Silicon Tracker [78] is the innermost sub-detector of CMS, and has active elements as close as 4.4cm to the interaction point [13]. The tracker has a length 5.8m, a diameter of 2.6m and covers a range in pseudorapidity of $|\eta| < 2.5$. At the center of the tracker are three rings of silicon pixels around the beam axis with two disks of silicon ~~pixels~~ pixel detectors to cap the rings. The pixel detector portion of the silicon tracker is comprised of 66×10^6 pixels. The silicon ~~pixels~~ pixel detectors are surrounded by silicon strips. The silicon strips are separated into 4 different sections: the Tracker Inner Barrel, the Tracker Inner Disk, the Tracker Outer Barrel, and the Tracker End Caps. The silicon strip detectors as a whole are comprised of 9.3×10^6 silicon strips. The high number of pixels and strips allow for the ability to distinguish and collect enough distinct points to reconstruct the path of the 1000 or so charged particles per bunch crossing expected at peak luminosity [13].

The amount of material present in the tracker is substantial enough to alter the path of particles




Figure 3.2: Layout of the silicon tracker with the pixels closest to the interaction point, marked with a black dot, and the strips segments beyond the pixels.

as they pass through the tracker. Fig. 3.3 shows the amount of material in the tracker as a function of radiation lengths (X_0). A radiation length is the mean distance a photon travels before interacting. The deflection angle set by the strength of the magnetic field determines the momentum resolution for high momentum particles. The momentum resolution for lower momentum tracks is limited by the loss of energy due to multiple scattering of these particles off the material of the detector. For UPC J/ψ , this is the primary factor contributing to the resolution of the reconstructed muon tracks.




Figure 3.3: Material budget in the tracker broken down by sub-detector(left) and category (right) [13].

3.2 ECAL

The next detector beyond the tracker is the electromagnetic calorimeter system, ECAL. The calorimeter system is made of 61,200 lead tungstate (PbWO_4) crystals in the central barrel and 7,324 in each of the two endcaps [13]. The barrel (EB) covers a pseudorapidity range $|\eta| < 1.479$ and has an $\eta - \phi$ segmentation of approximately 0.0174×0.0174 , depending slightly on the position of the fixed sized crystals. Lead tungstate is very dense giving the ECAL crystals a high number of radiation lengths within a short depth. The crystals of the barrel have a depth of 230 mm corresponding to $25.8 X_0$. The endcaps (EE) cover the pseudorapidity region $1.479 < |\eta| < 3$. In the endcap the crystals have an exposed area of $28.62 \times 28.62 \text{ mm}^2$, and a depth of 220 mm corresponding to $24.7 X_0$. The energy resolution of the ECAL as measured by test beam data can be seen in Figure 3.4. The fractional energy resolution reduces quickly above 10 GeV and levels off at about 0.5% for




Figure 3.4: The energy resolution of ECAL as a function of energy [13].

energies above 100 GeV. For J/ψ with rapidity near 2, the electron will carry energies around 6 GeV and can be resolved by the ECAL.

3.3 HCAL

The HCAL [79] like the ECAL has both a barrel (HB) and endcaps (HE). The pseudorapidity region $|\eta| < 1.3$ is covered by HB [13]. HB has an $\eta - \phi$ segmentation of 0.0897×0.0897 , and is 25 times more sparsely granulated than EB. HE covers the pseudorapidity region $1.3 < |\eta| < 3$. HE, like EE and the tracker endcaps, is aligned parallel to the beam axis resulting in granularity that changes with η . In the region $1.3 < |\eta| < 1.6$ HE has an $\eta - \phi$ segmentation of 0.0897×0.0897 . The $\eta - \phi$ cell size roughly doubles to 0.17×0.17 in the region $1.6 < |\eta| < 3$. The energy resolution of the barrel and endcaps can be seen in Figure 3.5. The thickness of the hadronic calorimeter is best described in interaction lengths, λ_I , which is the mean distance a hadron travels before it experiences a nuclear interaction. At $\eta = 0$ the barrel has a thickness 5.82 interaction lengths (λ_I), and increases as the path length through the material increases to $10.6 \lambda_I$ at $|\eta| = 1.3$.




Figure 3.5: The E_T resolution of HCAL as a function of $|\eta|$ and E_T [13].

In addition to HB and HE, HCAL has two additional calorimeters. Because the space between ECAL and the magnet is restricted to 1.18 m, an outer hadronic calorimeter section (HO) is placed beyond the magnet in the region $|\eta| < 1.3$ [13]. The main function of HO is to collect energy from the highest energy hadrons before they reach the muon system. HO is not used in this analysis, but does contribute to the material budget. To increase the total calorimetric coverage, HCAL also has a quartz fiber calorimeter (HF) in the forward region, $3 < |\eta| < 5$. For the majority of HF's

13 η rings the $\eta - \phi$ segmentation is 0.175×0.175 . In the lowest $|\eta|$ ring the segmentation is 0.111×0.175 in $\eta - \phi$. In the highest two $|\eta|$ rings the segmentation in ϕ is 0.349, with an η segmentation of 0.175 in the outer and 0.300 in the innermost ring. The longitudinal direction is effectively segmented by using both short and long fibers. The long fibers are sensitive to energy deposited throughout the calorimeter, i.e. to both the hadronic and electromagnetic parts of the shower. The short fibers are only sensitive to the energy deposited more than 22cm inside the calorimeter, which is mainly hadronic. This allows electromagnetic showers to be distinguished from purely hadronic showers [13].

The energy resolution for HF can be seen in Figure 3.5. As with the ECAL, the fractional resolution increases with energy. This is due to the random nature of shower development. At larger energies, the fluctuations in the shower of particles collected in the calorimeter tend to average out.

3.4 CASTOR

Very forward angles are covered at one end of CMS ($6.6 < \eta < 5.2$) by the CASTOR calorimeter [80]. CASTOR is 1.6m long and 0.6m in diameter and sits 14.4m away from the nominal interaction point. It is 1.6m long and 0.6m in diameter. CASTOR is a Cerenkov calorimeter made of quartz fibers/plates embedded in tungsten absorbers. The quartz and absorbers are inclined at 40° to the beam axis to maximize the amount of Cerenkov light. CASTOR is segmented into 16 ϕ -sectors and 14 z modules. The two front modules comprise the electromagnetic section of the calorimeter with a depth of 20 radiation lengths. Each of the electromagnetic channels has half the depth of the hadronic ones. The total thickness of CASTOR is about 10 interaction lengths.

3.5 ZDC

Beyond HF, the Zero Degree Calorimeters (ZDCs) [81] covers the very forward rapidity region. The ZDCs sit between the beam pipes on either side of the interaction point covering the area

around $\theta = 0$, $|\eta| > 8.3$. In heavy ion collisions the ZDC has the ability to measure neutral particles that do not participate in the collision [13]. This detector plays an important role in the analysis described in this thesis by measuring energy due to neutrons produced by photon-induced nuclear break. These measurements are used to identify events with asymmetric neutron emission, reducing the contribution to the sample ~~by~~of peripheral heavy-ion collisions. This will be discussed further in Chapter ??.

The ZDC has a total of 18 channels. Half of these 18 channels are on either side of the interaction point. The 9 channels on the side of CMS that correspond to positive η are denoted ZDC^+ , where as the 9 channels on the negative side are denoted ZDC^- . The 9 channels on each side are further sub-divided into an electro-magnetic (EM) section and a hadronic (HAD) section. The EM section is positioned in front of the HAD section with respect to the interaction point and is segmented transverse to the beam direction. The 5 EM sections are positioned in front to absorb the energy from electro-magnetically induced showers, which develop over a shorter distance than hadronically induced showers. The transverse segmentation allows for a measurement of the transverse shower width and the size of the beam spot at the ZDC. The HAD section is segmented in the direction of the beam and consists of 4 channels. The longitudinal segmentation allows for absorption of the full extended hadronic shower and the ability to measure the longitudinal shower shape.

Each of the 18 channels contains a tungsten target and quartz fibers. The dense tungsten target is used to initiate the shower. The quartz fibers shine Cerenkov light as the high momentum charged particles from the shower pass through it. the light from the quartz fibers is channeled to photo-multiplier tubes, one for each ZDC channel. Through a cascade of photon induced electrical discharges, the photo-multiplier converts the Cerenkov light to an electrical pulse.

This electrical pulse travels ~ 200 m down a coaxial cable from the LHC tunnel to the counting house in the CMS service cavern. There the electrical pulse is digitized by the Charge Integrator and Encoder (QIE). The QIE integrates the current each 25 ns. The charge is then mapped logarithmically to the 128 bits. These bits are sent across a small fiber optic cable to the HTR firmware

card. Here each 25 ns signal is stored in a 250 ns buffer, and the timing is synchronize with the rest of the detector to ensure the ZDC signal arrives at the central data acquisition system at the same time as the other sub detectors from the same collision.

3.6 Muons

The muon system [13] resides just outside of the superconducting magnet. It consists of three complementary systems: drift tubes (DT) in the barrel, cathode strip chambers (CSC) in the endcaps, and resistive plate chambers (RPC) in both the barrel and endcap regions [13]. Each of these gaseous detectors function in the same way. As the muon penetrates the gas volume electrons are knocked off of the gas atoms and these electrons are collected in the positively charged anode, whereas the ionized gas moves to the cathode. The DTs in the barrel and the CSCs in the endcap have better spatial precision relative to the RPCs, which are quicker and have more precise timing. The combination of the DTs and RPCs in the barrel and the CSCs and RPCs in the endcap allow for fast triggering and muon identification during data reconstruction.




Figure 3.6: The CMS muon system showing the four DT stations in the barrel (MB1-MB4), the four CSC stations in the endcap (ME1-ME4), and the RPC stations.

As seen in Fig. 3.6, the DTs reside only in the barrel, covering the region $|\eta| < 1.2$. Consisting of a total of 172,000 cells, the DT cells are collected into 250 chambers. The DT chambers are interwoven into the magnet field return yoke and are labeled by 5 segmentations in z , YB-2 to YB+2. Each z segment is divided into 12 ϕ segments labeled 1 at $\phi = 0$ and going to 12 rotating in positive ϕ with segments 4 and 10 containing 2 chambers. The segmentation in r is divided into four parts, MB1-MB4. Fig. 3.7 shows how each chamber is made of three super layers. Super layers $SL \Phi_1$ and $SL \Phi_2$ measure (r, ϕ) , whereas $SL \Theta$ measures z .




Figure 3.7: Schematic of the DT chambers and an individual DT cell.

The RPCs complement the DTs in the barrel and the CSCs in the endcap primarily for the purpose of triggering. Throughout the barrel and endcap there are a total of 1020 RPC modules,

480 in the barrel and 540 in the endcap. Denoted by the red lines in Fig. 3.6, the RPCs are mounted on both sides of the DTs in the barrel for the inner most layers, MB1 and MB2. For the outer two layers of the barrel, MB3 and MB4, a single layer of RPCs is mounted on the inner side of the DTs. Each corresponding DT segment has two RPCs modules, except in MB2 where some segments contain 3 RPC modules. The outer ring of the muon endcaps are instrumented with RPCs covering up to $|\eta| = 1.6$.




Figure 3.8: Schematic of a RPC cell.

There are a total of 468 CSC modules in the muon system endcap. For the sectors in z , ME2, ME3, and ME4 in the endcap, the modules in the ring closest to the beam cover 20° in ϕ . The modules in ME1 and the modules beyond the inner ring in ME2 and ME3 have a $10^\circ \phi$ segmentation. The 2 million wires of the CSCs are grouped into 400,000 channels and are powered with 9000 high voltage supplies. The resolution of the CSCs is on the order of 1mm for position measurements in the plane perpendicular to the beam and they are 99% efficient for muons that pass through all four sectors, ME1, ME2, ME3, and ME4. Because of the relatively low momentum of the muons originating from J/ψ in UPC events, the muons in the analysis discussed in this thesis are all in the range $1.6 < |\eta| < 2.4$ and rely on the CSCs for triggering.

The primary function of the muon systems is to trigger on and identify muons. The tracker is still the primary instrument for measuring the muons. Fig. 3.10 shows that the muon system only improves the momentum resolution for muons with momenta above 100 GeV. For UPC J/ψ events, where muons momenta are between 4-8 GeV, the muon system does not provide any advantage in terms of improved resolution. The muon system is however important in distinguishing which




Figure 3.9: Schematic of the CSC chambers and an individual CSC cell.

tracks are due to muons as opposed to other charged particles. Because of their much larger mass relative to electrons, muons emit less bremsstrahlung, or breaking radiation, as they penetrate the inner layers of the CMS on its way to the muon systems. Fig. 3.11 shows the amount of material traversed by particles traveling through CMS as a function of $|\eta|$. The total of nearly 10 interaction lengths between the interaction point and the muon chambers ensures that hadrons like charged pions, which almost exclusively decay to muons, are absorbed in the calorimeters before converting to muons. By eliminating backgrounds from both electrons and hadrons, the CMS muons system allows for identification of muons for both triggering and reconstruction.

The low-momentum nature of UPC physics creates complications due to the large amount of material between the interaction point and the muon systems. About 3 GeV of momentum is needed to reach the first layers of the muon system. In the rest frame of the J/ψ , the J/ψ equally shares its rest mass with its decay products creating 2 muons with momenta of about 1.5 GeV. For these daughter muons to reach the muon system, their parent J/ψ must be pushed to higher




Figure 3.10: The momentum resolution for muons using only the tracker and the whole muon system in the barrel (left) and end cap (right) [13].

momentum by the initial particles which created the J/ψ . For this reason, muons from UPC J/ψ s are only detected at higher η values. Understanding this momentum restriction of the muon system was a major focus of the analysis discussed in this thesis with details described in Section. 5.4.




Figure 3.11: The amount of material in CMS as a function of η in number of interaction lengths [13].

3.7 Trigger

The CMS trigger is two tiered [82, 83]. The L1 trigger is the lower level hardware based system. The High Level Trigger (HLT) is ~~a software base~~ software based and runs on a computer farm located at point 5.

The purpose of the L1 trigger is to make quick decisions about which events will be kept temporarily for further processing. It uses only information from the calorimeter and muon systems and is used to identify events where the tracker should be read out. The output from the sub-detectors is synchronized to ensure that the signal from each of the sub-detectors comes from the same collision.

If an event passes the L1 trigger, the data from all the sub-detectors, including the tracker are sent to the HLT computing farm. At this level the raw data from all the sub-detectors is unpacked and combined. The information from the calorimeters, muon system, and tracker can all be used to reconstruct basic physic objects in the HLT farm. For example, tracks can be associated with either ECAL energy clusters to form electron candidates, or tracks can be combined with hits in the muon system to create muon candidates. At the HLT, the whole detector is used to select events. The raw data from the events that survive the HLT are recorded permanently, those that do not are lost forever.

The HLT farm must always be ready to accept events from the L1 trigger. For this reason, the amount of computing time each HLT trigger path uses must be balanced. For more rare L1 triggers, which will occur at a lower rate, more complex reconstruction software can be used. Conversely, simpler, faster, methods must be used for more common high rate triggers. Because of this time constraint in the HLT farm, the reconstruction algorithms used for triggering ~~tended~~ tend to differ from the final reconstruction algorithms. In the HLT these algorithms are optimized for quickness, whereas the final reconstruction is optimized for precision and accuracy. By having the ability to spend different amounts of computing time on different L1 triggered events, the complexity of the event selection offered by the HLT is heightened.

The two tiered triggering system creates very low dead times while maintaining purity and

830 selectivity. The wide ~~gamete~~gamut of physics topics that are pursued by the CMS collaboration
831 are a testament to the effectiveness and versatility of the CMS two tiered triggering system. In
832 Chapter 4, the development of L1 triggers and HLT paths for selecting UPC events are discussed.

Chapter 4

UPC trigger development for CMS

In the 2011 LHC PbPb run the ~~rate~~cross section of UPC J/ψ production was about 10^5 times smaller than that of ordinary nucleus-nucleus collisions. To record these events dedicated UPC triggers were needed. Unlike most heavy-ion triggers, UPC triggers are optimized for low- p_T and low multiplicity events. For this reason trigger development specific to UPC events was required to carry out the analysis in this thesis.

The increase in collision rate of the LHC PbPb beams from 2010 to 2011 was nearly a factor of 15. To accommodate this increase in rate, the 2011 trigger scheme needed to be more selective than in 2010 where CMS could take any event which appeared to have a collision. The available bandwidth was allocated equally amongst the various heavy ion analysis groups to pursue as wide a physics program as possible. From this consideration, bandwidth limits were placed on the trigger rates for each analysis group's trigger package.

4.1 L1 trigger

The UPC L1 triggers were designed to study UPC J/ψ production via the dimuon and dielectron channels (see Section 3.7). To achieve this, the loosest muon and electron triggers were combined with a trigger on energy in the ZDCs and no activity in BSCs (BSC veto). Additional triggers were commissioned in case radiation damage during the run reduced the sensitivity of the BSCs. This

required no activity in HF (HF veto). These triggers are summarized in Table 4.1. The ECAL2 and ECAL5 triggers in Table 4.1 indicate a 2 and 5 GeV threshold on E_T measured in the ECAL. The MuonOpen trigger indicates that the trigger only requires a muon candidate in one of the three muon sub-systems and that there is no momentum threshold. ZDC in the trigger names indicate energy constant with at least one neutron. The sign on the ZDC label indicates which of the two ZDCs is required.

L1 trigger name	Rate (Hz)	Prescale	Id	Type
MuonOpen and (ZDC ⁺ or ZDC ⁻) and BSC veto	2.1	1	1	Physics
ECAL2 and (ZDC ⁺ or ZDC ⁻) and BSC veto	1.8	2	2	
ECAL5 and (ZDC ⁺ or ZDC ⁻) and BSC veto	0.3	1	3	
(ZDC ⁺ or ZDC ⁻)	35	1500	4	Monitor
MuonOpen and (ZDC ⁺ or ZDC ⁻) and HF veto	0	off	5	Backup
ECAL2 and (ZDC ⁺ or ZDC ⁻) and HF veto	0	off	6	
ECAL5 and (ZDC ⁺ or ZDC ⁻) and HF veto	0	off	7	

Table 4.1: List of 2011 L1 seeds.

The cumulative L1 trigger rate for all the UPC L1 trigger seeds was required to be no greater than 200 Hz. This requirement came from the need to keep the tracker read-out rate low. The trackers baseline voltage can fluctuate due to the high tracker hit multiplicities in PbPb collisions. In order to monitor the zero suppression of the tracker, the zero suppression algorithm was executed using the HLT computing farm rather than in the tracker firmware.

The (ZDC⁺ or ZDC⁻) was used to estimate the efficiency of the UPC electron and muon triggers. The factor that the rate is reduced by is called the prescale. A prescale of 2 for example means that half the triggers were accepted. If the prescale is set to 1, then whole trigger rate is accepted. The prescale factors for the triggers were set to balance the competing objectives of rate reduction and increasing the overlap between the monitoring and signal triggers.

4.2 HLT trigger

An event must pass the selection criteria of an HLT path in order to be recorded. As opposed to the L1 trigger, which has access only to information from calorimeters and muon chambers, the HLT has access to all of the CMS sub-detectors including the tracker. Reconstruction of a track in the pixel detector is used by the UPC trigger paths. The use of the pixel detector only, as opposed to using the whole tracker including the silicon strip detector, allows for quick track reconstruction saving computing cycles. The UPC triggers were required to have at least one reconstructed pixel track in order to reject backgrounds where no particles were reconstructed by the tracker. For the muon trigger in Table 4.2 the rate was reduced by nearly a factor of 4 compared to its L1 seed rate in Table 4.1.

HLT trigger	Rate (Hz)	L1 prescale	HLT prescale	L1 seed	Type
L1UPCMuon and Pixel Track	0.52	1	1	1	Physics
L1UPCECAL2 and Pixel Track	1.65	2	1	2	
L1UPCECAL5 and Pixel Track	0.26	1	1	3	
L1ZDCOr	3.6	1500	11	4	Monitor
L1ZDCOr and Pixel Track	2.8	1500	1	4	
L1UPCMuonHFVeto and Pixel Track	0	off	off	5	Backup
L1UPCECAL2HFVeto and Pixel Track	0	off	off	6	
L1UPCECAL5HFVeto and Pixel Track	0	off	off	7	

Table 4.2: List of 2011 HLT trigger.

The total HLT output for the UPC trigger package was limited to 20 Hz. The limiting factor for the HLT rate was the amount of disk space given to this analysis. To meet the bandwidth requirements and collect a significant sample of data for estimating efficiencies, the prescales were balanced with the goal of achieving at least 5% statistical precision on the efficiency measurements. As an example of the balancing of the prescales, the HLT ZDC trigger that did not require a pixel track was given a additional prescale factor of 11 on the HLT. The ZDC path that also required a pixel track on the HLT, which used the same L1 seed, was only prescaled at the L1. The prescale of 11 was set to ensure that at least 1000 of the pixel track ZDC triggers overlapped with the ZDC L1 only triggers so that efficiency of the pixel track requirement in the trigger could be estimated

886 from the tracks lost.

887 **4.3 Studies of 2011 PbPb data**

888 The UPC triggers for the 2011 PbPb run make several studies possible. Three such studies are
889 discussed below: $\gamma\gamma \rightarrow e^+e^-$, UPC interactions in peripheral nuclear collisions, and forward UPC
890 J/ψ using HF.

891 **4.3.1 High mass $\gamma\gamma \rightarrow e^+e^-$ in PbPb 2011**

892 This measurement would make use of the electron triggers and combine the current dimuon data
893 with dielectron data from the ECAL triggers. Because of the smaller mass of the electron, di-
894 electron production is slightly favored compared to dimuon production. STARlight predicts that
895 dielectron cross section is a factor of 2.5 higher in Xn break-up mode than for the dimuons channel
896 when looking at masses above 4 GeV. The ECAL is positioned just beyond the tracker, whereas
897 the muon system is the outermost sub-detector. Electrons would therefore be less effected by the
898 material budget, which is the main factor in reducing yields of reconstructed dimuon candidates.

899 The contribution from higher order diagrams can be explored by studying photoproduction
900 of dilepton pairs. Because the Pb nucleus has a charge 82 times higher than that of the proton,
901 the electromagnetic coupling is stronger. Thus higher order terms in the perturbative expansion
902 could potentially be more important for the Pb nucleus compared to the proton. Recent results
903 by ALICE favor very small contributions for higher order terms [74]. In addition, this analysis
904 provides a useful cross check to the UPC quarkonia analyses such as J/ψ by verifying the cross
905 section normalization.

906 **4.3.2 UPC hadronic overlap**

907 In the model calculations for UPC quarkonia photoproduction all hadronic interactions are rejected.
908 However, inclusive p_T spectra of J/ψ measured by ALICE in peripheral PbPb collisions show a

low momentum peak consistent with coherent photoproduction [84]. The ALICE spectra provide




Figure 4.1: Coherent excess in inclusive J/ψ p_T spectrum.

hints that UPC processes might also be present in peripheral nucleus-nucleus collisions at the LHC.

To study the overlap between photoproduction and hadronic production of quarkonium events, the inelastic sample and the UPC sample could both be used. The looseness of the rejection criteria to reject hadronic interactions, which uses the BSC detectors, leaves a significant overlap with peripheral hadronic collisions. The inclusive quarkonia sample from typical hadronic collisions can also be utilized. Coherent quarkonia photoproduction has a distinctive low p_T structure that can be used to identify photoproduced candidates in a sample that contains photoproduction combined with hadronic interactions. This measurement would open up the door to exploring the boundary between photoproduction and hadronic production.

4.3.3 UPC J/ψ with electrons in HF

As higher rapidities are explored both lower and higher momentum partons of the nucleus are probed. Because these two contributions to the UPC photoproduction cross section can be separated using neutron tagging in incoherent events, exploring higher dilepton rapidities becomes

attractive. HF extends to 5 in η , which is 2.6 units beyond the edge of the tracker. By combining hits in HF with tracks in the tracker, J/ψ from higher rapidities can be measured. When combined with neutron tagging of incoherently produced quarkonia, the current study can be extended to probe lower- x nuclear partons by identifying leptons in HF.

4.4 Trigger development for the LHC pPb Run

Specific UPC triggers were also developed for the pPb run in 2013. For this period of running a much higher total trigger rate was read out relative to 2011. The total rate allocated for UPC triggers at the L1 in 2013 was 5 kHz and 50 Hz at the HLT. This factor of 5 increase in HLT and factor of 25 in L1 bandwidth, allowed for a change in emphasis from the L1 to the HLT.

The basic strategy in 2013 was the same as in 2012, use the loosest available ECAL and muon L1 triggers to capture the lowest p_T electrons and muons possible and reject hadronic interactions. Because of the L1 bandwidth restrictions in 2011, both the ZDCs and the BCSs were used on the L1 to reduce rates. In 2013 only the muon and ECAL triggers were used on the L1 allowing for rejection of hadronic interactions through cuts on track multiplicity. In addition, a more sophisticated trigger using full dimuon reconstructed was developed to increase purity. The main advantage in this shift in strategy was a higher purity due to the increased sophistication of the reconstruction on the HLT. In addition, the cross section for photoproduction without regard to neutron emission is higher. Therefore, relaxing the requirement of activity in the ZDCs further increases measurable yields.

The HLT triggers in 2013 rejected hadronic interactions through counting tracks. For the five UPC trigger paths included in the HLT menu, three levels of reconstruction were done at the HLT.

- Pixel tracks were reconstructed from the inner pixel section of the silicon tracker alone, tracks were reconstructed using the full tracker using the strips as well, and full dimuon reconstruction was done using the tracker and muon detector.
- The least restrictive pixel track paths required at least one track reconstructed from the pixel

948 detector and less than 10 pixel tracks in the event.

- 949 • Full tracking paths were added on top of the pixel track paths and included an additional
950 requirement of one full track and less than 7 reconstructed tracks.
- 951 • The most restrictive path added to the pixel and full tracking paths and required reconstruc-
952 tion of dimuons with a mass between 2 and 12 GeV.

953 **4.4.1 J/ψ photoproduction in ultra-peripheral pPb collisions**

954 The CMS UPC triggers commissioned for the 2013 LHC pPb run will allow for further analysis of
955 J/ψ photoproduction. In ultra-peripheral pPb collisions, J/ψ photoproduction is dominated $\gamma - p$
956 interactions [85, 86]. The measurement would primarily probe the proton gluon densities. In
957 Eq. 2.13 the photon flux depends on the square of the number of protons in the parent nucleus, Z^2 .
958 However, the cross section of the target increases as the total number of nucleons to the two thirds
959 power, $A^{2/3}$. The much higher photon flux from the Pb-ion compensates for the decreased size of
960 the proton.

961 A pPb UPC J/ψ measurement will complement the measurements done at HERA [87, 88, 89],
962 and measurements done by ALICE [90]. CMS will contribute by adding additional kinematic
963 coverage and cover a unique range of γp center of mass energies, $W_{\gamma p}$. The difference in beam
964 energies and species at LHC versus HERA result in access to different $W_{\gamma p}$. ALICE and CMS
965 have different acceptance in J/ψ rapidity, which also translates to coverage of different $W_{\gamma p}$. In
966 addition, an excess in the UPC cross section compared to HERA measurements would indicate a
967 non-exclusive contribution to the pPb UPC J/ψ cross section. This measurement will enhance the
968 current understanding of the J/ψ photoproduction cross section as a function of the photon-proton
969 center of mass energy, $W_{\gamma p}$.

Chapter 5

Analysis

In this chapter the various parts of the analysis are explained. In Section 5.1, the simulations used to estimate the detector's ability to measure UPC processes are discussed. The selection of UPC events is detailed in Section 5.2. Extraction of the number of coherent J/ψ candidates is explained in Section 5.3. The determination of the detector's efficiency for measuring UPC events is explained in Section 5.4.

5.1 Physics generators and Monte Carlo simulations

Every physical measurement is the product of the underlying physics folded with the response of the detector used to do the measurement. In order to understand the underlying physical process, the detector's effect on the measurement must be understood and accounted for. As instruments become more and more complicated, the interplay among all of the many parts of the detector makes an analytic approach to the problem untenable. For this reason, the numerical technique of Monte Carlo (MC) simulation is often the most effective approach for describing detector effects.

MC simulations use random number generation to model the many statistical effects of particles interacting with different parts of the detector. First, particles are generated according to theoretical distributions. These particles are then propagated through a simulation of the detector. As the particles pass through the detector, random numbers are used to determine how these particles

interact with the materials of the detector based on the known properties of the material. In this way, the theoretical distributions are convolved with a realistic model of the detector's response. A more detailed picture of how the detector shapes the underlying distributions emerges with each successive event. The final goal of the MC simulation is to produce a set of events that accurately reproduce what would be measured if the theoretical input describes nature well.

5.1.1 STARlight and particle gun MC

In this thesis, two classes of generator input samples were used, STARlight [40, 41] and a particle gun. The STARlight samples correspond to the theoretical calculations described in Section 2.2, while the particle gun produces particles with a user defined transverse momentum distribution and rapidity distribution and isotropic decay to muon pairs in the J/ψ rest frame. For STARlight, three different physical processes were simulated: coherent J/ψ production, where the photon couples to the nucleus as a whole; incoherent J/ψ production; where the photon couples to a single nucleon within the nucleus, and photon-photon interactions, where the photons from the two nuclei interact with each other to produce a pair of oppositely charge muons. All three STARlight samples contain a μ^+ and μ^- in the final state.

Because STARlight is not integrated into the standard CMS software framework (CMSSW), a simulation software chain with 5 steps was developed. First, STARlight is run in the specified mode, and a single file is created for each physics process. In step 2, the STARlight output file is converted to the Les Houches Event (LHE) format [91], and the momentum of the parent J/ψ or the initial photon-photon pair is added to the record of each event. The event record produced by STARlight only contains the final state particles. To process the events in parallel, the STARlight files are subdivided in step 2, creating several LHE files from a single STARlight file. The LHE files are used as input to CMSSW.

Steps 3 to 5 take place within CMSSW. In step three the generated particles are propagated through the GEANT4 [92] detector simulation. This accounts for all the interactions with the detector and produces as output a format identical to the raw data that is recorded during data

1014 taking. Steps 4 and 5 are processed using the same software as in data taking. In step 4 the
1015 reconstruction software used during data taking is run on the output of the detector simulation.
1016 The output of the reconstruction is reduced to the information that is needed for the final analysis
1017 in the final step.

1018 The particle gun samples were created entirely within CMSSW. J/ψ mesons were created ac-
1019 cording to user defined p_T and rapidity distributions. The decay of J/ψ s to a $\mu^+\mu^-$ pair was
1020 simulated with a uniform decay distribution, corresponding to unpolarized J/ψ particles. As with
1021 the STARlight samples, these muons are propagated through the GEANT4 simulation [92] of the
1022 detector, and the raw data is produced. The remaining steps of running the reconstruction code and
1023 reducing the data to the final data format needed for the analysis are identical to the STARlight
1024 production.

1025 The momentum of the final state muons is the main driver of whether the candidate can be
1026 measured. One of the two daughter muons must have large enough momentum to fire the trigger,
1027 and both muons must have enough momentum to be reconstructed and tagged as a muon. There
1028 are at least 10 interaction lengths of material through which the muons must travel in order reach
1029 the muon chambers (see Fig. 3.11). This imposes an effective momentum threshold of about 7
1030 GeV in order for muons can fire the trigger.

1031 The p_T distribution and the polarization of the J/ψ s produced are the main factors controlling
1032 the momentum of the muon daughters, which vary for the different MC samples. The polarization
1033 effects how the momentum is shared between the daughters [93]. In the rest frame of the parent
1034 J/ψ , equal momentum is given to each daughter muon. However in the lab frame of the detector,
1035 the muon daughters which are emitted from transversely polarized J/ψ will tend to be emitted in
1036 the direction the J/ψ is traveling and will have unequal momenta. The daughter traveling in the
1037 direction of the J/ψ will have increased momentum, whereas the daughter traveling opposite to the
1038 J/ψ direction will have decreased momentum.

1039 In Fig. 5.1 the J/ψ s p_T from the STARlight generated coherent, and incoherent, and the dimuon
1040 p_T for photon-photon samples are compared. Both the coherent and the photon-photon samples are

concentrated at low p_T , and neither sample extends much beyond 0.15 GeV. The incoherent sample is peaked near 0.5 GeV and extends beyond 1 GeV. The two particle gun samples resemble the incoherent and coherent samples p_T distributions. The first sample has a Gaussian p_T distribution extending to approximately 0.15 GeV, whereas the second is flat in p_T up to 2 GeV.




Figure 5.1: Generator level rapidity (left) and p_T (right) distributions for the coherent (black), incoherent (red), and photon-photon process (green).

The particle gun samples are unpolarized, whereas the STARlight samples have transverse polarization. The cosine of the helicity angle, the angle between the J/ψ spin vector in the rest frame and the direction of the boost to the lab frame, are shown for the particle gun samples and the STARlight samples in Fig. 5.2. For the STARlight sample the helicity angle prefers to be either parallel or anti-parallel. However, the particle gun samples have no preferred direction.

Figures 5.3, 5.4, and 5.5 show the rapidity, ϕ , and p_T distributions of reconstructed coherent J/ψ from STARLight MC and data. All distributions are normalized to one. The mean of the STARLight rapidity distribution is slightly shift by ≈ 0.1 to higher rapidity. The ϕ , and p_T distributions from STARLight and data are both very similar.




Figure 5.2: The J/ψ polarization of the particle gun (red), coherent (blue), and incoherent samples are plotted as the cosine of the helicity angle.




Figure 5.3: Comparison of the of the dimuon rapidity distributions between coherent J/ψ MC sample and data.

5.2 Event selection

The unprecedented amounts of data produced by the LHC has made it possible to investigate novel physics processes like UPC J/ψ production. The data for this analysis were recorded during the 2011 LHC PbPb run. During this period, $150 \mu b^{-1}$ were recorded by the CMS detector,




Figure 5.4: Comparison of the of the dimuon ϕ distributions between coherent J/ψ MC sample and data.




Figure 5.5: Comparison of the of the dimuon p_T distributions between coherent J/ψ MC sample and data.

1058 corresponding to $> 10^9$ PbPb collisions. Of this, 143 ± 7 (syst) μb^{-1} of data were used in this
1059 analysis due to the ZDCs being temporally disconnected to test the Forward Shower Counters.

5.2.1 Data sets

The data were divided into three specially selected samples, Physics, Monitoring, and Zero bias, based on the triggers which recorded the events (see Table 5.1). By recording this hierarchy of samples, interesting events are selected with a much higher purity in the physics sample, while the zero bias and ZDC triggered samples allow for the investigation of the selection criteria. The purity, which is a measure of how many signal events relative to background events are in a sample, is obtained by using more selective triggers. Less selective triggers, those assigned to the monitoring and zero bias samples, were used to investigate to what extent signal events are lost because of the higher selectivity of physics triggers. These samples were recorded using subsets of the HLT triggers found in Table 4.2 of Chapter 4. The J/ψ events discussed in this thesis were obtained by analyzing the sample labeled in Table 5.1 as physics. A ZDC triggered monitoring sample was recorded for the sake of estimating efficiencies. Lastly, a zero bias sample was recorded for investigating the ZDC and the noise distributions of HF.

The physics sample containing the J/ψ signal was recorded by the muon trigger labeled “L1UPCMuon and Pixel Track” in Table 4.2. Because of the characteristically low momentum of UPC J/ψ as compared to J/ψ created by other physics processes, the loosest muon trigger was used. The noise trigger rate for the muon trigger alone was 50 Hz, but in coincidence with the BCS veto and the ZDC trigger the noise rate was below 2 Hz. By pairing the muon trigger with the ZDC on the L1 trigger, the noise contribution was reduced from the noise contribution from either of the two sub-detectors to the noise coincidence between the two sub-detectors. Contributions from hadronic interactions were reduced by the veto on the BSCs. This trigger was designed to balance reducing the rate with maximizing the efficiency, allowing for the data to be recorded without producing high rates that would have resulted in dead time for the detector.

In order to investigate the muon trigger and the other parts of the event selection, a monitoring sample was recorded by requiring energy consistent with at least one neutron in either of the ZDCs. Neutron production is a much more common process than the UPC J/ψ production. This process has cross sections on the order of 200 b compared to 10 mb predicted for J/ψ production. For this

Sample	Events	\mathcal{L}_{int}
Physics	346K	$143.3 \mu b^{-1}$
Monitor	1.1M	$31.6 mb^{-1}$
Zero Bias	8.8M	$580 b^{-1}$

Table 5.1: Integrated luminosities and number of events for the three samples used in this analysis.

reason, the rates of this trigger are much higher than the physics trigger, and only a small ~~sub-set~~ subset of these events are recorded. From this trigger the pixel track portion of the HLT trigger efficiency was estimated as well as the ZDC trigger efficiency, as will be described in Section 5.4.

In addition to the monitoring and physics sample, a zero bias sample was recorded to examine the ZDC neutron reconstruction and the HF noise distributions. The zero bias trigger fired every time both beams passed through CMS. Only 4 events out of every million triggered were recorded for this sample. This sample allowed for an unbiased measurement of the ZDC neutron threshold energies as discussed in Section ?? . Because the zero bias trigger only requires the presence of both LHC beams, the sample contains very few hadronic collisions. This allowed for a measurement of the electronic noise distribution of the HF detector, which is important to reducing contamination from hadronic interactions.

The integrated luminosity for each of the three samples is calculated by recording activity in HF [94]. The cross section for HF activity is measured from a van der Meer scan [95]. In this way, the amount of integrated luminosity for any running period is related to the activity in HF.

5.2.2 Event selection cuts

The analysis described in this thesis focuses on UPC J/ψ s decaying to muons. The trigger used for this analysis recorded 346841 events. A set of off-line cuts was applied to increase the relative contribution of UPC events to background processes. Two sets of event selection cuts were applied to reject background events. The first set rejects background from the beam. The second rejects events where hadronic collisions have occurred. Table 5.2 summarizes all the event selection cuts. Figure 5.6 shows a candidate event after applying all event selection cuts.

Cut type	Cut	Events
—	all triggered	346841
beam background rejection	good vertex requirement	340997
	beam halo muon rejection	302777
	cluster shape compatibility requirement	233590
hadronic interaction rejection	single-sided neutron requirement	149992
	two track requirement	32732
	HF signal rejection	5392
fake muon rejection	muon quality requirement	2047
kinematic cut	J/ψ mass requirement	696
	muon detectability cuts	567

Table 5.2: Effects of event selection cuts.




Figure 5.6: Event display of a UPC J/ψ candidate event.

To reject beam induced background the following cuts were applied:

- The reconstructed vertex must be within 2 cm in the transverse direction and 25 cm in the longitudinal direction. This cut ensures that reconstructed particles come from interactions between the two beams rather than event where one of the two beams interact with gas particles near the interaction point.
- Beam halo muons were rejected using the timing of the muon hits. The beam halo cut rejects events where muons surrounding the beam stream through the detector.

- Pixel cluster shape should be compatible with the vertex. This cut requires that energy deposited in the silicon tracker have a shape consistent with the primary vertex.

When applied after all other cuts these beam background cuts do not reject any UPC J/ψ candidates.

The second set of background rejection cuts were designed to reduce contamination from hadronic interactions.

- No more than 2 reconstructed tracks in the event.
- Maximum reconstructed hit energy in HF was required to be below the threshold for electronic noise. Nearly all hadronic interactions (about 98%) produce particles in the range $3 < |\eta| < 5$ covered by the HF detector. By requiring that the energy deposits in HF resemble noise, nearly all inelastic hadronic collisions are expected to be rejected.
- Energy in the ZDCs consistent with neutrons on only one side of the interaction point. In hadronic interactions both nuclei break-up. By requiring that ZDC only reconstruct neutrons on one side of the interaction point, hadronic interactions that produce neutrons on both sides were rejected.

Each of these cuts were designed to reject topologies produced by hadronic interactions. The effect of these cuts can be seen in Table 5.2 and are denoted hadronic interaction rejection.

To establish the HF noise thresholds, the noise distributions were measured in zero bias events. An offline selection of events with no reconstructed tracks was used to ensure that no collision had taken place. The HF noise threshold was defined as the cut that keeps 99% of the zero bias events. The noise distribution from this zero bias sample is compared to the physics sample and MC in Fig. 5.7.

The following standard muon quality cuts are applied:

- Tracker track matched with at least one muon segment (in any station) in both X and Y coordinates ($< 3 \sigma$).




Figure 5.7: Comparison of HF noise distributions in zero bias data, physics triggered data, and MC.

- Cut on number of tracker layers with hits > 5 .
- Number of pixel layers > 0 .
- The χ^2 per degrees of freedom of the track fit < 3 .
- Loose transverse and longitudinal impact parameter cuts, within 3 cm in the transverse direction and within 30 cm in the longitudinal direction with respect to the primary vertex.

These cuts are applied to reduce the number of fake muons and have been validated for other muon analyses [96].

5.3 Signal extraction

After all event selection cuts, the remaining events contain a combination of coherent J/ψ , incoherent J/ψ , and dimuons from the photon-photon process. Each process must be separated from the final mix. To achieve this, the invariant mass and p_T distributions are used to distinguish between the three processes. The photon-photon process is extended in invariant mass whereas the J/ψ is peaked strongly near 3.1 GeV. The dimuon transverse momentum distribution of the photon-photon and the coherent process have similar distributions, both are sharply peaked below 0.1 GeV,

whereas the incoherent process is more broadly distributed across an interval extending to nearly 1 GeV. The mass distribution was fit to separate the photon-photon process from the J/ψ process. The p_T distribution was used to separate the incoherent process from the photon-photon process, and the coherent process. In this way, a separate yield was extracted for all three processes.

The invariant mass distribution for opposite sign dimuons is shown in Fig. 5.8. A J/ψ signal is clearly visible together with tails at higher and lower mass due to the photon-photon process. A fit to the invariant mass distribution was performed using a Gaussian to account for the J/ψ signal and a first-order polynomial function for the photon-photon process. The extracted number of J/ψ candidates from this fit includes all J/ψ s in the mass window that passed the analysis cuts, i.e. both coherent and incoherent process contribute to yield from the mass fit. The p_T distribution is needed to separate the two different contributions to the J/ψ peak.




Figure 5.8: Mass fit to J/ψ using Gaussian for the signal and a first-order polynomial for the photon-photon continuum.

Figure 5.9 shows the p_T spectrum of the events plotted in Fig. 5.8. There is a clear coherent peak at $p_T = 60$ MeV followed by a broad distribution that with a mean near $p_T = 450$ MeV.

1167 To extract the contribution of coherent, incoherent and gamma-gamma processes in the data the
 1168 spectrum in Fig. 5.9 was fit to the sum of three MC templates corresponding to the final output
 1169 of the MC simulations for these three processes. The clear overlap of the coherent and photon-
 1170 photon process, and the clear separation of these two lower p_T processes from the incoherent
 1171 process is apparent. The shape of the p_T distribution for the coherent, incoherent, and photon-
 1172 photon process are taken from the final output of MC after applying all analysis cuts. In Fig.5.9,
 1173 the yield parameters that were fit were left unconstrained for all three process.




Figure 5.9: Fit to MC p_T templates.

1174 The shape of the photon-photon and coherent J/ψ process are very similar in transverse mo-
 1175 mentum. Accordingly, the contribution from the photon-photon process and the coherent process
 1176 are difficult to separate from the p_T distribution. The confidence contours in Fig. 5.10 from the
 1177 template fit in Fig. 5.9 demonstrate the strong anti-correlation between the coherent yield param-
 1178 eter, nCo , and the yield parameter for the photon-photon process, $nGamma$. Because of the anti-
 1179 correlation, the statistical uncertainty on nCo and $nGamma$ from the fit are larger than \sqrt{nCo} and
 1180 \sqrt{nGamma} expected from Poisson statistics. The information from the invariant mass and p_T dis-

1181 tributions were combined to break this correlation. Through this combination, the contribution to
 1182 the final yield from the three process was measured.




Figure 5.10: 68%, 95%, and 99% confidence contours from the p_T template fit.

1183 A simultaneous fit to the mass spectrum and p_T spectrum was preformed to utilize the mass
 1184 fits ability to distinguish the photon-photon process from the coherent and incoherent process all
 1185 while utilizing the p_T fits ability to separate the coherent and photon-photon processes from the
 1186 incoherent. Fig. 5.11 shows the result of the simultaneous fit. The simultaneous fit forces the
 1187 parameter $nGamma$ to both describe the photon-photon continuum present in the side bands of the
 1188 J/ψ mass peak as well the photon-photon contribution to the low- p_T part of the p_T spectrum. In
 1189 addition, the J/ψ yield from the mass fit is forced to equal the contribution from the incoherent
 1190 and coherent process in the fit to the p_T distribution. In this way, the correlation between the yield
 1191 parameters was broken, and the contribution from the three process were made independent of
 1192 each other.

1193 Fig. 5.12 shows the confidence contours for nCo and $nGamma$ from the simultaneous fit in
 1194 Fig. 5.11. The slope of the confidence contours in Fig. 5.12 is noticeably than in Fig. 5.10. The




Figure 5.11: Simultaneous fit to the mass and p_T spectra.

1195 contours for the simultaneous fit are also reduced compared to Fig. 5.10 with widths in nCo and
 1196 $nGamma$ similar to those expected from Poisson statistics. From the simultaneous fit, reasonable
 1197 statistical errors were obtained along with the yields for the three processes.




Figure 5.12: 68%, 95%, and 99% confidence contours from the simultaneous fit.

5.4 Efficiency determination

Each step of the triggering, event selection, and analysis has an associated efficiency that must be accounted for in the measurement of the J/ψ cross section. The ZDC trigger efficiency, the muon trigger efficiency, and the muon reconstruction efficiency are the two most significant contributors to the total efficiency measurement. The efficiency of the pixel track requirement, and the veto on activity in the BSCs from the trigger are also estimated but found to be consistent with fully efficient. The following section explains how each of these efficiencies was measured with a special emphasis on the ZDC trigger efficiency and the muon trigger and reconstruction efficiencies.

5.4.1 Muon efficiencies

The muon efficiencies were measured using a combination of MC and data based methods. The MC based measurement accounts for the detector acceptance and the efficiency of the muon quality cuts discussed in Section 5.2. The trigger efficiencies were measured in data using the tag and probe method [97], which is discussed below.

CMS has a limited acceptance for J/ψ s, particularly in the case of J/ψ s with low momentum like those produced in UPC events. To measure the acceptance of CMS for J/ψ s, reconstructed dimuon candidates were considered detectable if both reconstructed muon daughters fell into a detectability region in p_T and η . The muon detectability region was defined using the coherent J/ψ events obtained from STARlight. The efficiency for reconstructing single muons ϵ_{reco}^μ is defined by $\epsilon_{reco}^\mu = \frac{N_{reco}^\mu}{N_{gen}^\mu}$, where N_{reco}^μ is the number reconstructed muons obtained after the full CMS detector simulation that passed the standard muon quality cuts, and N_{gen}^μ is the number of generated muons from STARlight. Fig. 5.13 shows the efficiency for reconstructing single muons from coherent J/ψ events. To avoid the edges of the detectors acceptance, all reconstructed muons that fall into a $(p_T, |\eta|)$ bin that has an efficiency less than 20% were rejected. This condition defines the detectability region. The acceptance and reconstruction efficiency for reconstructing dimuons




Figure 5.13: Muon daughter detectability from coherent J/ψ

was calculated from MC using the following formula:

$$A \times \varepsilon = \frac{N_{det}(|y|)}{N_{gen}(|y|)}, \quad (5.1)$$

where N_{det} is the number of reconstructed dimuons where both daughters fall into the detectability region, and N_{gen} is the number of generated dimuons. From Eq. 5.1, the acceptance for J/ψ was calculated as a function of $|y|$, and p_T (see Fig. 5.14).

The “tag and probe method” is a data driven approach used to measure the trigger efficiency of the muon daughters from J/ψ decays. In this method there are three categories of daughter muons. *Tag muons* are high quality muons. *Passing probes* are reconstructed muons that match the muon trigger, while *failing probes* do not. Each dimuon will have one daughter classified as a tag and the other as a probe. From here three invariant mass histograms are studied. One histogram is created from all pairs. The second comes from pairs where the probe is a passing probe. The last histogram comes from pairs where the probe fails to fulfill the trigger, it is a failing probe. By matching the




Figure 5.14: Dimuon acceptance from coherent J/ψ .

tag to the trigger, the probe is unbiased by the trigger and the efficiency can be measured by fitting the three mass histograms.

Because the trigger efficiency depends on the p_T and $|\eta|$ of the muon, one set of three histograms for each $(p_T, |\eta|)$ bin of the probe is created. To extract the single muon trigger efficiency ϵ_{trig}^μ , each set of invariant mass histograms were simultaneously fit. The signal was fit using a Crystal Ball function [98], and the background was fit to an exponential. The Crystal Ball parameters were simultaneously fit to all three histograms. The exponential function was fit to the failing and passing probe histograms separately. Because the background shapes are in principle different for the two samples, the efficiency is driven by this difference.

Fig. 5.15 shows the fit of the three sets of pairs. This fit was done for each bin of the probes p_T and η . The efficiency from the fits in each bin are shown in Fig. 5.16.

The dimuon trigger efficiency ϵ_{trig}^{dimuon} was calculated from the single muon efficiencies using the following equation:

$$\epsilon_{trig}^{dimuon} = 1 - (1 - \epsilon_{trig}^{\mu_1})(1 - \epsilon_{trig}^{\mu_2}), \quad (5.2)$$

where $\epsilon_{trig}^{\mu_1}$ is the tag and probe efficiency of the first dimuon daughter, and $\epsilon_{trig}^{\mu_2}$ is the efficiency of the second muon daughter. In Eq. 5.2 the probability of at least one daughter firing the




Figure 5.15: Fits to tag and probe pairs in the J/ψ mass region for pairs with a probe $2 < |\eta| < 2.2$ and $1.55 < p_T < 1.8$ GeV.

trigger is calculated by subtracting one from the probability that neither daughter fires the trigger, thus giving the dimuon trigger efficiency.

The average dimuon trigger efficiency for each dimuon $|y|$ bin was calculated by averaging the efficiency of dimuon candidates in each bin. The dimuon trigger efficiency ranges from about 50% to 90%. As expected the J/ψ trigger efficiency increases with rapidity since the longitudinal momentum of the J/ψ is given by $p_Z = M_{J/\psi} \cdot \sinh(y)$. Thus J/ψ mesons at forward rapidity distribute more momentum to their daughter muons which therefore have a greater chance of punching through into the muon chamber. The average trigger efficiency was multiplied by the acceptance and reconstruction efficiency from the MC to produce a total factor for both efficiency and acceptance.

The total combined efficiency and acceptance factor for coherent J/ψ between $2.0 \leq |y| \leq 2.2$




Figure 5.16: Muon trigger efficiencies in p_T and η bins from the tag and probe method.




Figure 5.17: The trigger efficiency from tag and probe averaged over candidates in each $|y|$ bin.




Figure 5.18: The acceptance times averaged trigger efficiency from tag and probe.

was found to be $\approx 5\%$. The acceptance factor of roughly 7% from the MC was found to be the main contributor to the total efficiency. The interplay of the polarization of the J/ψ and the material in detector drives down the efficiency by creating an effective momentum threshold for detection (see Section 5.1). The reconstruction efficiency of the daughters range between 20%-60% for muons in the defined detectability range. The trigger efficiency for the detectable muons ranges from 30%-80% depending on p_T .

5.4.2 ZDC trigger efficiency

As discussed in Section ??, the trigger labeled “L1ZDCOr and Pixel Track” in Table 4.2 was used to measure the ZDC trigger efficiency. This trigger required either a ZDC^+ or ZDC^- trigger, together with at least one pixel track. The veto on the BSC minimum bias trigger, as in the physics triggers, was applied offline. The BSC veto excludes events where BSCs from both sides of the interaction point are above threshold. This trigger was used in order to collect the most inclusive possible sample without using the minimum bias triggers designed to collect hadronic interactions.

ZDC Side	N_{events}	N_{trig}	ϵ_{ZDC}
ZDC ⁻	73028	71706	0.9819 ± 0.0037
ZDC ⁺	76132	71859	0.9439 ± 0.0035

Table 5.3: ZDC trigger efficiencies for ZDC reconstruction method 1 and 2

This ZDC triggered sample suffers from a trigger bias. For example, a sample triggered by ZDC⁺ would always produce a ZDC⁺ trigger efficiency of one. To avoid this, a similar technique to tag and probe was used. Each event is either tagged as triggered by ZDC⁺ or triggered by the ZDC⁻. The ZDC⁺ trigger efficiency is measured from the ZDC⁻ tagged sample, and vice versa.

To estimate the efficiency, the number of events with energy in ZDC⁺ greater than the single neutron threshold, N_{events} , was measured. From this set of events, the number of events that also fire the ZDC⁺, N_{trig} , was measured. The ratio between the number of single neutron events that fired the trigger and all single neutron events was taken as the estimate of trigger efficiency. The same procedure was applied for each side of the ZDC. The trigger efficiency was found to be 98% for ZDC⁻ and 94% for ZDC⁺.

Chapter 6

Results and summary

In the previous two chapters the analysis steps to measure the coherent J/ψ photoproduction were given. In this chapter, the measured cross section for this process is given in Section 6.1. Comparisons to the theoretical models are discussed. In addition, the rapidity correlation between UPC J/ψ and forward neutrons are presented and compared to model calculations in Section 6.2. Finally, a summary of the complete thesis is given.

6.1 Coherent J/ψ cross section

The coherent J/ψ cross section is calculated using the following formula:

$$\frac{d\sigma_{coh}^{J/\psi}}{dy}(Xn0n) = \frac{N_{coh}^{J/\psi}}{\mathcal{L} \cdot (2 \cdot \Delta y) \cdot BR}, \quad (6.1)$$

where $N_{coh}^{J/\psi}$ is the corrected number of coherent J/ψ candidates, \mathcal{L} is the integrated luminosity used for this analysis, Δy is the width of the rapidity interval, and BR is the branching ratio for J/ψ to $\mu^+\mu^-$. To obtain $N_{coh}^{J/\psi}$ the following formula was used:

$$N_{coh}^{J/\psi}(Xn0n) = \frac{N_{yield}^{J/\psi}}{(A \times \epsilon) \cdot \epsilon_{trig}^{ZDC} \cdot \epsilon_{trig}^{J/\psi}}, \quad (6.2)$$

$N_{yield}^{J/\psi}$	65 ± 12	$N_{coh}^{J/\psi}$	1213 ± 240
$A \times \varepsilon$	0.07	\mathcal{L}	143
ε_{trig}^{ZDC}	0.96	BR	0.0593
$\varepsilon_{trig}^{J/\psi}$	0.79 ± 0.05	Δy	0.2
$N_{coh}^{J/\psi}(Xn0n)$	1213 ± 240	$\frac{d\sigma_{coh}^{J/\psi}}{dy}(Xn0n)$	356 ± 71 (stat) $^{+43}_{-50}$ (syst) μb

Table 6.1: The measured quantities used to calculate the corrected number of coherent J/ψ , $N_{coh}^{J/\psi}(Xn0n)$ (right), and the quantities used to calculate the cross section for UPC J/ψ photo-production, $\frac{d\sigma_{coh}^{J/\psi}}{dy}(Xn0n)$ (left).

where $N_{yield}^{J/\psi}$ is the J/ψ yield with a $p_T < 0.15$ GeV, $A \times \varepsilon$ is the combined acceptance and efficiency, ε_{trig}^{ZDC} is the ZDC triggering efficiency for detecting at least one neutron emitted in the forward region, and $\varepsilon_{trig}^{J/\psi}$ is the J/ψ trigger efficiency measured by the “tag and probe” method. Each of these quantities can be found in Table 6.1.

The coherent J/ψ cross section $\frac{d\sigma_{coh}^{J/\psi}}{dy}(Xn0n)$ requires single-sided neutron emission, Xn0n. Previous measurements carried by the ALICE Collaboration measured the cross section without any neutron emission requirement. In order to compare to their results, the cross section is scaled by the factor 5.06 to account for the increase in cross section. This value was obtained from STARlight. The resulting cross section is $1.80 \pm 0.37(\text{stat})^{+0.22}_{-0.25}$ (syst) mb.

Figure 6.1 shows the measured cross section compared to the ALICE data points, as well as STARlight, AB and LTA models (see Chapter 2). The measured cross section is in agreement with previous measurements, and favors the AB-EPS09 model. Given the measured cross section is significantly below STARlight, nuclear gluon shadowing plays an important role for $x \sim 10^{-2}$.

6.2 UPC J/ψ -neutron rapidity correlation

In addition to the production cross section, neutron dependent J/ψ photoproduction can be studied in order to better understand the separation between coherent and incoherent photoproduction. Because of the inclusion of the ZDC in the trigger, the data for this thesis is well suited for studying the correlations between the J/ψ and neutron rapidities. The difference between neutron emission




Figure 6.1: Coherent J/ψ photoproduction cross section measured by ALICE and CMS including feed-down and rescaled for nuclear break-up.

in coherent and incoherent production was studied in the past [58], and recent studies have explored what is possible within the acceptance of CMS [59]. In this section, the ratio between J/ψ s produced with neutrons in the same side of the detector and opposite side of the detector will be shown and compared with theory.

Figure 6.2 shows the rapidity distribution of UPC J/ψ that are accompanied by neutron detection on only one side of the interaction point. The candidates are divided by p_T and the side of the interaction point the neutron was detected in. Low- p_T corresponds to the $p_T < 0.15$ GeV, and high p_T corresponds to $0.15 \text{ GeV} < p_T < 1 \text{ GeV}$. The filled black circles correspond to candidates that are produced in conjunction with a neutron in ZDC^- , whereas the open squares indicate a neutron in ZDC^+ . In Fig. 6.2, there is a clear preference for the neutron and J/ψ to be detected in the same side of the detector for the high- p_T candidates, however the low- p_T candidates do not show a strong preference.

The ratio between candidates with neutrons on the opposite side of the detector and candidates with neutrons on the same side of the detector, $R_{opp/same}$, is measured. $R_{opp/same}$ is shown as a




Figure 6.2: Rapidity distribution of J/ψ in the case of the events having the neutron in negative and positive rapidity for the $p_T < 0.15$ GeV J/ψ (left), $0.15 < p_T < 1.0$ GeV J/ψ (right) and dimuons from $\gamma\gamma$ sample (bottom).

function of the candidate p_T and compared with theory from [59] in Figure 6.3. As the p_T of the candidates increases beyond 0.3 GeV in Fig. 6.3, a transition from J/ψ that are uncorrelated with neutron production to J/ψ candidates with tend to be created in the same direction as the detected neutron.

The transition shown in Fig. 6.3 can be explained by the different neutron production processes that occur in coherent versus incoherent photoproduction [58]. In coherent photoproduction, the photon couples elastically to the nucleus as a whole. In this case the nucleus will typically stay intact. Neutron emission then occurs though exchange of an additional photon. The target struck by this additional photon need not be the same nucleus that was the target of the photoproduction process. For this reason, the neutron and J/ψ in coherent production are not expected to be correlated. Neutron emission in incoherent production is due to the excitation of the nucleus that occurs when the photon couples to an individual nucleon within the nucleus. The excited nucleus then decays by neutron emission. In this case, the target nucleus for both the J/ψ and the neutron are the same, and ~~there for~~ therefor correlated.

The correlation demonstrated in Fig. 6.3 can potentially offer a means of separating the two




Figure 6.3: Ratio between the transverse momentum distribution of the J/ψ when J/ψ and neutron have the opposite direction and the transverse momentum distribution of the J/ψ when J/ψ and neutron have the same direction.

terms of Eq.2.28, which probe different values of x . In incoherent events, because the neutron indicates the direction in which the target was traveling, the momentum of the photon can be deduced. If the photon carries higher longitudinal momentum than the gluons with which it interacts, the photoproduced J/ψ will be pushed by the photon in the direction opposite the emitted neutron. This would correspond to the low- x contribution to Eq.2.28. When the gluons within the nucleus are of higher longitudinal momentum, the J/ψ will travel in the direction of the target and the emitted neutron. This gives the high- x contribution.

The measurement of $R_{opp/same}$ shown in Fig. 6.3 agrees qualitatively with the predictions of [59]. This indicates that neutron tagging can both aid in separating the coherent from the incoherent process, and potentially separate the low- x and high- x contribution to UPC photoproduction cross sections at high rapidity values. However, this qualitative agreement requires additional theoretical

and experimental work for quantitative confirmation.

6.3 Summary

As physicists' understanding of the QGP has deepened over the past 30 years of doing experimental heavy ion physics, the questions surrounding the QGP have shifted from the confirmation of creation of a deconfined state to understanding the properties of the state that is created. It appears that the QGP is a hot nearly viscosity free fluid of strongly coupled quarks and gluons. The control measurements from dAu collisions at RHIC, and pPb collisions at the LHC have shown signs of collective behavior such as flow, which made these results difficult to interpret. Additional knowledge of the initial state of the colliding nuclei is needed in order to fully understand the QGP signal seen in PbPb and AuAu collisions. UPC events can provide this needed knowledge. This thesis contributes to the understanding of the initial state through the measurement of the UPC J/ψ photoproduction cross section.

Ultra-peripheral collisions are a clean probe of the initial state. In UPC J/ψ photoproduction, the nuclei interact through the electromagnetic force precluding the possibility of creating a collective medium. The theoretical models of coherent UPC J/ψ photoproduction model these electromagnetic interactions by combining a semi-classical calculation of the photon flux with a variety of phenomenological and QCD based calculations of the nuclear gluon density. The Weisäcker-Williams approximation [49] is used to calculate the flux of photons that surround the colliding nuclei. The interaction of these photons with the nucleus is either calculated through a nuclear modification of the proton gluon density [39, 99], or by using the Glauber model approach that consists in modeling the nucleus as a collection of nucleons and scaling the nucleon photoproduction cross sections from e-p collisions [40]. Photoproduction cross sections from UPC events can determine at what energy scale the Glauber based method breaks down. For the gluon density based calculations, there is a wide discrepancy between the predictions, and photoproduction cross sections constrain which gluon density models are viable.

The analysis in this thesis consists of three major components, development of a trigger, estimation of efficiency, and measurement of signal events. The trigger development involved designing a trigger based on rate estimates from past data that ensured a sample that could be used for both measuring the signal and estimate the efficiency of the trigger, reconstruction, and event selection. The number of J/ψ candidates was measured by first applying a set of event selection cuts that rejected background events such as hadronic collisions and beam gas collisions, then fitting the remaining events to templates from simulation to separate the three remaining physics processes, the coherent, incoherent, and photon-photon processes. The efficiencies for each part of the trigger were measured from data. The acceptance and reconstruction efficiency were estimated from MC. The cross section was calculated by combining the efficiency with the measured luminosity and number of coherent J/ψ . The statistical uncertainties were taken from the template fit. The systematic uncertainties were estimated by varying the method used on each component of the analysis.

The UPC J/ψ photoproduction cross section in the $Xn0n$ break-up mode, $\frac{d\sigma_{co}^{J/\psi}}{dy}$ ($Xn0n$), was found to be 356 ± 71 (stat) $^{+43}_{-50}$ (syst) μb . When rescaled by a factor of 5.06 to account for the difference of break-up mode between the measurement in this thesis and the ALICE result in [73, 74], the result of 1.80 ± 0.37 (stat) $^{+0.22}_{-0.25}$ (syst) mb was found to be consistent with the predictions in [39]. The calculation in [39] is also favored by the ALICE measurements. Of the gluon distributions used in [39], a gluon distribution with moderately strong gluon shadowing, EPS09 [57], is consistent with both the results from this thesis and the previous ALICE results. This indicates that at the scale of the mass of the J/ψ the nucleus gluon density is significantly suppressed compared to the gluon densities of a nucleon. At this scale the nucleus can not be represented as a collection of nucleons as in the Glauber like model described in [40]. In addition, the J/ψ -neutron rapidity correlation shown in Fig. 6.3 by the $R_{opp/same}$ ratio demonstrates the potential to separate the low- x and high- x contributions to the photoproduction cross section.

The measurements in this thesis confirm the ability to increase the knowledge of the initial state through the exploration of UPC events. This confirmation opens the door to additional measure-

1404 ments in this growing field of UPC research. A whole host of measurements will be possible with
1405 the data already recorded and with the data that will be recorded in the coming years by CMS and
1406 the other LHC experiments. UPC photoproduction measurements probing the initial state of the
1407 nucleus will play a crucial role in the heavy-ion physics community as plans for building a new
1408 electron-ion collider come into focus.

Bibliography

- [1] E. Witten, “Cosmic Separation of Phases,” *Phys.Rev.* **D30** (1984) 272–285.
- [2] R. Vogt, “Quark-Gluon Plasma Signatures,”.
- [3] L. Evans and P. Bryant, “LHC Machine,” *JINST* **3** (2008) S08001.
- [4] M. Gyulassy and M. Plumer, “Jet Quenching in Dense Matter,” *Phys.Lett.* **B243** (1990) 432–438.
- [5] T. Matsui and H. Satz, “ J/ψ Suppression by Quark-Gluon Plasma Formation,” *Phys.Lett.* **B178** (1986) 416.
- [6] **PHENIX Collaboration** Collaboration, A. Adare *et al.*, “Cold Nuclear Matter Effects on J/ψ Yields as a Function of Rapidity and Nuclear Geometry in Deuteron-Gold Collisions at $\sqrt{s_{NN}} = 200$ GeV,” *Phys.Rev.Lett.* **107** (2011) 142301, arXiv:1010.1246 [nucl-ex].
- [7] A. Baltz, G. Baur, D. d’Enterria, L. Frankfurt, F. Gelis, *et al.*, “The Physics of Ultraperipheral Collisions at the LHC,” *Phys.Rept.* **458** (2008) 1–171, arXiv:0706.3356 [nucl-ex].
- [8] C. A. Bertulani, S. R. Klein, and J. Nystrand, “Physics of ultra-peripheral nuclear collisions,” *Ann.Rev.Nucl.Part.Sci.* **55** (2005) 271–310, arXiv:nucl-ex/0502005 [nucl-ex].
- [9] S. Margetis, K. Safarik, and O. Villalobos Baillie, “Strangeness production in heavy-ion collisions,” *Ann.Rev.Nucl.Part.Sci.* **50** (2000) 299–342.

- [10] F. Karsch, E. Laermann, and A. Peikert, “Quark mass and flavor dependence of the QCD phase transition,” *Nucl.Phys.* **B605** (2001) 579–599, arXiv:hep-lat/0012023 [hep-lat].
- [11] S. Hands, “The Phase diagram of QCD,” *Contemp.Phys.* **42** (2001) 209–225, arXiv:physics/0105022 [physics.ed-ph].
- [12] “The Frontiers of Nuclear Science, A Long Range Plan,” arXiv:0809.3137 [nucl-ex].
- [13] **ALICE Collaboration** Collaboration, K. Aamodt *et al.*, “The ALICE experiment at the CERN LHC,” *JINST* **3** (2008) S08002.
- [14] **ATLAS Collaboration** Collaboration, G. Aad *et al.*, “The ATLAS Experiment at the CERN Large Hadron Collider,” *JINST* **3** (2008) S08003.
- [15] S. Chatrchyan *et al.*, “The cms experiment at the cern lhc,” *Journal of Instrumentation* **3** no. 08, (2008) S08004. <http://stacks.iop.org/1748-0221/3/i=08/a=S08004>.
- [16] **LHCb Collaboration** Collaboration, J. Alves, A. Augusto *et al.*, “The LHCb Detector at the LHC,” *JINST* **3** (2008) S08005.
- [17] **CMS Collaboration** Collaboration, S. Chatrchyan *et al.*, “Multiplicity and transverse momentum dependence of two- and four-particle correlations in pPb and PbPb collisions,” *Phys.Lett.* **B724** (2013) 213–240, arXiv:1305.0609 [nucl-ex].
- [18] **ALICE Collaboration** Collaboration, B. Abelev *et al.*, “Long-range angular correlations on the near and away side in p -Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV,” *Phys.Lett.* **B719** (2013) 29–41, arXiv:1212.2001 [nucl-ex].
- [19] G. Roland, K. Safarik, and P. Steinberg, “Heavy-ion collisions at the LHC,” *Prog.Part.Nucl.Phys.* **77** (2014) 70–127.
- [20] **CMS Collaboration** Collaboration, S. Chatrchyan *et al.*, “Measurement of the pseudorapidity and centrality dependence of the transverse energy density in pb-pb

collisions at $\sqrt{s_{NN}}=2.76$ tev,” *Phys. Rev. Lett.* **109** (Oct, 2012) 152303.

<http://link.aps.org/doi/10.1103/PhysRevLett.109.152303>.

- [21] **PHENIX Collaboration** Collaboration, S. S. Adler *et al.*, “Systematic studies of the centrality and $\sqrt{s_{NN}}$ dependence of the $dE_T/d\eta$ and $dn_{ch}/d\eta$ in heavy ion collisions at midrapidity,” *Phys. Rev. C* **71** (Mar, 2005) 034908.

<http://link.aps.org/doi/10.1103/PhysRevC.71.034908>.

- [22] J. D. Bjorken, “Highly relativistic nucleus-nucleus collisions: The central rapidity region,” *Phys. Rev. D* **27** (Jan, 1983) 140–151.

<http://link.aps.org/doi/10.1103/PhysRevD.27.140>.

- [23] **PHENIX Collaboration** Collaboration, A. Adare *et al.*, “Enhanced production of direct photons in Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV and implications for the initial temperature,” *Phys. Rev. Lett.* **104** (Mar, 2010) 132301.

<http://link.aps.org/doi/10.1103/PhysRevLett.104.132301>.

- [24] M. Wilde, “Measurement of direct photons in pp and pb–pb collisions with {ALICE},” *Nuclear Physics A* **904–905** no. 0, (2013) 573c – 576c.

<http://www.sciencedirect.com/science/article/pii/S0375947413001954>. The Quark Matter 2012 Proceedings of the {XXIII} International Conference on Ultrarelativistic Nucleus-Nucleus Collisions.

- [25] **STAR Collaboration** Collaboration, K. Ackermann *et al.*, “Elliptic flow in Au + Au collisions at $(S(NN))^{1/2} = 130$ GeV,” *Phys.Rev.Lett.* **86** (2001) 402–407, arXiv:nucl-ex/0009011 [nucl-ex].

- [26] **CMS Collaboration** Collaboration, S. Chatrchyan *et al.*, “Measurement of the elliptic anisotropy of charged particles produced in pbpb collisions at $\sqrt{s_{NN}} = 2.76$ tev,” *Phys. Rev. C* **87** (Jan, 2013) 014902. <http://link.aps.org/doi/10.1103/PhysRevC.87.014902>.

- [27] J.-Y. Ollitrault, “Anisotropy as a signature of transverse collective flow,” *Phys.Rev.* **D46** (1992) 229–245.
- [28] P. Sorensen, “Elliptic Flow: A Study of Space-Momentum Correlations In Relativistic Nuclear Collisions,” arXiv:0905.0174 [nucl-ex].
- [29] H. Song, S. A. Bass, U. Heinz, T. Hirano, and C. Shen, “200 *a* gev Au + Au collisions serve a nearly perfect quark-gluon liquid,” *Phys. Rev. Lett.* **106** (May, 2011) 192301.
<http://link.aps.org/doi/10.1103/PhysRevLett.106.192301>.
- [30] N. Borghini, P. M. Dinh, and J.-Y. Ollitrault, “Flow analysis from cumulants: A Practical guide,” arXiv:nucl-ex/0110016 [nucl-ex].
- [31] N. Borghini, R. Bhalerao, and J. Ollitrault, “Anisotropic flow from Lee-Yang zeroes: A Practical guide,” *J.Phys.* **G30** (2004) S1213–S1216, arXiv:nucl-th/0402053 [nucl-th].
- [32] R. Bhalerao, N. Borghini, and J. Ollitrault, “Genuine collective flow from Lee-Yang zeroes,” *Phys.Lett.* **B580** (2004) 157–162, arXiv:nucl-th/0307018 [nucl-th].
- [33] M. Ryskin, “Diffractive J / psi electroproduction in LLA QCD,” *Z.Phys.* **C57** (1993) 89–92.
- [34] S. J. Brodsky, L. Frankfurt, J. Gunion, A. H. Mueller, and M. Strikman, “Diffractive lepton production of vector mesons in QCD,” *Phys.Rev.* **D50** (1994) 3134–3144, arXiv:hep-ph/9402283 [hep-ph].
- [35] C. von Weizsacker, “Radiation emitted in collisions of very fast electrons,” *Z.Phys.* **88** (1934) 612–625.
- [36] E. Williams, “Nature of the high-energy particles of penetrating radiation and status of ionization and radiation formulae,” *Phys.Rev.* **45** (1934) 729–730.
- [37] M. Klasen, “Hard photoproduction at hera,” hep-ph/0702052.

- [38] A. Adeluyi and C. A. Bertulani, “Gluon distributions in nuclei probed at energies available at the CERN large hadron collider,” *Phys. Rev. C* **84** (Aug, 2011) 024916.
<http://link.aps.org/doi/10.1103/PhysRevC.84.024916>.
- [39] A. Adeluyi and T. Nguyen, “Coherent photoproduction of ψ and Υ mesons in ultraperipheral pPb and PbPb collisions at the CERN Large Hadron Collider at $\sqrt{s_{NN}} = 5$ TeV and $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys. Rev. C* **87** (Feb, 2013) 027901, arXiv:1302.4288 [nucl-th]. <http://link.aps.org/doi/10.1103/PhysRevC.87.027901>.
- [40] S. R. Klein and J. Nystrand, “Exclusive vector meson production in relativistic heavy ion collisions,” *Phys. Rev. C* **60** (Jun, 1999) 014903.
<http://link.aps.org/doi/10.1103/PhysRevC.60.014903>.
- [41] <http://starlight.hepforge.org>.
- [42] L. Frankfurt, V. Guzey, and M. Strikman, “Leading Twist Nuclear Shadowing Phenomena in Hard Processes with Nuclei,” *Phys.Rept.* **512** (2012) 255–393, arXiv:1106.2091 [hep-ph].
- [43] V. Rebyakova, M. Strikman, and M. Zhalov, “Coherent rho and J/psi photoproduction in ultraperipheral processes with electromagnetic dissociation of heavy ions at RHIC and LHC,” *Phys.Lett.* **B710** (2012) 647–653, arXiv:1109.0737 [hep-ph].
- [44] V. Goncalves and M. Machado, “Vector Meson Production in Coherent Hadronic Interactions: An update on predictions for RHIC and LHC,” *Phys.Rev.* **C84** (2011) 011902, arXiv:1106.3036 [hep-ph].
- [45] A. Cisek, W. Schafer, and A. Szczurek, “Exclusive coherent production of heavy vector mesons in nucleus-nucleus collisions at LHC,” *Phys.Rev.* **C86** (2012) 014905, arXiv:1204.5381 [hep-ph].

- [46] T. Lappi and H. Mantysaari, “ J/ψ production in ultraperipheral Pb+Pb and p +Pb collisions at energies available at the CERN Large Hadron Collider,” *Phys.Rev.* **C87** no. 3, (2013) 032201, arXiv:1301.4095 [hep-ph].
- [47] J. Hufner, B. Kopeliovich, and A. B. Zamolodchikov, “Inelastic J/ψ photoproduction off nuclei: Gluon enhancement or double color exchange?,” *Z.Phys.* **A357** (1997) 113–120, arXiv:nucl-th/9607033 [nucl-th].
- [48] J. D. Jackson, *Classical electrodynamics*. Wiley, New York, NY, 3rd ed. ed., 1999.
<http://cdsweb.cern.ch/record/490457>.
- [49] E. Fermi, “On the Theory of Collisions between Atoms and Electrically Charged Particles,” in *Electromagnetic Probes of Fundamental Physics*, W. Marciano and S. White, eds., pp. 243–252. Sept., 2003. hep-th/0205086.
- [50] C. A. Brau, *Modern Problems in Classical Electrodynamics*. Oxford, New York, NY, 1st ed. ed., 2004.
- [51] T. Bauer, R. Spital, D. Yennie, and F. Pipkin, “The Hadronic Properties of the Photon in High-Energy Interactions,” *Rev.Mod.Phys.* **50** (1978) 261.
- [52] M. L. Miller, K. Reygers, S. J. Sanders, and P. Steinberg, “Glauber modeling in high energy nuclear collisions,” *Ann.Rev.Nucl.Part.Sci.* **57** (2007) 205–243, arXiv:nucl-ex/0701025 [nucl-ex].
- [53] S. Donnachie, H. G. Dosch, O. Nachtmann, and P. Landshoff, “Pomeron physics and QCD,” *Camb.Monogr.Part.Phys.Nucl.Phys.Cosmol.* **19** (2002) 1–347.
- [54] J. Pumplin, D. Stump, J. Huston, H. Lai, P. M. Nadolsky, *et al.*, “New generation of parton distributions with uncertainties from global QCD analysis,” *JHEP* **0207** (2002) 012, arXiv:hep-ph/0201195 [hep-ph].

- [55] A. Martin, W. Stirling, R. Thorne, and G. Watt, “Parton distributions for the LHC,”
Eur.Phys.J. **C63** (2009) 189–285, arXiv:0901.0002 [hep-ph].
- [56] K. J. Eskola, H. Paukkunen, and C. A. Salgado, “An Improved global analysis of nuclear
parton distribution functions including RHIC data,” *JHEP* **0807** (2008) 102,
arXiv:0802.0139 [hep-ph].
- [57] K. Eskola, H. Paukkunen, and C. Salgado, “EPS09: A New Generation of NLO and LO
Nuclear Parton Distribution Functions,” *JHEP* **0904** (2009) 065, arXiv:0902.4154
[hep-ph].
- [58] M. Strikman, M. Tverskoy, and M. Zhalov, “Neutron tagging of quasielastic J/ψ
photoproduction off nucleus in ultraperipheral heavy ion collisions at RHIC energies,”
Phys.Lett. **B626** (2005) 72–79, arXiv:hep-ph/0505023 [hep-ph].
- [59] V. Guzey, M. Strikman, and M. Zhalov, “Disentangling coherent and incoherent J/ψ
photoproduction on nuclei by neutron tagging in ultraperipheral ion collisions at the LHC,”
arXiv:1312.6486 [hep-ph].
- [60] H. Emling, “Electromagnetic excitation of the two phonon giant dipole resonance,”
Prog.Part.Nucl.Phys. **33** (1994) 729–786.
- [61] E. de Passos, M. Hussein, L. Canto, and B. Carlson, “The Mean energy, strength and width
of triple giant dipole resonances,” *Phys.Rev.* **C65** (2002) 034326, arXiv:nucl-th/0108002
[nucl-th].
- [62] I. A. Pshenichnov, J. P. Bondorf, I. N. Mishustin, A. Ventura, and S. Masetti, “Mutual heavy
ion dissociation in peripheral collisions at ultrarelativistic energies,” *Phys. Rev. C* **64** (Jul,
2001) 024903. <http://link.aps.org/doi/10.1103/PhysRevC.64.024903>.
- [63] M. Goldhaber and E. Teller, “On nuclear dipole vibrations,” *Phys.Rev.* **74** (1948) 1046–1049.
- [64] H. Steinwedel and J. H. D. Jensen, “,” *Zeits. f. Naturforsch* **5a** (1950) 343.

- [65] P. Chomaz, “Collective excitations in nuclei,”
- [66] A. J. Baltz, C. Chasman, and S. N. White, “Correlated forward–backward dissociation and neutron spectra as a luminosity monitor in heavy-ion colliders,” *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment* **417** no. 1, (1998) 1 – 8.
<http://www.sciencedirect.com/science/article/pii/S0168900298005750>.
- [67] I. A. Pshenichnov, I. N. Mishustin, J. P. Bondorf, A. S. Botvina, and A. S. Iljinov, “Particle emission following coulomb excitation in ultrarelativistic heavy-ion collisions,” *Phys. Rev. C* **60** (Sep, 1999) 044901. <http://link.aps.org/doi/10.1103/PhysRevC.60.044901>.
- [68] M. Chiu, A. Denisov, E. Garcia, J. Katzy, A. Makeev, M. Murray, and S. White, “Measurement of mutual coulomb dissociation in $\sqrt{s_{NN}} = 130\text{GeV}$ $au + au$ collisions,” *Phys. Rev. Lett.* **89** (Jun, 2002) 012302.
<http://link.aps.org/doi/10.1103/PhysRevLett.89.012302>.
- [69] **STAR Collaboration** Collaboration, G. Agakishiev and others, “ ρ^0 photoproduction in auau collisions at $\sqrt{s_{NN}} = 62.4\text{ gev}$ measured with the star detector,” *Phys. Rev. C* **85** (Jan, 2012) 014910. <http://link.aps.org/doi/10.1103/PhysRevC.85.014910>.
- [70] **STAR Collaboration** Collaboration, C. Adler and others, “Coherent ρ^0 production in ultraperipheral heavy-ion collisions,” *Phys. Rev. Lett.* **89** (Dec, 2002) 272302.
<http://link.aps.org/doi/10.1103/PhysRevLett.89.272302>.
- [71] **STAR Collaboration** Collaboration, B. I. Abelev *et al.*, “ ρ^0 photoproduction in ultraperipheral relativistic heavy ion collisions at $\sqrt{s_{NN}} = 200\text{ gev}$,” *Phys. Rev. C* **77** (Mar, 2008) 034910. <http://link.aps.org/doi/10.1103/PhysRevC.77.034910>.
- [72] S. Afanasiev *et al.*, “Photoproduction of and of high mass in ultra-peripheral au-au collisions at $\sqrt{s_{NN}}$,” *% em Physics Letters B* **679** no. 4, (2009) 321 – 329.
<http://www.sciencedirect.com/science/article/pii/S0370269309008983>.

- [73] **ALICE Collaboration** Collaboration, B. Abelev *et al.*, “Coherent J/ψ photoproduction in ultra-peripheral Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Phys.Lett.* **B718** (2013) 1273–1283, arXiv:1209.3715 [nucl-ex].
- [74] **ALICE Collaboration** Collaboration, E. Abbas *et al.*, “Charmonium and e^+e^- pair photoproduction at mid-rapidity in ultra-peripheral Pb-Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV,” *Eur.Phys.J.* **C73** (2013) 2617, arXiv:1305.1467 [nucl-ex].
- [75] V. Guzey, E. Kryshen, M. Strikman, and M. Zhalov, “Evidence for nuclear gluon shadowing from the ALICE measurements of pbbp ultraperipheral exclusive production,” *Physics Letters B* **726** no. 1–3, (2013) 290 – 295.
<http://www.sciencedirect.com/science/article/pii/S0370269313006825>.
- [76] **CMS Collaboration** Collaboration, G. Bayatian *et al.*, “CMS technical design report, volume II: Physics performance,” *J.Phys.* **G34** (2007) 995–1579.
- [77] **CMS Collaboration** Collaboration, “Addendum to the CMS ECAL technical design report: Changes to the CMS ECAL electronics,”.
- [78] **CMS Collaboration** Collaboration, “Addendum to the CMS tracker TDR,”.
- [79] **CMS HCAL Collaboration** Collaboration, G. Baiaian *et al.*, “Design, Performance, and Calibration of CMS Hadron-Barrel Calorimeter Wedges,”.
- [80] V. Andreev, X. Aslanoglou, A. Azman, M. Bakirci, S. Basegmez, *et al.*, “Performance studies of a full-length prototype for the CASTOR forward calorimeter at the CMS experiment,” *Eur.Phys.J.* **C67** (2010) 601–615.
- [81] **CMS Collaboration** Collaboration, O. A. Grachov *et al.*, “Status of zero degree calorimeter for CMS experiment,” *AIP Conf.Proc.* **867** (2006) 258–265, arXiv:nucl-ex/0608052 [nucl-ex].

- [82] **CMS Collaboration** Collaboration, S. Dasu *et al.*, “CMS. The TriDAS project. Technical design report, vol. 1: The trigger systems,”.
- [83] **CMS Collaboration** Collaboration, P. Sphicas, “CMS: The TriDAS project. Technical design report, Vol. 2: Data acquisition and high-level trigger,”.
- [84] A. Lardeux, “ J/ψ production in Pb-Pb collisions at $\sqrt{s_{NN}}=2.76$ TeV in the ALICE experiment,” *J.Phys.Conf.Ser.* **446** (2013) 012042.
- [85] L. Frankfurt, M. Strikman, and M. Zhalov, “Elastic and large t rapidity gap vector meson production in ultraperipheral proton-ion collisions,” *Phys.Lett.* **B640** (2006) 162–169, arXiv:hep-ph/0605160 [hep-ph].
- [86] V. Guzey and M. Zhalov, “Rapidity and momentum transfer distributions of coherent J/ψ photoproduction in ultraperipheral pPb collisions at the LHC,” *JHEP* **1402** (2014) 046, arXiv:1307.6689 [hep-ph].
- [87] **ZEUS Collaboration** Collaboration, S. Chekanov *et al.*, “Exclusive photoproduction of J/ψ mesons at HERA,” *Eur.Phys.J.* **C24** (2002) 345–360, arXiv:hep-ex/0201043 [hep-ex].
- [88] **H1 Collaboration** Collaboration, A. Aktas *et al.*, “Elastic J/ψ production at HERA,” *Eur.Phys.J.* **C46** (2006) 585–603, arXiv:hep-ex/0510016 [hep-ex].
- [89] **H1 Collaboration** Collaboration, C. Alexa *et al.*, “Elastic and Proton-Dissociative Photoproduction of J/ψ Mesons at HERA,” *Eur.Phys.J.* **C73** (2013) 2466, arXiv:1304.5162 [hep-ex].
- [90] **ALICE Collaboration** Collaboration, B. B. Abelev *et al.*, “Exclusive J/ψ photoproduction off protons in ultra-peripheral p-Pb collisions at $\sqrt{s_{NN}}=5.02$ TeV,” arXiv:1406.7819 [nucl-ex].

- [91] J. Alwall, A. Ballestrero, P. Bartalini, S. Belov, E. Boos, *et al.*, “A Standard format for Les Houches event files,” *Comput.Phys.Commun.* **176** (2007) 300–304, arXiv:hep-ph/0609017 [hep-ph].
- [92] J. Allison *et al.*, “Geant4 developments and applications,” *Nuclear Science, IEEE Transactions on* **53** no. 1, (Feb, 2006) 270–278.
- [93] P. Faccioli, C. Lourenco, J. Seixas, and H. K. Wohri, “Towards the experimental clarification of quarkonium polarization,” *Eur.Phys.J.* **C69** (2010) 657–673, arXiv:1006.2738 [hep-ph].
- [94] **CMS Collaboration** Collaboration, S. Chatrchyan *et al.*, “Measurement of CMS Luminosity,”.
- [95] S. van der Meer, “Calibration of the Effective Beam Height in the ISR,”.
- [96] S. Chatrchyan *et al.*, “ J/ψ and $\psi(2s)$ production in pp collisions at $\sqrt{s} = 7$ tev,” *Journal of High Energy Physics* **2012** no. 2, (2012) .
<http://dx.doi.org/10.1007/JHEP02%282012%29011>.
- [97] T. C. collaboration, “Performance of cms muon reconstruction in pp collision events at $\sqrt{s} = 7$ tev,” *Journal of Instrumentation* **7** no. 10, (2012) P10002.
<http://stacks.iop.org/1748-0221/7/i=10/a=P10002>.
- [98] J. Gaiser, “Charmonium Spectroscopy From Radiative Decays of the J/ψ and ψ' ,”.
- [99] V. Rebyakova, M. Strikman, and M. Zhalov, “Coherent ρ and J/Ψ photoproduction in ultraperipheral processes with electromagnetic dissociation of heavy ions at RHIC and LHC,” *Phys. Letters B* **710** no. 4–5, (2012) 647 – 653.
<http://www.sciencedirect.com/science/article/pii/S0370269312003152>.