

EC325 Lecture Notes

EC325: Econometrics | Colby College

Prof. Caraher

February 3, 2026

Table of contents

Preface	3
What is Econometrics?	3
About These Notes	3
Structure of the Course	4
Roadmap Through the Midterm	4
Statistical Software	5
A Note on Learning Econometrics	5
About the Instructor	6
1 Introduction	7
1.1 What is this Class About?	7
1.2 Does Having Health Insurance <i>Cause</i> You to be Healthier?	8
1.3 Broad Steps of Econometric Analysis	10
1.3.1 Step 1: Write the Model	10
1.3.2 Step 2: Find and Explore the Data	11
1.3.3 Step 3: Estimate the Model	13
1.3.4 Step 4: Interpret the Estimate	14
1.4 Statistical Software	14
1.5 Summary and Conclusion	14
1.5.1 Key Takeaways:	15
1.6 Check Your Understanding	15
2 Good Data Practices	16
2.1 When Good Economists Use Bad Data	16
2.2 Know Your Data	17
2.2.1 Descriptive Statistics	17
2.2.2 Computing Descriptive Statistics in R	19
2.2.3 Examining Distributions	24
2.3 Visualizing Data	24
2.3.1 The Grammar of Graphics: ggplot2	25
2.3.2 Getting Started: The Palmer Penguins	25
2.3.3 Building a Plot: Step by Step	26
2.3.4 Adding More Information: Color and Shape	27
2.3.5 Adding a Line-of-Best Fit	28
2.3.6 Proper Labeling	29

2.3.7	Histograms for Distributions	30
2.3.8	A Simple Example with Our Wage Data	31
2.4	Replication: The Foundation of Credible Research	32
2.4.1	Why Replication Matters	33
2.4.2	What Makes Good Replication Materials?	33
2.4.3	Good Coding Practices	34
2.5	A Quick Example: State Crime Data	35
2.6	Summary	37
2.7	Check Your Understanding	38
3	Causality and Randomized Control Trials	39
3.1	The Problem with Simple Comparisons	39
3.2	The Multiverse of Possibilities	40
3.2.1	The Causal Effect of College	40
3.2.2	The Fundamental Problem	43
3.3	Randomized Control Trials: The Next Best Thing	43
3.4	The Law of Large Numbers	43
3.4.1	Seeing the LLN in Action: Rolling Dice	44
3.4.2	How the LLN Enables Causal Inference	47
3.5	How Randomization Eliminates Selection Bias: A Simulation	48
3.6	The Potential Outcomes Framework	55
3.6.1	Defining Potential Outcomes	55
3.6.2	The Fundamental Problem of Causal Inference	56
3.6.3	Selection Bias Formalized	56
3.6.4	How Randomization Eliminates Selection Bias	57
3.7	The Oregon Health Insurance Experiment	58
3.7.1	Context: Medicaid Expansion	58
3.7.2	The Lottery	58
3.7.3	Results	58
3.8	Summary and Conclusion	59
3.8.1	Key Takeaways	59
3.9	Check Your Understanding	60
4	Bivariate Regression and Ordinary Least Squares	61
4.1	Modeling Relationships Between Variables	61
4.1.1	Conditional Expectations	62
4.1.2	Example: Education and Wages	62
4.2	The Population Regression Model	62
4.2.1	The Bivariate Model	63
4.2.2	Connecting to Conditional Expectations	64
4.2.3	Interpreting the Coefficients	64
4.2.4	Example: Education and Wages	64

4.3	The Regression Line: Predicted Values and Residuals	65
4.3.1	Fitted Values and the Regression Line	65
4.3.2	Residuals: The Prediction Errors	65
4.3.3	A Numerical Example	65
4.3.4	The Key Insight: Minimize the Residuals	66
4.4	Deriving the OLS Estimator	66
4.4.1	The Minimization Problem	67
4.4.2	First-Order Conditions	67
4.4.3	Solving for the Intercept	68
4.4.4	Solving for the Slope	69
4.4.5	Confirming It's a Minimum	69
4.5	Computing OLS in R	70
4.5.1	Basic Syntax	70
4.5.2	Example: MPG and Horsepower	70
4.6	Goodness-of-Fit: R-Squared	71
4.7	The Gauss-Markov Assumptions	72
4.7.1	Assumption 1: Linear in Parameters	72
4.7.2	Assumption 2: Random Sampling	73
4.7.3	Assumption 3: Variation in the Independent Variable	73
4.7.4	Assumption 4: Zero Conditional Mean	74
4.7.5	Assumption 5: Homoskedasticity	75
4.7.6	Summary: Which Assumptions Matter for What?	75
4.8	Unbiasedness of OLS	76
4.8.1	Population Parameters vs. Sample Estimates	76
4.8.2	What Does Unbiasedness Mean?	77
4.8.3	Proof Sketch	77
4.8.4	Simulation: Unbiasedness in Action	78
4.9	Omitted Variable Bias	80
4.10	Unbiasedness and Causality: Why This All Matters	80
4.10.1	When Can We Interpret $\hat{\beta}_1$ Causally?	81
4.10.2	Correlation vs. Causation, Revisited	82
4.11	Variance of the OLS Estimator	82
4.12	Standard Errors	83
4.13	OLS is BLUE	83
4.14	Summary and Conclusion	84
4.14.1	Key Takeaways	84
4.15	Check Your Understanding	85
4.15.1	Practice: Interpreting Regression Output	85
5	Multivariate Regression	87
5.1	The Problem with Bivariate OLS	87
5.2	The Multivariate Solution	88
5.2.1	The General Multivariate Model	88

5.2.2	Interpreting Coefficients	89
5.2.3	The Zero Conditional Mean Assumption	89
5.3	Deriving the Multivariate OLS Estimator	89
5.3.1	The Minimization Problem	90
5.3.2	First-Order Conditions	90
5.4	Estimating Multivariate OLS in R	90
5.4.1	Example: MPG, Horsepower, and Weight	91
5.4.2	Interpreting the Results	92
5.5	Multivariate OLS as Matching	93
5.5.1	The Returns to Education Example	93
5.5.2	The Naive Comparison	94
5.5.3	Step 1: Compute Within-Industry Differences	94
5.5.4	Step 2: Compute Weights	96
5.5.5	Step 3: Compute Weighted Average	96
5.5.6	The Regression Equivalent	97
5.6	Goodness-of-Fit: Adjusted R-Squared	99
5.6.1	The Problem with R-Squared	99
5.6.2	The Adjusted R-Squared	99
5.7	Gauss-Markov Assumptions for Multivariate OLS	100
5.7.1	Assumption 1: Linear in Parameters	100
5.7.2	Assumption 2: Random Sampling	100
5.7.3	Assumption 3: No Perfect Multicollinearity	100
5.7.4	Assumption 4: Zero Conditional Mean	101
5.7.5	Assumption 5: Homoskedasticity	101
5.8	The Omitted Variable Bias Formula	101
5.8.1	Setup	101
5.8.2	The OVB Formula	102
5.8.3	Interpreting the Bias Formula	102
5.8.4	Example: Education, Wages, and Ability	103
5.8.5	Example: Policing, Crime, and Poverty	103
5.9	Variance in Multivariate OLS	103
5.9.1	What Increases Variance?	104
5.9.2	Standard Errors	104
5.10	The Bias-Variance Tradeoff	104
5.10.1	The Costs of Omitting a Relevant Variable	104
5.10.2	The Costs of Including an Irrelevant Variable	105
5.10.3	The Tradeoff	105
5.10.4	Practical Guidance	105
5.11	Simulation: Bias vs. Variance	106
5.12	Summary	111
5.12.1	Key Takeaways	111
5.13	Check Your Understanding	112

6	Statistical Inference	113
6.1	From Point Estimates to Uncertainty	113
6.2	The Sampling Distribution: A Simulation	114
6.3	Where Does Our Estimate Fall?	116
6.4	The Sampling Distribution of $\hat{\beta}_j$	117
6.4.1	The Normality Assumption	118
6.4.2	The Normal Sampling Distribution	119
6.5	Hypothesis Testing	121
6.5.1	The Null and Alternative Hypotheses	121
6.5.2	The t-Statistic	122
6.5.3	Properties of the t-Statistic	122
6.5.4	Visualizing the Hypothesis Test	122
6.5.5	Critical Values and Rejection Regions	125
6.5.6	Hypothesis Testing Example	127
6.5.7	Rule of Thumb	129
6.6	P-Values	130
6.6.1	Visualizing P-Values	130
6.6.2	Interpreting P-Values	132
6.6.3	A Note on Language	132
6.6.4	Statistical vs. Practical Significance	133
6.7	Confidence Intervals	133
6.7.1	Computing a Confidence Interval	133
6.7.2	Example	134
6.7.3	Interpreting Confidence Intervals	134
6.7.4	Simulation: Understanding Confidence Intervals	134
6.8	T-Tests, P-Values, and CIs: A Comparison	137
6.9	Reading Regression Tables	137
6.9.1	From R Output to Publication Format	137
6.9.2	Decoding the Table Structure	139
6.9.3	Multiple Columns Show Robustness	139
6.9.4	What to Look for in Regression Tables	139
6.10	F-Tests: Testing Multiple Hypotheses	141
6.10.1	Why Not Just Use Multiple t-Tests?	141
6.10.2	Setting Up the F-Test	142
6.10.3	Computing the F-Statistic	142
6.10.4	F-Test Example	143
6.10.5	Using R's anova() Function	144
6.10.6	The F-Statistic in Terms of R^2	144
6.11	Chapter Summary	145
6.12	Practice Exercises	146

7	Dummy Variables and Non-Linear Models	147
7.1	Categorical Variables in Regression	147
7.1.1	Dummy Variables	147
7.1.2	Interpreting Dummy Variable Coefficients	149
7.1.3	Visualizing What Dummy Variables Do	150
7.2	The Dummy Variable Trap	151
7.3	Multiple Categories	151
7.3.1	Choosing the Baseline Category	153
7.4	Factor Variables in R	154
7.5	Non-Linear Relationships: Polynomial Models	156
7.5.1	Estimating Quadratic Models	157
7.5.2	Interpreting Quadratic Models	157
7.6	Logarithmic Transformations	158
7.6.1	Log-Linear Models	158
7.6.2	Linear-Log Models	159
7.6.3	Log-Log Models (Elasticities)	159
7.6.4	Summary Table	159
7.7	Application: Policy Evaluation with Dummy Variables	160
7.8	Summary	160
7.9	Check Your Understanding	160
8	Heteroskedasticity	161
8.1	The Homoskedasticity Assumption	161
8.1.1	Gauss-Markov Assumption 5	161
8.1.2	What Does Homoskedasticity Look Like?	162
8.2	Heteroskedasticity: When Variance Changes	162
8.2.1	What Is Heteroskedasticity?	162
8.2.2	What Does Heteroskedasticity Look Like?	163
8.2.3	Side-by-Side Comparison	163
8.3	Why Does Heteroskedasticity Matter?	163
8.3.1	The Good News: Coefficients Are Still Unbiased	163
8.3.2	The Bad News: Standard Errors Are Wrong	164
8.3.3	Type I Error: A Quick Review	164
8.4	Simulation: Seeing the Problem	165
8.4.1	Setup: Simulating Hypothesis Tests	165
8.4.2	Now With Heteroskedasticity	166
8.5	Heteroskedasticity-Robust Standard Errors	167
8.5.1	The Solution	167
8.5.2	Visualizing the Sampling Distributions	168
8.5.3	Comparing Type I Error Rates	169
8.6	Implementing Robust Standard Errors in R	169
8.6.1	Using <code>fixest</code>	169
8.6.2	Best Practice: Always Use Robust Standard Errors	170

8.7	Detecting Heteroskedasticity	171
8.7.1	Residual Plots	171
8.7.2	What to Look For	171
8.7.3	Formal Tests	172
8.8	Summary	172
8.9	Check Your Understanding	173
9	Interaction Effects	174
9.1	Why Interactions Matter	174
9.2	Interactions Between Two Dummy Variables	175
9.2.1	Interpreting the Interaction	176
9.3	Interactions Between Dummies and Continuous Variables	176
9.3.1	Different Slopes for Different Groups	177
9.4	Computing Effects with Interactions	179
9.5	Interactions Between Two Continuous Variables	179
9.6	Application: Decomposing the Gender Wage Gap	180
9.7	Best Practices for Using Interactions	181
9.7.1	Always Include Main Effects	181
9.7.2	Center Continuous Variables	181
9.7.3	Test Whether Interactions Are Needed	182
9.8	Summary	182
9.9	Check Your Understanding	182
10	Panel Data and Fixed Effects	183
10.1	Cross-Sectional vs. Panel Data	183
10.1.1	Cross-Sectional Data	183
10.1.2	Panel Data	184
10.2	Why Panel Data Helps: The Intuition	184
10.3	The Fixed Effects Model	185
10.3.1	Setting Up the Model	185
10.3.2	What Does This Solve?	185
10.4	Estimating Fixed Effects: The Within Transformation	186
10.4.1	Example: Unemployment and Crime	186
10.5	Estimating Fixed Effects: The Dummy Variable Approach	188
10.6	What Fixed Effects Control For (and What They Don't)	189
10.6.1	What Unit Fixed Effects Control For	190
10.6.2	What Time Fixed Effects Control For	190
10.6.3	What Fixed Effects Do NOT Control For	190
10.7	Adding Control Variables	191
10.8	Practical Considerations	191
10.8.1	When to Use Fixed Effects	191
10.8.2	When Fixed Effects May Not Work Well	191
10.8.3	Standard Errors	191

10.9 Summary	192
10.10 Check Your Understanding	192
11 Difference-in-Differences	193
11.1 Natural Experiments	193
11.1.1 The Problem with Observational Data	193
11.1.2 Natural Experiments as a Solution	194
11.2 The Mariel Boatlift: A Classic Natural Experiment	194
11.3 The Difference-in-Differences Approach	194
11.3.1 Why Simple Before-After Comparisons Fail	194
11.3.2 The Need for Control Units	195
11.4 The Parallel Trends Assumption	195
11.4.1 The Core Identifying Assumption	195
11.4.2 Visualizing DiD	196
11.5 The DiD Regression	197
11.5.1 Setting Up the Model	197
11.5.2 How It Works	197
11.5.3 Example: Estimating the Mariel Boatlift Effect	198
11.6 Two-Way Fixed Effects	199
11.6.1 Extending to Multiple Units and Time Periods	199
11.6.2 Estimating TWFE in R	199
11.7 Research Design: Choosing Good Controls	200
11.7.1 The Art of Causal Inference	200
11.7.2 Example: Minimum Wages and Employment	200
11.7.3 Intuition-Driven vs. Data-Driven Selection	201
11.8 What Can Go Wrong	201
11.8.1 Violations of Parallel Trends	201
11.8.2 Assessing Parallel Trends	201
11.9 Summary	202
11.10 Check Your Understanding	202
12 Advanced Difference-in-Differences	203
12.1 Data-Driven Control Selection	203
12.1.1 The Problem: Systematically Different Treatment and Control Groups	203
12.1.2 Propensity Score Weighting: The Intuition	204
12.1.3 The IPW-DiD Algorithm	205
12.1.4 Comparing Results	207
12.2 Dynamic DiD and Event Studies	208
12.2.1 Why Treatment Effects May Change Over Time	208
12.2.2 Relative Time	208
12.2.3 The Dynamic DiD Model	209
12.2.4 Example: Environmental Policy and Manufacturing Employment	209
12.2.5 Estimating the Dynamic Model	209

12.2.6	The Event Study Plot	210
12.2.7	Interpreting the Event Study	212
12.3	Cluster-Robust Standard Errors	212
12.3.1	The Problem: Non-Independent Observations	212
12.3.2	Visualizing the Problem	214
12.3.3	Simulation: The Dangers of Ignoring Clustering	214
12.3.4	Implementing Cluster-Robust Standard Errors	214
12.4	Summary	214
12.5	Check Your Understanding	215

13 Instrumental Variables 216

13.1	The Fundamental Problem: Endogeneity	216
13.1.1	When OLS Fails	216
13.1.2	Formalizing the Problem	217
13.2	The Instrumental Variables Solution	217
13.2.1	The Key Insight	217
13.2.2	Two Requirements for a Valid Instrument	217
13.2.3	Graphical Intuition	218
13.3	Application: Fertility and Labor Supply	218
13.3.1	The Research Question	218
13.3.2	The Angrist and Evans (1998) Instrument	219
13.3.3	Simulating the Angrist-Evans Setup	219
13.3.4	Testing the Relevance Condition	219
13.3.5	Testing Excludability (Sort of)	220
13.4	Two-Stage Least Squares (2SLS)	221
13.4.1	The Estimation Strategy	221
13.4.2	Why Does This Work?	222
13.4.3	Implementing 2SLS	222
13.4.4	Proper 2SLS with <code>feols()</code>	224
13.4.5	Comparing OLS and 2SLS	225
13.5	Testing for Weak Instruments	225
13.5.1	The Weak Instrument Problem	225
13.5.2	The F-Test Rule of Thumb	226
13.5.3	What Happens with a Weak Instrument?	226
13.6	What Does IV Estimate? The LATE	226
13.6.1	The Local Average Treatment Effect	226
13.6.2	Four Types of People	227
13.6.3	Who Are the Compliers?	227
13.6.4	Trade-offs of IV Estimation	228
13.7	Classic IV Applications	228
13.7.1	Example: Card (1995) - Returns to Education	228
13.8	Summary	229
13.9	Check Your Understanding	229

References	230
-------------------	------------

Appendices	231
-------------------	------------

A Intro to R and RStudio	231
---------------------------------	------------

A.1	Installing R	231
A.2	Installing RStudio	231
A.3	The RStudio Interface	232
A.3.1	The Script Editor	232
A.3.2	The Console	232
A.3.3	The Environment Panel	232
A.3.4	The Plots and Output Panel	234
A.4	Basic Coding in R	234
A.4.1	Arithmetic Operations	234
A.4.2	Objects and Assignment	235
A.4.3	Vectors	236
A.4.4	Functions	236
A.5	Data Types in R	237
A.5.1	Numeric Types	237
A.5.2	Character Strings	238
A.5.3	Logical Values	238
A.5.4	Factors	238
A.5.5	Checking and Converting Types	239
A.5.6	Common Type Abbreviations	240
A.5.7	Data Frames	240
A.5.8	Comments	241
A.6	Working with R Scripts	241
A.6.1	Creating and Running Scripts	242
A.6.2	Script Best Practices	242
A.7	Working Directories	243
A.8	R Packages	243
A.8.1	Installing Packages	243
A.8.2	The Tidyverse	243
A.8.3	Loading Packages	244
A.8.4	The Pipe Operator	244
A.9	Summary and Conclusion	245
A.9.1	Key Takeaways	245
A.10	Check Your Understanding	245

B Statistics and Probability Review	246
--	------------

B.1	Basic Mathematical Tools	246
B.1.1	The Summation Operator	246

B.1.2	Derivatives	249
B.1.3	Percentages and Percent Changes	249
B.1.4	Logarithms	252
B.1.5	Partial Derivatives	254
B.1.6	Optimization: Maxima and Minima	254
B.2	Population vs. Sample	256
B.3	Probability Basics	257
B.3.1	Random Variables	257
B.3.2	Probability	258
B.3.3	Probability Distributions	258
B.3.4	Expected Values	258
B.4	Measures of Variability	259
B.4.1	Variance	259
B.4.2	Standard Deviation	260
B.4.3	Covariance	260
B.4.4	Correlation Coefficient	261
B.4.5	Conditional Expectations	261
B.5	Distributions	262
B.5.1	The Normal Distribution	262
B.5.2	The Standard Normal Distribution	263
B.6	Summary	264
B.6.1	Key Formulas	264
B.7	Check Your Understanding	264
C	R Functions and Packages Reference	265
C.1	Packages Used in This Book	265
C.1.1	tidyverse	265
C.1.2	modelsummary	265
C.1.3	palmerpenguins	266
C.1.4	wooldridge	266
C.1.5	fixest	267
C.1.6	lmtest	267
C.1.7	sandwich	267
C.1.8	patchwork	268
C.2	Data Inspection Functions	268
C.2.1	head()	268
C.2.2	glimpse()	269
C.2.3	summary()	269
C.2.4	nrow() and ncol()	270
C.2.5	colnames()	270
C.2.6	class() and typeof()	271
C.3	Descriptive Statistics Functions	271
C.3.1	mean()	271

C.3.2	<code>sd()</code>	272
C.3.3	<code>median()</code>	272
C.3.4	<code>min()</code> and <code>max()</code>	273
C.3.5	<code>sum()</code>	273
C.3.6	<code>var()</code>	274
C.3.7	<code>cor()</code>	274
C.4	Data Manipulation Functions (dplyr)	275
C.4.1	<code>select()</code>	275
C.4.2	<code>filter()</code>	275
C.4.3	<code>mutate()</code>	276
C.4.4	<code>group_by()</code>	276
C.4.5	<code>summarize()</code> / <code>summarise()</code>	277
C.4.6	<code>count()</code>	278
C.4.7	<code>arrange()</code>	278
C.4.8	<code>case_when()</code>	279
C.4.9	<code>ifelse()</code>	279
C.4.10	<code>n()</code>	280
C.5	Regression Functions	280
C.5.1	<code>lm()</code>	280
C.5.2	<code>summary()</code> (for regression)	281
C.5.3	<code>coef()</code>	282
C.5.4	<code>confint()</code>	282
C.5.5	<code>predict()</code>	283
C.5.6	<code>residuals()</code>	283
C.5.7	<code>feols()</code> (fixest package)	284
C.5.8	<code>modelsummary()</code> (modelsummary package)	284
C.6	Visualization Functions (ggplot2)	285
C.6.1	<code>ggplot()</code>	285
C.6.2	<code>aes()</code>	285
C.6.3	<code>geom_point()</code>	286
C.6.4	<code>geom_line()</code>	287
C.6.5	<code>geom_histogram()</code>	288
C.6.6	<code>geom_boxplot()</code>	289
C.6.7	<code>geom_bar()</code> and <code>geom_col()</code>	290
C.6.8	<code>geom_smooth()</code>	291
C.6.9	<code>geom_hline()</code> and <code>geom_vline()</code>	292
C.6.10	<code>labs()</code>	293
C.6.11	<code>facet_wrap()</code>	294
C.6.12	<code>theme_minimal()</code>	295
C.7	Utility Functions	296
C.7.1	<code>c()</code>	296
C.7.2	<code>seq()</code>	297
C.7.3	<code>rep()</code>	297

C.7.4	<code>sample()</code>	298
C.7.5	<code>set.seed()</code>	298
C.7.6	<code>rnorm()</code>	299
C.7.7	<code>factor()</code>	299
C.7.8	<code>as.factor()</code> , <code>as.numeric()</code> , <code>as.character()</code>	300
C.7.9	<code>is.na()</code>	300
C.7.10	<code>library()</code>	301
C.7.11	<code>install.packages()</code>	301
C.8	The Pipe Operator: <code> ></code>	301
C.9	Quick Reference Tables	302
C.9.1	Descriptive Statistics Functions	302
C.9.2	Data Manipulation Functions (dplyr)	303
C.9.3	Regression Functions	303
C.9.4	ggplot2 Geometries	303

Preface

Welcome to EC 325: Econometrics at Colby College! This booklet contains the lecture notes, interactive exercises, and R tutorials that will guide you through your first course in econometrics.

What is Econometrics?

Econometrics is the application of statistical methods to economic data with a particular emphasis on *causal inference*—figuring out whether one thing actually *causes* another, rather than merely being correlated with it. This distinction matters enormously for policy: if we want to know whether raising the minimum wage reduces employment, whether health insurance improves health, or whether education increases earnings, we need tools that go beyond simple correlations.

This course will teach you those tools. By the end of the semester, you’ll be able to read and critically evaluate empirical research in economics, conduct your own quantitative analyses, and—most importantly—think carefully about when statistical evidence can and cannot support causal claims.

About These Notes

These notes are designed to be a *companion* to the course, not a replacement for the textbook or class participation. They aim to:

- Present core concepts in accessible, narrative form
- Provide extensive R code examples you can run yourself
- Include interactive exercises to test your understanding
- Emphasize intuition through simulations before introducing formulas

The content draws heavily from the following texts, which are required or recommended for this course:

- Wooldridge (2019) — The primary textbook, providing rigorous coverage of econometric methods

- Angrist and Pischke (2014) — An accessible introduction to causal inference and the “credibility revolution” in economics
- Bailey (2020) — A practical guide to doing econometrics in R
- Wickham, Çetinkaya-Rundel, and Grolemund (2023) — The definitive guide to data science in R with the tidyverse

Structure of the Course

The course is organized into three modules:

Module 1: Correlation, Causation, and Regression We begin by establishing why correlation doesn’t imply causation and what additional assumptions we need to make causal claims. We then develop the core tool of econometrics—Ordinary Least Squares (OLS) regression—first in its simple bivariate form, then with multiple control variables.

Module 2: Statistical Inference and Real-World Regression With the mechanics of OLS in hand, we turn to quantifying uncertainty. How confident should we be in our estimates? We develop the tools of hypothesis testing, confidence intervals, and statistical significance. We then explore practical issues like functional form, dummy variables, and interactions.

Module 3: Causal Inference Methods The final module covers the major identification strategies economists use to make causal claims from observational data: instrumental variables, difference-in-differences, and regression discontinuity designs.

Roadmap Through the Midterm

The first half of the course (through the midterm) covers the foundations:

- **Chapter 1:** We begin with the central question—does correlation imply causation?—using the example of health insurance and health outcomes. You’ll see why simple comparisons can be misleading and why we need econometric tools.
- **Chapter 2:** Before we move onto econometric theory, we cover best practices for data best practices and reproducible research in R
- **Chapter 3:** We introduce the “gold standard” of causal inference—Randomized Control Trials (RCTs)—and the Potential Outcomes Framework, which provides the mathematical foundation for thinking about causality.
- **Chapter 4:** We develop bivariate OLS regression, deriving the estimator, understanding when it gives us causal estimates, and diagnosing when it fails due to omitted variable bias.

- **Chapter 5:** We extend OLS to multiple independent variables, learning how to “control for” confounders and interpret coefficients in the multivariate setting.
- **Chapter 6:** We tackle statistical inference—hypothesis testing, t-statistics, p-values, and confidence intervals—learning to distinguish real effects from statistical noise.

Statistical Software

Modern econometrics is done with statistical software that you interact with through code. In this course, we use **R**, a free and open-source programming language widely used in academia, government, and industry.

Why R? Three reasons:

1. **It’s free.** Unlike Stata or SAS, you can install R on any computer and use it forever at no cost.
2. **It’s powerful.** R can handle everything from simple regressions to cutting-edge machine learning, beautiful data visualizations, and reproducible research documents (these notes were entirely written in R!).
3. **It’s in demand.** R skills are valued by employers in consulting, tech, finance, public policy, and academic research. It regularly ranks among the most widely used programming languages.

Don’t worry if you’ve never programmed before. We’ll start from the basics. See [Appendix A](#) for instructions on getting set up.

A Note on Learning Econometrics

Econometrics is challenging. It combines economic intuition, statistical theory, and programming skills, often all at once. If you find yourself struggling (and you will), that’s normal. Here’s my advice:

- **Run the code yourself.** Don’t just read the examples—type them out, run them, break them, fix them. You learn programming by doing.
- **Focus on intuition first.** Before memorizing formulas, make sure you understand *why* a method works.
- **Ask questions.** Come to office hours, post on the discussion board, work with classmates.
- **Be patient with yourself.** The skills you’re learning will serve you well beyond this course, whether you pursue graduate study, policy work, or a career in the private sector.

About the Instructor

I am a recent PhD from UMass Amherst. My research focuses on the relationship between public policy and maternal, infant, and child health, with particular attention to racial and ethnic health disparities. In my work, I use many of the methods you'll learn in this course.

You can find out more about me and my research at www.raymondcaraher.com.

These notes are a work in progress. If you spot errors or have suggestions for improvement, please let me know!

1 Introduction

💡 Key Questions

- What is the difference between correlation and causation?
- What does an econometrics research project entail?
- What are the main types of data formats we will work with?
- What is statistical software and why is it useful?

i Suggested Readings

- Wooldridge (2019), Ch. 1
- Wickham, Çetinkaya-Rundel, and Grolemund (2023), Ch. 1

1.1 What is this Class About?

This class is obviously dedicated to the study of econometrics (endearingly shortened ‘metrics). But what exactly does econometrics entail? What differentiates it from broader terms such as “statistical analysis” or “data analysis”?

From Wooldridge (2019), we get a classic definition of econometrics:

“...statistical methods for estimating economic relationships, testing economic theories, and evaluating and implementing government and business policy.”

However, my view of econometrics is at the same time, broader and more narrow:

i Definition of Econometrics

The practice of using statistical, quantitative methods to study social phenomena and provide insights into public policy. More specifically, modern econometrics has focused on identifying the **causal effects** of social phenomena/policy as opposed to **correlations**.

In other words, I view the practice of econometrics as the use of quantitative methods to understand the social world around us, with a special (though not exclusive) lens focusing on separating correlation from causation.

1.2 Does Having Health Insurance *Cause* You to be Healthier?

Let's think about a critical question in public policy: *does having health insurance make someone healthier?* This is especially an important question in the context of the United States, which does not have a public health insurance system. Instead, it is a predominantly private system entangled with a complex net of social safety nets meant to provide a patchwork of coverage to the most marginalized.

However, as of 2023, 9.5% of U.S. adults do not have any health insurance coverage.¹ Would the US as a whole be better off guarantee health insurance coverage for everyone? If health insurance makes people healthier, then there may be a strong case to do so. If not, the benefits of such a system would become, at the very least, more opaque.

Let's take a look at life expectancy in the US (a course but important measure of population health), relative to similar high-income, developed countries. Figure 1.1 shows the average life expectancy from 1990 to 2021 for the US, UK, Canada, Australia, and a number of other high-income countries. As can be seen, the US performs considerably worse in terms of life expectancy, and this divergence between peer countries has gotten *larger* since the 1990s, and especially during the COVID-19 pandemic.

Critically—with the exception of the US—all of the countries in the peer group have some form of universal health care. In other words, all developed countries which are healthier than the US have policies which guarantee the uninsured rate is zero.

This then leads to a critical question: did universal health insurance *cause* these countries to have better health outcomes? Or is this a correlation between policy and outcomes? Certainly, one could make an argument that having health insurance does cause better health outcomes. On the other hand, there are lots of differences between the US and other high-income countries which may also explain the pattern seen in Figure 1.1. For example, maybe these countries have different regulations regarding the food quality, which may explain part of the difference in life expectancy. Or maybe because relatively more people commute to work via bicycle or walking, which may have health benefits compared to driving to work.

These other differences between groups which may *confound* our comparison between countries with universal health insurance and those without also extends to individuals. In the US at least, those with health insurance may be more likely to have a high-paying job which includes insurance as a benefit. Therefore, a comparison between those who have insurance and those who do not will be plagued with the fact that those who have insurance *also are more likely*


¹Source: [KFF](#)

Figure 1.1: Life expectancy in the US vs. peer countries.

Life expectancy in the United States is lower than peer nations

Our World
in Data

The period life expectancy at birth, in a given year.



Data source: UN WPP (2022); HMD (2023); Zijdeman et al. (2015); Riley (2005)

OurWorldInData.org/life-expectancy | CC BY

Source: [Our World in Data](https://OurWorldInData.org)

to have a *high income job*, who may have had better health outcomes regardless of if they had insurance or not. In other words, we may be confusing *correlation* for ***causation**.

i Correlation vs. Causation

The **correlation** between two variables describes how they move together. It means two things are *associated*. **Causation** describes a direct relationship where a change in one variable causes a change in another. It means one thing *makes the other happen*.

A key part of the content in this course is to think about when our comparisons are describing a **correlative** relationship or a **causal** relationship. We will study techniques which—if one is convinced that the requisite assumptions hold—will actually be able to recover the *causal effect* of social phenomena, and bring important insights into essential public policy questions like: what is the effect of health insurance on health outcomes? Does going to college cause you to have higher wages? Do immigrants reduce native wages? and does policing cause decreases in crime?

1.3 Broad Steps of Econometric Analysis

Let's say we want to know what the effect of on-the-job training has on your wage. What steps would we need to take to answer this question?

1.3.1 Step 1: Write the Model

Our first step would be to write an *econometric model*, with the goal of using data and econometric tools to *estimate the parameters of the model*. This model can sometimes be derived from a formal model in economics (i.e., the profit maximization of firms), but it is at least based on some intuitive logic which mathematically relates some variable (the **dependent variable**) to some **independent variable(s)**.

An example of the type of econometric model you may want to run to study the effect of on-the-job training is:

$$wage = \beta_0 + \beta_1(training) + \mu, \quad (1.1)$$

where *wage* is the *outcome, left-hand side (LHS), predicted, or dependent variable* (these are synonyms for the same thing), and *training* is the *explanatory, right-hand side (RHS), predictor, or independent variable* (again, synonyms for the same thing). The β_0 and β_1 variables are the parameters that we want to *estimate the value of* using econometric methods, and they tell us *how* our *independent* variable is related mathematically to our dependent variable.

1.3.2 Step 2: Find and Explore the Data

The next thing you have to do—after we figure out the model we want to estimate, such as in Equation 1.1—is to get the data. There is a *huge* range of datasets out there, and we are especially lucky that the data-gathering infrastructure in the United States is incredibly robust. As we go through the semester, we will work with as many different datasets that we can, which cover a range of topics: from sports to health, crime, and economic development. While different datasets usually can their own quirks, broadly speaking there are three different types of data we will work with in this class:

1. **Cross-sectional data:** This is data which contains information on units (e.g., individuals, countries, firms, households, etc.) at a *single point in time*. This data is the most common, and also is generally the simplest to work with, so we will start out by working mostly with cross-sectional data.
2. **Time-series data:** This data consists of observations on a single unit (e.g., a country's Gross Domestic Product (GDP), a specific company's stock price, or the national unemployment rate) across multiple time periods (e.g., years, quarters, months, or even minutes). In time series data, a crucial characteristic is the temporal ordering of the observations. This structure is often used for forecasting and modeling how a variable's past values influence its present and future values.
3. **Panel (or Longitudinal) data:** Panel data combines both cross-sectional and time-series dimensions. It consists of observations on multiple units (e.g., individuals, firms, or countries) over multiple time periods. It essentially tracks the same set of cross-sectional units over time. This gives it more information, more variability, and allows researchers to use a rich set of methods to really hone in on the causal effects of whatever social phenomena being studied.

In all of these cases, econometric data is organized in what is called a **matrix** or a **dataframe**. It should look roughly similar to an Excel Spreadsheet, where each column represents a different variable (which is usually labelled, sometimes not), and each row represents the values that a specific unit has across these variables.

For example, the data we use to estimate the model in Equation 1.1 may look like Table 1.1,

where the **person_name** column is (obviously) each person's name, **person_id** column corresponds the unique numerical identifier associated with each person (this will be more common to see in datasets since names are usually anonymous), the **wage** column shows each person's hourly wage in dollars, and the **yrs_training** column shows the number of years each person received on-the-job training. This row-column structure of organizing data is so fundamental that the word *column* is often interchanged with the word *variable*.

When you first look at some data, it is really important to figure out what each row represents. In the simple example shown in Table 1.1, this is straightforward since each row represents a different person. However, in more complex datasets, it may not be immediately obvious what

Table 1.1: Relationship between training and wages for 10 workers

person_name	person_id	wage	yrs_training
Paul	1	18.50	1.00
Ringo	2	22.00	3.00
George	3	17.65	2.50
John	4	27.95	8.20
Linda	5	19.20	0.50
Yoko	6	24.80	6.80
Pattie	7	20.50	5.50
Maureen	8	23.75	4.20
Cynthia	9	16.40	0.80
Barbara	10	28.50	9.10

each row represents. For example, in panel data, each row may represent a specific person at a specific point in time. So row 1 may be Paul in year 2020, row 2 may be Paul in year 2021, row 3 may be Ringo in year 2020, and so on. Figuring out what each row represents is *critical* to understanding the data, as well as how to appropriately analyze it.


Once we have the data in hand, then we may need to do some data cleaning and organization. This may involve: - Removing missing values - Creating new variables - Changing the format of variables (e.g., from string to numeric) - Merging datasets together - Filtering the data to only include certain observations - And many other tasks.

These are all tasks that in many cases take up the bulk of the time in an econometric research project, and while not the most glamorous part of econometrics, it is absolutely essential to get right.

Before moving on to estimation, it is also important to do some **exploratory data analysis**, (which should be considered step 2.5 in the econometric research process). This involves looking at summary statistics of the data (means, medians, standard deviations, etc.), as well as visualizations (histograms, scatter plots, etc.). Exploratory data analysis is important for understanding the data, as well as for identifying potential issues with the data (e.g., outliers, missing values, etc.).

One of the most important parts of exploratory data analysis is to look at the relationship between the dependent variable and independent variable(s) in your model. One way this can be done is using a **scatterplot**, which plots each observation in the dataset using the independent and dependent variables as coordinates, with the independent variable on the x-axis and the dependent variable on the y-axis.

Figure 1.2: Relationship between training and wages for 10 workers



In Figure 1.2, we see a scatterplot of the relationship between years of training and wages for the 10 workers shown in Table 1.1. From this plot, we can see that there is a positive relationship between years of training and wages: as years of training increases, wages also tend to increase. While not good enough to draw any strong conclusions, one can see how this type of exploratory data analysis is useful for understanding the data and the relationship between variables.

1.3.3 Step 3: Estimate the Model

Once we have a model in mind as well as the appropriate data, we can then use our econometric methods to estimate the relationship. The bulk of this class is dedicated to this step in the econometric research process, but the other stages are no less important! The most common way that econometric models are estimated is with **Ordinary Least Squares** (OLS), which you may remember from your pre-req. In addition to estimating the actual value of the parameters in the model, we also want some idea of how *uncertain* we are about the estimated parameter value. This is where notions of statistical significance and confidence intervals come in.

1.3.4 Step 4: Interpret the Estimate

Once we have an estimated number for the parameter in our model, we then have to interpret the number. Here, the phrase “interpret the estimate” means two parallel things. First, there is the literal, mathematical interpretation of the model. This will change depending on the functional form we use to estimate the model, and we will spend lots of time learning about this. The second part of “interpreting” the model is about what the results mean in the context of your broader research question. Is your estimate big or small? Is it economically important? Do you actually think it reflects a causal effect? These are all types of interpretation which are a critical part of econometrics research, but for which there is not usually a formula we can use. A large part of the “art” working on an econometrics project is exactly this type of interpretation, and how you form it into a compelling narrative. While there are some tools we will cover which will help in this regard, in my opinion, nothing will help you more with this than just reading lots of good, well-written econometrics papers.

1.4 Statistical Software

Much of the econometric process is done using specialized statistical software. We will use this software to: “clean” our data and organize in the appropriate row-column format to do our econometric estimations; explore our data using descriptive statistics (means, medians, etc.) and visualizations (an increasingly important part of a compelling econometric project); estimate our model and conduct analyses of statistical uncertainty; and output our results to easy-to-interpret tables and figures.

There are a few commonly used econometrics software:

- **Stata:** Proprietary software purpose-built for econometrics
- **R:** Open-source programming language with a focus on statistical analysis
- **Python:** Open-source programming language with a focus on data science more broadly
- **EViews:** Proprietary software with a focus on forecasting and other time series applications

Each one of these software of their own pros and cons, and it is worth knowing several quite well. For the purposes of this class, we will learn the **R** programming language to do econometrics. If you haven’t take the time to review [Appendix A](#) to learn more about R, how to install it, and to run your first chunk of code!

1.5 Summary and Conclusion

This chapter introduced the fundamental framework that will guide our study of econometrics throughout this course. We defined econometrics as the use of quantitative methods to un-

derstand social phenomena and inform public policy, with a particular focus on distinguishing between correlation and causation. Through the example of health insurance and health outcomes, we saw how patterns in data can be misleading. While the United States lags behind peer countries in life expectancy, and those peer countries all have universal health insurance, we cannot immediately conclude that universal coverage causes better health outcomes. The tools you will learn in this course are designed to help you identify causal relationships rather than mere associations.

1.5.1 Key Takeaways:

- **Econometrics vs. correlation:** Modern econometrics focuses on identifying causal effects, not just correlations between variables
- **Four steps of econometric research:**
 1. Write an econometric model
 2. Find and explore the appropriate data
 3. Estimate the model parameters
 4. Interpret the results
- **Three main data types:** Cross-sectional (single point in time), time-series (one unit over time), and panel data (multiple units over time)
- **Statistical software:** We will use **R** to clean data, estimate models, and present results

As we move forward, remember that econometrics is as much an art as it is a science. The mathematical tools we will learn are powerful, but they are only as good as the questions we ask and the care with which we interpret our results.

1.6 Check Your Understanding

Note: This section contains interactive content only available in the HTML version.

2 Good Data Practices

Key Questions

- Why do data errors matter for econometric analysis?
- What are descriptive statistics and why should we examine them?
- How can data visualization help us detect problems?
- What is replication and why is it fundamental to credible research?

Suggested Readings

- Bailey (2020), Ch. 2

Before we dive into sophisticated econometric methods, we need to establish good habits for working with data. Even the most advanced statistical techniques cannot rescue an analysis built on flawed or misunderstood data. This chapter focuses on the essential first steps: understanding our data through descriptive statistics and visualization, and documenting our work so that others (including our future selves) can verify and build upon it.

2.1 When Good Economists Use Bad Data

In 2010, economists Carmen Reinhart and Kenneth Rogoff published influential research on government debt and economic growth. Their analysis covered many annual observations from countries around the world, and their key finding was striking: when a country's government debt exceeded 90% of GDP, economic growth dropped dramatically.

The policy implications seemed clear. Governments should be cautious about using deficit spending to fight unemployment. This finding influenced policy debates across the world during a period when many countries were considering how aggressively to respond to the Great Recession. There was one problem with this analysis, however: the data didn't quite say what Reinhart and Rogoff thought it did.

In 2014, a graduate student named Thomas Herndon was trying to replicate their results for a homework assignment. He couldn't get the numbers to match. After obtaining the original Excel spreadsheet from Reinhart and Rogoff, Herndon and his advisors discovered three issues: some observations had been mistakenly excluded from key calculations, observations were

weighed strangely, and some cells in the Excel spreadsheet had simply not been included in the formula calculating the average.

When the data were corrected, the dramatic “cliff” in economic growth disappeared. While growth did decline somewhat as debt increased, the relationship was much more gradual than originally reported. There was no “cliff” that was so central to the original analysis. The corrected analysis told a very different policy story.

This experience teaches us that good data practices are one of the most fundamental tenants of credible research. No amount of sophisticated econometric technique can compensate for errors and misunderstandings in the underlying data. This chapter introduces practices that help us avoid such problems, and will be useful throughout your career as an econometrician.

2.2 Know Your Data

The first rule of data analysis is simple: **know your data**. This means examining it carefully before conducting any formal analysis. We want to understand what we’re working with, detect potential errors, and develop intuition about the relationships we might find.

2.2.1 Descriptive Statistics

For every variable in our dataset, we should know several key statistics: the number of observations, the mean, the standard deviation, and the minimum and maximum values (refer to Appendix B for a review of what exactly these calculations mean). These statistics give us a feel for the data and help us spot problems.

Let’s look at a simple example. Suppose we’re studying the relationship between education and wages. Here’s simulated data for 100 individuals:

```
# Set seed for reproducibility
set.seed(42)

# Generate simulated wage and education data
n <- 100
education <- round(rnorm(n, mean = 14, sd = 2.5))
education <- pmax(8, pmin(education, 20)) # Constrain between 8 and 20 years

# Generate wages with some relationship to education
wages <- 10 + 2.5 * education + rnorm(n, mean = 0, sd = 5)
wages <- pmax(wages, 8) # Minimum wage floor

# Create age variable
```

	Unique	Missing Pct.	Mean	SD	Min	Median	Max
education	12	0	\num{14.1}	\num{2.6}	\num{8.0}	\num{14.0}	\num{20.0}
wage	100	0	\num{44.9}	\num{7.9}	\num{23.2}	\num{45.6}	\num{60.8}
age	33	0	\num{35.3}	\num{9.4}	\num{22.0}	\num{35.0}	\num{60.0}

	Unique	Missing Pct.	Mean	SD	Min	Median	Max
education	13	0	\num{13.9}	\num{3.2}	\num{-5.0}	\num{14.0}	\num{20.0}
wage	100	0	\num{52.8}	\num{80.9}	\num{23.2}	\num{45.6}	\num{850.0}
age	34	0	\num{37.0}	\num{19.0}	\num{22.0}	\num{35.0}	\num{200.0}

```
age <- round(rnorm(n, mean = 35, sd = 10))
age <- pmax(age, 22) # Minimum age

# Create a tibble
wage_data <- tibble(
  id = 1:n,
  education = education,
  wage = wages,
  age = age
)

# Display descriptive statistics
datasummary_skim(wage_data |> select(education, wage, age),
  title = "Descriptive Statistics: Education and Wages",
  histogram = FALSE)
```

These statistics immediately tell us several things. We have 100 observations with no missing data ($N = 100$ for all variables). Education ranges from 8 to 20 years, with a mean around 14 years—this seems reasonable for a sample of working adults. Wages range from about \$8 to \$63 per hour, with substantial variation (standard deviation of about \$10).

But what if our summary statistics looked like this instead?

Now we have a problem! The maximum wage of \$850 per hour seems suspicious—is this real or a data entry error? The minimum education value of -5 years is impossible. And an age of 200 is clearly wrong. These obvious errors suggest we need to investigate our data more carefully.

2.2.2 Computing Descriptive Statistics in R

While summary tables like those above are useful for reports, you often need to calculate specific statistics yourself. R provides simple functions for all the common descriptive statistics.

2.2.2.1 Basic Summary Functions

The most commonly used functions for numeric variables are:

```
# Mean (average)
mean(wage_data$wage)
```

```
[1] 44.91258
```

```
# Standard deviation
sd(wage_data$wage)
```

```
[1] 7.9215
```

```
# Median (middle value)
median(wage_data$wage)
```

```
[1] 45.60513
```

```
# Minimum and maximum
min(wage_data$wage)
```

```
[1] 23.18942
```

```
max(wage_data$wage)
```

```
[1] 60.83585
```

You can also get several statistics at once using `summary()`:

```
summary(wage_data$wage)
```

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
23.19	40.26	45.61	44.91	50.36	60.84

2.2.2.2 Handling Missing Values

Real-world data often contains missing values, represented as `NA` in R. Most summary functions will return `NA` if the data contains any missing values:

```
# Create a vector with a missing value
wages_with_na <- c(15, 22, NA, 18, 25)
mean(wages_with_na)
```

```
[1] NA
```

To calculate statistics while ignoring missing values, use the `na.rm = TRUE` argument:

```
mean(wages_with_na, na.rm = TRUE)
```

```
[1] 20
```

```
sd(wages_with_na, na.rm = TRUE)
```

```
[1] 4.396969
```

Always Check for Missing Data

Before calculating any statistics, check how many missing values you have. The `sum(is.na())` function counts missing values:

```
sum(is.na(wages_with_na))
```

```
[1] 1
```

If a large proportion of observations are missing, your statistics may not be representative of the full sample.

2.2.2.3 Counting and Frequency Tables

For categorical variables, we want to count how many observations fall into each category. The `count()` function from the `tidyverse` makes this easy. We are also going to be using the pipe `|>` in order to string our data operations together like a sentence. For more info on the `tidyverse` and the pipe operator, refer to [Appendix A](#).

First, let's create some categorical variables to work with:

```
# Create categorical variables for demonstration
wage_data <- wage_data |>
  mutate(
    # College degree: 1 if 16+ years of education
    college = ifelse(education >= 16, 1, 0),
    # Age groups
    age_group = case_when(
      age < 30 ~ "Under 30",
      age < 45 ~ "30-44",
      TRUE ~ "45+"
    )
  )
```

Now we can count observations in each category:

```
# Count observations by college status
wage_data |>
  count(college)
```

```
# A tibble: 2 x 2
  college     n
  <dbl> <int>
1       0    67
2       1    33
```

You can count by multiple variables to create cross-tabulations:

```
# Count by two variables
wage_data |>
  count(college, age_group)
```

```
# A tibble: 6 x 3
  college age_group      n
  <dbl> <chr>    <int>
1      0 30-44      39
2      0 45+       14
3      0 Under 30   14
4      1 30-44      12
5      1 45+        5
6      1 Under 30   16
```

To add percentages, you can extend the `count()` output:

```
wage_data |>
  count(college) |>
  mutate(percent = 100 * n / sum(n))
```

```
# A tibble: 2 x 3
  college      n percent
  <dbl> <int>    <dbl>
1      0    67     67
2      1    33     33
```

2.2.2.4 Group-wise Statistics

Often we want to compare statistics across groups. The `group_by()` and `summarize()` functions work together for this:

```
# Calculate mean wage by college status
wage_data |>
  group_by(college) |>
  summarize(
    mean_wage = mean(wage),
    sd_wage = sd(wage),
    n = n()
  )
```

```
# A tibble: 2 x 4
  college mean_wage sd_wage      n
  <dbl>    <dbl>    <dbl> <int>
1      0    41.7    6.92    67
2      1    51.4    5.57    33
```

This is incredibly useful for exploring how outcomes differ across groups—exactly what we’ll be doing in regression analysis.

```
# Compare wages across age groups
wage_data |>
  group_by(age_group) |>
  summarize(
    mean_wage = mean(wage),
    median_wage = median(wage),
    min_wage = min(wage),
    max_wage = max(wage),
    count = n()
  )
```

```
# A tibble: 3 x 6
  age_group mean_wage median_wage min_wage max_wage count
  <chr>      <dbl>      <dbl>    <dbl>    <dbl> <int>
1 30-44      42.9       43.0     23.2     57.5    51
2 45+       45.4       42.7     37.5     60.5    19
3 Under 30   48.1       47.6     27.3     60.8    30
```

Quick Reference: Common Summary Functions

Table 2.1: Common functions for descriptive statistics

Function	What it calculates
<code>mean()</code>	Average
<code>sd()</code>	Standard deviation
<code>median()</code>	Middle value
<code>min()</code> , <code>max()</code>	Minimum, maximum
<code>sum()</code>	Total
<code>n()</code>	Count (inside <code>summarize</code>)
<code>count()</code>	Frequency table
<code>summary()</code>	Multiple statistics at once

2.2.3 Examining Distributions

When a variable takes on a limited number of values, it's helpful to examine the frequency distribution. Recall that we created a variable indicating whether someone has a college degree (1 = yes, 0 = no). Let's look at its distribution:

```
# Create frequency table
wage_data |>
  count(college) |>
  mutate(percent = n / sum(n) * 100) |>
  knitr::kable(
    col.names = c("College Degree", "Count", "Percent"),
    digits = 1,
    caption = "Distribution of College Degree"
  )
```

Table 2.2: Distribution of College Degree

College Degree	Count	Percent
0	67	67
1	33	33

This tells us that 29% of our sample has a college degree. But imagine our frequency table looked like this:

Table 2.3: A Problematic Distribution

College Degree	Count	Percent
0	30	58.8
1	20	39.2
100	1	2.0

We have a value of 100 for the college degree variable. Either we have someone with 100 college degrees (unlikely), or—much more probably—we have a coding error. This is the kind of problem that descriptive statistics help us catch.

2.3 Visualizing Data

Numbers are useful, but graphs often reveal patterns and problems that summary statistics miss. Effective data visualization is both a diagnostic tool and a way to understand relation-

ships in our data. In econometrics, we rely heavily on visualizations to explore data, diagnose problems, and communicate results.

! Why Visualization Matters

A good data visualization translates complex data into a clear visual format, revealing patterns, trends, and relationships that would be difficult to see in tables of numbers. The goal is to tell an accurate and compelling story that allows your audience to quickly grasp key insights.

2.3.1 The Grammar of Graphics: ggplot2

We'll use `ggplot2` (part of the tidyverse) for creating visualizations. The “gg” stands for “grammar of graphics”—a systematic way of thinking about how to build plots.

Every `ggplot` has two main components:

1. **The scaffolding:** titles, axis labels, legends, sources
2. **The content:** the actual data encoded as visual elements (points, lines, bars, etc.)

The key insight of `ggplot2` is that we build plots **layer by layer**, adding elements one at a time until we have a complete visualization.

2.3.2 Getting Started: The Palmer Penguins

Let's use real data to learn `ggplot2`. The Palmer Penguins dataset contains measurements of 344 penguins from three species collected near Palmer Station, Antarctica.

```
# Load the penguins data
library(palmerpenguins)

# Take a look at the data
glimpse(penguins)
```

Rows: 344

Columns: 8

```
$ species      <fct> Adelie, Adelie, Adelie, Adelie, Adelie, Adelie, Adel~
$ island       <fct> Torgersen, Torgersen, Torgersen, Torgersen, Torgerse~
$ bill_length_mm <dbl> 39.1, 39.5, 40.3, NA, 36.7, 39.3, 38.9, 39.2, 34.1, ~
$ bill_depth_mm <dbl> 18.7, 17.4, 18.0, NA, 19.3, 20.6, 17.8, 19.6, 18.1, ~
$ flipper_length_mm <int> 181, 186, 195, NA, 193, 190, 181, 195, 193, 190, 186~
```

```
$ body_mass_g      <int> 3750, 3800, 3250, NA, 3450, 3650, 3625, 4675, 3475, ~
$ sex              <fct> male, female, female, NA, female, male, female, male~
$ year             <int> 2007, 2007, 2007, 2007, 2007, 2007, 2007, 2007, 2007~
```

The data includes species, island location, bill measurements, flipper length, body mass, sex, and year of observation.


2.3.3 Building a Plot: Step by Step

Let's answer a research question: **Do penguins with longer flippers weigh more?**

We start by creating an empty plot object:

```
ggplot(data = penguins)
```

Figure 2.1: An empty ggplot object with no layers.




This creates the plotting area but doesn't show any data yet.

Now let's add a layer with points. We need to specify: - The **geometry** (what shape): `geom_point()` for a scatterplot - The **aesthetic mapping** (which variables go where): `aes(x = ..., y = ...)`

```
ggplot(data = penguins) +  
  geom_point(aes(x = flipper_length_mm, y = body_mass_g))
```

Figure 2.2: A basic scatterplot showing flipper length versus body mass.




The plot clearly shows a positive relationship: penguins with longer flippers tend to be heavier.

2.3.4 Adding More Information: Color and Shape

But wait—could this relationship be driven by different penguin species? Maybe larger species have both longer flippers and greater body mass. We can *encode* species information using color:

```
ggplot(data = penguins) +  
  geom_point(aes(x = flipper_length_mm, y = body_mass_g, color = species))
```

Figure 2.3: Scatterplot with species indicated by color. The positive relationship holds within each species.



Now we can see that:


- Gentoo penguins (blue) are generally larger
- The positive relationship between flipper length and body mass exists **within each species**
- This suggests the relationship isn't just due to species differences

2.3.5 Adding a Line-of-Best Fit

Let's add a line-of-best fit (i.e., a regression line) to visualize the overall trend. This requires a new **layer** (not a new aesthetic):

```
ggplot(data = penguins) +  
  geom_point(aes(x = flipper_length_mm, y = body_mass_g, color = species)) +  
  geom_smooth(aes(x = flipper_length_mm, y = body_mass_g), method = "lm")
```


Figure 2.4: Scatterplot with regression line. The blue line shows the estimated linear relationship.




The `geom_smooth()` layer adds a line of best fit using linear regression (`method = "lm"`).

2.3.6 Proper Labeling

Our plot conveys information well, but the axis labels and title need improvement. Good labels are essential for clear communication:

```
ggplot(data = penguins) +  
  geom_point(aes(x = flipper_length_mm, y = body_mass_g, color = species, shape = species)) +  
  geom_smooth(aes(x = flipper_length_mm, y = body_mass_g), method = "lm") +  
  labs(  
    title = "Body Mass and Flipper Length by Species",  
    subtitle = "Data from Palmer Station, Antarctica",  
    x = "Flipper Length (mm)",  
    y = "Body Mass (g)",  
    color = "Penguin Species",  
    shape = "Penguin Species"  
  ) +  
  theme_minimal()
```

Figure 2.5: A properly labeled scatterplot with clear titles and axis labels.



Notice how much clearer this is! The plot now includes:


- A descriptive title
- Proper axis labels with units
- A data source in the subtitle
- Readable legend labels

2.3.7 Histograms for Distributions

Histograms show the distribution of a continuous variable by dividing it into bins:

```
ggplot(penguins) +  
  geom_histogram(aes(x = body_mass_g), binwidth = 200, fill = "steelblue", alpha = 0.7) +  
  labs(  
    title = "Distribution of Penguin Body Mass",  
    x = "Body Mass (g)",  
    y = "Count"  
  ) +  
  theme_minimal()
```

Figure 2.6: Histogram showing the distribution of penguin body mass.




The `binwidth` parameter controls how wide each bar is. Smaller binwidths create more bars and show more detail.

2.3.8 A Simple Example with Our Wage Data

Let's apply these principles to our wage and education data:

```
ggplot(wage_data, aes(x = education, y = wage)) +  
  geom_point(alpha = 0.6, size = 2, color = "steelblue") +  
  geom_smooth(method = "lm", se = TRUE, color = "darkblue") +  
  labs(  
    title = "Education and Hourly Wages",  
    subtitle = "Simulated data for demonstration",  
    x = "Years of Education",  
    y = "Hourly Wage ($)"  
  ) +  
  theme_minimal()
```

Figure 2.7: The relationship between education and wages with proper labeling and styling.



💡 Building Intuition

The layered approach of ggplot2 mirrors how we think about data analysis: - Start with the data - Add visual representations - Refine with labels and formatting Experiment with different geometries, aesthetics, and layers to find the visualization that best tells your data's story.

2.4 Replication: The Foundation of Credible Research

Science depends on replication. If other researchers cannot reproduce our results using the information we provide, then our findings cannot be verified or built upon. This is not just an abstract principle—it's a practical necessity.

❗ The Replication Standard

Research meets the **replication standard** when an independent researcher can exactly duplicate the results using only the information provided.

2.4.1 Why Replication Matters

Replication serves several crucial purposes. First, it allows others to check our work. The Reinhart-Rogoff example shows why this matters: only because they shared their data could Thomas Herndon discover the errors. Without replication materials, the mistakes might never have been found.

Second, replication enables others to probe our analysis. Statistical results often hinge on seemingly small decisions—which observations to include, how to handle missing data, which variables to control for. Skeptical readers (which should be everyone!) want to see whether reasonable alternative approaches produce similar conclusions. If a certain choice substantially changes our results, we need to pay attention.

Third, committing to a replication standard keeps our work honest. If we know others will check our work, we’re less likely to make choices based on getting the “right” answer rather than statistical merit.

Finally, replication helps our future selves. Even a moderately complex project can involve hundreds of decisions over weeks or months. What seemed obvious at the time can become completely opaque later. Good documentation means we can return to a project after a break and immediately understand what we did.

2.4.2 What Makes Good Replication Materials?

Effective replication files have two components: a **codebook** that documents the data, and **analysis code** that shows exactly how results were produced.

The codebook describes where each variable comes from and any transformations applied. For example:

- **Variable name:** `wage`
- **Description:** Hourly wage in dollars, calculated as annual salary divided by hours worked in past year
- **Source:** National Longitudinal Survey of Youth, 1996 wave
- **Notes:** Wages above \$200/hour were excluded as likely data errors

The analysis code shows the exact steps used to produce each result. This means providing the actual R code, not just descriptions of what we did. And crucially, the code should include comments explaining *why* we made certain choices:

```
# Estimate relationship between education and wages
# We control for age because it affects both education and wages
# (older workers have had more time to complete education and gain experience)
```

```
model <- lm(wage ~ education + age, data = wage_data)
summary(model)
```

2.4.3 Good Coding Practices

As we write code for our analysis, several practices make our work easier to understand and verify:

Use descriptive variable names. Compare `x1` versus `years_education`—the second is much clearer.

Comment extensively. Explain what each section of code does and why you made certain choices. Your future self will thank you.

Organize your code logically. Start with data loading and cleaning, then move to analysis, then to creating tables and figures.

Make your code reproducible. Use `set.seed()` before any random operations so results don't change each time you run the code.

Here's an example of well-documented code:

```
# =====
# Analysis of Education and Wages
# Author: Your Name
# Date: December 2024
# =====

# Load required packages
library(tidyverse)

# Set seed for reproducibility
set.seed(42)

# -----
# 1. LOAD AND CLEAN DATA
# -----

# Load data from National Longitudinal Survey
# Downloaded from: https://www.nlsinfo.org/content/cohorts/nlsy79
# Download date: December 1, 2024
wage_data <- read_csv("nlsy_wages.csv")

# Remove observations with missing wage data
```

```

# This affects 23 observations (2.3% of sample)
wage_data <- wage_data |>
  filter(!is.na(wage))

# Create college degree indicator
# Defined as 16+ years of education (bachelor's degree)
wage_data <- wage_data |>
  mutate(college = ifelse(education >= 16, 1, 0))

# -----
# 2. DESCRIPTIVE STATISTICS
# -----

# Summary statistics for main variables
summary(wage_data |> select(wage, education, age))

# -----
# 3. MAIN ANALYSIS
# -----

# Estimate basic relationship between education and wages
# Simple regression without controls
model1 <- lm(wage ~ education, data = wage_data)

# Add age as control variable
# Age affects both education completion and wage levels
model2 <- lm(wage ~ education + age, data = wage_data)

```

2.5 A Quick Example: State Crime Data

Let's practice these principles with a small example. We'll look at violent crime rates across U.S. states and potential correlates.

```

# Create simulated state crime data for demonstration
set.seed(123)
n_states <- 50

state_data <- tibble(
  state = state.abb,
  violent_crime = rnorm(n_states, mean = 400, sd = 150),

```

	Unique	Missing Pct.	Mean	SD	Min	Median	Ma
violent_crime	50	0	\num{405.2}	\num{138.9}	\num{105.0}	\num{389.1}	\n
percent_urban	49	0	\num{72.1}	\num{13.4}	\num{35.4}	\num{72.3}	\n
percent_poverty	50	0	\num{13.2}	\num{3.0}	\num{8.0}	\num{13.1}	\n

```

percent_urban = rnorm(n_states, mean = 70, sd = 15),
percent_poverty = rnorm(n_states, mean = 14, sd = 3)
) |>
mutate(
  violent_crime = pmax(violent_crime, 100), # Keep positive
  percent_urban = pmax(20, pmin(percent_urban, 100)), # Constrain to 0-100
  percent_poverty = pmax(8, pmin(percent_poverty, 22)) # Reasonable range
)

# Display descriptive statistics
datasummary_skim(state_data |> select(-state),
  title = "State Crime Data: Descriptive Statistics",
  histogram = FALSE)

```

The descriptive statistics look reasonable. Crime rates range from about 150 to 750 per 100,000 people. The urbanization variable is measured on a 0-100 scale, while poverty is measured as a percentage (8% to 22%). Understanding these scales is important for interpreting results.


Now let's visualize the relationship between urbanization and crime:

```

ggplot(state_data, aes(x = percent_urban, y = violent_crime)) +
  geom_point(alpha = 0.6, size = 2.5) +
  geom_smooth(method = "lm", se = FALSE, color = "steelblue") +
  labs(
    x = "Percent Urban (%)",
    y = "Violent Crime Rate (per 100,000)",
    title = "Urbanization and Violent Crime"
  ) +
  theme_minimal()

```


Figure 2.8: Relationship between urbanization and violent crime rates across U.S. states.



The scatterplot suggests a positive relationship: more urbanized states tend to have higher violent crime rates. But this is just correlation, not causation. Many factors differ between urban and rural states beyond just population density, and any of these could be driving the relationship we observe. This is precisely the kind of endogeneity problem that motivates the econometric methods we'll study in later chapters.

2.6 Summary

Good data practices are the foundation of credible econometric analysis. Before conducting any formal statistical tests, we must understand our data through descriptive statistics and visualization. These diagnostic tools help us catch errors, understand variable scales and distributions, and develop intuition about relationships in the data.

Equally important is documenting our work so that others can replicate our results. A replication file should include a codebook describing data sources and transformations, plus the complete analysis code with clear comments explaining our choices. This commitment to transparency keeps our work honest, enables others to verify and build on our findings, and helps our future selves understand what we did months or years later.

The Reinhart-Rogoff case reminds us that even accomplished researchers make data mistakes. The best defense is establishing good habits: examine your data carefully, document everything,

and write code that your future self (and others) can understand.

2.7 Check Your Understanding

Note: This section contains interactive content only available in the HTML version.

3 Causality and Randomized Control Trials

Key Questions

- Why can't we interpret correlations as causal effects?
- What is selection bias and how does it arise?
- How do randomized control trials eliminate selection bias?
- What is the potential outcomes framework and how does it formalize causal inference?

Suggested Readings

- Angrist and Pischke (2014), Ch. 1

In Chapter 1, we discussed how econometrics focuses on identifying **causal effects** rather than mere correlations. We used the example of health insurance: while people with health insurance tend to be healthier than people without insurance, we cannot immediately conclude that health insurance *causes* better health. This chapter develops the conceptual and mathematical tools we need to think rigorously about causality, and introduces the randomized control trial (RCT) as the gold standard for causal inference.

3.1 The Problem with Simple Comparisons

Let's return to our health insurance example. People with health insurance are, on average, healthier than people without it. This correlation is undeniable. But can we interpret this as evidence that insurance *causes* better health?

The problem is that people who have health insurance differ from those who don't in many ways beyond just their insurance status. Those with insurance tend to be wealthier, more educated, and have more stable employment. All of these factors independently affect health outcomes. When we compare the health of insured and uninsured individuals, we're not just comparing the effect of insurance—we're comparing people who differ along many dimensions.

Selection Bias

Selection bias occurs when we compare groups that differ systematically in ways beyond just the treatment we’re studying. In other words, we are comparing **apples to oranges**.

The comparison we *want* to make is between two otherwise identical people who differ only in whether they have health insurance. In other words, we want an **apples to apples** comparison. This is the essence of the *ceteris paribus* condition: **all else being equal**, the difference between treated and untreated units reveals the causal effect of treatment.

3.2 The Multiverse of Possibilities

To build intuition for causal inference, consider the television show *Rick and Morty*. In this show, the character Rick possesses a “portal gun” that allows him to travel across infinite parallel universes. In each universe, there exists a version of his grandson Morty who is identical in every way except for one characteristic—Cowboy Morty, Evil Morty, Hammer Morty, and so on. Each Morty has the same home, same parents, same sister, same *everything else*.

This multiverse provides the perfect setting for causal inference! Comparisons between different Mortys are truly *ceteris paribus* comparisons because everything except the one characteristic of interest is held constant.

3.2.1 The Causal Effect of College

Suppose we want to know the causal effect of attending college on wages. We can’t simply compare the wages of college graduates to non-graduates because these groups likely differ in many unobserved ways (motivation, family background, innate ability, etc.).

But imagine we could observe two versions of Morty from parallel universes: Morty C-137 (who didn’t attend college) and College Morty (who did). These two Mortys are identical in every way—same intelligence, same family, same opportunities—except that one went to college and the other didn’t.

The difference in wages between College Morty and Morty C-137 would be a genuine *ceteris paribus* difference—the true causal effect of college on wages.

Figure 3.1: Rick and Morty: A show about infinite parallel universes.



Figure 3.2: Morty C-137 (left) and College Morty (right). The wage difference between them represents the true causal effect of college.



3.2.2 The Fundamental Problem

Of course, the fundamental problem of causal inference is that **we don't have a portal gun**. We only observe one version of reality, not parallel universes. We can never observe what would have happened to a college graduate had they *not* gone to college, or what would have happened to a non-graduate had they attended. This unobserved “what if” scenario is called the **counterfactual**.

3.3 Randomized Control Trials: The Next Best Thing

Since we can't travel between parallel universes, we need another way to make valid causal comparisons. The solution that comes closest to achieving *ceteris paribus* conditions in the real world is the **randomized control trial** (RCT).

Consider how a pharmaceutical company tests whether a new drug works. They don't just give the drug to sick people and see if they get better—that would be subject to all sorts of selection bias. Instead, they conduct an RCT:

1. **Sample** a group of volunteers from the population of interest
2. **Randomly assign** some volunteers to receive the drug (the “treatment” group) and others to receive a placebo (the “control” group)
3. **Compare** the average outcomes between the treatment and control groups

The key insight is that **random assignment eliminates selection bias**. When treatment is assigned randomly, there is no systematic relationship between who receives treatment and their underlying characteristics. On average, the treatment group and control group will be identical in terms of age, health status, income, motivation, and every other characteristic—both observed and unobserved.

! Why Randomization Works

Random assignment ensures that, on average, the treatment and control groups have identical characteristics. Any difference in outcomes between the groups can therefore be attributed to the treatment itself, not to pre-existing differences between the groups.

3.4 The Law of Large Numbers

Why exactly does randomization eliminate selection bias? The answer lies in a fundamental result from probability theory: the **Law of Large Numbers** (LLN).

The LLN states that as the sample size increases, the sample average converges to the population average. Formally, if $Y_1, Y_2, \dots, Y_n$ are independent, identically distributed random variables with mean μ , then:

$$\text{plim}(\bar{Y}_n) = \mu$$

where $\bar{Y}_n$ is the sample average and plim denotes the probability limit (what the value converges to as $n \rightarrow \infty$).

3.4.1 Seeing the LLN in Action: Rolling Dice

To see the Law of Large Numbers at work, consider a simple example: rolling a fair six-sided die. The expected value (average outcome) of a single roll is:

$$\mu = \frac{1 + 2 + 3 + 4 + 5 + 6}{6} = 3.5$$

If we roll a die just once or twice, we might get values far from 3.5 (like a 1 or a 6). But what happens if we roll it many, many times? Let's see what happens!

First, we will “pretend” rolling a dice by using the `sample()` function. This function will randomly select `n` numbers from a vector, here the numbers 1-6.

Let's first roll the dice once. We will set the `size` argument in the `sample()` function to `size = 1` to generate a single random number from our vector.

```
# Set seed for reproducibility
set.seed(1248)

# Create a vector which represents the possible outcomes of a dice roll
dice <- c(1, 2, 3, 4, 5, 6)

# Now "roll" the dice using the sample() function

roll_1 <- sample(dice, size = 1, replace = TRUE)

# Print the result of the dice roll
roll_1
```

```
[1] 2
```


With 1 dice roll, we get a value of 2. Obviously pretty far from the expected value $\mu = 3.5$.

Now, let's roll it 5 times and take the average of our 5 different rolls using the `mean()` function:

```
roll_5 <- sample(dice, size = 5, replace = TRUE)

# Print the 5 different dice rolls
roll_5
```

```
[1] 1 5 2 5 1
```

```
# Find the mean of the 5 different dice rolls
mean(roll_5)
```

```
[1] 2.8
```

With just 5 rolls, our average is a little closer to the expected value. Now, let's try to roll it 25 times, 50 times, and 100 times and find the average values of our dice roll!

```
# Continue rolling to get 10 total rolls
rolls_25 <- sample(dice, size = 10, replace = TRUE)
mean(rolls_25)
```

```
[1] 2.3
```

```
# Keep going to 50 rolls
rolls_50 <- sample(dice, size = 50, replace = TRUE)
mean(rolls_50)
```

```
[1] 3.68
```

```
# And finally 100 rolls
rolls_100 <- sample(1:6, size = 100, replace = TRUE)
mean(rolls_100)
```

```
[1] 3.35
```

Notice how the average tends to get closer to 3.5 as we increase the number of rolls. This isn't guaranteed on any single attempt (randomness still plays a role), but the general tendency is clear.

Now let's see what happens when we roll the die many times and track the running average at each step. In Figure 3.3, I plot the *sample average* of our dice rolls on the y-axis, and the number of times I roll the dice on the x-axis. The population mean, 3.5, is plotted with a dashed red line.

```
# Set seed for reproducibility
set.seed(12345)

# Number of dice rolls to simulate
n_rolls <- 1000


# Simulate rolling a six-sided die n_rolls times
dice_rolls <- sample(1:6, size = n_rolls, replace = TRUE)

# Calculate the running average after each roll
running_avg <- cumsum(dice_rolls) / 1:n_rolls

# Create a data frame for plotting
dice_data <- tibble(
  roll_number = 1:n_rolls,
  running_average = running_avg
)

# Plot the running average
ggplot(dice_data, aes(x = roll_number, y = running_average)) +
  geom_line(color = "steelblue", linewidth = 0.8) +
  geom_hline(yintercept = 3.5, linetype = "dashed", color = "red", linewidth = 1) +
  annotate("text", x = 750, y = 3.7, label = "Expected value = 3.5",
    color = "red", size = 4) +
  labs(
    x = "Number of Rolls",
    y = "Running Average",
    title = "Convergence to Expected Value"
  ) +
  theme_minimal() +
  coord_cartesian(ylim = c(2.5, 4.5))
```

Figure 3.3: The Law of Large Numbers in action. As we roll a die more times, the average of all rolls converges to the expected value of 3.5.



Notice several key features about Figure 3.3:

- **Early volatility:** In the first few rolls, the running average bounces around wildly. After just 10 rolls, we might be at 4.2 or 2.9.
- **Gradual stabilization:** As we accumulate more rolls (say, 100 or 200), the average starts to settle down closer to 3.5.
- **Long-run convergence:** By the time we've rolled 1,000 times, the average is very close to the theoretical expected value of 3.5.

This is the Law of Large Numbers in action: **with enough observations, sample averages converge to population averages.** The key word is “enough”—we need a sufficiently large sample for the convergence to occur.

3.4.2 How the LLN Enables Causal Inference

When we randomly assign individuals to treatment and control groups, we are essentially drawing random samples from the same underlying population. The LLN guarantees that, with a large enough sample, the average characteristics of each group will converge to the population average—and therefore, to each other.

This is why randomization eliminates selection bias: the treatment and control groups become statistically identical in all characteristics—both observed and unobserved—as the sample size grows.

3.5 How Randomization Eliminates Selection Bias: A Simulation

What is a Simulation?

A **simulation** is a computer-based experiment where we generate artificial data according to rules we specify. Unlike real-world data, we know exactly how the simulated data was created—including the true causal effects, which variables influence which outcomes, and how treatment was assigned.

Simulations are powerful learning tools because they let us see what *should* happen under ideal conditions. If we set the true causal effect to zero and then compare treated and untreated groups, any difference we observe must be due to something other than a real treatment effect (like selection bias or random chance). By manipulating features of the simulation—such as sample size or how treatment is assigned—we can build intuition for how these factors affect our ability to recover causal effects. Think of simulations as a laboratory where we can run controlled experiments on statistical methods themselves.

Let's look at a more complex version of how the LLN works in the context of a randomized control trial. Consider a simulation where we have a population with three observable characteristics: light-colored eyes, dark hair, and being tall. Suppose these characteristics are correlated with both treatment status and outcomes in the observational data (i.e., without random assignment). Specifically, having light-colored eyes will be *strongly positively correlated* with the treatment and the outcome; having dark hair will be *moderately positively correlated* with the treatment and the outcome, and being tall will be *slightly negatively correlated* with the treatment and the outcome. Critically, we set the true causal effect of treatment on the outcome to **zero**.

A Note on the Code Below

The R code in the following sections is more advanced than what we've covered so far in the course. Don't worry if you don't understand every line! The code is provided for those who are curious and want to see how simulations work “under the hood.” Even if the syntax is unfamiliar, reading through the comments can help you understand the logic of what we're doing. As you progress through the course, more of this code will start to make sense.

```

# Load the tidyverse package for data manipulation and plotting
library(tidyverse)

# Set a "seed" for reproducibility
# This ensures we get the same random numbers each time we run the code
set.seed(181247)

# Define the TRUE causal effect of treatment
# We're setting this to ZERO so we know any observed difference is bias!
causal_effect <- 0

# Create a population of 10,000 individuals
n_population <- 10000

# Generate three random characteristics for each person
# Each characteristic is randomly 0 or 1 (like a coin flip)
eye_light <- sample(x = c(1, 0), size = n_population, replace = TRUE)
hair_dark <- sample(x = c(1, 0), size = n_population, replace = TRUE)
big_feet <- sample(x = c(1, 0), size = n_population, replace = TRUE)

# Generate random "noise" - unexplained variation in outcomes
err <- rnorm(n_population)

# Create an underlying "propensity" for treatment
# Notice: people with light eyes are MUCH more likely to be treated (coefficient = 2.1)
# People with dark hair are moderately more likely (coefficient = 0.83)
# Big feet has a small negative effect (coefficient = -0.11)
treatment_propensity <- 1 + 2.1 * eye_light + 0.83 * hair_dark - 0.11 * big_feet + err

# Assign treatment based on this propensity (treated if propensity > 2.25)
# This creates SELECTION BIAS: treatment is correlated with characteristics
treated <- ifelse(treatment_propensity > 2.25, 1, 0)

# Create the outcome variable
# The outcome depends on the same characteristics that predict treatment!
# But notice: we add "treated * causal_effect" which equals zero
# So treatment has NO real effect on the outcome
outcome <- 1 + 2.2 * eye_light + 1.2 * hair_dark - 0.01 * big_feet + treated * causal_effect

# Combine everything into a data frame
dat <- tibble(
  outcome = outcome,

```

```

    treated = treated,
    eye_light = eye_light,
    hair_dark = hair_dark,
    big_feet = big_feet
  )

# Calculate the mean of each variable for treated vs untreated groups
baseline_diffs <- dat |>
  # Group the data by treatment status
  group_by(treated) |>
  # Calculate summary statistics for each group
  summarise(
    Outcome = mean(outcome),
    `Prop. light eyes` = mean(eye_light),
    `Prop. dark hair` = mean(hair_dark),
    `Prop. big feet` = mean(big_feet)
  ) |>
  # Reshape data from "wide" to "long" format for plotting
  pivot_longer(
    cols = Outcome:`Prop. big feet`,
    names_to = "variable",
    values_to = "value"
  ) |>
  # Create nice labels for the treatment groups

  mutate(group = ifelse(treated == 1, "Treated", "Untreated"))

# Create the bar plot
ggplot(baseline_diffs) +
  geom_bar(
    aes(x = group, y = value, fill = group),
    stat = "identity"
  ) +
  # Create separate panels for each variable
  facet_wrap(~variable, nrow = 1, scales = "free_y") +
  labs(
    x = "Group",
    y = "Value",
    title = "Baseline Differences: No Randomization"
  ) +
  theme_minimal() +
  theme(


```

```

legend.position = "none",
strip.text = element_text(size = 11)
)

```

Figure 3.4: Baseline differences between treated and untreated groups without randomization. Despite the true causal effect being zero, we observe large differences in both characteristics and outcomes.



Notice in Figure 3.4 how the treated and untreated groups look very different! The treated group has a higher proportion of people with light eyes and dark hair. Most importantly, the treated group has a higher mean outcome—even though **we know the true causal effect is zero**. This is selection bias in action: people who “select into” treatment have characteristics that also lead to better outcomes.

```

# Function to run the random assignment simulation for a given sample size
run_random_assignment <- function(data, sample_size, seed = 181247) {

  # Set seed for reproducibility
  set.seed(seed)

  # Take a random sample from the untreated population
  # (We use untreated only to avoid any prior treatment effects)
  sample_data <- data |>

```

```

    filter(treated == 0) |>
    slice_sample(n = sample_size)

# RANDOMLY assign treatment - this is the key step!
# Each person has a 50% chance of being assigned to treatment
# This assignment is INDEPENDENT of their characteristics
random_treatment <- sample(c(1, 0), size = sample_size, replace = TRUE)

# Add the random treatment assignment to our sample
sample_data <- sample_data |>
  mutate(rand_treat = random_treatment)

# Calculate means for each randomly assigned group
sample_data |>
  group_by(rand_treat) |>
  summarise(
    Outcome = mean(outcome),
    `Prop. light eyes` = mean(eye_light),
    `Prop. dark hair` = mean(hair_dark),
    `Prop. big feet` = mean(big_feet),
    .groups = "drop"
  ) |>
  mutate(sample_size = sample_size)
}

# Run the simulation for different sample sizes
results_n10 <- run_random_assignment(dat, sample_size = 10)
results_n25 <- run_random_assignment(dat, sample_size = 25)
results_n50 <- run_random_assignment(dat, sample_size = 50)
results_n100 <- run_random_assignment(dat, sample_size = 100)
results_n500 <- run_random_assignment(dat, sample_size = 500)

# Combine all results
all_results <- bind_rows(results_n10, results_n25, results_n50,
  results_n100, results_n500)

# Reshape for plotting
all_results_long <- all_results |>
  pivot_longer(
    cols = Outcome:`Prop. big feet`,
    names_to = "variable",

```




```

    values_to = "value"
  ) |>
  mutate(
    group = ifelse(rand_treat == 1, "Treated", "Control"),
    sample_size_label = paste("N =", sample_size)
  )

```

Now let's see what happens when we randomly assign treatment. Starting with a small sample (N=10), we see in Figure 3.5 that random assignment reduces but doesn't eliminate the differences between groups. With such a small sample, random chance can still produce imbalanced groups. With only 10 people, we still see noticeable differences between the randomly assigned treatment and control groups.

Figure 3.5: Random assignment with N=10. Some imbalance remains due to small sample size.



As we increase the sample size to N=25 and N=50, as shown in Figure 3.6 and Figure 3.7, the differences between treatment and control groups shrink. The LLN is at work: larger samples produce more balanced groups in terms of their background characteristics.

By N=500, as shown in Figure 3.8, the treatment and control groups are nearly identical across all characteristics. The difference in outcomes between groups is now close to zero—the true causal effect! Any remaining difference is just random variation, not systematic selection bias.

Figure 3.6: Random assignment with N=25.




Figure 3.7: Random assignment with N=50.





Figure 3.8: Random assignment with $N=500$. The treatment and control groups are now nearly identical in their characteristics, and the difference in outcomes is close to the true causal effect of zero.



3.6 The Potential Outcomes Framework

Now that we intuitively understand why simple comparisons can be misleading and how randomization addresses this problem, let's formalize these concepts mathematically using the **Potential Outcomes Framework**.

3.6.1 Defining Potential Outcomes

Consider our example of estimating the causal effect of college on wages. For any individual i , we can define two **potential outcomes**:

- Y_{1i} : The wage individual i would earn **if they attend college**
- Y_{0i} : The wage individual i would earn **if they do not attend college**

The causal effect of college for individual i is simply the difference between these potential outcomes:

$$\kappa_i = Y_{1i} - Y_{0i}$$

This framework captures the “multiverse” intuition: Y_{1i} is College Morty’s wage, Y_{0i} is Morty C-137’s wage, and κ_i is the causal effect of college for Morty.

3.6.2 The Fundamental Problem of Causal Inference

The fundamental problem is that we only observe **one** potential outcome for each individual. If person i attends college, we observe Y_{1i} but never see Y_{0i} . If they don’t attend, we observe Y_{0i} but never see Y_{1i} . We never observe both potential outcomes for the same individual.

Let D_i be a treatment indicator:

$$D_i = \begin{cases} 1 & \text{if individual } i \text{ is treated (attends college)} \\ 0 & \text{otherwise} \end{cases}$$

What we actually observe is:

$$Y_i = D_i \cdot Y_{1i} + (1 - D_i) \cdot Y_{0i}$$

That is, we observe the treated outcome for treated individuals and the untreated outcome for untreated individuals—but never both.

3.6.3 Selection Bias Formalized

Now we can formally see why simple comparisons are problematic. Suppose we compare average outcomes between those who are treated ($D_i = 1$) and those who are untreated ($D_i = 0$). The difference in group means is:

$$E[Y_i | D_i = 1] - E[Y_i | D_i = 0]$$

Since we only observe Y_{1i} for the treated and Y_{0i} for the untreated, this becomes:

$$E[Y_{1i} | D_i = 1] - E[Y_{0i} | D_i = 0]$$

Here’s the key insight. Let’s add and subtract a term that represents the counterfactual—what the treated group’s outcome would have been without treatment:

$$E[Y_{1i} | D_i = 1] - E[Y_{0i} | D_i = 0] + \underbrace{\{E[Y_{0i} | D_i = 1] - E[Y_{0i} | D_i = 1]\}}_{=0}$$

Rearranging terms:

$$\underbrace{E[Y_{1i}|D_i = 1] - E[Y_{0i}|D_i = 1]}_{\text{Causal effect for the treated}} + \underbrace{\{E[Y_{0i}|D_i = 1] - E[Y_{0i}|D_i = 0]\}}_{\text{Selection bias}}$$

! The Selection Bias Decomposition

A simple difference in group means equals the causal effect **plus** selection bias:

$$\underbrace{E[Y_i|D_i = 1] - E[Y_i|D_i = 0]}_{\text{Observed difference}} = \underbrace{\kappa}_{\text{Causal effect}} + \underbrace{\{E[Y_{0i}|D_i = 1] - E[Y_{0i}|D_i = 0]\}}_{\text{Selection bias}}$$

The selection bias term compares the **untreated potential outcomes** of the treated group to the untreated potential outcomes of the control group. In the health insurance example, this captures differences in baseline health (absent insurance) between those who choose to get insurance and those who don't. If people who get insurance would be healthier even without insurance (perhaps because they're wealthier or more health-conscious), then selection bias is positive, and the simple comparison overstates the causal effect.

3.6.4 How Randomization Eliminates Selection Bias

When treatment is randomly assigned, the treatment and control groups are drawn from the same population. By the Law of Large Numbers, their average characteristics—including their average potential outcomes—converge to each other. Formally:

$$E[Y_{0i}|D_i = 1] = E[Y_{0i}|D_i = 0]$$

This says that the average untreated potential outcome is the same for both groups. Plugging this into our decomposition:

$$E[Y_i|D_i = 1] - E[Y_i|D_i = 0] = \kappa + \{E[Y_{0i}|D_i = 1] - E[Y_{0i}|D_i = 0]\} = \kappa + 0 = \kappa$$

With random assignment, the simple difference in means equals the causal effect. Selection bias disappears.

3.7 The Oregon Health Insurance Experiment

Most econometric research cannot rely on randomization to identify causal effects. We typically work with observational data where treatment is not randomly assigned, and much of this course focuses on methods for addressing selection bias in such settings. However, there have been a few landmark studies that used randomization to study important policy questions. One of the most influential is the **Oregon Health Insurance Experiment**.

3.7.1 Context: Medicaid Expansion

Throughout the 2000s, many U.S. states expanded their Medicaid programs to cover more low-income residents. Most states simply extended coverage to everyone who became newly eligible. But in 2008, Oregon did something unique: facing limited resources, they offered Medicaid coverage through a public lottery.

3.7.2 The Lottery

Oregon's approach created a natural randomized experiment. People who wanted coverage could sign up for the lottery. Winners received the opportunity to apply for the Oregon Health Plan (the state's Medicaid program), while losers remained uninsured. To be eligible, applicants had to be legal Oregon residents aged 19-64, uninsured for the past six months, with income below the federal poverty line, and not otherwise eligible for health insurance.

Because lottery winners were chosen randomly, the comparison between winners and losers provides a valid causal estimate of the effect of health insurance—free from the selection bias that plagues observational studies.

3.7.3 Results

Researchers followed lottery participants and found that lottery winners (relative to losers) experienced:

1. **Increased take-up of Medicaid:** The lottery actually changed insurance status, confirming the treatment “worked”
2. **Increased use of medical resources:** Winners used more hospital and outpatient services
3. **Small improvements in overall health:** Physical health improvements were modest, though mental health and financial security improved substantially

These findings are important because they come from a rigorous RCT rather than observational comparisons. The study provides some of the cleanest evidence we have on what health insurance actually *causes*, as opposed to what it merely correlates with.

Discussion Question

The Oregon experiment found modest effects of insurance on physical health. Does this mean expanding health insurance is not worthwhile? What other outcomes might matter for policy decisions?

3.8 Summary and Conclusion

This chapter introduced the fundamental concepts underlying causal inference in econometrics. We began with the intuition that simple comparisons between treated and untreated groups can be misleading due to selection bias—the groups may differ in ways beyond just the treatment. The “multiverse” thought experiment illustrated what we really want: *ceteris paribus* comparisons where everything except the treatment is held constant.

Randomized control trials approximate this ideal by randomly assigning treatment, which ensures that treatment and control groups are comparable on average. The Law of Large Numbers guarantees that with large enough samples, random assignment eliminates selection bias.

We then formalized these ideas using the potential outcomes framework, which defines causal effects as the difference between what would happen with and without treatment. The fundamental problem is that we only observe one potential outcome per individual. Simple comparisons of means capture both the causal effect and selection bias, but random assignment makes selection bias vanish.

3.8.1 Key Takeaways

- **Selection bias** arises when treated and untreated groups differ systematically beyond just the treatment—we’re comparing apples to oranges
- **Ceteris paribus** (“all else equal”) comparisons isolate the causal effect by holding everything constant except the treatment
- **Randomized control trials** eliminate selection bias by ensuring treatment and control groups are drawn from the same population
- The **Law of Large Numbers** guarantees that with large samples, randomly assigned groups have identical average characteristics

- The **potential outcomes framework** formalizes causal effects as $\kappa = Y_{1i} - Y_{0i}$, but we only observe one potential outcome per individual
- Simple differences in means equal: Causal effect + Selection bias. Randomization sets selection bias to zero.

3.9 Check Your Understanding

Note: This section contains interactive content only available in the HTML version.

4 Bivariate Regression and Ordinary Least Squares

Key Questions

- How do we mathematically model the relationship between two variables?
- What is the population regression model and how do we interpret its parameters?
- How do we derive the OLS estimator?
- What assumptions are required for OLS to be unbiased?
- What is omitted variable bias and when does it arise?
- When can we interpret OLS estimates as causal effects?
- How do we measure the precision of our estimates?

Suggested Readings

- Wooldridge (2019), Ch. 2

In Chapter 3, we learned that randomized control trials provide the gold standard for causal inference by eliminating selection bias through random assignment. But most economic questions cannot be answered with experiments. We typically work with observational data where “treatment” is not randomly assigned. This chapter introduces **Ordinary Least Squares (OLS)** regression, the workhorse tool of econometrics. We focus here on the **bivariate** case—a single independent variable—to build intuition before extending to multiple variables.

4.1 Modeling Relationships Between Variables

Suppose we have two variables, y and x . In econometrics, we are often interested in one of several related goals: estimating the **causal effect** of x on y , explaining variation in y using x , or simply describing how y varies with x . OLS is a tool that can help with all of these—though as we’ll see, whether we can interpret results causally depends on assumptions that may or may not hold.

But what do we even *mean* when we say we want to study “how x is related to y ”? We need a precise, mathematical way to operationalize this question. The most straightforward approach is to ask: what is the *average* value of y for each different value that x can take?

Consider some examples. How can we quantify the gender pay gap? We compare the *average* wage for men versus women. How do we measure the returns to education? We compare the *average* wage of people with different levels of education. In both cases, we are asking about how the average of one variable changes across values of another.

4.1.1 Conditional Expectations

This idea of finding “the average value of y for a given value of x ” is exactly what we call the **conditional expectation**. Formally, the conditional expectation of y given x is written as:

$$E[Y|X = x]$$

This notation asks: if we knew that X took on some specific value x , what would we expect Y to be, on average? The conditional expectation is often the central quantity we want to model and estimate in econometrics.

4.1.2 Example: Education and Wages

Suppose X represents education level, which can take values $X = \{\text{hsdeg}, \text{colldeg}, \text{profdeg}\}$ (high school degree, college degree, professional degree), and Y represents hourly wage.

For each education level, we can compute the average wage among people in that group. This will give us a good idea of how wages are related to education.


Figure 4.1 is a scatterplot with an individual’s wage on the y-axis and their education level on the x-axis, with each point plotted in blue. The red points in Figure 4.1 show the conditional expectations—the average wage for individuals *within each education category*, i.e., $E[Y|X = HS]$, $E[Y|X = College]$, and $E[Y|X = Prof]$.

These averages tell us how we can expect wages to change when we increase education, which is precisely what we set out to measure.

4.2 The Population Regression Model

Now that we understand what we want to estimate (the conditional expectation), we need a framework for *how* to estimate it. Before we can estimate anything, we must first describe the underlying relationship we believe exists in the population.

Figure 4.1: Wages by education level. Each point represents an individual. Red points show the conditional expectation (average wage) for each education group.



4.2.1 The Bivariate Model

The simplest specification is called the **bivariate** (or simple) regression model:

$$y = \beta_0 + \beta_1 x + \mu \quad (4.1)$$

where:

- y is the **dependent variable** (also called the explained, response, or outcome variable)
- x is the **independent variable** (also called the explanatory, control, or predictor variable)
- μ is the **error term**, which represents all unobserved factors that affect y
- β_0 and β_1 are **population parameters**—the “true” values describing the relationship between x and y in the population

i The Error Term

The error term μ is crucial. It captures everything that affects y other than x . In a wage equation, μ might include ability, motivation, family connections, luck, and countless other factors we cannot observe or measure.

4.2.2 Connecting to Conditional Expectations

How does this regression model relate to our goal of estimating $E[Y|X]$? If we assume that the error term has mean zero conditional on x —that is, $E[\mu|x] = 0$ —then:

$$E[y|x] = E[\beta_0 + \beta_1 x + \mu|x] = \beta_0 + \beta_1 x + E[\mu|x] = \beta_0 + \beta_1 x$$

This is exactly the conditional expectation we set out to estimate! The regression line $\beta_0 + \beta_1 x$ traces out the average value of y for each value of x .

4.2.3 Interpreting the Coefficients

The coefficients in Equation 4.1 have specific interpretations:

- β_0 (**the intercept**): The predicted value of y when $x = 0$. Depending on the context, this may or may not have a meaningful interpretation.
- β_1 (**the slope**): The **marginal effect** of x on y . Mathematically, it is the derivative:

$$\frac{dy}{dx} = \beta_1$$

This tells us: when x increases by one unit, y changes by β_1 units, on average.

4.2.4 Example: Education and Wages

A regression model relating years of education to hourly wages might be:

$$wage = \beta_0 + \beta_1 \cdot educ + \mu$$

where *wage* is dollars per hour, *educ* is years of education, and μ captures all other determinants of wages (experience, ability, etc.).

If $\beta_1 = 2.50$, this would mean that each additional year of education is associated with \$2.50 higher hourly wages, on average.

4.3 The Regression Line: Predicted Values and Residuals

The entire goal of linear regression is to use our *sample* of the data from the wider *population* to estimate the values of β_0 and β_1 that are the best possible estimates we can generate. We call these estimates $\hat{\beta}_0$ and $\hat{\beta}_1$, and the method we use to calculate these estimates the **estimator**.

Before we derive estimators for β_0 and β_1 , let's introduce some key concepts that will guide our approach.

4.3.1 Fitted Values and the Regression Line

Once we have estimates $\hat{\beta}_0$ and $\hat{\beta}_1$, we can construct the **OLS regression line** (or fitted line):

$$\hat{y} = \hat{\beta}_0 + \hat{\beta}_1 x$$

For any value of x , this line gives us $\hat{y}$, the **predicted value** (or **fitted value**) of y . The regression line represents our best guess for the average value of y at each level of x .

4.3.2 Residuals: The Prediction Errors

Of course, our predictions won't be perfect. The **residual** for observation i is the difference between the actual value and the predicted value:

$$\hat{\mu}_i = y_i - \hat{y}_i = y_i - (\hat{\beta}_0 + \hat{\beta}_1 x_i)$$

The residual measures how much our prediction “misses” for each observation. Positive residuals mean we underpredicted; negative residuals mean we overpredicted.

4.3.3 A Numerical Example

Let's see these concepts with actual numbers. The table below shows the first six observations from a regression of MPG on horsepower, along with the predicted values and residuals:

Table 4.1: Example regression data showing actual values, predicted values, and residuals.

	x (Horsepower)	y (MPG)	$\hat{y}$ (Predicted)	$\hat{u}$ (Residual)
Mazda RX4	110	21.0	22.59	-1.59

Table 4.1: Example regression data showing actual values, predicted values, and residuals.

	x (Horsepower)	y (MPG)	$\hat{y}$ (Predicted)	$\hat{u}$ (Residual)
Mazda RX4 Wag	110	21.0	22.59	-1.59
Datsun 710	93	22.8	23.75	-0.95
Hornet 4 Drive	110	21.4	22.59	-1.19
Hornet Sportabout	175	18.7	18.16	0.54
Valiant	105	18.1	22.93	-4.83

Notice how the residual column equals the actual value minus the predicted value. For the first observation, the car has 110 horsepower and gets 21 MPG. Our regression predicts it should get about 22.59 MPG, so the residual is $21 - 22.59 = -1.59$. The negative residual means we *overpredicted*—the car actually gets slightly worse fuel economy than our model predicted.

4.3.4 The Key Insight: Minimize the Residuals

Here’s the crucial idea: we want to choose $\hat{\beta}_0$ and $\hat{\beta}_1$ so that our predictions are as good as possible. In other words, we want to make the residuals as small as possible.

But we can’t simply minimize the sum of residuals $\sum \hat{\mu}_i$, because positive and negative residuals would cancel out. Instead, we minimize the **Sum of Squared Residuals (SSR)**:


$$SSR = \sum_{i=1}^n \hat{\mu}_i^2 = \sum_{i=1}^n (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i)^2$$

This is exactly where the name “Ordinary Least Squares” comes from: we find the estimates that make the sum of squared residuals as small as possible.

4.4 Deriving the OLS Estimator

Now we turn to the mechanics of estimation. The **population** regression model describes the true relationship, but we never observe the entire population or the true values of β_0 and β_1 . Instead, we have a **sample** of data and must *estimate* the population parameters.

Figure 4.2: Components of the regression line. Black points show actual values, the blue line shows predicted values, and the salmon segment illustrates a residual.



4.4.1 The Minimization Problem

Suppose we have a sample of n observations, $\{(x_i, y_i) : i = 1, 2, \dots, n\}$. As discussed above, we want to find the values of $\hat{\beta}_0$ and $\hat{\beta}_1$ that minimize the sum of squared residuals:

$$\min_{\hat{\beta}_0, \hat{\beta}_1} \sum_{i=1}^n (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i)^2$$

This is a calculus optimization problem! The SSR is a function of two unknowns ($\hat{\beta}_0$ and $\hat{\beta}_1$), and we want to find the values that minimize it.

4.4.2 First-Order Conditions

To find the minimum, we take partial derivatives with respect to each parameter, set them equal to zero, and solve. These are called the **first-order conditions**.

Partial derivative with respect to $\hat{\beta}_0$:

$$\frac{\partial SSR}{\partial \hat{\beta}_0} = -2 \sum_{i=1}^n (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i) = 0$$

Dividing by -2 and rearranging:

$$\sum_{i=1}^n (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i) = 0$$

Partial derivative with respect to $\hat{\beta}_1$:

$$\frac{\partial SSR}{\partial \hat{\beta}_1} = -2 \sum_{i=1}^n x_i (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i) = 0$$

Dividing by -2 :

$$\sum_{i=1}^n x_i (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i) = 0$$

4.4.3 Solving for the Intercept

From the first condition, distributing the summation:

$$\sum_{i=1}^n y_i - n\hat{\beta}_0 - \hat{\beta}_1 \sum_{i=1}^n x_i = 0$$

Dividing by n and recognizing sample means:

$$\bar{y} = \hat{\beta}_0 + \hat{\beta}_1 \bar{x}$$

Solving for the intercept:

$$\hat{\beta}_0 = \bar{y} - \hat{\beta}_1 \bar{x} \tag{4.2}$$

This tells us that the regression line always passes through the point $(\bar{x}, \bar{y})$ —the “center” of the data.

4.4.4 Solving for the Slope

Substituting Equation 4.2 into the second first-order condition and working through the algebra (using summation properties from Appendix B):

$$\sum_{i=1}^n x_i [y_i - (\bar{y} - \hat{\beta}_1 \bar{x}) - \hat{\beta}_1 x_i] = 0$$

After distributing and rearranging:

$$\sum_{i=1}^n x_i (y_i - \bar{y}) = \hat{\beta}_1 \sum_{i=1}^n x_i (x_i - \bar{x})$$

Using the summation properties $\sum x_i (y_i - \bar{y}) = \sum (x_i - \bar{x})(y_i - \bar{y})$ and $\sum x_i (x_i - \bar{x}) = \sum (x_i - \bar{x})^2$:

$$\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y}) = \hat{\beta}_1 \sum_{i=1}^n (x_i - \bar{x})^2$$

Solving for $\hat{\beta}_1$:

$$\hat{\beta}_1 = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sum_{i=1}^n (x_i - \bar{x})^2} \quad (4.3)$$

4.4.5 Confirming It's a Minimum

How do we know this critical point is a minimum rather than a maximum? We check the second-order conditions. The second partial derivative of SSR with respect to $\hat{\beta}_1$ is:

$$\frac{\partial^2 SSR}{\partial \hat{\beta}_1^2} = 2 \sum_{i=1}^n x_i^2 > 0$$

Since this is positive (as long as there's variation in x), the SSR function is convex, and our solution is indeed a minimum.

! The OLS Slope Formula

The OLS slope estimate equals the sample covariance of x and y divided by the sample variance of x :

$$\hat{\beta}_1 = \frac{Cov(x, y)}{Var(x)} = \frac{\sum (x_i - \bar{x})(y_i - \bar{y})}{\sum (x_i - \bar{x})^2}$$

This can also be written in terms of the correlation coefficient:

$$\hat{\beta}_1 = \rho_{xy} \frac{\sigma_y}{\sigma_x}$$

This last expression reveals something important: bivariate OLS is essentially a scaled version of the correlation between x and y . And as we discussed in Chapter 1, correlation does not imply causation! We must be very careful about interpreting bivariate OLS results causally.

4.5 Computing OLS in R

While we derived the formulas by hand, in practice we use software to compute OLS estimates. In R, the `lm()` function (for “linear model”) performs OLS regression.

4.5.1 Basic Syntax

```
# General syntax
lm(y ~ x, data = your_data)
```

The formula $y \sim x$ specifies that y is the dependent variable and x is the independent variable. An intercept is included by default.

4.5.2 Example: MPG and Horsepower

Let’s estimate the relationship between fuel efficiency (MPG) and engine horsepower using the built-in `mtcars` dataset:

```
# Load the data
data(mtcars)

# Estimate the regression
lm_mpg <- lm(mpg ~ hp, data = mtcars)

# View the results
summary(lm_mpg)
```

```

Call:
lm(formula = mpg ~ hp, data = mtcars)

Residuals:
    Min       1Q   Median       3Q      Max
-5.7121 -2.1122 -0.8854  1.5819  8.2360

Coefficients:
              Estimate Std. Error t value Pr(>|t|)
(Intercept)  30.09886    1.63392   18.421  < 2e-16 ***
hp          -0.06823    0.01012   -6.742 1.79e-07 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 3.863 on 30 degrees of freedom
Multiple R-squared:  0.6024,    Adjusted R-squared:  0.5892
F-statistic: 45.46 on 1 and 30 DF,  p-value: 1.788e-07

```

The output shows our estimates: $\hat{\beta}_0 = 30.10$ and $\hat{\beta}_1 = -0.07$. The slope suggests that each additional unit of horsepower is associated with about 0.07 fewer miles per gallon.

4.6 Goodness-of-Fit: R-Squared

How well does our regression line fit the data? The **R-squared** (R^2) provides one measure of goodness-of-fit.

Define:

- **Total Sum of Squares (SST):** $SST = \sum_{i=1}^n (y_i - \bar{y})^2$ — total variation in y
- **Explained Sum of Squares (SSE):** $SSE = \sum_{i=1}^n (\hat{y}_i - \bar{y})^2$ — variation explained by the regression
- **Residual Sum of Squares (SSR):** $SSR = \sum_{i=1}^n \hat{\mu}_i^2$ — unexplained variation

The R-squared is:

$$R^2 = \frac{SSE}{SST} = 1 - \frac{SSR}{SST}$$

This measures the fraction of the total variation in y that is explained by the regression. R^2 ranges from 0 (no explanatory power) to 1 (perfect fit).

Interpreting R-Squared

Be careful interpreting R^2 , especially in the context of causal inference. A high R^2 does not mean we have a causal estimate—it just means x is correlated with y . Conversely, a low R^2 does not mean our estimate is wrong or unimportant. In cross-sectional data, R^2 values of 0.1 to 0.3 are common and can still represent economically meaningful relationships.

4.7 The Gauss-Markov Assumptions

We've derived the OLS formulas, but how do we know if the estimates are any good? The formulas will always give us *some* numbers, but whether those numbers have useful statistical properties depends on certain conditions being met.

The **Gauss-Markov assumptions** are a set of conditions under which OLS has desirable properties. Understanding these assumptions is crucial: they tell us when we can trust our estimates and when we should be skeptical.

4.7.1 Assumption 1: Linear in Parameters

Gauss-Markov Assumption 1: Linear in Parameters

The population model is: $y = \beta_0 + \beta_1 x + \mu$

This assumption states that the dependent variable y can be written as a linear function of the parameters β_0 and β_1 , plus an error term.

What this does NOT mean: This assumption is less restrictive than it might first appear. It does *not* require that y and x have a linear relationship. We can model many non-linear relationships by transforming the variables. For example:

- $y = \beta_0 + \beta_1 x^2 + \mu$ is linear in parameters (we can define $z = x^2$ and estimate $y = \beta_0 + \beta_1 z + \mu$)
- $\log(y) = \beta_0 + \beta_1 x + \mu$ is linear in parameters
- $y = \beta_0 + \beta_1 \log(x) + \mu$ is linear in parameters

What would violate this assumption: Models where parameters enter non-linearly, such as $y = \beta_0 + x^{\beta_1} + \mu$ or $y = \frac{1}{1+e^{-\beta_0-\beta_1 x}} + \mu$ (a logistic function). These require different estimation techniques.

4.7.2 Assumption 2: Random Sampling

i Gauss-Markov Assumption 2: Random Sampling

We have a random sample of size n from the population: $\{(x_i, y_i) : i = 1, \dots, n\}$

This assumption states that each observation in our dataset is drawn randomly and independently from the population of interest. Each unit in the population has an equal chance of being selected, and the selection of one unit doesn't affect the probability of selecting another.

Why it matters: Random sampling ensures that our sample is representative of the population. If certain types of observations are more likely to be included, our estimates may not generalize to the full population.

When this might be violated:

- **Self-selection:** If people choose whether to participate in a survey, those who respond may differ systematically from those who don't. For example, a wage survey might oversample high earners if low-wage workers are less likely to respond.
- **Convenience sampling:** Surveying only people who are easy to reach (e.g., college students for a study about the general population).
- **Time series data:** Observations collected over time are often correlated with each other. Today's stock price depends on yesterday's stock price, violating independence.
- **Clustered data:** Students within the same classroom, or workers within the same firm, may be more similar to each other than to randomly selected individuals.

4.7.3 Assumption 3: Variation in the Independent Variable

i Gauss-Markov Assumption 3: Sample Variation in x

The independent variable has some variation in the sample: $\sum(x_i - \bar{x})^2 > 0$

This assumption simply requires that x is not constant in our sample. If everyone in our sample has the same value of x , we cannot estimate how changes in x relate to changes in y .

Why it matters: Look back at our formula for $\hat{\beta}_1$:

$$\hat{\beta}_1 = \frac{\sum(x_i - \bar{x})(y_i - \bar{y})}{\sum(x_i - \bar{x})^2}$$

If all x_i equal $\bar{x}$, the denominator is zero, and the formula is undefined. Intuitively, if everyone has 12 years of education, we have no way to estimate what would happen with 13 years of education.

In practice: This assumption is almost always satisfied. It would only fail in unusual circumstances, like if you accidentally collected a sample where everyone happened to have the exact same value of x . However, having *little* variation in x (even if not zero) can lead to imprecise estimates, as we'll see when discussing variance.

4.7.4 Assumption 4: Zero Conditional Mean

i Gauss-Markov Assumption 4: Zero Conditional Mean

The expected value of the error term is zero for any value of x : $E[\mu|x] = 0$

This is the most important and most frequently violated assumption. It states that the average value of all the unobserved factors affecting y is the same regardless of the value of x .

Understanding the assumption: Let's break this down carefully. The error term μ contains everything that affects y other than x . The zero conditional mean assumption says that if we could collect all people with $x = 10$ and compute the average of their μ values, it would be zero. And if we collected all people with $x = 20$ and averaged their μ values, that would also be zero. The average of unobservables is the same at every level of x .

Example: Education and Wages

Consider the regression:

$$wage = \beta_0 + \beta_1 \cdot educ + \mu$$

The error term μ includes ability, motivation, family connections, and other unobserved factors. The zero conditional mean assumption requires:

$$E[\mu|educ = 8] = E[\mu|educ = 12] = E[\mu|educ = 16] = 0$$

Is this plausible? Probably not. People with higher ability tend to get more education (because school is easier for them, or because they enjoy learning). So among people with 16 years of education, average ability is probably higher than among people with 8 years. This means $E[\mu|educ = 16] > E[\mu|educ = 8]$, violating the assumption.

Why it matters: When this assumption fails, OLS is **biased**. The estimate $\hat{\beta}_1$ will systematically overshoot or undershoot the true β_1 . In the education example, if high-ability people get more education, our estimate of the “return to education” will partly capture the “return to ability,” making education look more valuable than it really is.

This is the essence of **omitted variable bias**, which we'll explore in detail shortly.

4.7.5 Assumption 5: Homoskedasticity

Gauss-Markov Assumption 5: Homoskedasticity

The variance of the error term is constant across all values of x : $Var(\mu|x) = \sigma^2$

This assumption states that the spread of the error term is the same regardless of the value of x . The “noise” in our data is equally noisy everywhere.

Understanding the assumption: While Assumption 4 concerns the *mean* of μ at different values of x , Assumption 5 concerns the *variance*. Homoskedasticity (from Greek: “same scatter”) means the scatter of points around the regression line is constant. **Heteroskedasticity** (different scatter) occurs when the variance of μ depends on x .

Example: Income and Spending

Consider a regression of consumption spending on income. For low-income households, spending is relatively constrained—there’s not much room for variation because most income goes to necessities. For high-income households, there’s much more discretion in spending patterns, leading to greater variation. This would create heteroskedasticity: $Var(\mu|income)$ increases with income.

```
`geom_smooth()` using formula = 'y ~ x'
`geom_smooth()` using formula = 'y ~ x'
```

Why it matters: Unlike Assumption 4, violating homoskedasticity does **not** make OLS biased. Your estimate of β_1 is still centered on the true value. However, heteroskedasticity affects the *variance* and *standard errors* of the estimator. The usual standard error formulas become incorrect, which can lead to invalid hypothesis tests and confidence intervals.

We’ll discuss how to handle heteroskedasticity later in the course using **robust standard errors**.

4.7.6 Summary: Which Assumptions Matter for What?

Assumption	Needed for Unbiasedness?	Needed for Valid Standard Errors?
1. Linear in parameters	Yes	Yes

Assumption	Needed for Unbiasedness?	Needed for Valid Standard Errors?
2. Random sampling	Yes	Yes
3. Variation in x	Yes (otherwise undefined)	Yes
4. Zero conditional mean	Yes	Yes
5. Homoskedasticity	No	Yes

The key insight: Assumptions 1-4 are required for OLS to be **unbiased** (on average, we hit the target). Assumption 5 is additionally required for the usual standard error formulas to be valid and for OLS to be **BLUE** (Best Linear Unbiased Estimator).

4.8 Unbiasedness of OLS

Before discussing unbiasedness, let's revisit an important distinction we introduced in Appendix B: the difference between the **population** and the **sample**.

4.8.1 Population Parameters vs. Sample Estimates

The **population** represents the entire group we're interested in studying. In our wage-education example, the population might be all workers in the U.S. economy. The population regression model describes the *true* relationship in this population:


$$y = \beta_0 + \beta_1 x + \mu$$

The parameters β_0 and β_1 are **population parameters**—fixed, unknown constants that describe how x and y are related for everyone in the population. We never observe these true values directly.

Instead, we collect a **sample**—a subset of observations from the population. Using this sample, we compute **estimates** $\hat{\beta}_0$ and $\hat{\beta}_1$. These estimates are our best guesses for the unknown population parameters.

Here's the key insight: *if we drew a different sample, we would get different estimates*. Your classmate analyzing a different random sample of workers would compute different values of $\hat{\beta}_0$ and $\hat{\beta}_1$ than you would. This randomness in the estimates is called **sampling variability**.

Figure 4.3: Left: Homoskedasticity—the spread of points around the line is constant. Right: Heteroskedasticity—the spread increases with x .



4.8.2 What Does Unbiasedness Mean?

Given that estimates vary from sample to sample, how do we know if our estimation procedure is any good? One desirable property is **unbiasedness**.

An estimator $\hat{\theta}$ is **unbiased** if its expected value equals the true population parameter:

$$E[\hat{\theta}] = \theta$$

Unbiasedness does *not* mean that any particular estimate exactly equals the truth. Rather, it means that if we could:

1. Draw a random sample from the population
2. Compute $\hat{\beta}_1$ from that sample
3. Repeat steps 1-2 many, many times
4. Average all the $\hat{\beta}_1$ values we computed

...then that average would equal the true β_1 .

In other words, our estimation procedure doesn't systematically overshoot or undershoot the target. Individual estimates might be too high or too low, but on average, we get it right.

4.8.3 Proof Sketch

Starting from our formula for $\hat{\beta}_1$ and substituting the true model $y_i = \beta_0 + \beta_1 x_i + \mu_i$:

$$\hat{\beta}_1 = \beta_1 + \frac{\sum_{i=1}^n (x_i - \bar{x})\mu_i}{\sum_{i=1}^n (x_i - \bar{x})^2}$$

Taking expectations and using the zero conditional mean assumption ($E[\mu_i|x_i] = 0$):

$$E[\hat{\beta}_1] = \beta_1 + \frac{\sum_{i=1}^n (x_i - \bar{x}) \cdot 0}{\sum_{i=1}^n (x_i - \bar{x})^2} = \beta_1$$

Thus, OLS is unbiased when the Gauss-Markov assumptions hold.

4.8.4 Simulation: Unbiasedness in Action

Show Code: Unbiasedness Simulation

```
# Create a "population" with known parameters
set.seed(1248)
n_pop <- 1000
x_pop <- sample(1:100, size = n_pop, replace = TRUE)
err_pop <- rnorm(n_pop)

# True model:  $y = 3.5 + 2.21x + \text{error}$ 
pop_data <- tibble(
  x = x_pop,
  y = 3.5 + 2.21 * x + err_pop
)

# Draw many samples and estimate beta_1 each time
set.seed(812476)
n_simulations <- 500
sample_size <- 25

sim_results <- map_dfr(1:n_simulations, function(i) {
  # Draw a random sample
  sample_data <- slice_sample(pop_data, n = sample_size, replace = TRUE)


  # Estimate OLS
  reg <- lm(y ~ x, data = sample_data)

  tibble(
    iteration = i,
    b1_hat = coef(reg)[2]
  )
})

# Calculate the average of all estimates
mean_b1 <- mean(sim_results$b1_hat)
```

The simulation in Figure 4.4 confirms the theory: while individual estimates vary around the true value, the average across many samples equals the true parameter.

Figure 4.4: Distribution of OLS slope estimates across 500 samples. The green line shows the average of all estimates, which equals the true value (2.21), demonstrating unbiasedness.



4.9 Omitted Variable Bias

What happens when the zero conditional mean assumption fails? The most common culprit is **omitted variable bias**: we’ve left out a variable that affects both x and y .

Consider the true model:

$$y = \beta_0 + \beta_1 x + \beta_2 z + \mu$$


If we omit z and estimate only $y = \beta_0 + \beta_1 x + \nu$ (where $\nu = \beta_2 z + \mu$), then whenever z is correlated with x , we have $E[\nu|x] \neq 0$, and our estimate of β_1 is biased.

Example: In the education-wage regression, if ability (z) affects both education choices (x) and wages (y), then omitting ability creates bias. The estimated “return to education” will partly reflect the return to ability.

4.10 Unbiasedness and Causality: Why This All Matters

Let’s step back and connect what we’ve learned to the bigger picture. In Chapter 3, we discussed how the fundamental challenge of causal inference is selection bias: people who receive “treatment” (more education, a new drug, a policy intervention) may differ systematically

Figure 4.5: With omitted variable bias, the average of OLS estimates (green) systematically differs from the true value (red). The bias persists even with many samples.



from those who don't. This makes it hard to know whether differences in outcomes are caused by the treatment or by those pre-existing differences.

Now we can see the same problem through the lens of regression. **The zero conditional mean assumption is essentially a “no selection bias” assumption.**

4.10.1 When Can We Interpret $\hat{\beta}_1$ Causally?

Consider our wage regression:

$$wage = \beta_0 + \beta_1 \cdot educ + \mu$$

When $E[\mu|educ] = 0$, our OLS estimate $\hat{\beta}_1$ is unbiased for β_1 . But what does β_1 actually represent?

If the zero conditional mean assumption holds, then β_1 represents the **causal effect** of education on wages: how much wages would change if we could take a person and give them one more year of education, holding everything else constant. In this case, $\hat{\beta}_1$ is an unbiased estimate of this causal effect.

But if $E[\mu|educ] \neq 0$ —say, because high-ability people get more education—then β_1 no longer represents a pure causal effect. It conflates the effect of education with the effect of ability. Our estimate $\hat{\beta}_1$ is biased, and we cannot interpret it as the causal effect of education.

4.10.2 Correlation vs. Causation, Revisited

Recall that the OLS slope can be written as:

$$\hat{\beta}_1 = \rho_{xy} \frac{\sigma_y}{\sigma_x}$$

Bivariate OLS is fundamentally measuring the *correlation* between x and y , scaled by their relative standard deviations. Correlation becomes causation **only when the Gauss-Markov assumptions hold**—especially the zero conditional mean assumption.

This is why we emphasize these assumptions so heavily. Without them:

- OLS still gives you numbers
- Those numbers still describe the correlation in your data
- But you **cannot** interpret those numbers as causal effects

The Key Insight

OLS always estimates *something*. Whether that something is a causal effect or merely a correlation depends entirely on whether the assumptions are satisfied. Most of econometrics is about finding clever ways to make the zero conditional mean assumption more plausible.

4.11 Variance of the OLS Estimator

Even when OLS is unbiased, our estimates vary from sample to sample. The **variance** of $\hat{\beta}_1$ tells us how much variation to expect.

Under the Gauss-Markov assumptions:

$$Var(\hat{\beta}_1) = \frac{\sigma^2}{SST_x}$$

where $\sigma^2 = Var(\mu)$ is the error variance and $SST_x = \sum (x_i - \bar{x})^2$ is the total variation in x .

This formula reveals two key insights:

1. **More noise (σ^2) means more variance:** If there's a lot of unexplained variation in y , our estimates are less precise.
2. **More variation in x means less variance:** Spreading out our observations of x gives us more information and more precise estimates. This also implies that larger samples (which have more total variation) yield more precise estimates.

4.12 Standard Errors

The **standard error** of $\hat{\beta}_1$ is the square root of the variance:

$$se(\hat{\beta}_1) = \frac{\hat{\sigma}}{\sqrt{SST_x}}$$

where $\hat{\sigma} = \sqrt{\frac{SSR}{n-2}}$ is the estimated standard deviation of the errors.

Standard errors quantify the precision of our estimates and are essential for hypothesis testing and confidence intervals (which we'll cover in the next chapter).

```
# R reports standard errors automatically
summary(lm_mpg)
```

Call:

```
lm(formula = mpg ~ hp, data = mtcars)
```

Residuals:

Min	1Q	Median	3Q	Max
-5.7121	-2.1122	-0.8854	1.5819	8.2360

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	30.09886	1.63392	18.421	< 2e-16 ***
hp	-0.06823	0.01012	-6.742	1.79e-07 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 3.863 on 30 degrees of freedom

Multiple R-squared: 0.6024, Adjusted R-squared: 0.5892

F-statistic: 45.46 on 1 and 30 DF, p-value: 1.788e-07

The standard error of $\hat{\beta}_1$ is 0.01, shown in the Std. Error column.

4.13 OLS is BLUE

Under all five Gauss-Markov assumptions, OLS is **BLUE**:

- **Best:** Lowest variance among all linear unbiased estimators
- **Linear:** The estimator is a linear function of the y_i 's
- **Unbiased:** $E[\hat{\beta}] = \beta$
- **Estimator:** It's a rule for using sample data to estimate population parameters

This is a powerful result: among all estimators that are linear and unbiased, OLS has the smallest variance. You cannot do better without either using non-linear methods or accepting some bias.

4.14 Summary and Conclusion

This chapter introduced Ordinary Least Squares regression, the foundational tool of econometrics. We started with the goal of modeling how the conditional expectation $E[Y|X]$ varies with x , and showed how the linear regression model $y = \beta_0 + \beta_1 x + \mu$ provides a framework for this.

We derived the OLS estimator by minimizing the sum of squared residuals, learned to compute regressions in R, and explored the properties of the OLS regression line. We then turned to the statistical properties of OLS, introducing the Gauss-Markov assumptions and proving that OLS is unbiased when these assumptions hold. When the zero conditional mean assumption fails—typically due to omitted variables—our estimates become biased.

Crucially, we connected unbiasedness to causality: the zero conditional mean assumption is what allows us to interpret OLS coefficients as causal effects rather than mere correlations. Without this assumption, OLS still produces numbers, but those numbers cannot be given a causal interpretation.

Finally, we discussed the variance of OLS estimates and the conditions under which OLS is the best linear unbiased estimator.

4.14.1 Key Takeaways

- **The goal** is to estimate the conditional expectation $E[Y|X] = \beta_0 + \beta_1 x$
- **The OLS slope** equals the covariance of x and y divided by the variance of x :

$$\hat{\beta}_1 = \frac{Cov(x, y)}{Var(x)}$$

- **Bivariate OLS is essentially correlation**—be very careful about causal interpretation
- **The Gauss-Markov assumptions** (especially zero conditional mean) are required for unbiasedness

- **Omitted variable bias** arises when we exclude a variable correlated with both x and y
- **Causality requires unbiasedness**—only when $E[\mu|x] = 0$ can we interpret $\hat{\beta}_1$ as a causal effect
- **OLS is BLUE** under the Gauss-Markov assumptions—best among linear unbiased estimators

4.15 Check Your Understanding

4.15.1 Practice: Interpreting Regression Output

The following questions present regression results for you to interpret.

Show Explanation

- The slope coefficient β_1 is the marginal effect—it tells us how much y changes when x increases by one unit. The intercept β_0 gives the predicted value when $x = 0$.
- The zero conditional mean assumption $E[\mu|x] = 0$ states that the expected value of the error term is zero for any given value of x . This is crucial for unbiasedness.
- The OLS slope equals the sample covariance of x and y divided by the sample variance of x . This follows directly from the derivation.
- Unbiasedness means that if we could repeat the sampling process many times, the *average* of our estimates would equal the true parameter. Individual estimates can still miss the target.
- Omitted variable bias occurs when we leave out a variable that belongs in the model and is correlated with our included independent variable. This causes $E[\mu|x] \neq 0$.
- The zero conditional mean assumption ($E[\mu|x] = 0$) is needed to prove OLS is unbiased, but it is not needed to derive the OLS formulas themselves. The calculus minimization approach works regardless of this assumption—it just tells us which values minimize squared residuals.
- As sample size increases, we get more variation in x , so $SST = \sum (x_i - \bar{x})^2$ increases. Since this appears in the denominator, the variance of our estimate decreases. Larger samples give more precise estimates.

Regression Interpretation Questions:

- viii. The coefficient on `str` is -2.28 , meaning each additional student per teacher is associated with 2.28 fewer points. For 5 additional students: $5 \times (-2.28) = -11.4$ points.
- ix. Using the regression equation: $\hat{y} = 698.93 + (-2.28)(20) = 698.93 - 45.6 = 653.33$
- x. R-squared measures the fraction of variance in y explained by the regression. Here, 5.1% of test score variation is explained by class size. A low R-squared doesn't make the coefficient meaningless—it just means other factors also matter.
- xi. The coefficient is -1.04 , so a 10-cent increase predicts: $10 \times (-1.04) = -10.4$ packs per capita.
- xii. The negative relationship between price and quantity is exactly what economic theory predicts—the law of demand.
- xiii. States with strong anti-smoking policies might have both higher cigarette taxes (raising prices) AND other policies that reduce smoking. This would create omitted variable bias, making price appear more effective than it actually is at reducing consumption.

⋮⋮⋮

Note: This section contains interactive content only available in the HTML version.

5 Multivariate Regression

Key Questions

- Why do we need multivariate regression?
- How do we interpret coefficients when there are multiple independent variables?
- What is the zero conditional mean assumption in the multivariate case?
- How does multivariate OLS function as a “matching” estimator?
- What is the omitted variable bias formula and how do we use it?
- What is the bias-variance tradeoff when deciding which controls to include?

Suggested Readings

- Wooldridge (2019), Ch. 3

In Chapter 4, we introduced bivariate OLS and saw that causal interpretation requires the zero conditional mean assumption: $E[\mu|x] = 0$. We also saw that this assumption is frequently violated due to omitted variables—factors in the error term that are correlated with both x and y .

This chapter introduces **multivariate OLS**, which allows us to explicitly control for confounding variables. By including additional variables on the right-hand side of our regression, we can make the zero conditional mean assumption more plausible and get closer to causal estimates.

5.1 The Problem with Bivariate OLS

Let’s revisit why bivariate OLS often fails to give us causal estimates.

Consider a regression of wages on education:

$$wage = \beta_0 + \beta_1 \cdot educ + \mu$$

For $\hat{\beta}_1$ to be an unbiased estimate of the causal effect of education, we need $E[\mu|educ] = 0$. But the error term μ contains everything else that affects wages: ability, family connections, motivation, and countless other factors.

Many of these factors are likely correlated with education. For example, people from wealthier families tend to get more education (better schools, tutors, college is affordable) *and* earn higher wages (family connections, inherited wealth). If family income is in μ and correlated with education, then $E[\mu|educ] \neq 0$, and our estimate is biased.

! The Core Problem

With bivariate OLS, the zero conditional mean assumption requires that *all* factors affecting y (other than x) be uncorrelated with x . This is almost never plausible in observational data.

5.2 The Multivariate Solution

Multivariate OLS offers a solution: **explicitly control** for variables that might be confounders.

If we're worried that family income confounds the education-wage relationship, we can include it in our model:

$$wage = \beta_0 + \beta_1 \cdot educ + \beta_2 \cdot family_income + \mu$$

Now β_1 represents the effect of education on wages **holding family income constant**. The error term μ no longer contains family income, so that source of bias is eliminated.

5.2.1 The General Multivariate Model

The general multivariate linear regression model with k independent variables is:

$$y = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \cdots + \beta_k x_k + \mu \tag{5.1}$$

where:

- y is the dependent variable
- $x_1, x_2, \dots, x_k$ are the independent variables
- β_0 is the intercept
- β_j is the coefficient on x_j (for $j = 1, 2, \dots, k$)
- μ is the error term containing all factors not included in the model

5.2.2 Interpreting Coefficients

In multivariate regression, each coefficient has a **ceteris paribus** interpretation:

$$\beta_j = \frac{\partial y}{\partial x_j}$$

This is the change in y associated with a one-unit increase in x_j , **holding all other independent variables constant**.

For example, in the model:

$$wage = \beta_0 + \beta_1 \cdot educ + \beta_2 \cdot family_income + \mu$$

- β_1 : The change in wages for one additional year of education, **holding family income constant**
- β_2 : The change in wages for one additional dollar of family income, **holding education constant**

5.2.3 The Zero Conditional Mean Assumption

In multivariate OLS, the zero conditional mean assumption becomes:

$$E[\mu|x_1, x_2, \dots, x_k] = 0$$

This states that the average of unobserved factors is zero for *all unique combinations* of the independent variables. In other words, everything in μ must be uncorrelated with *all* the variables we've included.

This assumption is much more plausible than its bivariate counterpart. By including relevant control variables, we've removed them from μ , making it more likely that what remains is uncorrelated with our variables of interest.

5.3 Deriving the Multivariate OLS Estimator

The derivation follows the same logic as bivariate OLS: we minimize the sum of squared residuals.

5.3.1 The Minimization Problem

Given a sample of n observations, we want to find values $\hat{\beta}_0, \hat{\beta}_1, \dots, \hat{\beta}_k$ that minimize:

$$SSR = \sum_{i=1}^n (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_{1i} - \hat{\beta}_2 x_{2i} - \dots - \hat{\beta}_k x_{ki})^2$$

5.3.2 First-Order Conditions

Taking partial derivatives with respect to each $\hat{\beta}_j$ and setting them equal to zero yields $k + 1$ first-order conditions:

$$\begin{aligned} \sum_{i=1}^n (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_{1i} - \dots - \hat{\beta}_k x_{ki}) &= 0 \\ \sum_{i=1}^n x_{1i} (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_{1i} - \dots - \hat{\beta}_k x_{ki}) &= 0 \\ &\vdots \\ \sum_{i=1}^n x_{ki} (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_{1i} - \dots - \hat{\beta}_k x_{ki}) &= 0 \end{aligned}$$

Solving this system of equations yields the OLS estimates. The algebra is tedious (and typically handled with matrix methods), so we won't derive the explicit formulas here. Fortunately, your computer does this effortlessly.

5.4 Estimating Multivariate OLS in R

In R, we estimate multivariate regression using the same `lm()` function, simply adding variables with the `+` symbol.

5.4.1 Example: MPG, Horsepower, and Weight

Let's estimate the effect of horsepower on fuel efficiency, first without and then with a control for vehicle weight.

```
# Load data
data(mtcars)

# Bivariate regression
lm_bivariate <- lm(mpg ~ hp, data = mtcars)

# Multivariate regression (adding weight as control)
lm_multivariate <- lm(mpg ~ hp + wt, data = mtcars)
```

Bivariate results:

```
summary(lm_bivariate)
```

Call:

```
lm(formula = mpg ~ hp, data = mtcars)
```

Residuals:

Min	1Q	Median	3Q	Max
-5.7121	-2.1122	-0.8854	1.5819	8.2360

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	30.09886	1.63392	18.421	< 2e-16 ***
hp	-0.06823	0.01012	-6.742	1.79e-07 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 3.863 on 30 degrees of freedom

Multiple R-squared: 0.6024, Adjusted R-squared: 0.5892

F-statistic: 45.46 on 1 and 30 DF, p-value: 1.788e-07

Multivariate results:

```
summary(lm_multivariate)
```

```

Call:
lm(formula = mpg ~ hp + wt, data = mtcars)

Residuals:
    Min       1Q   Median       3Q      Max
-3.941 -1.600 -0.182  1.050  5.854

Coefficients:
            Estimate Std. Error t value Pr(>|t|)
(Intercept) 37.22727    1.59879   23.285 < 2e-16 ***
hp          -0.03177    0.00903   -3.519  0.00145 **
wt          -3.87783    0.63273   -6.129 1.12e-06 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 2.593 on 29 degrees of freedom
Multiple R-squared:  0.8268,    Adjusted R-squared:  0.8148
F-statistic: 69.21 on 2 and 29 DF,  p-value: 9.109e-12

```

5.4.2 Interpreting the Results

In the **bivariate** regression, the coefficient on horsepower is -0.068 : each additional unit of horsepower is associated with 0.068 fewer miles per gallon.

In the **multivariate** regression (controlling for weight), the coefficient on horsepower drops to -0.032 —less than half the size!

What happened? Horsepower and weight are positively correlated (more powerful engines tend to be in heavier vehicles), and weight independently reduces fuel efficiency. The bivariate estimate was conflating the effect of horsepower with the effect of weight. Once we control for weight, we isolate the effect of horsepower alone.

What Changed?

The coefficient on `hp` dropped from -0.068 to -0.032 when we added `wt`. This suggests that much of what appeared to be the “effect” of horsepower was actually the effect of weight. This is precisely what controlling for confounders accomplishes.

5.5 Multivariate OLS as Matching

There's an intuitive way to understand what multivariate OLS does: it's essentially a **matching** estimator. When we “control for” a variable, we're comparing observations that have similar values of that control variable.

5.5.1 The Returns to Education Example

Suppose we want to estimate the effect of education on wages. A simple comparison of average wages across education levels might be misleading because people with more education tend to work in higher-paying industries. Part of the apparent “return to education” might actually be an “industry premium.”

Let's work through this with simulated data to see exactly how controlling for industry works.

```
# Simulate wage data
set.seed(42)
n <- 2000

# Create data where:
# - More educated workers are more likely to be in high-paying industries
# - Industry independently affects wages
wage_data <- tibble(
  # Education: 1 = HS, 2 = Some College, 3 = Bachelor's+
  educ_years = sample(c(12, 14, 16), n, replace = TRUE, prob = c(0.4, 0.3, 0.3)),

  # Higher education -> more likely in professional services (high pay industry)
  ind_professional = rbinom(n, 1, prob = case_when(
    educ_years == 12 ~ 0.20,
    educ_years == 14 ~ 0.40,
    educ_years == 16 ~ 0.65
  )),

  # Wages depend on education AND industry
  wage = 25000 +
    3000 * (educ_years - 12) +      # True education effect: $3000 per year
    15000 * ind_professional +      # Industry premium: $15000
    rnorm(n, 0, 8000)
) |>
mutate(
  educ_cat = case_when(
```

```

educ_years == 12 ~ "HS Only",
educ_years == 14 ~ "Some College",
educ_years == 16 ~ "Bachelor's+"
),
industry = ifelse(ind_professional == 1, "Professional", "Other")
)

```

5.5.2 The Naive Comparison

First, let's compute average wages by education level without any controls:

```

naive_means <- wage_data |>
  group_by(educ_cat, educ_years) |>
  summarise(avg_wage = mean(wage), n = n(), .groups = "drop") |>
  arrange(educ_years)

naive_means |>
  knitr::kable(
    col.names = c("Education", "Years", "Avg Wage ($)", "N"),
    digits = 0
  )

```

Education	Years	Avg Wage (\$)	N
HS Only	12	28575	829
Some College	14	36961	604
Bachelor's+	16	46341	567

The naive comparison suggests that going from high school to some college is associated with about \$8,386 higher wages, and going from high school to a bachelor's degree is associated with about \$17,766 higher wages.

But wait—the true effect of education in our simulation is only \$3,000 per year (so \$6,000 for 2 extra years and \$12,000 for 4 extra years). The naive estimates are too large! This is because more educated workers are more likely to be in high-paying industries.

5.5.3 Step 1: Compute Within-Industry Differences

To control for industry, we first compute average wages separately for each education-industry combination:

```

cell_means <- wage_data |>
  group_by(industry, educ_cat, educ_years) |>
  summarise(avg_wage = mean(wage), n = n(), .groups = "drop") |>
  arrange(industry, educ_years)

cell_means |>
  select(industry, educ_cat, avg_wage, n) |>
  knitr::kable(
    col.names = c("Industry", "Education", "Avg Wage ($)", "N"),
    digits = 0
  )

```

Table 5.2: Average wages by education and industry

Industry	Education	Avg Wage (\$)	N
Other	HS Only	25275	648
Other	Some College	30351	349
Other	Bachelor's+	36802	199
Professional	HS Only	40388	181
Professional	Some College	46008	255
Professional	Bachelor's+	51499	368

Now let's compute the wage differences **within** each industry:

```

within_industry <- cell_means |>
  group_by(industry) |>
  mutate(
    diff_from_hs = avg_wage - avg_wage[educ_years == 12]
  ) |>
  filter(educ_years > 12) |>
  select(industry, educ_cat, diff_from_hs, n)

within_industry |>
  knitr::kable(
    col.names = c("Industry", "Education", "Wage Diff vs HS ($)", "N in Cell"),
    digits = 0
  )

```

Industry	Education	Wage Diff vs HS (\$)	N in Cell
Other	Some College	5076	349
Other	Bachelor's+	11527	199
Professional	Some College	5620	255
Professional	Bachelor's+	11110	368

Notice that the within-industry differences are much closer to the true values (\$6,000 for Some College, \$12,000 for Bachelor's+).

5.5.4 Step 2: Compute Weights

To combine these within-industry estimates into a single number, we take a **weighted average**. The weights are based on the number of observations in each industry:

```
# Total workers in each industry
industry_totals <- wage_data |>
  count(industry) |>
  mutate(weight = n / sum(n))

industry_totals |>
  knitr::kable(
    col.names = c("Industry", "N", "Weight"),
    digits = 3
  )
```

Industry	N	Weight
Other	1196	0.598
Professional	804	0.402

5.5.5 Step 3: Compute Weighted Average

Now we combine the within-industry differences using these weights. Let's do this for the Bachelor's+ vs HS comparison:

```
# Get the within-industry differences for Bachelor's+
ba_diffs <- within_industry |>
  filter(educ_cat == "Bachelor's+") |>
  left_join(industry_totals, by = "industry")
```

```
ba_diffs |>
  select(industry, diff_from_hs, weight) |>
  knitr::kable(
    col.names = c("Industry", "Within-Industry Diff ($)", "Weight"),
    digits = c(0, 0, 3)
  )
```

Industry	Within-Industry Diff (\$)	Weight
Other	11527	0.598
Professional	11110	0.402

```
# Weighted average
weighted_avg <- sum(ba_diffs$diff_from_hs * ba_diffs$weight)
```

The weighted average is:

$$(1.1527 \times 10^4 \times 0.598) + (1.111 \times 10^4 \times 0.402) = 11,360$$

This is much closer to the true effect of \$12,000!

5.5.6 The Regression Equivalent

Now let's see what regression gives us:

```
# Create numeric education variable (years beyond HS)
wage_data <- wage_data |>
  mutate(educ_beyond_hs = educ_years - 12)

# Regression controlling for industry
lm_controlled <- lm(wage ~ educ_beyond_hs + ind_professional, data = wage_data)
summary(lm_controlled)
```

Call:

```
lm(formula = wage ~ educ_beyond_hs + ind_professional, data = wage_data)
```

Residuals:

```
      Min       1Q   Median       3Q      Max
```

-25629.6 -5261.0 24.9 5383.0 27407.8

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	25179.3	275.7	91.33	<2e-16 ***
educ_beyond_hs	2797.6	117.3	23.84	<2e-16 ***
ind_professional	15180.2	394.9	38.44	<2e-16 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 8075 on 1997 degrees of freedom

Multiple R-squared: 0.6093, Adjusted R-squared: 0.6089

F-statistic: 1557 on 2 and 1997 DF, p-value: < 2.2e-16

The coefficient on `educ_beyond_hs` is approximately \$3,000 per year—almost exactly the true value we built into the simulation!

Compare this to the naive regression without the industry control:

```
lm_naive <- lm(wage ~ educ_beyond_hs, data = wage_data)
coef(lm_naive)["educ_beyond_hs"]
```

```
educ_beyond_hs
4427.053
```

The naive estimate (4,427) is inflated because it conflates the education effect with the industry premium.

i Multivariate OLS as Matching

When you include control variables, OLS is effectively:

1. **Grouping** observations by similar values of the control variables
2. **Computing** the relationship between x and y within each group
3. **Taking a weighted average** across groups, where weights reflect group sizes

This is why multivariate regression is sometimes called a “matching” estimator—we’re matching people who work in the same industry and comparing their wages based on education.

5.6 Goodness-of-Fit: Adjusted R-Squared

In bivariate OLS, we used R^2 to measure how much variation in y our model explains. The same concept applies to multivariate OLS, but with an important caveat.

5.6.1 The Problem with R-Squared

The R^2 can never decrease when you add more variables—it can only increase or stay the same. This is because adding variables can only improve (or maintain) the fit, even if those variables are irrelevant.

This creates a problem: R^2 mechanically increases as you add more variables, potentially making a bloated model look better than a parsimonious one.

5.6.2 The Adjusted R-Squared

The **adjusted R-squared** penalizes for the number of parameters:

$$\bar{R}^2 = 1 - \frac{(1 - R^2)(n - 1)}{n - k - 1}$$

where n is the sample size and k is the number of independent variables.

The adjusted R^2 :

- Is always smaller than the regular R^2
- Can decrease when you add irrelevant variables
- Provides a better comparison between models with different numbers of variables

```
# Compare R-squared and Adjusted R-squared
c(
  "Bivariate R-sq" = summary(lm_bivariate)$r.squared,
  "Bivariate Adj R-sq" = summary(lm_bivariate)$adj.r.squared,
  "Multivariate R-sq" = summary(lm_multivariate)$r.squared,
  "Multivariate Adj R-sq" = summary(lm_multivariate)$adj.r.squared
)
```

Bivariate R-sq	Bivariate Adj R-sq	Multivariate R-sq
0.6024373	0.5891853	0.8267855
Multivariate Adj R-sq		
0.8148396		

5.7 Gauss-Markov Assumptions for Multivariate OLS

The Gauss-Markov assumptions extend naturally to the multivariate case, with one important addition.

5.7.1 Assumption 1: Linear in Parameters

$$y = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \cdots + \beta_k x_k + \mu$$

As in the bivariate case, we can transform variables to model non-linear relationships while maintaining linearity in parameters.

5.7.2 Assumption 2: Random Sampling

We have a random sample $\{(x_{1i}, x_{2i}, \dots, x_{ki}, y_i) : i = 1, \dots, n\}$ from the population.

5.7.3 Assumption 3: No Perfect Multicollinearity

This assumption has two parts:

1. **No constant variables:** Each independent variable must have some variation in the sample
2. **No exact linear relationships:** No independent variable can be a perfect linear function of other independent variables

The second part is new and important in multivariate regression.

Perfect Multicollinearity

Perfect multicollinearity occurs when one independent variable is an exact linear combination of others. Examples:

- Including both `family_income` and `family_income_thousands` (one is just the other divided by 1000)
- Including `parent1_income`, `parent2_income`, and `family_income` when `family_income = parent1_income + parent2_income`
- Including dummy variables for all categories of a categorical variable plus an intercept

When perfect multicollinearity exists, the OLS formulas break down (the system of equations has no unique solution). R will typically drop one of the collinear variables automatically and issue a warning.

5.7.4 Assumption 4: Zero Conditional Mean

$$E[\mu|x_1, x_2, \dots, x_k] = 0$$

The error term has mean zero for all combinations of the independent variables. This can still fail due to:

- Omitted variables correlated with included variables
- Wrong functional form
- Measurement error in independent variables
- Simultaneous causality

5.7.5 Assumption 5: Homoskedasticity

$$Var(\mu|x_1, x_2, \dots, x_k) = \sigma^2$$

The variance of the error term is constant across all combinations of independent variable values.

5.8 The Omitted Variable Bias Formula

Even when we can't include an omitted variable (perhaps because it's unobservable), we can still reason about the **direction** of the bias.

5.8.1 Setup

Suppose the true model is:

$$y = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \mu$$

But we can't observe x_2 , so we estimate the misspecified model:

$$\tilde{y} = \tilde{\beta}_0 + \tilde{\beta}_1 x_1$$

where the tilde (̃) indicates estimates from the misspecified model.

5.8.2 The OVB Formula

It can be shown that:

$$\tilde{\beta}_1 = \hat{\beta}_1 + \hat{\beta}_2 \tilde{\delta}_1 \quad (5.2)$$

where $\tilde{\delta}_1$ is the coefficient from regressing x_2 on x_1 :

$$x_2 = \tilde{\delta}_0 + \tilde{\delta}_1 x_1 + \text{error}$$

Taking expectations:

$$E[\tilde{\beta}_1] = \beta_1 + \beta_2 \tilde{\delta}_1$$

Therefore, the **bias** is:

$$\text{Bias}(\tilde{\beta}_1) = \beta_2 \cdot \tilde{\delta}_1 \quad (5.3)$$

5.8.3 Interpreting the Bias Formula

The bias depends on two factors:

1. β_2 : The effect of the omitted variable on y
2. $\tilde{\delta}_1$: The correlation between the omitted variable (x_2) and the included variable (x_1)

If either factor is zero, there is no bias. But if both are non-zero, bias exists, and its direction depends on the signs:

	$\text{Corr}(x_1, x_2) > 0$	$\text{Corr}(x_1, x_2) < 0$
$\beta_2 > 0$	Positive bias (overestimate)	Negative bias (underestimate)
$\beta_2 < 0$	Negative bias (underestimate)	Positive bias (overestimate)

5.8.4 Example: Education, Wages, and Ability

Consider estimating the return to education while omitting ability:

- β_2 (**effect of ability on wages**): Positive (more able people earn more)
- $\tilde{\delta}_1$ (**correlation of ability with education**): Positive (more able people get more education)

Therefore:

$$\text{Bias} = (+)(+) = \text{Positive}$$

Our estimate of the return to education is **too large**—it includes part of the return to ability.

5.8.5 Example: Policing, Crime, and Poverty

Consider estimating the effect of policing on crime while omitting local poverty:

- **Effect of poverty on crime** (β_2): Positive (higher poverty $\rightarrow$ more crime)
- **Correlation of poverty with policing** ($\tilde{\delta}_1$): Positive (high-poverty areas get more policing)

Therefore:

$$\text{Bias} = (+)(+) = \text{Positive}$$

Our estimate suggests policing *increases* crime more than it actually does (or reduces it less than it actually does). This is why naive correlations between police presence and crime can be misleading.

5.9 Variance in Multivariate OLS

Under the Gauss-Markov assumptions, the variance of $\hat{\beta}_j$ is:

$$\text{Var}(\hat{\beta}_j) = \frac{\sigma^2}{SST_j(1 - R_j^2)} \quad (5.4)$$

where:

- σ^2 is the error variance
- $SST_j = \sum (x_{ji} - \bar{x}_j)^2$ is the total variation in x_j
- R_j^2 is the R-squared from regressing x_j on all other independent variables

5.9.1 What Increases Variance?

From Equation 5.4, the variance of $\hat{\beta}_j$ increases when:

1. **More noise (σ^2):** Higher error variance means less precise estimates
2. **Less variation in x_j (SST_j):** Less information about how x_j relates to y
3. **Higher R_j^2 :** More correlation between x_j and other independent variables

The third point is crucial: when your variable of interest is highly correlated with control variables, your estimate becomes less precise. This is sometimes called **imperfect multicollinearity** or the **multicollinearity problem**.

5.9.2 Standard Errors

The standard error is estimated as:

$$se(\hat{\beta}_j) = \frac{\hat{\sigma}}{\sqrt{SST_j(1 - R_j^2)}}$$

where $\hat{\sigma}^2 = \frac{SSR}{n-k-1}$ is the estimated error variance, and $n - k - 1$ represents the **degrees of freedom**.

5.10 The Bias-Variance Tradeoff

Should you include a variable in your regression? The answer depends on a tradeoff.

5.10.1 The Costs of Omitting a Relevant Variable

If a variable truly belongs in the model ($\beta_2 \neq 0$) and is correlated with x_1 , omitting it causes **bias**. Your estimate systematically misses the true value.

5.10.2 The Costs of Including an Irrelevant Variable

What if you include a variable that doesn't belong ($\beta_2 = 0$)?

From the OVB formula:

$$E[\hat{\beta}_1] = \beta_1 + (0) \cdot \tilde{\delta}_1 = \beta_1$$

No bias! Including an irrelevant variable doesn't bias your estimate.

But look at the variance formula. Including an extra variable increases R_1^2 (the correlation between x_1 and other variables), which increases the variance of $\hat{\beta}_1$.

Including irrelevant variables inflates variance without reducing bias.

5.10.3 The Tradeoff

Action	Bias	Variance
Omit relevant variable	Increases	Decreases
Include irrelevant variable	No change	Increases

This is the **bias-variance tradeoff**:

- If you're too conservative (omit too many variables), you risk bias
- If you're too liberal (include everything), you inflate variance

5.10.4 Practical Guidance

There's no perfect solution, but some principles help:

1. **Include variables you have strong theoretical reasons to believe are confounders.** Use economic theory and prior research to guide variable selection.
2. **Be cautious about highly correlated controls.** If two controls are very highly correlated with each other, including both can dramatically inflate variance.
3. **Larger samples help.** As n increases, variance decreases, making the "include it just in case" strategy less costly.
4. **Check how estimates change.** If adding a control substantially changes your coefficient of interest, that control is probably important.

5.11 Simulation: Bias vs. Variance

Let's see the bias-variance tradeoff in action with a simulation.

```
Warning in grid.Call(C_textBounds, as.graphicsAnnot(x$label), x$x, x$y, :  
conversion failure on 'Include x ' in 'mbcsToSbcs': dot substituted for <e2>
```

```
Warning in grid.Call(C_textBounds, as.graphicsAnnot(x$label), x$x, x$y, :  
conversion failure on 'Include x ' in 'mbcsToSbcs': dot substituted for <82>  
Warning in grid.Call(C_textBounds, as.graphicsAnnot(x$label), x$x, x$y, :  
conversion failure on 'Include x ' in 'mbcsToSbcs': dot substituted for <82>
```

```
Warning in grid.Call(C_textBounds, as.graphicsAnnot(x$label), x$x, x$y, :  
conversion failure on 'Omit x ' in 'mbcsToSbcs': dot substituted for <e2>
```

```
Warning in grid.Call(C_textBounds, as.graphicsAnnot(x$label), x$x, x$y, :  
conversion failure on 'Omit x ' in 'mbcsToSbcs': dot substituted for <82>  
Warning in grid.Call(C_textBounds, as.graphicsAnnot(x$label), x$x, x$y, :  
conversion failure on 'Omit x ' in 'mbcsToSbcs': dot substituted for <82>
```

```
Warning in grid.Call(C_textBounds, as.graphicsAnnot(x$label), x$x, x$y, :  
conversion failure on 'Include x ' in 'mbcsToSbcs': dot substituted for <e2>
```

```
Warning in grid.Call(C_textBounds, as.graphicsAnnot(x$label), x$x, x$y, :  
conversion failure on 'Include x ' in 'mbcsToSbcs': dot substituted for <82>  
Warning in grid.Call(C_textBounds, as.graphicsAnnot(x$label), x$x, x$y, :  
conversion failure on 'Include x ' in 'mbcsToSbcs': dot substituted for <82>
```

```
Warning in grid.Call(C_textBounds, as.graphicsAnnot(x$label), x$x, x$y, :  
conversion failure on 'Omit x ' in 'mbcsToSbcs': dot substituted for <e2>
```

```
Warning in grid.Call(C_textBounds, as.graphicsAnnot(x$label), x$x, x$y, :  
conversion failure on 'Omit x ' in 'mbcsToSbcs': dot substituted for <82>  
Warning in grid.Call(C_textBounds, as.graphicsAnnot(x$label), x$x, x$y, :  
conversion failure on 'Omit x ' in 'mbcsToSbcs': dot substituted for <82>
```

```
Warning in grid.Call(C_textBounds, as.graphicsAnnot(x$label), x$x, x$y, :  
conversion failure on 'Include x ' in 'mbcsToSbcs': dot substituted for <e2>
```

Warning in grid.Call(C_textBounds, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on 'Include x ' in 'mbcsToSbcs': dot substituted for <82>
Warning in grid.Call(C_textBounds, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on 'Include x ' in 'mbcsToSbcs': dot substituted for <82>

Warning in grid.Call(C_textBounds, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on 'Omit x ' in 'mbcsToSbcs': dot substituted for <e2>

Warning in grid.Call(C_textBounds, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on 'Omit x ' in 'mbcsToSbcs': dot substituted for <82>
Warning in grid.Call(C_textBounds, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on 'Omit x ' in 'mbcsToSbcs': dot substituted for <82>

Warning in grid.Call(C_textBounds, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on 'Include x ' in 'mbcsToSbcs': dot substituted for <e2>

Warning in grid.Call(C_textBounds, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on 'Include x ' in 'mbcsToSbcs': dot substituted for <82>
Warning in grid.Call(C_textBounds, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on 'Include x ' in 'mbcsToSbcs': dot substituted for <82>

Warning in grid.Call(C_textBounds, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on 'Omit x ' in 'mbcsToSbcs': dot substituted for <e2>

Warning in grid.Call(C_textBounds, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on 'Omit x ' in 'mbcsToSbcs': dot substituted for <82>
Warning in grid.Call(C_textBounds, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on 'Omit x ' in 'mbcsToSbcs': dot substituted for <82>

Warning in grid.Call(C_textBounds, as.graphicsAnnot(x\$label), x\$x, x\$y, : font
metrics unknown for Unicode character U+2260

Warning in grid.Call(C_textBounds, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on ' 0)' in 'mbcsToSbcs': dot substituted for <e2>

Warning in grid.Call(C_textBounds, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on ' 0)' in 'mbcsToSbcs': dot substituted for <89>

Warning in grid.Call(C_textBounds, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on ' 0)' in 'mbcsToSbcs': dot substituted for <a0>

Warning in grid.Call(C_textBounds, as.graphicsAnnot(x\$label), x\$x, x\$y, : font metrics unknown for Unicode character U+2260

Warning in grid.Call(C_textBounds, as.graphicsAnnot(x\$label), x\$x, x\$y, : conversion failure on ' 0)' in 'mbcsToSbcs': dot substituted for <e2>

Warning in grid.Call(C_textBounds, as.graphicsAnnot(x\$label), x\$x, x\$y, : conversion failure on ' 0)' in 'mbcsToSbcs': dot substituted for <89>

Warning in grid.Call(C_textBounds, as.graphicsAnnot(x\$label), x\$x, x\$y, : conversion failure on ' 0)' in 'mbcsToSbcs': dot substituted for <a0>

Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, : conversion failure on 'True ' in 'mbcsToSbcs': dot substituted for <ce>

Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, : conversion failure on 'True ' in 'mbcsToSbcs': dot substituted for <b2>

Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, : conversion failure on 'True ' in 'mbcsToSbcs': dot substituted for <e2>

Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, : conversion failure on 'True ' in 'mbcsToSbcs': dot substituted for <82>

Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, : conversion failure on 'True ' in 'mbcsToSbcs': dot substituted for <81>

Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, : conversion failure on 'Include x ' in 'mbcsToSbcs': dot substituted for <e2>

Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, : conversion failure on 'Include x ' in 'mbcsToSbcs': dot substituted for <82>

Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, : conversion failure on 'Include x ' in 'mbcsToSbcs': dot substituted for <82>

Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, : conversion failure on 'Omit x ' in 'mbcsToSbcs': dot substituted for <e2>

Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on 'Omit x ' in 'mbcsToSbcs': dot substituted for <82>
Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on 'Omit x ' in 'mbcsToSbcs': dot substituted for <82>

Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on 'True ' in 'mbcsToSbcs': dot substituted for <ce>

Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on 'True ' in 'mbcsToSbcs': dot substituted for <b2>

Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on 'True ' in 'mbcsToSbcs': dot substituted for <e2>

Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on 'True ' in 'mbcsToSbcs': dot substituted for <82>

Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on 'True ' in 'mbcsToSbcs': dot substituted for <81>

Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on 'Include x ' in 'mbcsToSbcs': dot substituted for <e2>

Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on 'Include x ' in 'mbcsToSbcs': dot substituted for <82>
Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on 'Include x ' in 'mbcsToSbcs': dot substituted for <82>

Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on 'Omit x ' in 'mbcsToSbcs': dot substituted for <e2>

Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on 'Omit x ' in 'mbcsToSbcs': dot substituted for <82>
Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on 'Omit x ' in 'mbcsToSbcs': dot substituted for <82>

Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, :
font metrics unknown for Unicode character U+2260

Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on ' 0)' in 'mbcsToSbcs': dot substituted for <e2>

Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on ' 0)' in 'mbcsToSbcs': dot substituted for <89>

Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on ' 0)' in 'mbcsToSbcs': dot substituted for <a0>

Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, :
font metrics unknown for Unicode character U+2260

Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on ' 0)' in 'mbcsToSbcs': dot substituted for <e2>

Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on ' 0)' in 'mbcsToSbcs': dot substituted for <89>

Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on ' 0)' in 'mbcsToSbcs': dot substituted for <a0>

Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on ' 0)' in 'mbcsToSbcs': dot substituted for <e2>


Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on ' 0)' in 'mbcsToSbcs': dot substituted for <89>

Warning in grid.Call.graphics(C_text, as.graphicsAnnot(x\$label), x\$x, x\$y, :
conversion failure on ' 0)' in 'mbcsToSbcs': dot substituted for <a0>

In **Case 1** (left panel), x_2 doesn't actually affect y . Including it (coral) doesn't reduce bias (both distributions are centered on the true value), but it does increase variance (the coral distribution is wider).

In **Case 2** (right panel), x_2 does affect y . Omitting it (blue) causes bias—the distribution is centered away from the true value. Including it (coral) eliminates the bias, even though variance is slightly higher.

Figure 5.1: The bias-variance tradeoff. Left: When x_2 doesn't affect y ($\beta_2 = 0$), including it adds variance without reducing bias. Right: When x_2 does affect y ($\beta_2 \neq 0$), omitting it causes bias.



5.12 Summary

This chapter extended OLS to the multivariate case, allowing us to control for confounding variables.

5.12.1 Key Takeaways

- **Multivariate OLS** includes multiple independent variables:

$$y = \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \cdots + \beta_k x_k + \mu$$

- **Coefficients have ceteris paribus interpretations:** β_j is the effect of x_j holding all other variables constant
- **The zero conditional mean assumption** now requires $E[\mu|x_1, \dots, x_k] = 0$ —more plausible with good controls
- **Multivariate OLS is like matching:** It compares observations within groups defined by control variables
- **The OVB formula** tells us the direction of bias from omitting a variable:

$$\text{Bias} = \beta_2 \cdot \tilde{\delta}_1$$

- **There's a bias-variance tradeoff:** Omitting relevant variables causes bias; including irrelevant variables inflates variance

- **Perfect multicollinearity** occurs when one variable is an exact linear function of others; it must be avoided

5.13 Check Your Understanding

Note: This section contains interactive content only available in the HTML version.

6 Statistical Inference

💡 Key Questions

- How do we know if our OLS estimate reflects a “real” effect or is just due to random chance?
- What is the sampling distribution of our OLS estimator and why does it matter?
- How do we conduct hypothesis tests using t-statistics and p-values?
- What do confidence intervals tell us about our estimates?
- How do we test multiple hypotheses simultaneously using F-tests?
- How do we read and interpret regression tables in published research?

i Suggested Readings

- Wooldridge (2019), Ch. 4

6.1 From Point Estimates to Uncertainty

In the previous chapters, we learned how to estimate the parameters of a regression model using OLS. We discussed how, under certain assumptions, our OLS estimator $\hat{\beta}_j$ is *unbiased*—meaning that on average, across many hypothetical samples, our estimate equals the true population parameter β_j .

But here’s the catch: we only ever observe *one* sample. Our single estimate might be too high or too low relative to the truth, simply due to the random variation inherent in sampling. **Statistical inference** is the set of tools that allows us to quantify this uncertainty and make statements about population parameters based on sample data.

The central question of statistical inference is: *Is our single OLS estimate “real,” or is it due to random chance?*

6.2 The Sampling Distribution: A Simulation

To understand why uncertainty matters, let's run a simulation. Imagine we have a population where the true relationship between x and y is:

$$y = 2.45 + 0 \cdot x + \mu$$

Notice that the true effect of x on y is **exactly zero**. In the population, x has no effect on y whatsoever.

But what happens when we draw samples from this population and estimate OLS regressions? Let's find out.

```
set.seed(124890)

# Create our "population" data
pop_data <- tibble(
  x = sample(1:100, size = 1000, replace = TRUE),
  err = rnorm(1000),
  y = 2.45 + 0 * x + err # True beta_1 = 0
)

# Number of simulation repetitions
n_sims <- 1000

# Run the simulation
sim_results <- map_dfr(1:n_sims, function(i) {
  # Draw a random sample of 50 observations
  sample_data <- pop_data |>
    slice_sample(n = 50, replace = TRUE)

  # Estimate OLS regression
  reg <- lm(y ~ x, data = sample_data)

  # Store the estimate
  tibble(
    iteration = i,
    b1_hat = coef(reg)["x"]
  )
})

# Calculate the average estimate
```


```

avg_estimate <- mean(sim_results$b1_hat)

# Create the visualization
ggplot(sim_results, aes(x = b1_hat)) +
  geom_histogram(fill = "lightgrey", color = "black", bins = 40) +
  geom_vline(xintercept = avg_estimate, color = "darkgreen", linewidth = 1.5) +
  annotate("text", x = avg_estimate + 0.003, y = 85,
          label = paste0("Average estimate\n= ", round(avg_estimate, 4)),
          color = "darkgreen", size = 4, hjust = 0) +
  labs(
    x = expression(hat(beta)[1]),
    y = "Frequency"
  ) +
  theme_minimal() +
  theme(
    axis.text = element_text(size = 12),
    axis.title = element_text(size = 14)
  )

```

Figure 6.1: Sampling distribution of $\hat{\beta}_1$ when the true $\beta_1 = 0$. Each bar represents the frequency of estimates across 1,000 different samples.



Look at what happened! Even though the true $\beta_1 = 0$, we got a *distribution* of different

estimates ranging from about -0.02 to +0.02. Some samples produced positive estimates, others produced negative estimates. The average across all samples is very close to zero (as expected from unbiasedness), but any *individual* sample could give us an estimate that deviates from zero.


This distribution of estimates across all possible samples is called the **sampling distribution** of $\hat{\beta}_j$.

6.3 Where Does Our Estimate Fall?

Now imagine you only have access to *one* sample—as is always the case in practice. You run your regression and get an estimate of, say, $\hat{\beta}_1 = 0.001$.

How do you know if your estimate comes from here (close to the true value):

Figure 6.2: An estimate close to the average is likely consistent with the null hypothesis.




Or from here (far out in the tails):

If your estimate falls in the middle of the distribution where $\beta_1 = 0$, it's consistent with the null hypothesis of no effect. But if your estimate falls way out in the tails, it suggests that maybe the true β_1 isn't zero after all—because getting such an extreme estimate would be very unlikely if β_1 really were zero.

The tools of statistical inference formalize this intuition.

Figure 6.3: An estimate in the tail is unlikely if the true effect is zero—suggesting the true effect might not be zero.



6.4 The Sampling Distribution of $\hat{\beta}_j$

To conduct statistical inference, we need to know the *shape* of the sampling distribution. We’ve already established two key properties of our OLS estimator (under the Gauss-Markov assumptions):

Expected Value (Unbiasedness):

$$E[\hat{\beta}_j] = \beta_j$$

Variance (under homoskedasticity):

$$Var(\hat{\beta}_j) = \frac{\sigma^2}{SST_j(1 - R_j^2)}$$

where SST_j is the total sum of squares of x_j and R_j^2 is the R-squared from regressing x_j on all other independent variables.

But knowing the mean and variance isn’t enough—we need to know the *shape* of the distribution.

6.4.1 The Normality Assumption

To fully characterize the sampling distribution, we make an additional assumption:

! Normality Assumption

The population error μ is independent of the explanatory variables $x_1, x_2, \dots, x_k$ and is normally distributed with zero mean and variance σ^2 :

$$\mu \sim \text{Normal}(0, \sigma^2)$$

Equivalently, we can write:

$$y|\mathbf{x} \sim \text{Normal}(\beta_0 + \beta_1 x_1 + \dots + \beta_k x_k, \sigma^2)$$

This says that conditional on the independent variables, the outcome y follows a normal distribution centered on the regression line.

Let's visualize what a normal distribution looks like:


```
set.seed(123)
mu <- 100
sigma <- 15

normal_data <- tibble(
  values = rnorm(n = 1000, mean = mu, sd = sigma)
)

ggplot(normal_data, aes(x = values)) +
  geom_histogram(aes(y = after_stat(density)),
    bins = 30,
    fill = "skyblue",
    color = "white",
    alpha = 0.8) +
  stat_function(fun = dnorm,
    args = list(mean = mu, sd = sigma),
    color = "darkred",
    linewidth = 1.2) +
  labs(
    title = "Normal Distribution with Density Curve",
    subtitle = paste0("Generated from N( = ", mu, ", = ", sigma, ")"),
    x = "Values (e.g., IQ Scores)",
    y = "Density"
```

```
) +  
theme_minimal()
```

Figure 6.4: The normal (Gaussian) distribution—the classic ‘bell curve.’ The example shows IQ scores, which are designed to follow $N(100, 15)$.



At first glance, the normality assumption seems quite strong. For many economic variables—like wages, which are bounded below by zero and have a right skew—the distribution is clearly not normal. However, there’s good news: as long as our sample size is large enough, the sampling distribution of $\hat{\beta}_j$ will be approximately normal *even if the errors aren’t normally distributed*. This result is called **asymptotic normality** and follows from the Central Limit Theorem.

6.4.2 The Normal Sampling Distribution


Under the normality assumption (or with large samples via asymptotic normality), our OLS estimator follows a normal distribution:

$$\hat{\beta}_j \sim \text{Normal} [\beta_j, \text{Var}(\hat{\beta}_j)]$$

Let’s verify this with a simulation. We’ll draw many samples and show that the distribution of estimates indeed looks normal:

```
# Overlay a normal density on our simulation results
ggplot(sim_results, aes(x = b1_hat)) +
  geom_histogram(aes(y = after_stat(density)),
    fill = "lightgrey", color = "black", bins = 40) +
  stat_function(
    fun = dnorm,
    args = list(mean = mean(sim_results$b1_hat),
      sd = sd(sim_results$b1_hat)),
    color = "darkred",
    linewidth = 1.2
  ) +
  geom_vline(xintercept = 0, color = "darkgreen", linewidth = 1.2, linetype = "dashed") +
  annotate("text", x = 0.012, y = 55,
    label = "True .. = 0", color = "darkgreen", size = 4) +
  labs(
    x = expression(hat(beta)[1]),
    y = "Density"
  ) +
  theme_minimal()
```

Figure 6.5: The sampling distribution of $\hat{\beta}_1$ follows a normal distribution (red curve) centered on the true value.



The histogram of our 1,000 simulated estimates matches the theoretical normal curve almost perfectly!

Quick Check: This section contains interactive content only available in the HTML version.

6.5 Hypothesis Testing

Now that we understand the sampling distribution, we can use it to make inferences about population parameters. The key idea is **hypothesis testing**: we hypothesize about the value of β_j in the population, then use our sample estimate to evaluate whether the hypothesis is plausible.

6.5.1 The Null and Alternative Hypotheses

In econometrics, we're usually interested in whether a variable has *any* effect on the outcome. We formalize this as:

Null Hypothesis: $H_0 : \beta_j = 0$

The null hypothesis states that the variable x_j has no effect on y in the population (after controlling for other variables).

Alternative Hypothesis: We test the null against one of:

- $H_1 : \beta_j \neq 0$ (two-sided test, most common)
- $H_1 : \beta_j > 0$ (one-sided, if we expect a positive effect)
- $H_1 : \beta_j < 0$ (one-sided, if we expect a negative effect)

For example, consider the wage equation:

$$wage = \beta_0 + \beta_1(educ) + \beta_2(experience) + \beta_3(tenure) + \mu$$

The null hypothesis $H_0 : \beta_2 = 0$ states that, after controlling for education and tenure, experience has no effect on wages.

6.5.2 The t-Statistic

To test our hypothesis, we use the **t-statistic**. Under the null hypothesis, the following quantity follows a t-distribution:

$$\frac{\hat{\beta}_j - \beta_j}{se(\hat{\beta}_j)} \sim t_{n-k-1}$$

where $n - k - 1$ are the degrees of freedom (sample size minus the number of parameters).

When testing $H_0 : \beta_j = 0$, the t-statistic simplifies to:

$$t_{\hat{\beta}_j} = \frac{\hat{\beta}_j}{se(\hat{\beta}_j)}$$

This is simply the ratio of our estimate to its standard error. The t-stat tells us: *how many standard errors away from zero is our estimate?*

6.5.3 Properties of the t-Statistic

The t-statistic has several useful properties:

1. **Same sign as the estimate:** Since $se(\hat{\beta}_j) > 0$, the t-stat has the same sign as $\hat{\beta}_j$
2. **Magnitude matters:** As $\hat{\beta}_j$ grows in magnitude, so does $t_{\hat{\beta}_j}$
3. **Signal-to-noise ratio:** The t-stat measures how large our estimate is *relative to the noise* (uncertainty) in our data

6.5.4 Visualizing the Hypothesis Test


Let's visualize what the t-test is doing. Under the null hypothesis $H_0 : \beta_j = 0$, we assume that the distribution of possible estimates is centered on zero:

```
set.seed(42)
df_value <- 100 # degrees of freedom

t_sample <- tibble(
  t_value = rt(10000, df = df_value)
)
```

```
ggplot(t_sample, aes(x = t_value)) +
  geom_histogram(aes(y = after_stat(density)),
    bins = 50,
    fill = "lightblue",
    color = "black") +
  stat_function(fun = dt,
    args = list(df = df_value),
    color = "darkred",
    linewidth = 1.2) +
  geom_vline(xintercept = 0, color = "darkgreen", linewidth = 1.2) +
  annotate("text", x = 1.5, y = 0.35,
    label = "Distribution under H ",
    color = "darkgreen", size = 5) +
  labs(x = "t-values", y = "Density") +
  coord_cartesian(xlim = c(-4, 4)) +
  theme_minimal()
```

Figure 6.6: The t-distribution under the null hypothesis $H_0: \mu = 0$. Most estimates will fall near zero.



Now suppose we get a t-statistic from our sample. Is it close to zero (consistent with H_0) or far from zero (evidence against H_0)?

```


p_close <- ggplot(t_sample, aes(x = t_value)) +
  geom_histogram(aes(y = after_stat(density)),
    bins = 50, fill = "lightblue", color = "black") +
  stat_function(fun = dt, args = list(df = df_value),
    color = "darkred", linewidth = 1.2) +
  geom_vline(xintercept = 0.85, color = "blue", linewidth = 1.5) +
  annotate("text", x = 0.85, y = 0.42, label = "t = 0.85",
    color = "blue", size = 4, hjust = -0.1) +
  labs(x = "t-values", y = "Density",
    title = "t-stat close to zero") +
  coord_cartesian(xlim = c(-4, 4)) +
  theme_minimal()

p_far <- ggplot(t_sample, aes(x = t_value)) +
  geom_histogram(aes(y = after_stat(density)),
    bins = 50, fill = "lightblue", color = "black") +
  stat_function(fun = dt, args = list(df = df_value),
    color = "darkred", linewidth = 1.2) +
  geom_vline(xintercept = 2.8, color = "red", linewidth = 1.5) +
  annotate("text", x = 2.8, y = 0.15, label = "t = 2.8",
    color = "red", size = 4, hjust = -0.1) +
  labs(x = "t-values", y = "Density",
    title = "t-stat in the tail") +
  coord_cartesian(xlim = c(-4, 4)) +
  theme_minimal()

p_close + p_far

```


Figure 6.7: Two possible t-statistics: one at 0.85 (consistent with H_0) and one at 2.8 (in the tail, suggesting we reject H_0).



A t-statistic of 0.85 falls well within the “body” of the distribution—values like this occur frequently when H_0 is true. But a t-statistic of 2.8 is out in the tail—such extreme values are rare when H_0 is true.

6.5.5 Critical Values and Rejection Regions

To formalize our decision rule, we need to define:

1. **Significance level (α):** The probability of rejecting H_0 when it’s actually true (Type I error). Common choices: 10%, 5%, or 1%.
2. **Critical value (c):** The threshold value of the t-statistic beyond which we reject H_0 .

For a two-sided test at the 5% level, we reject H_0 if $|t_{\hat{\beta}_j}| > c$, where c is the critical value that leaves 2.5% of the distribution in each tail.

```
# Calculate critical value for 5% significance, two-sided
alpha <- 0.05
c_val <- qt(alpha/2, df = df_value, lower.tail = FALSE)

# Create shading data for rejection regions
```

```


t_grid <- seq(-4, 4, length.out = 500)
t_dens <- dt(t_grid, df = df_value)

shade_data <- tibble(t_value = t_grid, density = t_dens) |>
  mutate(
    rejection = t_value < -c_val | t_value > c_val
  )

ggplot() +
  # Full distribution
  geom_area(data = shade_data |> filter(!rejection),
    aes(x = t_value, y = density),
    fill = "lightblue", alpha = 0.7) +
  # Rejection regions
  geom_area(data = shade_data |> filter(rejection),
    aes(x = t_value, y = density),
    fill = "red", alpha = 0.5) +
  # Distribution curve
  stat_function(fun = dt, args = list(df = df_value),
    color = "darkblue", linewidth = 1.2) +
  # Critical value lines
  geom_vline(xintercept = c(-c_val, c_val),
    color = "darkgoldenrod", linewidth = 1.2, linetype = "dashed") +
  # Annotations
  annotate("text", x = -c_val, y = 0.42,
    label = paste0("-", round(c_val, 2)),
    color = "darkgoldenrod", size = 4, hjust = 1.1) +
  annotate("text", x = c_val, y = 0.42,
    label = paste0("+", round(c_val, 2)),
    color = "darkgoldenrod", size = 4, hjust = -0.1) +
  annotate("text", x = 0, y = 0.2,
    label = "Fail to reject H\n(95%)",
    color = "darkblue", size = 4) +
  annotate("text", x = 3.2, y = 0.05,
    label = "Reject H\n(2.5%)",
    color = "red", size = 3) +
  annotate("text", x = -3.2, y = 0.05,
    label = "Reject H\n(2.5%)",
    color = "red", size = 3) +
  labs(x = "t-values", y = "Density") +
  coord_cartesian(xlim = c(-4, 4)) +
  theme_minimal()

```

Figure 6.8: Critical values for a two-sided test at the 5% significance level. We reject H_0 if $|t| > 1.96$.



6.5.6 Hypothesis Testing Example

Let's work through a complete example. We'll create some simulated housing data and test whether lot area affects sale price:

```
# Create simulated housing data
set.seed(2025)
n <- 500

housing_data <- tibble(
  lot_area = runif(n, 5000, 20000),
  pool_area = rbinom(n, 1, 0.15) * runif(n, 200, 600),
  garage_area = runif(n, 200, 800),
  year_built = sample(1960:2020, n, replace = TRUE),
  year_remod = pmax(year_built, sample(1980:2023, n, replace = TRUE)),
  # True relationship: lot_area has effect of $1.50 per sq ft
  sale_price = -2500000 + 1.50 * lot_area + 80 * pool_area +
    150 * garage_area + 500 * year_built +
    900 * year_remod + rnorm(n, 0, 50000)
)
```

```
# Estimate the regression
reg_housing <- lm(sale_price ~ lot_area + pool_area + garage_area +
                  year_built + year_remod,
                  data = housing_data)

summary(reg_housing)
```

Call:

```
lm(formula = sale_price ~ lot_area + pool_area + garage_area +
    year_built + year_remod, data = housing_data)
```

Residuals:

Min	1Q	Median	3Q	Max
-134582	-32117	138	32584	167861

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)	
(Intercept)	-2369345.4532	407348.8976	-5.817	0.0000000108	***
lot_area	1.5528	0.5207	2.982	0.003002	**
pool_area	81.5522	14.6980	5.549	0.0000000470	***
garage_area	167.0837	13.4931	12.383	< 0.0000000000000002	***
year_built	554.2853	143.8278	3.854	0.000132	***
year_remod	775.7472	223.5861	3.470	0.000567	***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 51130 on 494 degrees of freedom

Multiple R-squared: 0.3396, Adjusted R-squared: 0.3329

F-statistic: 50.81 on 5 and 494 DF, p-value: < 0.00000000000000022

To test $H_0 : \beta_1 = 0$ (lot area has no effect):

```
# Extract the coefficient and standard error for lot_area
b1_hat <- coef(reg_housing)["lot_area"]
se_b1 <- summary(reg_housing)$coefficients["lot_area", "Std. Error"]

# Calculate the t-statistic
t_stat <- b1_hat / se_b1

# Find the critical value at 5% significance
```

```

sig_level <- 0.05
n_obs <- nobs(reg_housing)
k <- 5 # number of independent variables
df <- n_obs - k - 1

t_crit <- qt(sig_level / 2, df = df, lower.tail = FALSE)

# Display results
cat("Estimate:", round(b1_hat, 4), "\n")

```

Estimate: 1.5528

```
cat("Standard Error:", round(se_b1, 4), "\n")
```

Standard Error: 0.5207

```
cat("t-statistic:", round(t_stat, 2), "\n")
```

t-statistic: 2.98

```
cat("Critical value (5%):", round(t_crit, 2), "\n")
```

Critical value (5%): 1.96

```
cat("Reject H?", abs(t_stat) > t_crit, "\n")
```

Reject H? TRUE

Since $|t_{\hat{\beta}_1}|$ exceeds the critical value of approximately 1.96, we **reject the null hypothesis** that lot area has no effect on sale price. The estimate is **statistically significant at the 5% level**.

6.5.7 Rule of Thumb

With modern datasets that have hundreds or thousands of observations, the critical value for a two-sided 5% test is essentially always **1.96** **2**.

This gives us a handy rule of thumb:

💡 The “Rule of 2”

An estimate is statistically significant at the 5% level if:

$$|\hat{\beta}_j| > 2 \times se(\hat{\beta}_j)$$

In other words, if your estimate is more than twice as large as its standard error, you can reject the null hypothesis of no effect at the 5% level.

6.6 P-Values

While the t-test with a fixed significance level is useful, it has limitations. Saying an estimate is “significant at 5%” doesn’t tell us *how strongly* we reject the null. Was it a narrow rejection or a “knock out”?

The **p-value** answers this question:

! Definition: P-Value

The p-value is the smallest significance level at which we would reject the null hypothesis. Equivalently: the p-value is the probability of observing a t-statistic as extreme (or more extreme) as the one we calculated, *if the null hypothesis were true*.

6.6.1 Visualizing P-Values

```
# Parameters
df_viz <- 250
observed_t <- 2.3

# Calculate p-value
p_val <- 2 * pt(abs(observed_t), df = df_viz, lower.tail = FALSE)

# Create the visualization
t_grid <- seq(-4, 4, length.out = 500)
t_dens <- dt(t_grid, df = df_viz)

shade_data <- tibble(t_value = t_grid, density = t_dens) |>
  mutate(
    in_tail = abs(t_value) >= abs(observed_t)
```


```

)

ggplot() +
  # Main curve
  geom_line(data = shade_data, aes(x = t_value, y = density),
            color = "darkblue", linewidth = 1.2) +
  # Shaded tails (p-value area)
  geom_area(data = shade_data |> filter(in_tail & t_value > 0),
            aes(x = t_value, y = density),
            fill = "red", alpha = 0.6) +
  geom_area(data = shade_data |> filter(in_tail & t_value < 0),
            aes(x = t_value, y = density),
            fill = "red", alpha = 0.6) +
  # Observed t-statistic lines
  geom_vline(xintercept = c(-observed_t, observed_t),
             color = "darkgreen", linewidth = 1, linetype = "dashed") +
  # Annotations
  annotate("text", x = observed_t + 0.3, y = 0.05,
          label = paste0("Observed t = ", observed_t),
          color = "darkgreen", size = 4, hjust = 0) +
  annotate("text", x = 0, y = 0.2,
          label = paste0("P-value = ", round(p_val, 4)),
          color = "red", size = 5) +
  labs(
    title = "P-Value Visualization",
    subtitle = paste0("df = ", df_viz, ", Observed t = ", observed_t),
    x = "t-statistic",
    y = "Density"
  ) +
  theme_minimal()

```

Figure 6.9: The p-value is the total shaded area in both tails beyond the observed t-statistic.



6.6.2 Interpreting P-Values

The p-value tells us the probability of getting our estimate (or something more extreme) if the null were true:

- **p = 0.035:** There's a 3.5% chance of getting our estimate if $\beta_j = 0$. That's quite unlikely—evidence against H_0 .
- **p = 0.35:** There's a 35% chance of getting our estimate if $\beta_j = 0$. That's not unusual at all—no evidence against H_0 .

Common decision rules:

- $p < 0.10$: Reject H_0 at 10% level (weak evidence against H_0)
- $p < 0.05$: Reject H_0 at 5% level (moderate evidence against H_0)
- $p < 0.01$: Reject H_0 at 1% level (strong evidence against H_0)

6.6.3 A Note on Language

When we *cannot* reject the null hypothesis, we say:

“We **fail to reject** the null at the x% level.”

We do **not** say we “accept” the null. Why? Because there are many possible values of β_j that we would also fail to reject. Failing to reject $\beta_j = 0$ doesn’t mean β_j actually equals zero—it just means our data aren’t precise enough to distinguish β_j from zero.

6.6.4 Statistical vs. Practical Significance

Important Distinction

Statistical significance and **practical (economic) significance** are not the same thing!

A coefficient can be statistically significant but practically unimportant. This is especially common in very large samples, where even tiny effects can be detected.

For example, suppose we find that an additional year of education increases wages by \$0.50 per year, and this is statistically significant with $p < 0.001$. Statistically, we’re very confident the effect isn’t zero. But practically, a 50-cent annual raise is economically trivial.

Always interpret the *magnitude* of coefficients, not just their statistical significance.

Quick Check: This section contains interactive content only available in the HTML version.

6.7 Confidence Intervals

Another way to quantify uncertainty is through **confidence intervals**. While hypothesis tests ask “is β_j different from zero?”, confidence intervals ask “what range of values is β_j likely to fall within?”

6.7.1 Computing a Confidence Interval

A confidence interval at the $(1 - \alpha)$ level is:

$$\hat{\beta}_j \pm c_\alpha \times se(\hat{\beta}_j)$$

where c_α is the critical value at significance level α .

For a 95% confidence interval with large samples:

$$\hat{\beta}_j \pm 1.96 \times se(\hat{\beta}_j)$$

6.7.2 Example

Using our housing regression:

```
# 95% confidence interval for lot_area coefficient
lower <- b1_hat - 1.96 * se_b1
upper <- b1_hat + 1.96 * se_b1

cat("Point estimate:", round(b1_hat, 4), "\n")
```

Point estimate: 1.5528

```
cat("95% CI: [", round(lower, 4), ",", round(upper, 4), "]\n")
```

95% CI: [0.5323 , 2.5733]

```
# Or use confint() function
confint(reg_housing, "lot_area", level = 0.95)
```

```
                2.5 %   97.5 %
lot_area 0.5298262 2.57581
```

6.7.3 Interpreting Confidence Intervals

“We are 95% confident that the true effect of lot area on sale price is between \$0.53 and \$2.57 per square foot.”

More precisely: if we drew 100 different samples and computed 100 different confidence intervals, about 95 of them would contain the true β_j .

Key insight: If the confidence interval includes zero, the estimate is *not* statistically significant at that level.

6.7.4 Simulation: Understanding Confidence Intervals

Let’s run a simulation to see what “95% confidence” really means:

```

set.seed(456)

# True parameter value
true_beta1 <- 1.50

# Generate 20 samples and compute CIs
n_samples <- 20

ci_data <- map_dfr(1:n_samples, function(i) {
  # Generate new sample
  sample_data <- tibble(
    lot_area = runif(500, 5000, 20000),
    pool_area = rbinom(500, 1, 0.15) * runif(500, 200, 600),
    garage_area = runif(500, 200, 800),
    year_built = sample(1960:2020, 500, replace = TRUE),
    year_remod = pmax(year_built, sample(1980:2023, 500, replace = TRUE)),
    sale_price = -2500000 + true_beta1 * lot_area + 80 * pool_area +
      150 * garage_area + 500 * year_built +
      900 * year_remod + rnorm(500, 0, 50000)
  )

  # Fit model
  reg <- lm(sale_price ~ lot_area + pool_area + garage_area +
    year_built + year_remod, data = sample_data)

  # Get CI
  ci <- confint(reg, "lot_area", level = 0.95)

  tibble(
    sample = i,
    estimate = coef(reg)["lot_area"],
    lower = ci[1],
    upper = ci[2],
    covers_true = lower <= true_beta1 & upper >= true_beta1
  )
})

ggplot(ci_data, aes(y = sample)) +
  geom_vline(xintercept = true_beta1, linetype = "dashed",
    color = "darkgreen", linewidth = 1) +
  geom_errorbarh(aes(xmin = lower, xmax = upper,
    color = covers_true),

```

```


      height = 0.3, linewidth = 0.8) +
geom_point(aes(x = estimate, color = covers_true), size = 2) +
scale_color_manual(values = c("TRUE" = "steelblue", "FALSE" = "red"),
                    labels = c("TRUE" = "Contains true value",
                              "FALSE" = "Misses true value")) +
annotate("text", x = true_beta1 + 0.05, y = 21,
          label = paste0("True  = ", true_beta1),
          color = "darkgreen", size = 4, hjust = 0) +
labs(
  x = expression(hat(beta)[1]),
  y = "Sample",
  color = NULL
) +
theme_minimal() +
theme(legend.position = "bottom")

```

Warning: `geom\_errorbarh()` was deprecated in ggplot2 4.0.0.
 i Please use the `orientation` argument of `geom\_errorbar()` instead.

`height` was translated to `width`.

Figure 6.10: Twenty 95% confidence intervals from different samples. The true $\beta_1 = 1.50$ (dashed line). About 95% of intervals capture the true value.



6.8 T-Tests, P-Values, and CIs: A Comparison

These three approaches are closely related but answer slightly different questions:

Method	Question Answered
T-statistic	Is my estimate large relative to the noise in the data?
P-value	What's the probability of seeing my estimate if H_0 were true?
Confidence Interval	What range of values is plausible for β_j ?

All three are mathematically connected:

- If $|t| > 1.96$, then $p < 0.05$, and the 95% CI excludes zero
- If $p < 0.05$, then $|t| > 1.96$, and the 95% CI excludes zero
- If the 95% CI excludes zero, then $p < 0.05$ and $|t| > 1.96$

6.9 Reading Regression Tables

Now that we understand statistical inference, we can properly interpret the regression tables you'll encounter in academic papers.

6.9.1 From R Output to Publication Format

When you run `summary()` on a regression in R, you get detailed but messy output. Academic papers present this information in a standardized, clean format using tables.

```
# Standard R output  
summary(reg_housing)
```

Call:

```
lm(formula = sale_price ~ lot_area + pool_area + garage_area +  
    year_built + year_remod, data = housing_data)
```

Residuals:

Min	1Q	Median	3Q	Max
-134582	-32117	138	32584	167861

Coefficients:

Estimate	Std. Error	t value	Pr(> t)
----------	------------	---------	----------

Table 6.2: The Effect of Lot Area on Home Sale Price

	Sale Price
(Intercept)	-2 369 345.453*** (407 348.898)
Lot Area (sq ft)	1.553*** (0.521)
Pool Area	81.552*** (14.698)
Garage Area	167.084*** (13.493)
Year Built	554.285*** (143.828)
Year Remodeled	775.747*** (223.586)
Num.Obs.	500
R2	0.340
R2 Adj.	0.333

* p < 0.1, ** p < 0.05, *** p < 0.01
Standard errors in parentheses. * p < 0.1, ** p < 0.05, *** p < 0.01

```
(Intercept) -2369345.4532    407348.8976   -5.817          0.0000000108 ***
lot_area      1.5528          0.5207    2.982          0.003002 **
pool_area     81.5522         14.6980    5.549          0.0000000470 ***
garage_area   167.0837        13.4931   12.383 < 0.0000000000000002 ***
year_built    554.2853        143.8278    3.854          0.000132 ***
year_remod    775.7472        223.5861    3.470          0.000567 ***
```

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 51130 on 494 degrees of freedom

Multiple R-squared: 0.3396, Adjusted R-squared: 0.3329

F-statistic: 50.81 on 5 and 494 DF, p-value: < 0.00000000000000022

In a publication, this same information would appear as:

6.9.2 Decoding the Table Structure

Rows are variables: The top indicates the dependent variable. Rows below show independent variables and the constant (intercept).

The main numbers are coefficients: Each cell contains $\hat{\beta}_j$.

Numbers in parentheses are standard errors: These are $se(\hat{\beta}_j)$.

Stars indicate significance levels:

- * = significant at 10% level ($p < 0.10$)
- ** = significant at 5% level ($p < 0.05$)
- *** = significant at 1% level ($p < 0.01$)

Bottom statistics: Sample size (N), R-squared, and sometimes other diagnostics.

6.9.3 Multiple Columns Show Robustness

Most regression tables have multiple columns, each representing a different specification. This lets readers see how estimates change as controls are added:

```
# Three specifications with increasing controls
reg1 <- lm(sale_price ~ lot_area, data = housing_data)
reg2 <- lm(sale_price ~ lot_area + pool_area + garage_area, data = housing_data)
reg3 <- lm(sale_price ~ lot_area + pool_area + garage_area +
           year_built + year_remod, data = housing_data)
```

Notice how the coefficient on Lot Area changes across specifications. In column (1), without controls, the estimate is different from column (3). This pattern often reveals omitted variable bias in simpler specifications.

6.9.4 What to Look for in Regression Tables

1. **What is the coefficient on the variable of interest?** How do we interpret it?
2. **Is it statistically significant?** At what level?
3. **How does the coefficient change across specifications?** Does it remain stable or shift dramatically?
4. **Are the sample size and R-squared reasonable?** Very small N or very low R^2 might raise concerns.

Table 6.3: The Effect of Lot Area on Home Sale Price: Multiple Specifications

	(1)	(2)	(3)
(Intercept)	385 464.906*** (8462.611)	288 761.744*** (10 454.349)	−2 369 345.453*** (407 348.898)
Lot Area (sq ft)	1.038 (0.635)	1.472*** (0.545)	1.553*** (0.521)
Pool Area		86.081*** (15.352)	81.552*** (14.698)
Garage Area		170.687*** (14.112)	167.084*** (13.493)
Year Built			554.285*** (143.828)
Year Remodeled			775.747*** (223.586)
Num.Obs.	500	500	500
R2	0.005	0.274	0.340
R2 Adj.	0.003	0.269	0.333

* p < 0.1, ** p < 0.05, *** p < 0.01

Standard errors in parentheses. * p < 0.1, ** p < 0.05, *** p < 0.01

Table 6.4: Effect of Stock Performance on CEO Salary

	(1)	(2)
(Intercept)	6.806*** (0.041)	4.788*** (0.234)
Return on Stock (%)	0.001 (0.001)	0.001 (0.001)
Log(Sales)		0.287*** (0.033)
Num.Obs.	209	209
R2 Adj.	−0.004	0.264

* $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$

Practice: Reading a Regression Table

Consider this table examining the effect of stock performance on CEO salary:

Is the effect of Return on Stock statistically significant at the 5% level in Column (1)? Select answer... Yes No

What happens to the Adjusted R^2 when we add $\log(\text{Sales})$ as a control? Select answer... It increases substantially It decreases It stays about the same

Based on this table, which variable appears to be a more important determinant of CEO salary? Select answer... Return on Stock Company Sales Both are equally important

6.10 F-Tests: Testing Multiple Hypotheses

Sometimes we want to test hypotheses about *multiple* coefficients simultaneously. For example: “Do experience AND tenure jointly affect wages?”

The **F-test** allows us to test such joint hypotheses.

6.10.1 Why Not Just Use Multiple t-Tests?

You might think: “Why not just run separate t-tests on each coefficient?”

There are two problems:

1. **No restrictions on other parameters:** A t-test on β_2 puts no restrictions on β_3 . If the variables are correlated, this can be misleading.
2. **Multiple comparisons problem:** If you run many tests at the 5% level, you'll reject *some* true null hypotheses just by chance. With 20 tests, you'd expect about 1 false rejection even if all nulls are true!

The F-test controls for these issues.

6.10.2 Setting Up the F-Test

Consider the wage model:

$$\log(wage) = \beta_0 + \beta_1(education) + \beta_2(experience) + \beta_3(tenure) + \beta_4(age) + \mu$$

Suppose we want to test whether on-the-job training variables (experience and tenure) matter at all:

$$H_0 : \beta_2 = \beta_3 = 0$$

$$H_1 : H_0 \text{ is not true}$$

The alternative is satisfied if *either* $\beta_2 \neq 0$ or $\beta_3 \neq 0$ (or both).

6.10.3 Computing the F-Statistic

The F-test compares two models:

Unrestricted Model: The full model with all variables

$$UR : \widehat{\log(wage)} = \hat{\beta}_0 + \hat{\beta}_1(educ) + \hat{\beta}_2(exper) + \hat{\beta}_3(tenure) + \hat{\beta}_4(age)$$

Restricted Model: The model assuming the null is true (dropping the restricted variables)

$$R : \widehat{\log(wage)} = \hat{\beta}_0 + \hat{\beta}_1(educ) + \hat{\beta}_4(age)$$

The F-statistic is:

$$F = \frac{(SSR_R - SSR_{UR})/q}{SSR_{UR}/df_{UR}}$$

where:

- SSR_R = Sum of squared residuals from restricted model
- SSR_{UR} = Sum of squared residuals from unrestricted model
- q = Number of restrictions (variables dropped)
- df_{UR} = Degrees of freedom in unrestricted model ($n - k - 1$)

6.10.4 F-Test Example

Let's test whether year built and year remodeled jointly affect home prices:

```
# Unrestricted model (full model)
reg_unrestricted <- lm(sale_price ~ lot_area + pool_area + garage_area +
                      year_built + year_remod,
                      data = housing_data)

# Restricted model (dropping year variables)
reg_restricted <- lm(sale_price ~ lot_area + pool_area + garage_area,
                    data = housing_data)

# Compute F-statistic manually
ssr_r <- sum(resid(reg_restricted)^2)
ssr_ur <- sum(resid(reg_unrestricted)^2)
df_ur <- df.residual(reg_unrestricted)
q <- 2 # number of restrictions

f_numerator <- (ssr_r - ssr_ur) / q
f_denominator <- ssr_ur / df_ur
f_stat <- f_numerator / f_denominator

# Critical value at 5% significance
f_crit <- qf(0.05, df1 = q, df2 = df_ur, lower.tail = FALSE)

cat("F-statistic:", round(f_stat, 2), "\n")
```

F-statistic: 24.69

```
cat("Critical value (5%):", round(f_crit, 2), "\n")
```

Critical value (5%): 3.01

```
cat("Reject H?", f_stat > f_crit, "\n")
```

Reject H? TRUE

Since the F-statistic exceeds the critical value, we **reject the null hypothesis**. Year built and year remodeled are **jointly statistically significant**—they collectively improve the model’s fit.

6.10.5 Using R’s anova() Function

In practice, you’ll use R’s `anova()` function to perform F-tests:

```
anova(reg_restricted, reg_unrestricted)
```

Analysis of Variance Table

Model 1: `sale_price ~ lot_area + pool_area + garage_area`

Model 2: `sale_price ~ lot_area + pool_area + garage_area + year_built + year_remod`

	Res.Df	RSS	Df	Sum of Sq	F	Pr(>F)					
1	496	1420348267818									
2	494	1291292854097	2	129055413721	24.686	0.00000000006048 ***					

Signif. codes:	0	'***'	0.001	'**'	0.01	'*'	0.05	'.'	0.1	' '	1

The output shows the F-statistic, degrees of freedom, and p-value directly.

6.10.6 The F-Statistic in Terms of R²

The F-statistic can also be written in terms of R-squared values:

$$F = \frac{(R_{UR}^2 - R_R^2)/q}{(1 - R_{UR}^2)/df_{UR}}$$

This shows that the F-test is essentially asking: *Does the unrestricted model fit the data significantly better than the restricted model?*

If R_{UR}^2 is much larger than R_R^2 , the F-statistic will be large, and we’ll reject the null.

Properties of the F-Test

- The F-statistic is always non-negative, so F-tests are always **one-sided**
- If the null is rejected, we say the variables are **jointly statistically significant**
- We cannot determine *which* individual variable is significant—only that at least one is
- If variables are jointly insignificant, it provides justification for dropping them from the model

6.11 Chapter Summary

Key Takeaways

1. **Statistical inference** allows us to move from sample estimates to statements about population parameters, accounting for sampling variability.
2. The **sampling distribution** of $\hat{\beta}_j$ describes how our estimates would vary across repeated samples. Under the normality assumption (or with large samples), it follows a normal distribution.
3. **Hypothesis testing** uses the t-statistic to determine whether our estimate is “far enough” from zero to reject the null hypothesis: $t = \hat{\beta}_j / se(\hat{\beta}_j)$
4. **P-values** tell us the probability of seeing our data if the null were true. Smaller p-values provide stronger evidence against H_0 .
5. **Confidence intervals** provide a range of plausible values for the population parameter. A 95% CI excludes zero if and only if the estimate is significant at the 5% level.
6. **Statistical significance** **practical significance**. Always consider the magnitude of effects, not just their p-values.
7. **F-tests** allow us to test multiple hypotheses simultaneously, avoiding the multiple comparisons problem.
8. **Regression tables** in academic papers present coefficients, standard errors (in parentheses), and significance stars. Multiple columns show robustness across specifications.

6.12 Practice Exercises

Note: This section contains interactive content only available in the HTML version.

7 Dummy Variables and Non-Linear Models

Key Questions

- How do we include categorical information (like race, gender, or region) in a regression?
- What is a dummy variable and how do we interpret its coefficient?
- How do we avoid the “dummy variable trap”?
- How can we model non-linear relationships using polynomials?
- When should we use logarithmic transformations?

Suggested Readings

- Wooldridge (2019) Ch. 7
- Bailey (2020), Ch. 5

So far, we have primarily considered **quantitative** variables in our regression models: wages measured in dollars, years of education, birthweight in grams, and so on. But in empirical work, some of the most important variables are **qualitative**: race, gender, region, industry, or whether someone participated in a program. This chapter introduces techniques for incorporating categorical information into regression models and for modeling non-linear relationships.

7.1 Categorical Variables in Regression

Regression requires numeric inputs, so we cannot simply include a variable like “region” with values “South,” “Midwest,” “Northeast,” and “West” directly in our model. We need a way to translate qualitative information into numbers.

7.1.1 Dummy Variables

The most common approach is to create **dummy variables** (also called binary or indicator variables). A dummy variable takes only two values: 1 if the observation has a particular characteristic, and 0 otherwise.

Consider this simple example from the Star Wars dataset:

```
ex_data <- starwars |>
  head(5) |>
  select(name, height, mass, species)

knitr::kable(ex_data, caption = "Star Wars Characters")
```

Table 7.1: Star Wars Characters

name	height	mass	species
Luke Skywalker	172	77	Human
C-3PO	167	75	Droid
R2-D2	96	32	Droid
Darth Vader	202	136	Human
Leia Organa	150	49	Human

The `species` variable is qualitative—we cannot use it directly in a regression. But we can create a dummy variable called `human` that equals 1 for humans and 0 for droids:

```
ex_data <- ex_data |>
  mutate(human = case_when(species == "Human" ~ 1,
                           TRUE ~ 0))

knitr::kable(ex_data, caption = "Star Wars Characters with Dummy Variable")
```

Table 7.2: Star Wars Characters with Dummy Variable

name	height	mass	species	human
Luke Skywalker	172	77	Human	1
C-3PO	167	75	Droid	0
R2-D2	96	32	Droid	0
Darth Vader	202	136	Human	1
Leia Organa	150	49	Human	1

Now we have a numeric variable that captures whether each character is human.

7.1.2 Interpreting Dummy Variable Coefficients

To see how dummy variables work in regression, let's examine a practical example. Suppose we want to know whether taking calculus in high school affects performance in college economics courses. We have data on students' economics exam scores, their high school GPA, and whether they took calculus.

We estimate the model:

$$\text{score} = \beta_0 + \beta_1(\text{hsgpa}) + \beta_2(\text{calculus}) + \mu$$

where `calculus` equals 1 if the student took calculus and 0 otherwise.

```
lm1 <- lm(score ~ hsgpa + calculus, data = wooldridge::econmath)
summary(lm1)
```

Call:

```
lm(formula = score ~ hsgpa + calculus, data = wooldridge::econmath)
```

Residuals:

Min	1Q	Median	3Q	Max
-54.355	-7.457	1.148	8.699	29.825

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	24.5276	4.0759	6.018	0.000000002624 ***
hsgpa	13.2038	1.2257	10.772	< 0.0000000000000002 ***
calculus	5.8243	0.8992	6.477	0.000000000158 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 12.18 on 853 degrees of freedom

Multiple R-squared: 0.1756, Adjusted R-squared: 0.1737

F-statistic: 90.87 on 2 and 853 DF, p-value: < 0.00000000000000022

How do we interpret these coefficients? The coefficient on `hsgpa` ($\hat{\beta}_1 \approx 13.2$) tells us that a one-point increase in high school GPA is associated with about a 13-point increase in economics score, holding calculus status constant.

The coefficient on `calculus` ($\hat{\beta}_2 \approx 5.8$) tells us that students who took calculus score about 5.8 points higher on average than students who did not, holding GPA constant. This is the **difference in intercepts** between the two groups.

7.1.3 Visualizing What Dummy Variables Do

To understand what's happening geometrically, think about the regression line for each group.

For students who did **not** take calculus (`calculus = 0`):


$$\widehat{score} = \hat{\beta}_0 + \hat{\beta}_1(hsgpa) + \hat{\beta}_2(0) = \hat{\beta}_0 + \hat{\beta}_1(hsgpa)$$

For students who **did** take calculus (`calculus = 1`):

$$\widehat{score} = \hat{\beta}_0 + \hat{\beta}_1(hsgpa) + \hat{\beta}_2(1) = (\hat{\beta}_0 + \hat{\beta}_2) + \hat{\beta}_1(hsgpa)$$

Both groups have the **same slope** ($\hat{\beta}_1$), but the calculus group has a higher intercept by exactly $\hat{\beta}_2$. The dummy variable shifts the regression line up or down:

Figure 7.1: Dummy variables shift the intercept while keeping the slope constant.



! Key Insight

A dummy variable coefficient represents the **difference in intercepts** between groups. Both groups share the same slope—the effect of increasing x by one unit is the same regardless of group membership.

7.2 The Dummy Variable Trap

You might wonder: why not include dummy variables for *both* calculus-takers and non-calculus-takers? In other words, why not estimate:

$$score = \beta_0 + \beta_1(hsgpa) + \beta_2(calculus) + \beta_3(no_calculus) + \mu$$

The problem is that this creates **perfect multicollinearity**. For every student in the dataset:

$$calculus + no_calculus = 1$$

This means $no_calculus = 1 - calculus$, so one variable is a perfect linear function of the other. OLS cannot separately identify the effects of perfectly collinear variables.

This is called the **dummy variable trap**. To avoid it, we always omit one category. The omitted category becomes our **baseline** or **reference group**, and all coefficients are interpreted relative to this baseline.

In our example, non-calculus-takers are the baseline. The coefficient $\hat{\beta}_2 = 5.8$ means calculus-takers score 5.8 points higher *than non-calculus-takers*.

7.3 Multiple Categories

What if our categorical variable has more than two values? Consider the Palmer Penguins data, where species can be Adelie, Chinstrap, or Gentoo.

```
set.seed(12487345)

ex_data <- penguins |>
  sample_n(5) |>
  select(body_mass_g, flipper_length_mm, sex, species)

knitr::kable(ex_data, caption = "Palmer Penguins Sample")
```

Table 7.3: Palmer Penguins Sample

body_mass_g	flipper_length_mm	sex	species
3450	193	female	Adelie
3475	187	male	Adelie
4650	184	male	Adelie

body_mass_g	flipper_length_mm	sex	species
3650	187	female	Chinstrap
4600	209	female	Gentoo

To include species in a regression, we create $g - 1$ **dummy variables** where g is the number of categories. With three species, we need two dummy variables:

```
ex_data <- ex_data |>
  mutate(species_Chinstrap = case_when(species == "Chinstrap" ~ 1, TRUE ~ 0),
         species_Gentoo = case_when(species == "Gentoo" ~ 1, TRUE ~ 0))

knitr::kable(ex_data, caption = "Penguins with Dummy Variables")
```

Table 7.4: Penguins with Dummy Variables

body_mass_g	flipper_length_mm	sex	species	species_Chinstrap	species_Gentoo
3450	193	female	Adelie	0	0
3475	187	male	Adelie	0	0
4650	184	male	Adelie	0	0
3650	187	female	Chinstrap	1	0
4600	209	female	Gentoo	0	1

Notice that Adelie penguins have 0 for both dummies—they are the omitted baseline category.

Now we can estimate the effect of body mass on flipper length, controlling for species:

$$\text{flipper_length} = \beta_0 + \beta_1(\text{body_mass}) + \beta_2(\text{Chinstrap}) + \beta_3(\text{Gentoo}) + \mu$$

```
penguin_data <- penguins |>
  mutate(species_Chinstrap = case_when(species == "Chinstrap" ~ 1, TRUE ~ 0),
         species_Gentoo = case_when(species == "Gentoo" ~ 1, TRUE ~ 0))

lm_penguin <- lm(flipper_length_mm ~ body_mass_g + species_Chinstrap + species_Gentoo,
                 data = penguin_data)

summary(lm_penguin)
```

```

Call:
lm(formula = flipper_length_mm ~ body_mass_g + species_Chinstrap +
    species_Gentoo, data = penguin_data)

Residuals:
    Min       1Q   Median       3Q      Max
-14.5455  -3.1845   0.1307   3.3533  17.5313

Coefficients:
              Estimate Std. Error t value Pr(>|t|)
(Intercept)    158.8602606    2.3865767   66.564 < 0.0000000000000002 ***
body_mass_g      0.0084021    0.0006339   13.255 < 0.0000000000000002 ***
species_Chinstrap  5.5974403    0.7882166    7.101  0.000000000000733 ***
species_Gentoo   15.6774699    1.0906591   14.374 < 0.0000000000000002 ***
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 5.395 on 338 degrees of freedom
(2 observations deleted due to missingness)
Multiple R-squared:  0.8541,    Adjusted R-squared:  0.8528
F-statistic: 659.4 on 3 and 338 DF,  p-value: < 0.00000000000000022

```

Interpretation:

- $\hat{\beta}_1 \approx 0.008$: A 1-gram increase in body mass is associated with a 0.008 mm increase in flipper length, holding species constant.
- $\hat{\beta}_2 \approx 5.6$: Chinstrap penguins have flippers about 5.6 mm longer than Adelie penguins of the same body mass.
- $\hat{\beta}_3 \approx 15.7$: Gentoo penguins have flippers about 15.7 mm longer than Adelie penguins of the same body mass.

7.3.1 Choosing the Baseline Category

How should we choose which category to omit? It depends on our research question.

If we only care about controlling for the categorical variable (not interpreting its coefficients), then the choice doesn't matter—the coefficient on our main variable of interest won't change.

If we do care about interpreting the dummy coefficients, choose a baseline that makes interpretation natural. For studying the gender wage gap, making “male” the baseline means the female coefficient directly measures the wage gap relative to men.

Let's verify that changing the baseline doesn't affect other coefficients:

```
# Switching baseline to Gentoo
penguin_data <- penguin_data |>
  mutate(species_Adelie = case_when(species == "Adelie" ~ 1, TRUE ~ 0))

lm_penguin2 <- lm(flipper_length_mm ~ body_mass_g + species_Chinstrap + species_Adelie,
  data = penguin_data)

# Compare body mass coefficients
cat("Body mass coefficient (Adelie baseline):", round(coef(lm_penguin)["body_mass_g"], 5), "
```

Body mass coefficient (Adelie baseline): 0.0084

```
cat("Body mass coefficient (Gentoo baseline):", round(coef(lm_penguin2)["body_mass_g"], 5))
```

Body mass coefficient (Gentoo baseline): 0.0084

The coefficient on body mass is identical regardless of which species is the baseline.

7.4 Factor Variables in R

Creating dummy variables manually is tedious, especially when categories are numerous. R's **factor** variables automate this process.

When you include a factor variable in a regression, R automatically creates the necessary dummy variables and omits one category:

```
# Convert species to factor
penguin_data <- penguins |>
  mutate(species_f = factor(species))

# R handles dummy creation automatically
lm_factor <- lm(flipper_length_mm ~ body_mass_g + species_f, data = penguin_data)
summary(lm_factor)
```

Call:

```
lm(formula = flipper_length_mm ~ body_mass_g + species_f, data = penguin_data)
```

Residuals:

Min	1Q	Median	3Q	Max
-14.5455	-3.1845	0.1307	3.3533	17.5313

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	158.8602606	2.3865767	66.564	< 0.0000000000000002 ***
body_mass_g	0.0084021	0.0006339	13.255	< 0.0000000000000002 ***
species_fChinstrap	5.5974403	0.7882166	7.101	0.000000000000733 ***
species_fGentoo	15.6774699	1.0906591	14.374	< 0.0000000000000002 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 5.395 on 338 degrees of freedom

(2 observations deleted due to missingness)

Multiple R-squared: 0.8541, Adjusted R-squared: 0.8528

F-statistic: 659.4 on 3 and 338 DF, p-value: < 0.0000000000000002

Notice the coefficients are identical to our manual approach. R chose “Adelie” as the baseline because it comes first alphabetically.

To change the baseline category, use the `relevel()` function:

```
# Make Gentoo the baseline
penguin_data <- penguin_data |>
  mutate(species_f = relevel(species_f, ref = "Gentoo"))

lm_factor2 <- lm(flipper_length_mm ~ body_mass_g + species_f, data = penguin_data)
summary(lm_factor2)
```

Call:

```
lm(formula = flipper_length_mm ~ body_mass_g + species_f, data = penguin_data)
```

Residuals:

Min	1Q	Median	3Q	Max
-14.5455	-3.1845	0.1307	3.3533	17.5313

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	174.5377305	3.2542422	53.634	< 0.0000000000000002 ***
body_mass_g	0.0084021	0.0006339	13.255	< 0.0000000000000002 ***
species_fAdelie	-15.6774699	1.0906591	-14.374	< 0.0000000000000002 ***

```
species_fChinstrap -10.0800297 1.1787380 -8.552 0.000000000000000428 ***
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```

Residual standard error: 5.395 on 338 degrees of freedom

(2 observations deleted due to missingness)

Multiple R-squared: 0.8541, Adjusted R-squared: 0.8528

F-statistic: 659.4 on 3 and 338 DF, p-value: < 0.00000000000000022


Now coefficients are relative to Gentoo penguins.

7.5 Non-Linear Relationships: Polynomial Models

So far we've assumed linear relationships between variables. But many economic relationships are non-linear. Consider the effect of work experience on wages: at low experience levels, additional experience substantially increases productivity and wages; but at very high experience levels, workers may become less productive as they age.

This suggests a **quadratic** relationship—wages increase with experience at first, then eventually decrease:

Figure 7.2: A quadratic model (orange) often fits experience-wage data better than a linear model (blue).



7.5.1 Estimating Quadratic Models

To estimate a quadratic model, we simply include both the variable and its square:

$$\log(\text{wage}) = \beta_0 + \beta_1(\text{exper}) + \beta_2(\text{exper}^2) + \mu$$

```
wage_data <- wooldridge::wage1 |>
  mutate(exper2 = exper^2)

lm_quad <- lm(log(wage) ~ exper + exper2, data = wage_data)
summary(lm_quad)
```

Call:

```
lm(formula = log(wage) ~ exper + exper2, data = wage_data)
```

Residuals:

Min	1Q	Median	3Q	Max
-2.05828	-0.33126	-0.05044	0.31806	1.47493

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	1.2952908	0.0494812	26.177	< 0.0000000000000002 ***
exper	0.0455341	0.0058595	7.771	0.0000000000000415 ***
exper2	-0.0009439	0.0001291	-7.312	0.0000000000009933 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 0.5041 on 523 degrees of freedom

Multiple R-squared: 0.104, Adjusted R-squared: 0.1006

F-statistic: 30.35 on 2 and 523 DF, p-value: 0.0000000000003377

7.5.2 Interpreting Quadratic Models

With a quadratic term, the marginal effect of x on y is no longer constant. Taking the derivative:

$$\frac{\partial y}{\partial x} = \beta_1 + 2\beta_2 x$$

The effect of experience **depends on your current level of experience.**

With our estimates ($\hat{\beta}_1 \approx 0.046$, $\hat{\beta}_2 \approx -0.0009$):

For a worker with **10 years** of experience:

$$0.046 + 2(-0.0009)(10) = 0.028 \approx 2.8\%$$

An additional year of experience increases wages by about 2.8%.

For a worker with **45 years** of experience:

$$0.046 + 2(-0.0009)(45) = -0.035 \approx -3.5\%$$

An additional year actually *decreases* wages—the worker is past the peak of the experience-wage relationship.

i Finding the Peak

The experience level where wages are maximized occurs where the marginal effect equals zero:

$$\beta_1 + 2\beta_2 x^* = 0 \implies x^* = -\frac{\beta_1}{2\beta_2}$$

With our estimates: $x^* = -\frac{0.046}{2(-0.0009)} \approx 25.6$ years.

7.6 Logarithmic Transformations

We've already seen that taking the log of the dependent variable changes how we interpret coefficients. Let's review the different cases systematically.

7.6.1 Log-Linear Models

When the dependent variable is logged but independent variables are not:

$$\ln(Y) = \beta_0 + \beta_1 X + \mu$$

Interpretation: A **one-unit** increase in X is associated with a $100 \times \beta_1$ **percent** change in Y .

This works because $\ln(Y_1) - \ln(Y_0) = \ln(Y_1/Y_0) \approx (Y_1 - Y_0)/Y_0$ for small changes.

⚠ Approximation Warning

The “multiply by 100” interpretation is an approximation that works well when β_1 is small (less than about 0.2 in absolute value). For larger coefficients, use the exact formula: percent change = $100 \times (e^{\beta_1} - 1)$.

7.6.2 Linear-Log Models

When independent variables are logged but the dependent variable is not:

$$Y = \beta_0 + \beta_1 \ln(X) + \mu$$

Interpretation: A **1 percent** increase in X is associated with a $\beta_1/100$ **unit** change in Y .

7.6.3 Log-Log Models (Elasticities)

When both sides are logged:

$$\ln(Y) = \beta_0 + \beta_1 \ln(X) + \mu$$

Interpretation: A **1 percent** increase in X is associated with a β_1 **percent** change in Y .

This is the **elasticity** of Y with respect to X —a unit-free measure of responsiveness.

7.6.4 Summary Table

Model	Equation	Interpretation of β_1
Linear-Linear	$Y = \beta_0 + \beta_1 X$	1-unit $\uparrow$ in $X \rightarrow \beta_1$ -unit change in Y
Log-Linear	$\ln(Y) = \beta_0 + \beta_1 X$	1-unit $\uparrow$ in $X \rightarrow (100 \times \beta_1)\%$ change in Y
Linear-Log	$Y = \beta_0 + \beta_1 \ln(X)$	1% $\uparrow$ in $X \rightarrow \beta_1/100$ unit change in Y
Log-Log	$\ln(Y) = \beta_0 + \beta_1 \ln(X)$	1% $\uparrow$ in $X \rightarrow \beta_1\%$ change in Y (elasticity)

7.7 Application: Policy Evaluation with Dummy Variables

Dummy variables are essential for evaluating policies and programs. Suppose we want to estimate the effect of a job training program on wages:

$$\ln(wage) = \beta_0 + \beta_1(training) + \beta_2(educ) + \beta_3(exper) + \mu$$

where `training` equals 1 if the person participated in the program and 0 otherwise.

The coefficient β_1 measures the **treatment effect**—the percentage difference in wages between program participants and non-participants, controlling for education and experience.

This is essentially a regression-based version of comparing treatment and control groups, where the control variables help us account for differences between those who did and didn't participate.

7.8 Summary

This chapter introduced two powerful extensions to basic regression: categorical variables and non-linear functional forms.

Dummy variables allow us to include qualitative information in regression models. The coefficient on a dummy variable represents the difference in intercepts between groups—the effect of having a particular characteristic relative to the baseline category. When a categorical variable has g categories, we include $g - 1$ dummy variables to avoid perfect multicollinearity (the dummy variable trap).

Polynomial models capture non-linear relationships by including squared or higher-order terms. With quadratic models, the marginal effect of x on y depends on the current level of x , allowing us to model relationships that increase and then decrease (or vice versa).

Logarithmic transformations change the interpretation of coefficients: log-linear models give percentage changes, linear-log models give the effect of percentage changes, and log-log models give elasticities. These transformations are especially useful when dealing with variables like wages or prices that are right-skewed and where percentage changes are more meaningful than absolute changes.

7.9 Check Your Understanding

Note: This section contains interactive content only available in the HTML version.

8 Heteroskedasticity

Key Questions

- What is homoskedasticity and why is it a Gauss-Markov assumption?
- What happens to OLS estimates when heteroskedasticity is present?
- How does heteroskedasticity affect our hypothesis tests?
- What are heteroskedasticity-robust standard errors and when should we use them?
- How can we detect heteroskedasticity in our data?

Suggested Readings

- Wooldridge (2019), Ch. 8

One of the key assumptions underlying OLS is that the variance of the error term is constant across all values of the explanatory variables. When this assumption fails, we have **heteroskedasticity**—and while our coefficient estimates remain unbiased, our standard errors become unreliable. This chapter explores what heteroskedasticity is, why it matters, and how to fix it.

8.1 The Homoskedasticity Assumption

8.1.1 Gauss-Markov Assumption 5

Recall the Gauss-Markov assumptions that make OLS the Best Linear Unbiased Estimator (BLUE). The fifth assumption is **homoskedasticity**:

Homoskedasticity

The error term μ has the **same variance** given any value of the explanatory variables:

$$\text{Var}(\mu|x_1, x_2, \dots, x_k) = \sigma^2$$


The variance is constant—it doesn't depend on x .

This assumption is crucial because it allows us to derive the correct formula for the variance of $\hat{\beta}$, which in turn gives us valid standard errors, t-statistics, and p-values.

8.1.2 What Does Homoskedasticity Look Like?

In a homoskedastic regression, the spread of observations around the regression line is **constant** across all values of x .

Figure 8.1: Homoskedastic data: the spread of points around the regression line is constant across all values of x .



Notice how the “cone” of data points has roughly the same width whether x is small or large. This is homoskedasticity in action.

8.2 Heteroskedasticity: When Variance Changes

8.2.1 What Is Heteroskedasticity?

Heteroskedasticity occurs when the variance of the error term changes with the level of the explanatory variable:


$$Var(\mu_i|x_i) = \sigma_i^2$$

The subscript i indicates that variance depends on observation i 's value of x .

8.2.2 What Does Heteroskedasticity Look Like?

The classic pattern is a “fan” or “cone” shape, where the spread of data increases (or decreases) as x changes:

Figure 8.2: Heteroskedastic data: the spread of points around the regression line increases with x , creating a ‘fan’ or ‘cone’ pattern.



This pattern is extremely common in economic data. For example:

- **Income and consumption:** Higher-income households have more variable consumption patterns
- **Firm size and profits:** Larger firms have more variable profit margins
- **Experience and wages:** More experienced workers have more variable wages


8.2.3 Side-by-Side Comparison

8.3 Why Does Heteroskedasticity Matter?

8.3.1 The Good News: Coefficients Are Still Unbiased

Here's the crucial point: **heteroskedasticity does NOT bias our coefficient estimates.** The unbiasedness of OLS depends only on Gauss-Markov assumptions 1-4, not on homoskedasticity.

Figure 8.3: Comparing homoskedastic (left) and heteroskedastic (right) data. The key difference is whether the spread of points changes with x .



So $E[\hat{\beta}] = \beta$ still holds. Our estimates are centered on the truth.

8.3.2 The Bad News: Standard Errors Are Wrong

However, the standard error formulas we've been using assume homoskedasticity. When this assumption is violated:

- The formula $SE(\hat{\beta}_1) = \frac{\hat{\sigma}}{\sqrt{SST_x}}$ is **invalid**
- Our computed standard errors will typically be **too small**
- This means t-statistics are **too large**
- And p-values are **too small**

The result? **We reject the null hypothesis too often**, even when it's true.

8.3.3 Type I Error: A Quick Review

! Type I Error

A **Type I error** (false positive) occurs when we reject a true null hypothesis. When we set $\alpha = 0.05$, we're accepting a 5% probability of making this error. If our test is working correctly, we should reject true null hypotheses exactly 5% of the time.

With heteroskedasticity and standard OLS standard errors, the actual Type I error rate can be much higher than 5%—sometimes 10%, 15%, or even higher.

8.4 Simulation: Seeing the Problem

8.4.1 Setup: Simulating Hypothesis Tests

Let's run a simulation to see exactly what goes wrong. We'll generate data where the **true effect is zero** ($\beta_1 = 0$), then see how often we incorrectly reject this true null hypothesis.

First, let's see what happens under **homoskedasticity** (where everything works correctly):

```
# Simulation parameters
n_sims <- 2000      # Number of simulations
n_obs <- 250        # Observations per sample
beta0_true <- 5     # True intercept
beta1_true <- 0     # True slope (NULL HYPOTHESIS IS TRUE!)

set.seed(42)

# Function for ONE simulation trial under HOMOSKEDASTICITY
run_homo_trial <- function(i) {
  x <- runif(n_obs, 1, 50)
  error <- rnorm(n_obs, mean = 0, sd = 10) # Constant variance
  y <- beta0_true + beta1_true * x + error

  model <- lm(y ~ x)

  # Get OLS p-value
  pval_ols <- summary(model)$coefficients["x", "Pr(>|t|)"]


  return(data.frame(pval_ols = pval_ols))
}

# Run simulations
homo_results <- map_dfr(1:n_sims, run_homo_trial)

# Calculate rejection rate (Type I error rate)
homo_reject_rate <- mean(homo_results$pval_ols < 0.05)
```

Under homoskedasticity, the Type I error rate is 5.7%—almost exactly 5%, as expected.

Figure 8.4: Under homoskedasticity, p-values are uniformly distributed when the null is true, and the rejection rate matches our significance level.



8.4.2 Now With Heteroskedasticity

Now let's see what happens when we have heteroskedasticity but use standard OLS standard errors:

```
# Function for ONE simulation trial under HETEROSKEDASTICITY
run_hetero_trial <- function(i) {
  x <- runif(n_obs, 1, 50)
  # Variance INCREASES with x (heteroskedasticity!)
  error <- rnorm(n_obs, mean = 0, sd = sqrt(0.2 * x^3))
  y <- beta0_true + beta1_true * x + error

  model <- lm(y ~ x)

  # Get both OLS and robust p-values
  res_ols <- coeftest(model)
  res_robust <- coeftest(model, vcov. = vcovHC(model, type = "HC1"))

  return(data.frame(
    pval_ols = res_ols["x", "Pr(>|t|)"],
    pval_robust = res_robust["x", "Pr(>|t|)"]
  ))
}
```

```


))
}

# Run simulations
hetero_results <- map_dfr(1:n_sims, run_hetero_trial)

# Calculate rejection rates
ols_reject_rate <- mean(hetero_results$pval_ols < 0.05)
robust_reject_rate <- mean(hetero_results$pval_robust < 0.05)

```

Figure 8.5: Under heteroskedasticity, standard OLS p-values pile up near zero, causing excessive rejections. Robust standard errors fix this problem.



The difference is stark! With standard OLS standard errors (left panel), we reject the true null hypothesis 10.6% of the time—more than **twice** our intended 5% rate. This is a serious problem: we’re finding “significant” effects that don’t exist.

With heteroskedasticity-robust standard errors (right panel), the rejection rate returns to approximately 5%.

8.5 Heteroskedasticity-Robust Standard Errors

8.5.1 The Solution

The fix is remarkably simple: use a different formula for the standard errors that accounts for heteroskedasticity. These are called **heteroskedasticity-robust standard errors** (also known as White standard errors or Huber-White standard errors).

The standard OLS variance formula is:

$$Var(\hat{\beta}_1) = \frac{\sum_{i=1}^n \hat{u}_i^2 / (n-2)}{SST_x^2}$$


The heteroskedasticity-robust formula is:

$$Var(\hat{\beta}_1) = \frac{\sum_{i=1}^n (x_i - \bar{x})^2 \hat{u}_i^2}{SST_x^2}$$

The key difference is that robust standard errors **weight each squared residual by how far its x_i is from the mean**. This accounts for the fact that residuals may have different variances at different values of x .

8.5.2 Visualizing the Sampling Distributions

Figure 8.6: The actual sampling distribution (blue) compared to what standard OLS SEs predict (red dashed) vs. robust SEs (green). OLS SEs are too narrow; robust SEs match reality.




The figure shows that:

- The **actual** sampling distribution (blue) has heavier tails than OLS standard errors predict
- Standard OLS SEs (red dashed) assume a narrower distribution than reality
- Robust SEs (green) correctly capture the true variability

8.5.3 Comparing Type I Error Rates

Figure 8.7: Type I error rates across different scenarios. Robust standard errors maintain the correct 5% rate even with heteroskedasticity.



8.6 Implementing Robust Standard Errors in R

8.6.1 Using fixest

The easiest way to get robust standard errors is with the `fixest` package, which we'll use throughout this course:

```
# Generate some heteroskedastic data
set.seed(123)
example_data <- tibble(
  x = runif(200, 1, 50),
  y = 5 + 0.5 * x + rnorm(200, 0, sqrt(0.1 * x^2))
)

# Standard OLS (wrong SEs with heteroskedasticity)
model_ols <- feols(y ~ x, data = example_data)
```

	Standard SE	Robust SE
(Intercept)	5.353*** (1.216)	5.353*** (0.906)
x	0.482*** (0.042)	0.482*** (0.045)
Num.Obs.	200	200
R2	0.402	0.402

+ p < 0.1, * p < 0.05, ** p < 0.01, *** p < 0.001

```
# With heteroskedasticity-robust SEs
model_robust <- feols(y ~ x, vcov = "HC1", data = example_data)
```

```
# Compare the results
modelsummary(
  list("Standard SE" = model_ols, "Robust SE" = model_robust),
  stars = TRUE,
  gof_map = c("nobs", "r.squared")
)
```

Notice that:

- The **coefficients are identical** (unbiasedness is not affected)
- The **standard errors differ** (robust SEs are typically larger)
- This affects **t-statistics and p-values**

8.6.2 Best Practice: Always Use Robust Standard Errors

Best Practice

Always report heteroskedasticity-robust standard errors.

Why? Because:

1. If heteroskedasticity is present, robust SEs are correct and standard SEs are wrong
2. If homoskedasticity holds, robust SEs are still valid (just slightly less efficient)

You can't lose by using robust SEs, but you can lose badly by not using them.

8.7 Detecting Heteroskedasticity


8.7.1 Residual Plots

The best way to detect heteroskedasticity is to **look at your residuals**. Plot the residuals against the fitted values (or against each explanatory variable) and look for patterns:

```
# Fit model to heteroskedastic data
model <- lm(y ~ x, data = example_data)

# Add residuals and fitted values to data
example_data <- example_data |>
  mutate(
    fitted = fitted(model),
    resid = residuals(model)
  )
```

Figure 8.8: Residual plot showing the ‘fan’ pattern characteristic of heteroskedasticity. The spread of residuals increases with fitted values.



8.7.2 What to Look For

Homoskedasticity: Residuals form a random cloud with constant spread

Heteroskedasticity: Look for:

- Fan/cone shapes (spread increases or decreases)
- Patterns in the spread related to fitted values
- Different variability for different groups

8.7.3 Formal Tests

While visual inspection is usually sufficient, you can also use formal tests like the **Breusch-Pagan test**:

```
# Breusch-Pagan test  
bptest(model)
```

studentized Breusch-Pagan test

```
data: model  
BP = 29.91, df = 1, p-value = 0.00000004526
```

A small p-value (< 0.05) suggests heteroskedasticity is present. However, **don't rely solely on formal tests**—always look at your residual plots too.

8.8 Summary

Heteroskedasticity is a common issue in economic data where the variance of errors changes with the explanatory variables.

Key Points:

1. **Heteroskedasticity does NOT bias coefficients** - $E[\hat{\beta}] = \beta$ still holds
2. **Heteroskedasticity DOES invalidate standard errors** - leading to incorrect inference
3. **The main symptom is inflated Type I error rates** - we reject true nulls too often
4. **The solution is simple: use robust standard errors** - specify `vcov = "HC1"` in `feols()`
5. **Detect heteroskedasticity with residual plots** - look for fan/cone patterns

Best Practice: Always use heteroskedasticity-robust standard errors. They're valid whether or not heteroskedasticity is present, so there's no downside to using them.

8.9 Check Your Understanding

Note: This section contains interactive content only available in the HTML version.

9 Interaction Effects

💡 Key Questions

- What happens when the effect of one variable depends on the value of another?
- How do we include and interpret interaction terms in regression?
- What's the difference between interacting two dummy variables versus a dummy and a continuous variable?
- How do interactions allow for different slopes across groups?

📖 Suggested Readings

- Wooldridge (2019), Ch. 7.4
- Bailey (2020), Ch. 5.4

In the previous chapter, we saw how dummy variables allow different groups to have different intercepts—the regression lines for different groups are parallel but shifted up or down. But what if the *effect* of a variable differs across groups? What if the returns to education are higher for men than for women, or the effect of experience on wages depends on whether someone has a college degree?

Interaction terms allow us to model these situations. When we include an interaction between two variables, we allow the effect of one variable to depend on the value of another.

9.1 Why Interactions Matter

Consider the gender wage gap. We might estimate:

$$\ln(wage) = \beta_0 + \beta_1(female) + \beta_2(educ) + \mu$$

This model assumes that the effect of education on wages (β_2) is the same for men and women. The only difference is the intercept—women's wage line is shifted down by β_1 .

But what if education has different effects for men and women? Perhaps discrimination means women receive lower returns to their education, or perhaps women are overrepresented in fields

where education is particularly valuable. To test this, we need to allow the *slope* on education to differ by gender.

This is exactly what interaction terms do.

9.2 Interactions Between Two Dummy Variables

Let's start with the simplest case: interacting two dummy variables. Suppose we want to understand how being female and being married jointly affect wages.

We create an **interaction term** by multiplying the two dummy variables:

$$\ln(\text{wage}) = \beta_0 + \beta_1(\text{female}) + \beta_2(\text{married}) + \beta_3(\text{female} \times \text{married}) + \mu$$

The interaction term $\text{female} \times \text{married}$ equals 1 only for married women and 0 for everyone else.

```
wage_data <- wooldridge::wage1 |>
  mutate(female_married = female * married)

lm1 <- lm(log(wage) ~ female + married + female_married, data = wage_data)
summary(lm1)
```

Call:

```
lm(formula = log(wage) ~ female + married + female_married, data = wage_data)
```

Residuals:

Min	1Q	Median	3Q	Max
-2.0240	-0.3245	-0.0800	0.3155	1.6849

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	1.52081	0.05099	29.827	< 0.0000000000000002 ***
female	-0.13164	0.06680	-1.971	0.0493 *
married	0.42669	0.06155	6.932	0.0000000000123 ***
female_married	-0.37479	0.08571	-4.373	0.0000148080201 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 0.4728 on 522 degrees of freedom

Multiple R-squared: 0.2132, Adjusted R-squared: 0.2087
 F-statistic: 47.15 on 3 and 522 DF, p-value: < 0.00000000000000022

9.2.1 Interpreting the Interaction

To understand what each coefficient means, let's work through the predicted wages for each group:

Group	female	married	female $\times$ married	Predicted $\ln(\text{wage})$
Unmarried men	0	0	0	β_0
Unmarried women	1	0	0	$\beta_0 + \beta_1$
Married men	0	1	0	$\beta_0 + \beta_2$
Married women	1	1	1	$\beta_0 + \beta_1 + \beta_2 + \beta_3$

From our estimates:

- **Unmarried men** (baseline): $\ln(\text{wage}) = 1.57$
- **Unmarried women**: The coefficient on **female** ($\hat{\beta}_1 \approx -0.13$) tells us unmarried women earn about 13% less than unmarried men.
- **Married men**: The coefficient on **married** ($\hat{\beta}_2 \approx 0.42$) tells us married men earn about 42% more than unmarried men—a large “marriage premium.”
- **Married women**: We need to add all applicable coefficients: $-0.13 + 0.42 + (-0.38) = -0.09$, so married women earn about 9% less than unmarried men.

! The Interaction Effect

The coefficient on the interaction term ($\hat{\beta}_3 \approx -0.38$) tells us that the marriage premium is **different** for women than for men. While married men earn 42% more than unmarried men, the marriage “premium” for women is only $0.42 + (-0.38) = 0.04$, or about 4%—much smaller. Equivalently, the gender wage gap is larger for married workers than for unmarried workers.

9.3 Interactions Between Dummies and Continuous Variables

Perhaps more powerful is interacting a dummy variable with a continuous variable. This allows the **slope**—not just the intercept—to differ across groups.

Suppose we want to test whether men and women have different returns to education:

$$\ln(wage) = \beta_0 + \beta_1(female) + \beta_2(educ) + \beta_3(female \times educ) + \mu$$

The interaction term $female \times educ$ equals 0 for all men (since $female = 0$) and equals the education level for women.

```
wage_data <- wooldridge::wage1 |>
  mutate(female_educ = female * educ)

lm2 <- lm(log(wage) ~ female + educ + female_educ, data = wage_data)
summary(lm2)
```

Call:

```
lm(formula = log(wage) ~ female + educ + female_educ, data = wage_data)
```

Residuals:

Min	1Q	Median	3Q	Max
-2.02673	-0.27468	-0.03721	0.26221	1.34740

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	0.82595465	0.11805024	6.997	0.000000000000808 ***
female	-0.36006448	0.18542958	-1.942	0.0527 .
educ	0.07722794	0.00898753	8.593	< 0.0000000000000002 ***
female_educ	-0.00006409	0.01450349	-0.004	0.9965

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 0.4459 on 522 degrees of freedom

Multiple R-squared: 0.3002, Adjusted R-squared: 0.2962

F-statistic: 74.65 on 3 and 522 DF, p-value: < 0.00000000000000022

9.3.1 Different Slopes for Different Groups

Let's write out the predicted wage equations for each group:

For men ($female = 0$):

$$\ln(wage) = \beta_0 + \beta_2(educ) = 0.389 + 0.082(educ)$$

For women ($female = 1$):


$$\begin{aligned}\ln(wage) &= (\beta_0 + \beta_1) + (\beta_2 + \beta_3)(educ) = (0.389 - 0.227) + (0.082 + 0.006)(educ) \\ &= 0.162 + 0.088(educ)\end{aligned}$$

Now we can see:

- **Men** start with a higher intercept (0.389 vs 0.162)
- **Women** have a slightly higher return to education (8.8% vs 8.2% per year)

The interaction coefficient ($\hat{\beta}_3 \approx 0.006$) is not statistically significant here, suggesting we cannot reject that men and women have the same returns to education. But the approach illustrates how interactions allow different slopes.

Figure 9.1: With an interaction term, men and women can have different slopes (returns to education), not just different intercepts.



i Dummy Variables vs. Interactions: A Summary

- **Dummy variable alone:** Different intercepts, same slopes (parallel lines)
- **Interaction with continuous variable:** Different intercepts AND different slopes

9.4 Computing Effects with Interactions

When interactions are present, computing the effect of a variable requires care. The effect of one variable depends on the value of the other variable in the interaction.

From our model:

$$\ln(wage) = \beta_0 + \beta_1(female) + \beta_2(educ) + \beta_3(female \times educ)$$

The effect of education on wages is:

$$\frac{\partial \ln(wage)}{\partial educ} = \beta_2 + \beta_3(female)$$

- For men ($female = 0$): Effect = $\beta_2 = 0.082$ (8.2% per year)
- For women ($female = 1$): Effect = $\beta_2 + \beta_3 = 0.082 + 0.006 = 0.088$ (8.8% per year)

Similarly, the effect of being female depends on education level:

$$\frac{\partial \ln(wage)}{\partial female} = \beta_1 + \beta_3(educ)$$

At 12 years of education: Effect = $-0.227 + 0.006(12) = -0.155$ (-15.5%)

At 16 years of education: Effect = $-0.227 + 0.006(16) = -0.131$ (-13.1%)

The gender wage gap **narrows** as education increases (though the interaction isn't statistically significant in this example).

9.5 Interactions Between Two Continuous Variables

We can also interact two continuous variables. For example, perhaps the effect of experience on wages depends on education level—more educated workers might have steeper experience-wage profiles because they're in careers with more room for advancement.

$$\ln(wage) = \beta_0 + \beta_1(educ) + \beta_2(exper) + \beta_3(educ \times exper) + \mu$$

```
wage_data <- wooldridge::wage1 |>
  mutate(educ_exper = educ * exper)

lm3 <- lm(log(wage) ~ educ + exper + educ_exper, data = wage_data)
summary(lm3)
```

Call:
lm(formula = log(wage) ~ educ + exper + educ_exper, data = wage_data)

Residuals:

	Min	1Q	Median	3Q	Max
	-2.05543	-0.29549	-0.05335	0.31047	1.44788

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	0.1532232	0.1673326	0.916	0.3603
educ	0.1030365	0.0127368	8.090	0.00000000000000422 ***
exper	0.0132641	0.0060374	2.197	0.0285 *
educ_exper	-0.0002473	0.0004945	-0.500	0.6172

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 0.4617 on 522 degrees of freedom
Multiple R-squared: 0.2497, Adjusted R-squared: 0.2454
F-statistic: 57.91 on 3 and 522 DF, p-value: < 0.00000000000000022

The interaction coefficient ($\hat{\beta}_3 \approx 0.0003$) is small but positive and marginally significant. This suggests that the returns to experience are slightly higher for more educated workers.

The effect of experience is:

$$\frac{\partial \ln(wage)}{\partial exper} = \beta_2 + \beta_3(educ)$$

For someone with 12 years of education: $0.019 + 0.0003(12) = 0.023$ (2.3% per year)

For someone with 16 years of education: $0.019 + 0.0003(16) = 0.024$ (2.4% per year)

9.6 Application: Decomposing the Gender Wage Gap

Economists have long sought to understand why women earn less than men on average. A classic approach uses regression with interactions to **decompose** the wage gap into:

1. **Explained differences:** Gaps due to differences in education, experience, occupation, etc.
2. **Unexplained differences:** Gaps that remain after controlling for observable characteristics (potentially reflecting discrimination)

Consider a comprehensive model:

$$\ln(wage) = \beta_0 + \beta_1(female) + \sum_j \beta_j(HumanCapital_j) + \sum_k \beta_k(Occupation_k) + \mu$$

- The human capital variables (education, experience) capture differences in worker productivity
- The occupation dummies capture sorting into different types of jobs
- The coefficient β_1 on **female** represents the **unexplained** gap—the wage difference that remains after accounting for these factors

Research by Blau and Kahn (2017) uses this approach with rich panel data. They find that while the raw gender wage gap has narrowed considerably since 1980, a substantial unexplained gap remains. Human capital differences explain less of the gap over time (as women have caught up in education and experience), but occupational and industry segregation continues to be important.

9.7 Best Practices for Using Interactions

9.7.1 Always Include Main Effects

When including an interaction term, always include the main effects too:

Correct:

$$y = \beta_0 + \beta_1(A) + \beta_2(B) + \beta_3(A \times B)$$

Incorrect:

$$y = \beta_0 + \beta_3(A \times B)$$

Omitting main effects forces the interaction to capture both the main effects and the interaction, leading to misleading coefficients.

9.7.2 Center Continuous Variables

When interacting continuous variables, centering them (subtracting the mean) can make coefficients more interpretable. After centering, the main effect coefficient gives the effect at the mean of the other variable, rather than at zero.

9.7.3 Test Whether Interactions Are Needed

Not every model needs interactions. Test whether the interaction coefficient is statistically significant before interpreting it substantively. You can also use F-tests to jointly test multiple interaction terms.

9.8 Summary

Interaction terms extend the flexibility of regression models by allowing the effect of one variable to depend on another.

Interactions between dummy variables allow us to estimate separate effects for groups defined by multiple categories. The interaction coefficient captures the additional effect of belonging to both categories beyond what we'd expect from adding the individual effects.

Interactions between dummies and continuous variables allow different slopes for different groups. Without interactions, a dummy variable only shifts the intercept (parallel lines). With interactions, each group can have its own slope.

Interactions between continuous variables allow the marginal effect of one variable to vary with the level of another. This is useful when economic theory suggests complementarities or diminishing returns.

When interactions are present, interpreting coefficients requires care. The effect of a variable involved in an interaction depends on the value of the other variable(s) in the interaction. Always compute effects at meaningful values (like means or specific policy-relevant points).

9.9 Check Your Understanding

Note: This section contains interactive content only available in the HTML version.

10 Panel Data and Fixed Effects

💡 Key Questions

- What distinguishes panel data from cross-sectional data?
- How can observing units over time help us address omitted variable bias?
- What is a fixed effect and what does it control for?
- How do we estimate fixed effects models in practice?
- What types of confounders can fixed effects eliminate, and what can they not?

📖 Suggested Readings

- Wooldridge (2019), Ch. 13-14

Throughout this course, we have emphasized that omitted variable bias poses the central threat to causal inference. We can add control variables, but we can only control for what we can measure. What about unobserved factors like natural ability, motivation, local culture, or a city's history?

Panel data—observing the same units over multiple time periods—offers a powerful solution. By tracking how outcomes change within the same unit over time, we can control for all characteristics of that unit that remain constant, even if we cannot observe or measure them directly. This chapter introduces panel data methods, with a focus on **fixed effects** estimation.

10.1 Cross-Sectional vs. Panel Data

10.1.1 Cross-Sectional Data

Most datasets we have examined so far are **cross-sectional**: each unit (person, firm, city) is observed at a single point in time. Examples include housing prices in Ames, Iowa in 2010; penguin measurements from a single expedition; or wage data from one year of the Current Population Survey.

With cross-sectional data, we can describe populations, identify correlations, and use multivariate regression to control for observable confounders. But we face a fundamental limitation: we cannot control for unobserved characteristics that differ across units.

10.1.2 Panel Data

Panel data (also called longitudinal data) tracks the same units across multiple time periods. Examples include:

- NBA players observed across multiple seasons
- Employment and wages in states tracked quarterly over several years
- Crime rates in cities measured annually for a decade
- Stock prices for companies tracked daily

The key feature is that we observe the **same unit** at **different times**, allowing us to see how that unit changes when circumstances change.

10.2 Why Panel Data Helps: The Intuition

Consider estimating the effect of unemployment on crime. With cross-sectional data from a single year, we might compare cities with high unemployment to cities with low unemployment. But cities differ in countless ways: geography, demographics, policing strategies, local culture, economic history. Many of these factors affect both unemployment and crime, creating omitted variable bias.

Now imagine we observe the **same cities** over two years. Some cities experience rising unemployment; others see it fall. We can ask: when unemployment rises *within a city*, does crime also rise?

This **within-unit comparison** automatically controls for everything about that city that stays constant over time—its geography, its historical legacy, its baseline demographics. We don't need to measure these factors; we just need them to be stable across the time periods we observe.

! The Key Insight

Panel data allows us to control for **all time-invariant characteristics** of each unit, whether observed or unobserved. The variation we use for identification comes from changes *within* units over time, not comparisons *across* different units.

10.3 The Fixed Effects Model

10.3.1 Setting Up the Model

Let's formalize this intuition. Suppose we observe units $i = 1, 2, \dots, N$ over time periods $t = 1, 2, \dots, T$. Our outcome is y_{it} and our variable of interest is x_{it} .

The panel data model is:

$$y_{it} = \beta_0 + \beta_1 x_{it} + a_i + \tau_t + \mu_{it} \quad (10.1)$$

where:

- y_{it} is the outcome for unit i at time t
- x_{it} is the explanatory variable (which varies across units and time)
- a_i is the **unit fixed effect**—capturing all time-invariant characteristics of unit i
- τ_t is the **time fixed effect**—capturing shocks that affect all units equally in period t
- μ_{it} is the idiosyncratic error term

The unit fixed effect a_i is the crucial element. It absorbs everything about unit i that doesn't change over time: genetics, geography, institutional history, baseline culture. In our crime example, a_i captures each city's fixed characteristics that affect crime rates.

10.3.2 What Does This Solve?

Recall that omitted variable bias occurs when an unobserved factor is correlated with both x and y . In the standard cross-sectional regression:

$$y_i = \beta_0 + \beta_1 x_i + \mu_i$$

any time-invariant unobserved factor correlated with x gets absorbed into the error term, biasing our estimate of β_1 .

With fixed effects, those time-invariant factors are captured by a_i and explicitly controlled for. As long as the confounders don't change over time, they cannot bias our estimate.

10.4 Estimating Fixed Effects: The Within Transformation

How do we actually estimate Equation 10.1 when we don't observe a_i ?

One approach is the **within transformation** (also called the “demeaning” approach). For each unit i , we compute the average of each variable over time:

$$\bar{y}_i = \beta_0 + \beta_1 \bar{x}_i + a_i + \bar{\tau} + \bar{\mu}_i$$

where $\bar{y}_i = \frac{1}{T} \sum_{t=1}^T y_{it}$ is unit i 's average outcome across all time periods.

Now subtract this time-averaged equation from the original:

$$y_{it} - \bar{y}_i = \beta_1(x_{it} - \bar{x}_i) + (\tau_t - \bar{\tau}) + (\mu_{it} - \bar{\mu}_i)$$

Notice what happened: the a_i term **completely disappears!** Since a_i is constant over time, $a_i - a_i = 0$.

We can write this more compactly as:

$$\ddot{y}_{it} = \beta_1 \ddot{x}_{it} + \ddot{\tau}_t + \ddot{\mu}_{it}$$

where the double-dots indicate “time-demeaned” variables (deviations from unit means).

i Why “Within” Transformation?

This estimator is called the “within” transformation because the variation used to identify β_1 comes entirely from variation *within* each unit over time. Cross-sectional differences between units are eliminated by the demeaning process.

10.4.1 Example: Unemployment and Crime

Let's see this in action. We have data on crime rates and unemployment for 46 cities observed in two years (1982 and 1987).

First, let's see what happens with simple cross-sectional regression using just 1987 data:

```
crime_data <- wooldridge::crime2

# Cross-sectional regression using only 1987
reg_cross <- lm(crmrte ~ unem, data = filter(crime_data, year == 87))
summary(reg_cross)
```

Call:

```
lm(formula = crmrte ~ unem, data = filter(crime_data, year ==  
      87))
```

Residuals:

Min	1Q	Median	3Q	Max
-57.55	-27.01	-10.56	18.01	79.75

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	128.378	20.757	6.185	0.00000018 ***
unem	-4.161	3.416	-1.218	0.23

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 34.6 on 44 degrees of freedom

Multiple R-squared: 0.03262, Adjusted R-squared: 0.01063

F-statistic: 1.483 on 1 and 44 DF, p-value: 0.2297

The coefficient on unemployment is **negative** (-4.16), suggesting higher unemployment is associated with *lower* crime. This counterintuitive result likely reflects omitted variable bias: cities with low unemployment might also have other characteristics (wealth, strong institutions) that lead to lower crime.

Now let's use both years and apply the within transformation:

```
# Create city identifier  
crime_data <- crime_data |>  
  mutate(city_id = rep(1:46, each = 2))  
  
# Compute city-level means  
city_means <- crime_data |>  
  group_by(city_id) |>  
  summarize(  
    mean_crmrte = mean(crmrte),  
    mean_unem = mean(unem),  
    mean_d87 = mean(d87)  
  )  
  
# Merge and demean  
crime_data <- crime_data |>  
  left_join(city_means, by = "city_id") |>
```

```
mutate(
  crmrte_demean = crmrte - mean_crmrte,
  unem_demean = unem - mean_unem,
  time_demean = d87 - mean_d87
)

# Estimate on demeaned data
reg_within <- lm(crmrte_demean ~ unem_demean + time_demean, data = crime_data)
summary(reg_within)
```

Call:

```
lm(formula = crmrte_demean ~ unem_demean + time_demean, data = crime_data)
```

Residuals:

Min	1Q	Median	3Q	Max
-26.458	-6.384	0.000	6.384	26.458

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	0.0000000000000001116	1.039332417007331255	0.000	1.000000
unem_demean	2.217999508245958040	0.617247710094903201	3.593	0.000535 ***
time_demean	15.402203621534521716	3.306166769286674967	4.659	0.0000111 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 9.969 on 89 degrees of freedom

Multiple R-squared: 0.1961, Adjusted R-squared: 0.178

F-statistic: 10.85 on 2 and 89 DF, p-value: 0.00006059

Now the coefficient on unemployment is **positive** (2.22)! After controlling for all time-invariant city characteristics, we find that when unemployment rises within a city, crime rises too. This makes much more intuitive sense.

The within transformation eliminated the bias from unobserved city characteristics that were driving the negative cross-sectional correlation.

10.5 Estimating Fixed Effects: The Dummy Variable Approach

In practice, we rarely compute the within transformation manually. Instead, we use an equivalent approach: including **dummy variables** for each unit.

We estimate:

$$y_{it} = \beta_0 + \beta_1 x_{it} + \sum_{j=2}^N \gamma_j D_j + \sum_{s=2}^T \delta_s T_s + \mu_{it}$$

where D_j is a dummy variable equal to 1 if the observation is from unit j , and T_s is a dummy equal to 1 if the observation is from time period s . We omit one unit and one time period to avoid the dummy variable trap.

This approach is mathematically equivalent to the within transformation but more flexible: it's easier to add control variables, and we can see the estimated fixed effects if desired.

```
# Create factor variable for city and dummy for 1987
crime_data <- crime_data |>
  mutate(
    city_f = factor(city_id),
    year_87 = ifelse(year == 87, 1, 0)
  )

# Estimate with dummy variables
reg_fe <- lm(crmrte ~ unem + year_87 + city_f, data = crime_data)

# Show just the key coefficients (not all 46 city dummies)
summary(reg_fe)$coefficients[1:4, ]
```

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	51.48923	12.3457756	4.170595	0.0001404348
unem	2.21800	0.8778658	2.526581	0.0151893177
year_87	15.40220	4.7021169	3.275589	0.0020604685
city_f2	17.29171	14.1954509	1.218116	0.2296714584

The coefficient on `unem` (2.22) is identical to our within transformation estimate!

The dummy variable coefficients themselves are usually not of primary interest, but they can provide useful information. For instance, we can see which cities have unusually high or low crime rates after accounting for unemployment.

10.6 What Fixed Effects Control For (and What They Don't)

Fixed effects are powerful, but they're not magic. It's crucial to understand exactly what they do and don't control for.

10.6.1 What Unit Fixed Effects Control For

Unit fixed effects (α_i) absorb all characteristics of unit i that are **constant over time**:

- Geography and location
- Historical legacy and founding conditions
- Stable institutional features
- Baseline demographics (to the extent they don't change much)
- Measurement differences (e.g., how crime is recorded)
- "Culture" and other hard-to-measure factors

10.6.2 What Time Fixed Effects Control For

Time fixed effects (τ_t) absorb all factors that **affect all units equally** in a given period:

- Macroeconomic conditions (recessions, booms)
- National policy changes
- Seasonal patterns
- Technological changes affecting everyone

10.6.3 What Fixed Effects Do NOT Control For

Fixed effects cannot eliminate bias from factors that vary **both across units and over time** in ways correlated with the treatment:

- Time-varying confounders specific to certain units
- Differential trends (some units changing faster than others)
- Reverse causality (if y affects x within a period)

Fixed Effects No OVB

Fixed effects dramatically reduce omitted variable bias by controlling for time-invariant confounders. But they don't eliminate all possible bias. Unit-specific, time-varying confounders can still cause problems. Always think carefully about what might be changing differently across your units over time.

For example, in our crime analysis, fixed effects control for each city's baseline characteristics. But if some cities increased their police budgets while others cut them, and these budget changes are correlated with unemployment changes, we would still have omitted variable bias. We could address this by adding police spending as an explicit control variable.

10.7 Adding Control Variables

We can include additional time-varying control variables in fixed effects models:

$$y_{it} = \beta_1 x_{it} + \gamma z_{it} + a_i + \tau_t + \mu_{it}$$

where z_{it} is a control variable that changes over time within units.

This helps address the remaining source of potential bias: unit-specific, time-varying confounders. Of course, we can only control for variables we observe and measure.

10.8 Practical Considerations

10.8.1 When to Use Fixed Effects

Fixed effects are most valuable when:

1. You have genuine panel data (same units observed multiple times)
2. There is meaningful variation in x within units over time
3. Time-invariant confounders are a major concern
4. The treatment effect is expected to be relatively immediate

10.8.2 When Fixed Effects May Not Work Well

Fixed effects may be less useful when:

1. Little within-unit variation exists (everyone's x is stable over time)
2. The treatment effect takes a long time to materialize
3. Time-varying confounders are the main concern
4. You have very few time periods (less precise estimates)

10.8.3 Standard Errors

With panel data, observations within the same unit are often correlated over time. Standard OLS standard errors assume independence and can be too small. In practice, researchers typically use **clustered standard errors** at the unit level to account for this correlation. We'll discuss this more in later chapters.

10.9 Summary

Panel data—observing the same units over time—offers a powerful tool for causal inference. By tracking how outcomes change within units when circumstances change, we can control for all time-invariant characteristics of each unit, whether observed or not.

The **fixed effects estimator** implements this idea. It can be computed via the within transformation (demeaning the data) or equivalently by including dummy variables for each unit. The coefficient of interest is identified purely from within-unit variation over time.

Fixed effects control for:

- All time-invariant unit characteristics (observed and unobserved)
- All time-specific shocks affecting all units equally (when time fixed effects are included)

Fixed effects do NOT control for:

- Unit-specific, time-varying confounders
- Differential trends across units
- Reverse causality

Panel methods are not a complete solution to omitted variable bias, but they represent a major improvement over cross-sectional analysis when time-invariant confounders are a primary concern.

10.10 Check Your Understanding

Note: This section contains interactive content only available in the HTML version.

11 Difference-in-Differences

💡 Key Questions

- What is a natural experiment and how does it help with causal inference?
- What is the parallel trends assumption and why is it crucial for DiD?
- How do we estimate the average treatment effect on the treated (ATT) using DiD?
- What is two-way fixed effects and how does it relate to DiD?
- How do we choose good control units for a DiD design?

i Suggested Readings

- TBD

We have built up an increasingly powerful toolkit for causal inference. Randomized controlled trials remain the gold standard, but are often infeasible. Multivariate OLS allows us to control for observable confounders, and fixed effects extend this to control for unobserved time-invariant factors. But what if even fixed effects aren't enough?

This chapter introduces **difference-in-differences (DiD)**, a method that combines the logic of experiments with the practicality of observational data. Instead of relying solely on control *variables*, DiD uses control *units*—a comparison group that shows us what would have happened to the treated group in the absence of treatment.

11.1 Natural Experiments

11.1.1 The Problem with Observational Data

Suppose we want to estimate the effect of immigration on native workers' wages. A naive approach might compare wages in cities with high immigration to wages in cities with low immigration. But this comparison is confounded: immigrants tend to move to cities with strong economies, creating a spurious positive correlation between immigration and wages.

Even with control variables, we may miss important unobserved factors like a city's economic trajectory or its attractiveness to both immigrants and high-wage employers. Fixed effects help

with time-invariant confounders, but cannot address factors that change differently across cities over time.

11.1.2 Natural Experiments as a Solution

A **natural experiment** occurs when some external force—nature, policy, or historical accident—assigns units to treatment and control groups in a way that resembles random assignment.

Natural Experiments

A **natural experiment** is a study based on observational data where “treatment” is assigned by forces outside the researcher’s control, such as policy changes, geographic boundaries, or historical events. Natural experiments create conditions that *resemble* random assignment, allowing us to make causal inferences without actually randomizing.

The key insight is that if treatment assignment is “as good as random,” then comparing treated and control units can identify causal effects—just like in an RCT.

11.2 The Mariel Boatlift: A Classic Natural Experiment

One of the most famous natural experiments in economics involves the Mariel Boatlift. In 1980, Cuba experienced severe economic turmoil, and Fidel Castro announced that anyone who wished to leave could do so. Between April and October, approximately 125,000 Cubans departed from Mariel Harbor and arrived in Miami.

This mass migration had nothing to do with labor market conditions in Miami. It was driven entirely by Cuban politics and Miami’s geographic proximity to Cuba. In other words, the “treatment” of a sudden large increase in labor supply was essentially randomly assigned to Miami.

David Card’s seminal 1990 study used this natural experiment to estimate the effect of immigration on native wages. But how exactly do we use such an event to identify causal effects?

11.3 The Difference-in-Differences Approach

11.3.1 Why Simple Before-After Comparisons Fail

Card first compared wages in Miami before the boatlift (1979) to wages after (1980-1985):

City	Pre (1979)	Post (1980-1985)	Difference
Miami	1.85	1.83	-0.02

Wages fell slightly. Can we interpret this as the causal effect of immigration?

No. Wages might have fallen everywhere during this period due to recession or other macroeconomic factors. Looking only at the treated unit’s change over time confuses the treatment effect with broader trends.

11.3.2 The Need for Control Units

To isolate the treatment effect, we need to know what *would have happened* to Miami’s wages if the boatlift had never occurred. This **counterfactual** is inherently unobservable—we cannot see the parallel universe where Cuba didn’t open its borders.

But we can approximate it using **control units**: cities similar to Miami that were not affected by the boatlift. Card chose Atlanta, Los Angeles, Houston, and Tampa-St. Petersburg as controls, arguing they had similar demographics and economic trajectories to Miami.

City	Pre (1979)	Post (1980-1985)	Difference
Miami	1.85	1.83	-0.02
Controls	1.93	1.91	-0.02
Difference			0.00

The control cities experienced the *same* wage decline as Miami. This suggests the wage drop in Miami was not caused by the boatlift but by broader economic trends affecting all cities.

The **difference-in-differences** estimate is the change in Miami minus the change in controls:

$$(-0.02) - (-0.02) = 0.00$$

Card concluded that the Mariel Boatlift had essentially no effect on native wages—a surprising and influential finding.

11.4 The Parallel Trends Assumption

11.4.1 The Core Identifying Assumption

The DiD approach rests on one critical assumption:

! Parallel Trends Assumption

In the absence of treatment, the average outcome for the treated group would have followed the **same trend** as the average outcome for the control group.

Mathematically:

$$E[Y_{i,post} - Y_{i,pre}|Treated] = E[Y_{i,post} - Y_{i,pre}|Control]$$


in the counterfactual world where treatment never occurred.

This assumption is **not testable** directly because we never observe the treated group's counterfactual trend. However, we can assess its plausibility by examining whether treated and control groups followed similar trends *before* treatment began. If pre-treatment trends are parallel, it's more credible that they would have remained parallel absent treatment.

11.4.2 Visualizing DiD

The logic of DiD is best understood graphically:

Figure 11.1: The DiD estimate is the gap between the observed treatment outcome and the counterfactual (dashed line) based on the control group's trend.



The **Average Treatment Effect on the Treated (ATT)** is the vertical distance between:

1. The observed outcome for the treated group after treatment
2. The counterfactual: where the treated group *would have been* if it had followed the control group's trend

The dashed line represents this counterfactual—the treated group's hypothetical trajectory without treatment.

11.5 The DiD Regression

11.5.1 Setting Up the Model

We typically estimate DiD using regression. For the simple 2×2 case (two groups, two time periods), the model is:

$$y_{it} = \beta_0 + \beta_1 D_i + \beta_2 P_t + \delta(D_i \times P_t) + \mu_{it} \quad (11.1)$$

where:

- D_i is a **treatment group indicator** (1 if treated, 0 if control)
- P_t is a **post-period indicator** (1 if post-treatment, 0 if pre-treatment)
- $D_i \times P_t$ is the interaction term (1 only for treated units in the post period)
- δ is the **DiD estimate**—our estimate of the ATT

11.5.2 How It Works

To see why δ captures the DiD effect, consider the predicted outcomes for each group-period combination:

Group	Period	Predicted Outcome
Control	Pre	0
Control	Post	0 + 2
Treated	Pre	0 + 1
Treated	Post	0 + 1 + 2 +

Now compute the differences:

Change for treated group: (0 + 1 + 2 +) − (0 + 1) = 2 +

Change for control group: (0 + 2) − 0 = 2

DiD estimate: (2 +) − 2 =

The coefficient on the interaction term exactly equals the difference-in-differences!

11.5.3 Example: Estimating the Mariel Boatlift Effect

```
# Create the Card (1990) data
miami_wages <- tibble(
  city = "miami",
  year = 1979:1985,
  wages = c(1.85, 1.83, 1.85, 1.82, 1.82, 1.82, 1.82)
)

control_wages <- tibble(
  city = "control",
  year = 1979:1985,
  wages = c(1.93, 1.90, 1.91, 1.91, 1.90, 1.91, 1.92)
)

boatlift_data <- bind_rows(miami_wages, control_wages) |>
  mutate(
    post = ifelse(year >= 1980, 1, 0),
    treated = ifelse(city == "miami", 1, 0),
    post_treat = post * treated
  )

# Estimate DiD regression
did_reg <- feols(wages ~ treated + post + post_treat,
  vcov = "HC1",
  data = boatlift_data)

summary(did_reg)
```

OLS estimation, Dep. Var.: wages

Observations: 14

Standard-errors: Heteroskedasticity-robust

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	1.930000	0.00000100	1930000.000000	< 2.2e-16 ***
treated	-0.080000	0.00000100	-80000.000000	< 2.2e-16 ***
post	-0.021667	0.00331942	-6.527254	0.000066615 ***
post_treat	-0.001667	0.00628785	-0.265062	0.796345671

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
RMSE: 0.008522 Adj. R2: 0.947329

The coefficient on `post_treat` (approximately 0.01) is our DiD estimate. It suggests the Mariel Boatlift had essentially no effect on wages—consistent with Card’s original findings.

11.6 Two-Way Fixed Effects

11.6.1 Extending to Multiple Units and Time Periods

The simple 2×2 DiD works well for clean natural experiments with one treated unit and one control unit observed before and after. But what if we have many treated units, many control units, and many time periods?

The **Two-Way Fixed Effects (TWFE)** model extends DiD to these more complex settings:

$$y_{it} = \delta D_{it} + \alpha_i + \tau_t + \mu_{it} \quad (11.2)$$

where:

- D_{it} equals 1 if unit i is treated at time t , and 0 otherwise
- α_i are **unit fixed effects** (one for each unit)
- τ_t are **time fixed effects** (one for each time period)
- δ is the treatment effect

This model subsumes the simple DiD: unit fixed effects capture permanent differences between groups (like $\beta_1 D_i$), and time fixed effects capture common shocks to all units (like $\beta_2 P_t$).

11.6.2 Estimating TWFE in R

The `fixest` package makes TWFE estimation straightforward. The syntax `| unit + time` specifies the fixed effects:

```
# TWFE estimation
twfe_reg <- feols(wages ~ post_treat | city + year,
                  vcov = "HC1",
                  data = boatlift_data)

summary(twfe_reg)
```

```

OLS estimation, Dep. Var.: wages
Observations: 14
Fixed-effects: city: 2, year: 7
Standard-errors: Heteroskedasticity-robust
               Estimate Std. Error   t value Pr(>|t|)
post_treat -0.001667    0.006491 -0.256776  0.80758
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
RMSE: 0.00622      Adj. R2: 0.943875
               Within R2: 0.002193

```

The coefficient on `post_treat` is identical to our simple DiD regression—as expected, since TWFE and the 2×2 DiD model are equivalent in this simple case.

11.7 Research Design: Choosing Good Controls

11.7.1 The Art of Causal Inference

Your DiD estimate is only as good as your research design. The goal is to find control units that make the parallel trends assumption credible.

Good Controls → Good Counterfactual → Credible DiD Estimate

Bad Controls → Bad Counterfactual → Misleading DiD Estimate

11.7.2 Example: Minimum Wages and Employment

Consider estimating the effect of minimum wage increases on employment. One approach might compare all states that raised their minimum wage to all states that didn't.

But states that raise minimum wages are systematically different: they tend to be more politically liberal, have stronger labor markets, different industries, and many other characteristics that affect employment. These differences violate parallel trends—minimum-wage states may have been on different employment trajectories regardless of the policy.

Card and Krueger (1994) proposed a more compelling research design. When New Jersey raised its minimum wage in 1992 and neighboring Pennsylvania did not, they compared fast-food restaurants in counties *right next to each other* on opposite sides of the NJ-PA border.

The identifying assumption: restaurants separated only by a state border are similar in all relevant ways except the minimum wage policy. Geographic proximity makes parallel trends much more plausible than comparing NJ to distant states with very different economies.

11.7.3 Intuition-Driven vs. Data-Driven Selection

There are two broad approaches to choosing control units:

Intuition-driven selection relies on substantive knowledge to identify appropriate comparisons. Card’s choice of control cities for Miami was based on demographic and economic similarities. Card and Krueger’s border design was based on geographic proximity. The advantage is transparency and interpretability.

Data-driven selection uses algorithms to find control units that match the treated unit on pre-treatment outcomes. Methods like synthetic control construct a weighted average of potential controls that best reproduces the treated unit’s pre-treatment trajectory. The advantage is objectivity; the disadvantage is complexity and potential overfitting.

In practice, the best research designs often combine both approaches: use substantive knowledge to identify a reasonable set of potential controls, then verify that pre-treatment trends are indeed parallel.

11.8 What Can Go Wrong

11.8.1 Violations of Parallel Trends

The most serious threat to DiD is violation of the parallel trends assumption. This can happen when:

1. **Differential trends:** Treated and control units were already on different trajectories before treatment
2. **Anticipation effects:** Units change behavior before treatment actually begins
3. **Spillovers:** Treatment affects control units (e.g., workers move from treated to control areas)
4. **Composition changes:** The mix of units changes over time in ways correlated with treatment

11.8.2 Assessing Parallel Trends

While we cannot test parallel trends directly, we can assess plausibility by:

1. **Examining pre-treatment trends:** Do treated and control groups move together before treatment? Diverging pre-trends are a red flag.
2. **Placebo tests:** Estimate “effects” at times when no treatment occurred. Significant placebo effects suggest the design is flawed.
3. **Event studies:** Estimate treatment effects for each time period separately to visualize the dynamic pattern.

If pre-treatment trends are not parallel, the DiD estimate will be biased. The direction of bias depends on whether the treated group was trending upward or downward relative to controls.

11.9 Summary

Difference-in-differences is one of the most widely used methods for causal inference in economics and social science. It combines the intuition of experimental design with the practicality of observational data.

The key elements are:

1. **Natural experiments** provide plausibly exogenous variation in treatment assignment
2. **Control units** approximate what would have happened to treated units absent treatment
3. **The parallel trends assumption** requires that treated and control groups would have followed the same trajectory without treatment
4. **The DiD estimator** subtracts the control group's change from the treated group's change to isolate the treatment effect
5. **TWFE** extends DiD to multiple units and time periods using unit and time fixed effects

The credibility of any DiD analysis rests entirely on the parallel trends assumption. Careful research design—choosing appropriate control units and verifying parallel pre-trends—is essential for producing convincing causal estimates.

11.10 Check Your Understanding

Note: This section contains interactive content only available in the HTML version.

12 Advanced Difference-in-Differences

💡 Key Questions

- How can we use data to select better control units for DiD?
- What are propensity scores and how do they improve DiD estimates?
- How do we estimate treatment effects that change over time?
- What is an event study and how do we interpret it?
- Why do we need cluster-robust standard errors in panel data?

📖 Suggested Readings

- TBD

The basic difference-in-differences framework is powerful, but real-world applications often require more sophisticated tools. This chapter covers three important extensions: **data-driven control selection** using propensity scores, **dynamic DiD** for effects that evolve over time, and **cluster-robust standard errors** for proper inference with grouped data.

12.1 Data-Driven Control Selection


12.1.1 The Problem: Systematically Different Treatment and Control Groups

In the previous chapter, we emphasized that your DiD estimate is only as good as your counterfactual. When treated and control groups are systematically different, the parallel trends assumption becomes implausible.

Consider estimating the effect of the Affordable Care Act (ACA) Medicaid expansion on uninsured rates. States that expanded Medicaid are very different from states that didn't: they tend to be more politically liberal, have different demographics, different industries, and different baseline health care systems.

Using all non-expanding states as controls produces biased estimates because these states were on different trajectories *even before* the policy.

Figure 12.1: Without adjustment, treated and control states show diverging pre-trends, violating the parallel trends assumption.



The figure shows that even *before* the ACA expansion in 2014, treated states had steeper downward trends in uninsured rates. This pre-existing difference would be incorrectly attributed to the policy in a naive DiD analysis.

12.1.2 Propensity Score Weighting: The Intuition

The solution is to find control units that are *similar* to treated units. If we could compare only states that were “on the verge” of expanding but didn’t, we would have a much better counterfactual.

Propensity score methods formalize this intuition. The idea is:

1. Estimate each unit’s probability of being treated based on observable characteristics
2. Give more weight to control units that “look like” they should have been treated
3. Use these weights in the DiD regression

Control units with high propensity scores (high predicted probability of treatment) but who weren’t actually treated become our primary comparison group. They’re similar to treated units on observables, making parallel trends more plausible.

12.1.3 The IPW-DiD Algorithm

Inverse Propensity Score Weighted DiD (IPW-DiD) proceeds in three steps:

Step 1: Estimate the Propensity Score

We estimate a logistic regression predicting treatment status from pre-treatment characteristics:

$$P(D_i = 1|X) = G(\beta_0 + \beta_1 X_1 + \beta_2 X_2 + \dots + \beta_k X_k)$$

where $G()$ is the logistic function and the X variables are characteristics that predict treatment.

For the ACA example, we might include political leaning (Democratic vote share), region, and baseline poverty rates.

```
# Aggregate to state level for propensity score estimation
state_data <- panel_data |>
  filter(post == 0) |> # Use only pre-treatment data
  group_by(state_id, is_expanded) |>
  summarize(
    dem_share = mean(X1),
    region = mean(X2),
    poverty = mean(X3),
    .groups = 'drop'
  )

# Estimate propensity scores with logistic regression
ps_model <- glm(is_expanded ~ dem_share + region + poverty,
  data = state_data,
  family = binomial("logit"))

# Get predicted probabilities
state_data <- state_data |>
  mutate(ps_score = predict(ps_model, type = "response"))
```

Step 2: Calculate IPW Weights

The inverse propensity weights are:

$$IPW_i = D_i + (1 - D_i) \times \frac{\hat{P}_i}{1 - \hat{P}_i}$$

- For treated units ($D_i = 1$): $\text{weight} = 1$
- For control units ($D_i = 0$): $\text{weight} = \frac{\hat{P}_i}{1 - \hat{P}_i}$

Control units with high propensity scores get large weights; those with low propensity scores get small weights.

```
# Calculate IPW weights
state_data <- state_data |>
  mutate(
    ipw = case_when(
      is_expanded == 1 ~ 1,
      is_expanded == 0 ~ ps_score / (1 - ps_score)
    )
  )

# Show example weights
state_data |>
  select(state_id, is_expanded, ps_score, ipw) |>
  arrange(desc(ipw)) |>
  head(8) |>
  knitr::kable(digits = 2, caption = "Example IPW Weights")
```

Table 12.1: Example IPW Weights

state_id	is_expanded	ps_score	ipw
32	0	0.86	5.94
181	0	0.79	3.77
51	0	0.75	3.06
101	0	0.69	2.26
31	0	0.65	1.85
150	0	0.63	1.71
29	0	0.61	1.58
106	0	0.59	1.41

Step 3: Estimate Weighted TWFE

Finally, we estimate the standard TWFE DiD regression but using the IPW weights:


```
# Merge weights into panel data
panel_weighted <- panel_data |>
  left_join(select(state_data, state_id, ipw), by = "state_id")
```

```
# Naive (unweighted) DiD
did_naive <- feols(uninsured_rate ~ treat_active | state_id + year,
                  vcov = "HC1",
                  data = panel_data)

# IPW-weighted DiD
did_ipw <- feols(uninsured_rate ~ treat_active | state_id + year,
                 vcov = "HC1",
                 weights = ~ipw,
                 data = panel_weighted)
```

12.1.4 Comparing Results

Figure 12.2: IPW weighting restores parallel pre-trends by up-weighting control units similar to treated units.



The IPW-weighted trends are much more parallel in the pre-period! This gives us greater confidence in the parallel trends assumption.

Table 12.2: Comparing Naive and IPW-DiD Estimates (True effect = -5)

Method	Estimate	True Effect
Naive DiD (all controls equal weight)	-10.91	-5
IPW-Weighted DiD	-6.24	-5

The naive estimate is substantially biased (too negative) because it attributes pre-existing trends to the policy. The IPW-weighted estimate is much closer to the true effect of -5 percentage points.

! Key Insight

IPW-DiD works by reweighting the control group so that it looks more like the treatment group on observable characteristics. This makes parallel trends more plausible—but the method is only as good as your propensity score model. If important confounders are omitted, bias remains.

12.2 Dynamic DiD and Event Studies

12.2.1 Why Treatment Effects May Change Over Time

The basic DiD model estimates a single treatment effect that applies to all post-treatment periods. But many policies have effects that **evolve over time**:

- A job training program might have small initial effects that grow as workers gain experience
- A minimum wage increase might have immediate disemployment effects that fade as firms adjust
- An environmental regulation might have growing effects as compliance increases

Dynamic DiD allows us to estimate separate treatment effects for each time period, capturing how the policy impact unfolds.

12.2.2 Relative Time

Dynamic DiD models use **relative time**—the number of periods since treatment occurred—rather than calendar time. If treatment occurs in 2015:

Calendar Year	Relative Time
2012	-3
2013	-2
2014	-1
2015 (treatment)	0
2016	+1
2017	+2

This framing lets us estimate effects at each distance from treatment and, crucially, test whether pre-treatment “effects” are zero (as they should be if parallel trends holds).

12.2.3 The Dynamic DiD Model

The dynamic DiD model includes dummy variables for each relative time period:

$$y_{it} = \sum_{k \neq -1} \beta_k D_{it}^k + \alpha_i + \tau_t + \mu_{it}$$

where $D_{it}^k = 1$ if unit i is treated and period t is k periods from treatment.

We **omit** $k = -1$ (the period just before treatment) as the reference category. All coefficients are interpreted relative to this baseline.

12.2.4 Example: Environmental Policy and Manufacturing Employment


Suppose we want to estimate how an environmental regulation affects manufacturing employment in cities. We have data on 50 cities observed from 1995 to 2010, with 25 cities adopting the policy in 2005.

Let’s visualize the raw trends first:

The trends are parallel before 2005, then the treated cities experience a growing decline in manufacturing employment.

12.2.5 Estimating the Dynamic Model

Figure 12.3: Treatment and control cities show parallel pre-trends, then diverge after the policy in 2005.




```
# Estimate dynamic DiD (note: rel_t_treat_-1 is omitted as reference)
dyn_did <- feols(
  manufacturing_emp ~ `rel_t_treat_-10` + `rel_t_treat_-9` + `rel_t_treat_-8` +
    `rel_t_treat_-7` + `rel_t_treat_-6` + `rel_t_treat_-5` + `rel_t_treat_-4` +
    `rel_t_treat_-3` + `rel_t_treat_-2` + `rel_t_treat_0` + `rel_t_treat_1` +
    `rel_t_treat_2` + `rel_t_treat_3` + `rel_t_treat_4` + `rel_t_treat_5` |
    city_id + year,
  vcov = "HC1",
  data = env_data
)
```

12.2.6 The Event Study Plot

The best way to present dynamic DiD results is an **event study plot**. We plot each coefficient with its 95% confidence interval, with the x-axis showing relative time and a vertical line at the treatment date.

Figure 12.4: Event study plot showing treatment effects over time. Pre-treatment coefficients near zero support parallel trends; post-treatment coefficients show the growing negative effect.



12.2.7 Interpreting the Event Study

The event study plot reveals several important patterns:

1. **Pre-treatment coefficients ($k < -1$):** These cluster around zero with confidence intervals that include zero. This is strong evidence for parallel trends—before the policy, treated and control cities were on similar trajectories.
2. **Reference period ($k = -1$):** This is normalized to zero by construction.
3. **Post-treatment coefficients ($k \geq 0$):** These show the causal effect of the policy. The effect starts at about -100 in the adoption year and grows to about -500 by year 3, where it levels off.

i Why Pre-Trends Matter

The pre-treatment coefficients serve as a **placebo test**. If we found large, significant “effects” before the policy was implemented, that would suggest our control group is invalid—the groups were already diverging for reasons unrelated to the treatment.

12.3 Cluster-Robust Standard Errors

12.3.1 The Problem: Non-Independent Observations

Standard OLS assumes observations are independent. But in panel data, observations from the same unit across time are often correlated. Students in the same classroom share a teacher; workers in the same firm share management; residents of the same state share policies.

When errors are **correlated within clusters**, standard errors from regular OLS are too small. This leads to:

- T-statistics that are too large
- P-values that are too small
- Confidence intervals that are too narrow
- **Too many false positives** (Type I errors)

Figure 12.5: With clustered data, observations within groups are correlated. Regular standard errors don't account for this, leading to overconfidence.





Figure 12.6: Distribution of p-values from 500 simulations where the true effect is zero. Naive standard errors (left) produce too many small p-values; cluster-robust standard errors (right) are properly calibrated.



12.3.2 Visualizing the Problem

12.3.3 Simulation: The Dangers of Ignoring Clustering

Let's simulate what happens when we ignore clustering. We'll generate data where the **true treatment effect is zero**, so any “significant” result is a false positive.

The simulation shows that:

- With **naive standard errors**: We reject the null 61% of the time (should be 5%)
- With **cluster-robust standard errors**: We reject 6% of the time (close to 5%)

Ignoring clustering leads to a **massive inflation of false positives**!

12.3.4 Implementing Cluster-Robust Standard Errors

In `fixest`, cluster-robust standard errors are easy to implement using the `cluster` argument:

```
# Standard (naive) standard errors
model_naive <- feols(y ~ treatment | state + year,
                    vcov = "HC1",
                    data = my_data)

# Cluster-robust standard errors (clustered at state level)
model_cluster <- feols(y ~ treatment | state + year,
                      cluster = "state",
                      data = my_data)
```

Rule of Thumb: Cluster at the Treatment Level

Cluster your standard errors at the level where treatment is assigned. If policies are set at the state level, cluster by state. If a program is implemented at the school level, cluster by school.

12.4 Summary

This chapter covered three important extensions to basic DiD:

Propensity Score Weighting (IPW-DiD) improves the counterfactual by up-weighting control units that are similar to treated units. When treated and control groups differ systematically, naive DiD violates parallel trends; IPW helps restore balance. The algorithm involves estimating propensity scores, computing inverse probability weights, and using those weights in the TWFE regression.

Dynamic DiD and Event Studies allow treatment effects to vary over time. By estimating separate effects for each period relative to treatment, we can see whether effects grow, shrink, or remain stable. Crucially, the pre-treatment coefficients serve as a placebo test: if they're significantly different from zero, parallel trends is violated.

Cluster-Robust Standard Errors correct for correlated errors within groups. When observations within clusters (states, schools, firms) are correlated, standard OLS inference is overconfident, leading to inflated false positive rates. Clustering at the treatment level provides valid inference.

These tools are essential for credible DiD analysis in practice.

12.5 Check Your Understanding

Note: This section contains interactive content only available in the HTML version.

13 Instrumental Variables

💡 Key Questions

- What is endogeneity and why does it cause OLS to fail?
- What are instrumental variables and what makes an instrument valid?
- How does Two-Stage Least Squares (2SLS) work?
- What is the Local Average Treatment Effect (LATE)?
- How do we test whether an instrument is strong enough?

📖 Suggested Readings

- TBD

We've built up a powerful toolkit for causal inference: randomized experiments, multivariate regression, fixed effects, and difference-in-differences. But what happens when none of these methods are available?

Sometimes we have only cross-sectional data, treatment is clearly endogenous, we cannot randomize, and there's no natural experiment to exploit. In these situations, **instrumental variables (IV)** offers a path forward—if we can find the right instrument.

13.1 The Fundamental Problem: Endogeneity

13.1.1 When OLS Fails

Recall that OLS provides unbiased estimates when the zero conditional mean assumption holds: $E[\mu|X] = 0$. When this assumption fails—when X is correlated with the error term—we say X is **endogenous**, and OLS is biased.

Endogeneity arises from three main sources:

1. Omitted Variable Bias

We want to estimate the effect of education on wages, but ability affects both education choices and wages. Since we can't measure ability, it's absorbed into the error term, biasing our estimate.

2. Selection Bias

We want to estimate the effect of health insurance on health outcomes, but healthier people are more likely to purchase insurance. Health status drives insurance decisions, contaminating our comparison.

3. Reverse Causality

We want to estimate the effect of police on crime, but high-crime areas get more police deployed. The causation runs both ways, making it impossible to isolate the effect of policing.

13.1.2 Formalizing the Problem

Suppose we want to estimate:

$$y = \beta_0 + \beta_1 x + \mu$$

When $E[\mu|x] \neq 0$, the variable x is **endogenous**. OLS will be biased and inconsistent—no matter how large our sample, we won't converge to the true β_1 .

We cannot interpret the OLS coefficient as a causal effect.

13.2 The Instrumental Variables Solution

13.2.1 The Key Insight

The IV approach finds a variable Z that “shakes up” X in a way that's unrelated to the error term. Think of it as finding a source of **exogenous variation** in X —variation that comes from outside the system and isn't contaminated by the factors causing endogeneity.

13.2.2 Two Requirements for a Valid Instrument

For Z to be a valid instrument, it must satisfy two conditions:

! The Two IV Assumptions

1. **Relevance:** $Cov(Z, X) \neq 0$

The instrument must be correlated with the endogenous variable. This is **testable**—we can check whether Z predicts X .

2. **Exclusion Restriction:** $Cov(Z, \mu) = 0$


The instrument affects Y **only through** X . There is no direct effect of Z on Y . This is **not directly testable**—it requires economic reasoning and assumption.

13.2.3 Graphical Intuition

The IV strategy can be visualized as a causal diagram:

Figure 13.1: The instrumental variables strategy. Z affects Y only through X (exclusion restriction), while unobservables (μ) affect both X and Y , creating endogeneity.

The IV Strategy: Using Exogenous Variation



The key insight: X has two types of variation—“good” variation unrelated to μ , and “bad” variation correlated with μ . OLS uses both and is therefore biased. IV uses only the “good” variation induced by Z , giving us an unbiased estimate.

13.3 Application: Fertility and Labor Supply

13.3.1 The Research Question

What is the causal effect of having children on mothers’ labor force participation? This question is crucial for understanding gender wage gaps, designing parental leave policies, and evaluating childcare subsidies.

But we cannot randomize fertility! And the decision to have children is deeply endogenous:

- **Reverse causality:** Women may time children based on career factors
- **Selection:** Less career-oriented women may choose larger families
- **Omitted variables:** Unobserved preferences affect both fertility and work

13.3.2 The Angrist and Evans (1998) Instrument

In a landmark study, Angrist and Evans (1998) found a clever instrument for fertility: the **sex composition of the first two children**.

The instrument works because parents prefer mixed-sex children. If your first two children are the same sex (both boys or both girls), you're more likely to try for a third child to "get" the other sex.

Why is this relevant? ($Cov(Z, X) \neq 0$)

- Parents with same-sex children are more likely to have a third child
- This is an empirical fact we can verify

Why is this excludable? ($Cov(Z, \mu) = 0$)

- Sex of children is essentially random (about 50% boys, 50% girls)
- Hard to imagine why same-sex vs. mixed-sex would *directly* affect labor supply
- The only plausible channel is through fertility decisions

13.3.3 Simulating the Angrist-Evans Setup

Let's create simulated data that mirrors the Angrist-Evans setting to illustrate how IV works:

13.3.4 Testing the Relevance Condition


First, let's verify that same-sex siblings predict having more than two children:

```
# Check relevance: does samesex predict morekids?
relevance_check <- ae_sim |>
  group_by(samesex_lab) |>
  summarize(prop_morekids = mean(morekids))

relevance_check
```

```
# A tibble: 2 x 2
  samesex_lab prop_morekids
  <chr>       <dbl>
1 Mixed sex   0.196
2 Same sex    0.232
```

Figure 13.2: The relevance condition: parents with same-sex children are more likely to have a third child.




The difference is substantial—same-sex parents are about 7 percentage points more likely to have a third child. The instrument is relevant.

13.3.5 Testing Excludability (Sort of)

We cannot directly test the exclusion restriction, but we can check whether the instrument has a small *reduced-form* relationship with the outcome:

The difference is small (about 1-2 percentage points), consistent with the effect operating through fertility rather than directly. Of course, this doesn't *prove* excludability—we must rely on economic reasoning to argue there's no direct channel.

Figure 13.3: Checking excludability: the direct relationship between same-sex and labor supply is small, consistent with the effect operating through fertility.



13.4 Two-Stage Least Squares (2SLS)

13.4.1 The Estimation Strategy

IV models are typically estimated using **Two-Stage Least Squares (2SLS)**:

Stage 1 (First Stage): Regress the endogenous variable on the instrument:

$$X_i = \pi_0 + \pi_1 Z_i + \nu_i$$

Get predicted values: $\hat{X}_i = \hat{\pi}_0 + \hat{\pi}_1 Z_i$

These predicted values contain only the variation in X that comes from Z —the exogenous variation.

Stage 2 (Second Stage): Regress the outcome on the predicted values:


$$Y_i = \beta_0 + \beta_1 \hat{X}_i + \mu_i$$

The coefficient $\hat{\beta}_1^{2SLS}$ is our causal estimate.

13.4.2 Why Does This Work?

The first stage “purges” X of its problematic variation. Since Z is exogenous (by assumption), the predicted values $\hat{X}$ contain only exogenous variation. When we regress Y on $\hat{X}$, we’re using only this “clean” variation, giving us an unbiased estimate.

Figure 13.4: 2SLS uses only the variation in X that comes from the instrument Z , discarding the endogenous variation.



13.4.3 Implementing 2SLS

Let’s estimate the model step by step:

Step 1: The First Stage

```
# First stage: regress endogenous variable on instrument
first_stage <- lm(morekids ~ samesex, data = ae_sim)
summary(first_stage)
```

Call:

```
lm(formula = morekids ~ samesex, data = ae_sim)
```

Residuals:

Min	1Q	Median	3Q	Max
-0.2316	-0.2316	-0.1963	-0.1963	0.8037

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	0.196288	0.002585	75.931	<0.0000000000000002 ***
samesex	0.035260	0.003664	9.623	<0.0000000000000002 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 0.4096 on 49998 degrees of freedom

Multiple R-squared: 0.001849, Adjusted R-squared: 0.001829

F-statistic: 92.61 on 1 and 49998 DF, p-value: < 0.00000000000000022

The F-statistic on `samesex` is very large (over 100), indicating a strong instrument.

Step 2: Get Predicted Values

```
# Get predicted values from first stage
ae_sim <- ae_sim |>
  mutate(morekids_hat = predict(first_stage))

# Look at the predictions
ae_sim |>
  select(samesex, morekids, morekids_hat) |>
  head(10)
```

A tibble: 10 x 3

	samesex	morekids	morekids_hat
	<int>	<int>	<dbl>
1	1	0	0.232
2	0	0	0.196
3	0	1	0.196
4	1	0	0.232
5	1	0	0.232
6	1	0	0.232
7	1	0	0.232
8	1	0	0.232
9	0	0	0.196
10	0	0	0.196

Step 3: The Second Stage

```
# Second stage: regress outcome on predicted values
# NOTE: Standard errors from this approach are WRONG
second_stage_manual <- lm(mom_worked ~ morekids_hat, data = ae_sim)
coef(second_stage_manual)["morekids_hat"]
```

```
morekids_hat
-0.2002152
```

⚠ Standard Error Warning

The manual two-stage approach gives the correct coefficient, but the standard errors are **wrong** because they don't account for uncertainty in the first stage. Always use a proper 2SLS estimator that computes correct standard errors.

13.4.4 Proper 2SLS with `feols()`

The `fixest` package provides proper 2SLS estimation with correct standard errors:

```
# Proper 2SLS estimation
# Syntax: outcome ~ controls | fixed effects | endogenous ~ instrument
tsls <- feols(mom_worked ~ 1 | # Just intercept (no controls)
              0 | # No fixed effects
              morekids ~ samesex, # First stage
              data = ae_sim)

summary(tsls)
```


```
TSLS estimation - Dep. Var.: mom_worked
                Endo.      : morekids
                Instr.     : samesex
Second stage: Dep. Var.: mom_worked
Observations: 50,000
Standard-errors: IID

              Estimate Std. Error  t value  Pr(>|t|)
(Intercept)  0.621714   0.026008  23.90457 < 2.2e-16 ***
fit_morekids -0.200215   0.121213  -1.65176  0.09859 .
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
RMSE: 0.477825  Adj. R2: 3.111e-5
F-test (1st stage), morekids: stat = 92.6067, p < 2.2e-16 , on 1 and 49,998 DoF.
Wu-Hausman: stat = 1.1581, p = 0.281862, on 1 and 49,997 DoF.
```

13.4.5 Comparing OLS and 2SLS

```
# OLS for comparison (biased)
ols <- lm(mom_worked ~ morekids, data = ae_sim)
```

Figure 13.5: Comparing OLS (biased) and 2SLS (unbiased) estimates. OLS overstates the negative effect because career orientation confounds the relationship.



The OLS estimate is more negative than the true effect (-0.10) because it's contaminated by selection: women with lower career orientation both have more children *and* are less likely to work, making the correlation more negative than the causal effect.

The 2SLS estimate is much closer to the true effect because it uses only the exogenous variation from the same-sex instrument.

13.5 Testing for Weak Instruments

13.5.1 The Weak Instrument Problem

If the instrument is only weakly correlated with X , 2SLS performs poorly:

- Estimates become biased (toward OLS!)

- Standard errors become unreliable
- Inference is invalid

13.5.2 The F-Test Rule of Thumb

Stock and Yogo (2005) established a rule of thumb:

i Weak Instrument Test

The first-stage F-statistic should exceed **10** for the instrument to be considered strong. With $F < 10$, 2SLS estimates may be severely biased.

In our example, the first-stage F-statistic is well over 100—a very strong instrument.

13.5.3 What Happens with a Weak Instrument?

Let's see what happens when we use a weak (essentially random) instrument:

```
# Create a weak instrument (random noise)
set.seed(123)
ae_sim <- ae_sim |>
  mutate(weak_instrument = rnorm(n()))

# Try 2SLS with weak instrument
weak_iv <- feols(mom_worked ~ 1 | 0 | morekids ~ weak_instrument,
  data = ae_sim)
```


With a weak instrument, the estimate is wildly off and the confidence interval is enormous. Always check your first-stage F-statistic!

13.6 What Does IV Estimate? The LATE

13.6.1 The Local Average Treatment Effect

An important subtlety: IV does not estimate the same quantity as OLS or DiD. It estimates the **Local Average Treatment Effect (LATE)**—the causal effect for a specific subpopulation called **compliers**.

Figure 13.6: With a weak instrument (random noise), the 2SLS estimate becomes unreliable with huge standard errors.



13.6.2 Four Types of People

In the fertility example, we can categorize people by how their fertility responds to the instrument:

Type	Description
Always-takers	Would have 3+ kids regardless of sex composition
Never-takers	Would never have 3+ kids regardless of sex composition
Compliers	Have 3+ kids <i>because</i> first two were same-sex
Defiers	Have 3+ kids <i>only if</i> first two were different-sex

IV identifies the causal effect only for compliers—those whose treatment status was actually changed by the instrument.

13.6.3 Who Are the Compliers?

In the Angrist-Evans context, compliers are parents who:

- Wanted mixed-sex children strongly enough to try for a third

- Would have stopped at two kids if the first two were different sexes

This is **not**:

- All parents with 3+ children
- Parents who wanted large families regardless
- Parents who strongly preferred small families

13.6.4 Trade-offs of IV Estimation

Advantages	Disadvantages
Credible causal identification	Estimates LATE, not ATE
Handles endogeneity	Less generalizable
No need for RCT or natural experiment	Requires strong assumptions
Works with cross-sectional data	Often larger standard errors

13.7 Classic IV Applications

Instrumental variables have been used to study many important questions:

Study	Question	Instrument
Card (1995)	Returns to education	Distance to college
Acemoglu et al. (2001)	Do institutions cause growth?	Colonial settler mortality
Levitt (1997)	Effect of police on crime	Electoral timing
Angrist & Krueger (1991)	Returns to education	Quarter of birth

13.7.1 Example: Card (1995) - Returns to Education

Question: What is the causal effect of education on wages?

Problem: Ability is unobserved but affects both education and wages, biasing OLS upward.

Instrument: Distance to the nearest college. People who grew up closer to a college got more education.

Why relevant? Proximity reduces the cost of attending college.

Why excludable? Distance to a college shouldn't directly affect wages decades later (only through education).

Finding: Returns to education are actually *higher* than OLS suggests—the ability bias works in the opposite direction from what many expected.

13.8 Summary

Instrumental variables provide a powerful tool for causal inference when other methods fail. The key is finding a variable that affects treatment but has no direct effect on the outcome.

The two requirements:

1. **Relevance:** The instrument predicts the endogenous variable (testable via first-stage F-statistic > 10)
2. **Exclusion:** The instrument affects the outcome only through the treatment (requires economic reasoning—not directly testable)

Key insights:

- 2SLS uses only the exogenous variation in X induced by Z
- IV estimates the Local Average Treatment Effect (LATE) for compliers
- Weak instruments produce unreliable estimates—always check the first-stage F-statistic
- The exclusion restriction is crucial but untestable—think carefully about whether it holds

IV requires strong assumptions, but when those assumptions are credible, it provides a path to causal inference even in challenging observational settings.

13.9 Check Your Understanding

Note: This section contains interactive content only available in the HTML version.

References

- Angrist, Joshua D., and Jörn-Steffen Pischke. 2014. *Mastering 'Metrics: The Path from Cause to Effect*. Princeton University Press.
- Bailey, Michael A. 2020. *Real Econometrics: The Right Tools to Answer Important Questions*. Oxford University Press.
- Wickham, Hadley, Mine Çetinkaya-Rundel, and Garrett Golemund. 2023. *R for Data Science: Import, Tidy, Transform, Visualize, and Model Data*. O'Reilly Media, Incorporated.
- Wooldridge, Jeffrey M. 2019. *Introductory Econometrics: A Modern Approach*. 7th ed. Cengage Learning.

A Intro to R and RStudio

💡 Key Questions

- How do I install R and RStudio on my computer?
- What are the main components of the RStudio interface?
- How do I write and run R code?
- What are objects and how do I use them?
- How do I install and load R packages?

📖 Suggested Readings

- Wickham, Çetinkaya-Rundel, and Grolemund (2023), Ch. 1-3

This appendix will walk you through installing R and RStudio, understanding the RStudio interface, and writing your first lines of R code. By the end of this chapter, you will be ready to start working with data and running econometric analyses.

A.1 Installing R

To get started with R, you first need to download and install the base R software. R is available for free from CRAN (the Comprehensive R Archive Network). Navigate to <https://cloud.r-project.org/> and click the download link for your operating system (Windows, macOS, or Linux). Follow the installation instructions provided for your system.

Once the installation is complete, you will have R installed on your computer. However, we will not typically interact with R directly. Instead, we will use RStudio, an integrated development environment (IDE) that makes working with R much more convenient.

A.2 Installing RStudio

RStudio is a powerful IDE designed specifically for working with R. It provides a user-friendly interface with helpful features like syntax highlighting, code completion, and integrated help documentation. You can download RStudio Desktop for free from <https://posit.co/download/>

[rstudio-desktop/](#). The website should automatically detect your operating system and show you the appropriate download link. Click the link and follow the installation instructions.

! Only Use RStudio

After installing both R and RStudio, you will only ever need to open RStudio. RStudio runs R in the background, so you don't need to open the base R application separately. Feel free to delete the shortcut for base R from your desktop if you wish. When you open an R script file (`.R`) for the first time, be sure to set RStudio as the default program to open such files.

A.3 The RStudio Interface

When you first open RStudio, you will see an interface divided into several panels (or “panes”), as seen in Figure A.1. Each panel serves a different purpose in your workflow. Understanding these panels is essential for working efficiently in R.

A.3.1 The Script Editor

The Script Editor is where you write your R code in script files. A script file is simply a text file containing R code that can be saved, edited, and run again later. Working in script files is essential for reproducibility—you can always go back and see exactly what you did, make changes, and re-run your analysis. To open a new script file, go to File → New File → R Script, or use the keyboard shortcut `Cmd+Shift+N` (Mac) or `Ctrl+Shift+N` (Windows/Linux).


A.3.2 The Console

The Console is where R actually executes your code. When you run code from a script, it gets sent to the console line by line. You can also type code directly into the console and press Enter to run it immediately. However, code typed directly into the console is not saved, so this approach is generally only recommended for quick tests or exploratory work. Output from many functions, including regression results, will be printed to the console.

A.3.3 The Environment Panel

The Environment panel displays all the “objects” you have created in your current R session. These objects might include data frames, vectors, or stored values. You can see each object's name along with a brief summary of its contents. This panel is useful for keeping track of what data and variables you currently have available.

Figure A.1: The RStudio interface with its four main panels labeled.



A.3.4 The Plots and Output Panel

This panel serves multiple purposes. When you create visualizations (like scatter plots or histograms), they will appear here. This panel also contains tabs for browsing files on your computer, viewing installed packages, and accessing R’s built-in help documentation.

Customizing Your Layout

You can hide or show different panels under the “View” menu. This is useful when you want more screen space for a particular task, such as focusing on your script while writing code.

A.4 Basic Coding in R

Now that RStudio is set up, let’s write some code! We’ll start with simple operations in the console to get familiar with R’s syntax.

A.4.1 Arithmetic Operations

R can function as a sophisticated calculator. Try typing the following expressions in the console and pressing Enter:

```
2 + 2
```

```
[1] 4
```

```
20 * 5 / 87
```

```
[1] 1.149425
```

```
sin(pi / 2)
```

```
[1] 1
```

```
sqrt(6)
```

```
[1] 2.44949
```

As you can see, R can handle basic arithmetic as well as mathematical functions like `sin()` and `sqrt()`. The `pi` you see in the third example is a built-in constant in R representing the mathematical constant π .

A.4.2 Objects and Assignment

R is an “object-oriented language,” meaning that data, variables, and nearly everything else is stored as an “object.” You create objects using the assignment operator `<-`. The assignment operator takes whatever is on the right side and stores it in the name on the left side.

```
x <- 2 + 5
```

After running this code, the value 7 is now stored in an object called `x`. Notice that R didn’t print anything—it just stored the value. To see what’s stored in an object, type its name and press Enter:

```
x
```

```
[1] 7
```

You can use objects in subsequent calculations, just like variables in algebra:

```
x + 10
```

```
[1] 17
```

You can also update an object by reassigning it:

```
x <- x + 50  
x
```

```
[1] 57
```

Naming Objects

When creating object names, keep them short but descriptive. Object names cannot contain spaces. Some good examples include: `flight_data`, `reg_controls_1`, `acs_male`, or `X_mat`.

A.4.3 Vectors

A vector is the simplest type of object in R. Vectors contain a sequence of values of the same type (all numbers, all text, etc.). You create vectors using the `c()` function, which stands for “combine”:

```
primes <- c(2, 3, 5, 7, 11, 13)
primes
```

```
[1]  2  3  5  7 11 13
```

One powerful feature of R is that arithmetic operations on vectors are applied element-by-element:

```
primes * 2
```

```
[1]  4  6 10 14 22 26
```

```
primes - 1
```

```
[1]  1  2  4  6 10 12
```

You can also perform arithmetic between two vectors of the same length:

```
odds <- c(1, 3, 5, 7, 9, 11)
primes + odds
```

```
[1]  3  6 10 14 20 24
```

A.4.4 Functions

R has a large collection of built-in functions that perform specific tasks. Functions are called using the syntax: `function_name(argument1 = value1, argument2 = value2, ...)`.

For example, the `seq()` function generates a sequence of numbers:

```
bb <- seq(from = 1, to = 30, by = 3)
bb
```



```
[1] 1 4 7 10 13 16 19 22 25 28
```

Here, **from**, **to**, and **by** are the *arguments* of the function. We set **from** = 1 to start at 1, **to** = 30 to end at 30, and **by** = 3 to increment by 3.

Getting Help

To learn more about any function, type **?** followed by the function name in the console. For example, **?seq** will open the documentation for the **seq()** function.

A.5 Data Types in R

Every piece of data in R has a **type** that determines how it can be used. Understanding data types is essential because certain operations only work with certain types, and R will sometimes convert between types automatically in ways that can cause unexpected results.

A.5.1 Numeric Types

R has two main numeric types:

Integer (**int**): Whole numbers without decimal places. Created by adding **L** after a number:

```
my_integer <- 5L
typeof(my_integer)
```

```
[1] "integer"
```

Double (**dbl**): Numbers with decimal places (also called “floating point”). This is the default for most numbers:

```
my_double <- 5.7
typeof(my_double)
```

```
[1] "double"
```

In practice, the distinction between integers and doubles rarely matters for econometric analysis—R handles conversions automatically. When you see **<int>** or **<dbl>** in your data, both are simply “numbers.”

A.5.2 Character Strings

Character (chr): Text data, always enclosed in quotes:

```
my_name <- "Economics"  
typeof(my_name)
```

```
[1] "character"
```

Character strings are used for labels, names, and categorical data that hasn't been converted to a factor.

A.5.3 Logical Values

Logical (lgl): TRUE or FALSE values (can be abbreviated as T and F):

```
is_enrolled <- TRUE  
typeof(is_enrolled)
```

```
[1] "logical"
```

Logical values are often created by comparisons:

```
5 > 3
```

```
[1] TRUE
```

```
10 == 10 # Note: == tests equality, = is for assignment
```

```
[1] TRUE
```

Logical values are essential for filtering data (e.g., “keep only observations where income > 50000”).

A.5.4 Factors

Factor (fct): Categorical variables with a fixed set of possible values (called “levels”). Factors are crucial in regression because R treats them differently from numeric variables:

```
# Create a factor for education level
edu_level <- factor(c("High School", "College", "Graduate", "College", "High School"))
edu_level
```

```
[1] High School College      Graduate    College    High School
Levels: College Graduate High School
```

Notice how R shows the “Levels”—the complete set of possible categories. Factors are especially important for dummy variables in regression.

! Why Factors Matter

When you include a factor in a regression, R automatically creates dummy variables for each level. If you include a character variable, R may treat it as text rather than a categorical variable, leading to errors or unexpected results. We will talk more about this in section [Chapter 7](#).

A.5.5 Checking and Converting Types

You can check an object’s type with `typeof()` or `class()`:

```
class(edu_level)
```

```
[1] "factor"
```

Convert between types using `as.numeric()`, `as.character()`, `as.factor()`, etc.:

```
# Convert character to factor
major <- c("Economics", "Math", "Economics", "Physics")
major_factor <- as.factor(major)
major_factor
```

```
[1] Economics Math      Economics Physics
Levels: Economics Math Physics
```

```
# Convert numeric to character
numbers <- c(1, 2, 3)
as.character(numbers)
```

```
[1] "1" "2" "3"
```

A.5.6 Common Type Abbreviations

When working with data frames, you'll frequently see these abbreviations:

Table A.1: Common data type abbreviations in R

Abbreviation	Type	Description	Example
<dbl>	Double	Numbers with decimals	3.14, 2.5
<int>	Integer	Whole numbers	1, 42
<chr>	Character	Text strings	"hello", "NY"
<fct>	Factor	Categorical variable	factor("low", "med", "high")
<lgl>	Logical	TRUE/FALSE	TRUE, FALSE
<date>	Date	Calendar dates	as.Date("2024-01-15")

A.5.7 Data Frames

Data frames are the primary way that structured, tabular data is stored in R. A data frame is essentially a table where each column is a vector and each row is an observation. R comes with several built-in data frames for practice, including `mtcars`, which contains data on various car models.

Use `head()` to view the first few rows of a data frame:

```
head(mtcars)
```

	mpg	cyl	disp	hp	drat	wt	qsec	vs	am	gear	carb
Mazda RX4	21.0	6	160	110	3.90	2.620	16.46	0	1	4	4
Mazda RX4 Wag	21.0	6	160	110	3.90	2.875	17.02	0	1	4	4
Datsun 710	22.8	4	108	93	3.85	2.320	18.61	1	1	4	1
Hornet 4 Drive	21.4	6	258	110	3.08	3.215	19.44	1	0	3	1
Hornet Sportabout	18.7	8	360	175	3.15	3.440	17.02	0	0	3	2
Valiant	18.1	6	225	105	2.76	3.460	20.22	1	0	3	1

You can explore the structure of a data frame using several helpful functions:

```
nrow(mtcars)
```

```
[1] 32
```

```
ncol(mtcars)
```

```
[1] 11
```

```
colnames(mtcars)
```

```
[1] "mpg"  "cyl"  "disp" "hp"   "drat" "wt"   "qsec" "vs"   "am"   "gear"  
[11] "carb"
```

To access a specific column (variable) within a data frame, use the `$` operator:

```
head(mtcars$mpg)
```

```
[1] 21.0 21.0 22.8 21.4 18.7 18.1
```

Here, `mtcars$mpg` extracts the `mpg` column from the `mtcars` data frame as a vector. I wrapped it in `head()` to show only the first few values.

A.5.8 Comments

In R, anything preceded by the `#` symbol is treated as a comment and will not be executed. Comments are essential for documenting your code—explaining what each section does, leaving notes for yourself, or marking areas that need revision.

```
# Find the mean miles per gallon  
mean(mtcars$mpg)
```

```
[1] 20.09062
```

Good commenting habits make your code readable to others (and to your future self).

A.6 Working with R Scripts

While the console is useful for quick tests, you should do the vast majority of your work in script files. Scripts allow you to save your code, edit it easily, and reproduce your entire analysis by running the script from top to bottom.

A.6.1 Creating and Running Scripts

To create a new script, go to File → New File → R Script (or use Cmd/Ctrl + Shift + N). Write your code in the script editor, then use the following keyboard shortcuts to run it:

Cmd/Ctrl + Enter executes the current line (or highlighted selection) in the console. The cursor then automatically moves to the next line, making it easy to step through your code line by line.

Figure A.2: Example R script with cursor position shown. Pressing Cmd/Ctrl + Enter will run the complete command that creates the `not_cancelled` object, then move the cursor to the next statement.

```
library(dplyr)
library(nycflights13)

not_cancelled <- flights |>
  filter(!is.na(dep_delay), !is.na(arr_delay))

not_cancelled |>
  group_by(year, month, day) |>
  summarize(mean = mean(dep_delay))
```

Cmd/Ctrl + Shift + S runs the entire script from top to bottom. This is useful for checking that your complete analysis works correctly and produces reproducible results.

A.6.2 Script Best Practices

Following good practices when writing scripts will save you time and frustration. Always start your script with your name and a brief description of the script's purpose in comments. Load all required packages at the beginning of the script using `library()` so anyone reading your code knows what dependencies are needed. Never include `install.packages()` in your script—you only need to install packages once on your computer, and you don't want your script to install packages on someone else's machine without their permission. Save your scripts with short, descriptive names that contain no spaces, such as `assignment_3.R` or `ec_325_data_plotting.R`.

A.7 Working Directories

R uses a concept called the “working directory”—the folder on your computer where R looks for files you want to load and where it saves files you create. You can see your current working directory at the top of the console, or by running:

```
getwd()
```

A good workflow is to save your R script in a dedicated folder for this course (or a subfolder for each week’s work), put all related data files in the same folder, and then set your working directory to that location. In RStudio, go to Session → Set Working Directory → To Source File Location to automatically set the working directory to wherever your script is saved.

A.8 R Packages

R comes with a set of “base” packages that provide fundamental functionality. However, the true power of R lies in its vast ecosystem of user-contributed packages. These packages extend R’s capabilities for specialized tasks—from data visualization to complex statistical modeling.

A.8.1 Installing Packages

You only need to install a package once on your computer. Use the `install.packages()` function:

```
install.packages("tidyverse")
```

A.8.2 The Tidyverse

The tidyverse is a collection of R packages designed for data science. These packages share a common philosophy and work seamlessly together, making data manipulation and visualization more intuitive. Some of the most commonly used tidyverse packages include `dplyr` (data manipulation), `ggplot2` (visualization), and `readr` (reading data files).

Installing `tidyverse` installs all of these packages at once.

A.8.3 Loading Packages

Once a package is installed, you need to load it each time you start a new R session using `library()`:

```
library(tidyverse)
```

When you load the `tidyverse`, you'll see a message listing which packages were attached. You may also see messages about “conflicts”—this just means some `tidyverse` functions have the same name as functions in base R, and the `tidyverse` versions will take precedence. This is normal and expected behavior.

A.8.4 The Pipe Operator

One of the most useful features of the `tidyverse` is the **pipe operator**: `|>`. The pipe takes the result of whatever is on its left and passes it as the first argument to the function on its right.

Think of the pipe like a conjunction in a sentence. Just as “and” or “then” connect actions in English, the pipe connects operations in R. Instead of saying “Take the data, then filter it, then summarize it,” you write code that reads almost the same way:

```
data |>
  filter(year == 2020) |>
  summarize(mean_income = mean(income))
```

This code reads naturally from top to bottom: “Start with `data`, *then* filter to rows where year equals 2020, *then* summarize by calculating the mean income.”

Without the pipe, you would need to either nest functions inside each other (hard to read) or create intermediate objects at each step (clutters your environment). The pipe lets you write code that mirrors how you think about the analysis.

i Base R Pipe vs. Magrittr Pipe

R now has a built-in pipe (`|>`) that works without loading any packages. The `tidyverse` historically used `%>%` from the `magrittr` package, which you may see other people use. Both work similarly for most purposes. We'll use the base R pipe `|>` in this course.

! Remember the Difference

`install.packages()` downloads and installs a package to your computer (do this once).
`library()` loads an already-installed package into your current R session (do this every time you start R).

A.9 Summary and Conclusion

This appendix walked you through the essential first steps for working with R. You installed R and RStudio, learned how the RStudio interface is organized, and wrote your first R code. You now understand how to create and manipulate objects, work with vectors and data frames, write and run scripts, and install packages.

A.9.1 Key Takeaways

- **RStudio is your interface to R:** Always work in RStudio, not base R. The script editor, console, environment, and plots panels each serve distinct purposes in your workflow.
- **Scripts are essential for reproducibility:** Write your code in script files that you can save, edit, and re-run. Use comments to document what your code does.
- **Objects store everything in R:** Use the assignment operator `<-` to create objects. Vectors hold sequences of values; data frames hold structured tabular data.
- **Packages extend R's capabilities:** Install packages once with `install.packages()`, then load them each session with `library()`. The tidyverse is a particularly useful collection of packages for data science.
- **Working directories matter:** Set your working directory to the folder containing your script and data files to keep your analysis organized.

With these foundations in place, you are ready to start loading real data, creating visualizations, and running econometric analyses.

A.10 Check Your Understanding

Note: This section contains interactive content only available in the HTML version.

B Statistics and Probability Review

Key Questions

- What are the properties of the summation operator?
- What are the key rules for logarithms, and why do economists use logs so often?
- How do we interpret derivatives and partial derivatives?
- How do we find the maximum or minimum of a function?
- What is the difference between a population and a sample?
- What is a random variable and how do we describe its distribution?
- How do we measure central tendency, variability, and relationships between variables?
- What is the normal distribution and why is it important?

Suggested Readings

- Wooldridge (2019), Appendix A, B, and C

This appendix reviews the essential mathematical and statistical concepts that form the foundation for econometrics. If you've taken introductory statistics or calculus, much of this material will be familiar. However, econometrics uses these tools in specific ways, so it's worth reviewing them with an eye toward how they'll appear throughout the course.

B.1 Basic Mathematical Tools

B.1.1 The Summation Operator

The summation operator provides a compact way to write the sum of a sequence of numbers. If $\{x_i : i = 1, 2, \dots, n\}$ is a sequence of n numbers, then the sum of these numbers is written as:

$$x_1 + x_2 + \dots + x_n = \sum_{i=1}^n x_i$$

The symbol Σ (capital Greek letter sigma) tells us to add up the terms. The expression $i = 1$ below the sigma indicates that we start counting at $i = 1$, and the n above tells us to stop at $i = n$.

B.1.1.1 Properties of Summation

The summation operator has several useful properties that we will use frequently.

Property 1: Summing a constant. If we add a constant c to itself n times, we get n times the constant:

$$\sum_{i=1}^n c = nc$$

Property 2: Factoring out constants. We can pull constants outside the summation:

$$\sum_{i=1}^n cx_i = c \sum_{i=1}^n x_i$$

Property 3: Separating additive terms. Sums can be split across addition:

$$\sum_{i=1}^n (ax_i + by_i) = a \sum_{i=1}^n x_i + b \sum_{i=1}^n y_i$$

Common Mistakes with Summation

These properties do **not** extend to division or multiplication in the ways you might expect. We **cannot** break up divided expressions:

$$\sum_{i=1}^n \frac{x_i}{y_i} \neq \frac{\sum_{i=1}^n x_i}{\sum_{i=1}^n y_i}$$

We **cannot** break up multiplicative expressions:

$$\sum_{i=1}^n x_i^2 \neq \left(\sum_{i=1}^n x_i \right)^2$$

B.1.1.2 The Sample Mean

Given n numbers $\{x_i : i = 1, 2, \dots, n\}$, we can compute their **average** or **mean** value by adding them up and dividing by n . The average value, denoted $\bar{x}$ (read “x-bar”), is written as:

$$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$$

The average is an example of a **descriptive statistic**: a value that describes or summarizes a characteristic of a set of data points. In this case, the mean describes the **central tendency** of x_i —where the “middle” of the data lies.

B.1.1.3 Useful Summation Results

Two results involving summations will appear repeatedly in this course.

Result 1: Deviations from the mean sum to zero. The sum of the deviations $(x_i - \bar{x})$ from the mean always equals zero:

$$\sum_{i=1}^n (x_i - \bar{x}) = 0$$

This can be proven using the properties above:

$$\sum_{i=1}^n (x_i - \bar{x}) = \sum_{i=1}^n x_i - \sum_{i=1}^n \bar{x} = \sum_{i=1}^n x_i - n\bar{x} = n\bar{x} - n\bar{x} = 0$$

The last step follows because $\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$ implies $n\bar{x} = \sum_{i=1}^n x_i$.

Result 2: Sum of squared deviations. The sum of squared deviations can be rewritten as:

$$\sum_{i=1}^n (x_i - \bar{x})^2 = \sum_{i=1}^n x_i^2 - n\bar{x}^2$$

More generally, for two variables x and y :

$$\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y}) = \sum_{i=1}^n x_i y_i - n\bar{x}\bar{y}$$

These identities are useful for computing variances and covariances, which we will discuss shortly.

B.1.2 Derivatives

The derivative $\frac{dy}{dx}$ of a function $y = f(x)$ measures its **instantaneous rate of change**. It tells you how sensitive the function's output is to a small change in its input. In econometrics, derivatives help us interpret how changes in independent variables affect dependent variables.

Here are derivatives of some common functional forms in econometrics:

Function	Derivative
$y = \beta_0 + \beta_1 x$	$\frac{dy}{dx} = \beta_1$
$y = \beta_0 + \beta_1 x + \beta_2 x^2$	$\frac{dy}{dx} = \beta_1 + 2\beta_2 x$
$y = \beta_0 + \frac{\beta_1}{x}$	$\frac{dy}{dx} = -\frac{\beta_1}{x^2}$
$y = \beta_0 + \beta_1 \ln(x)$	$\frac{dy}{dx} = \frac{\beta_1}{x}$

Notice that in the linear case, the derivative is simply the coefficient β_1 —the effect of x on y is constant. In the other cases, the derivative depends on the value of x , meaning the effect of x on y varies depending on where you start.

B.1.3 Percentages and Percent Changes

Understanding percentages is essential for econometrics. Many economic variables are reported as percentages (unemployment rate, interest rates, inflation), and we frequently describe changes in percentage terms. Getting the language right matters—confusing “percent” with “percentage points” is a common source of errors.

B.1.3.1 Percentages

A **percentage** expresses a number as a fraction of 100. To convert a decimal to a percentage, multiply by 100:

$$0.25 = 25\%$$

To convert a percentage to a decimal, divide by 100:

$$25\% = 0.25$$

In econometrics, we typically work with decimals in our calculations and convert to percentages for interpretation.

B.1.3.2 Percent Change

The **percent change** (or **percentage change**) measures how much a variable changed relative to its initial value:

$$\text{Percent Change} = \frac{\text{New Value} - \text{Old Value}}{\text{Old Value}} \times 100\%$$

Or more compactly:

$$\% \Delta Y = \frac{Y_2 - Y_1}{Y_1} \times 100\%$$

Example: If wages increase from \$20 to \$22 per hour:

$$\% \Delta \text{wage} = \frac{22 - 20}{20} \times 100\% = \frac{2}{20} \times 100\% = 10\%$$

Wages increased by **10 percent**.

Percent Change vs. Percentage Point Change

These are **not** the same thing! This distinction trips up many students.

- **Percent change:** The change relative to the starting value
- **Percentage point change:** The arithmetic difference between two percentages

Example: If the unemployment rate rises from 4% to 5%:

- The **percentage point change** is: $5\% - 4\% = 1$ percentage point
- The **percent change** is: $\frac{5-4}{4} \times 100\% = 25\%$

Both statements are correct:

- “Unemployment rose by 1 percentage point”
- “Unemployment rose by 25 percent”

They mean very different things! In news and policy discussions, this distinction matters enormously.

B.1.3.3 When to Use Which

Use **percentage points** when:

- Describing changes in variables already measured as percentages (interest rates, unemployment rates, test scores out of 100)
- You want to describe the absolute change in the percentage

Use **percent change** when:

- Describing changes in levels (wages, prices, GDP, population)
- You want to describe the relative or proportional change

Example: An interest rate cut from 5% to 4% is:

- A 1 percentage point decrease
- A 20% decrease (both are correct, but mean different things)
- ~~A 1% decrease~~ (this is ambiguous and often wrong)

B.1.3.4 Calculating Values After Percent Changes

If Y increases by $p\%$, the new value is:

$$Y_{new} = Y_{old} \times (1 + p/100)$$

If Y decreases by $p\%$, the new value is:

$$Y_{new} = Y_{old} \times (1 - p/100)$$

Example: If a \$50 item increases in price by 20%:

$$Price_{new} = 50 \times (1 + 0.20) = 50 \times 1.20 = \$60$$

B.1.3.5 Compounding Percent Changes

Percent changes don't add—they compound. If wages increase by 10% and then by another 10%, the total increase is **not** 20%.

$$\text{Final} = \text{Initial} \times 1.10 \times 1.10 = \text{Initial} \times 1.21$$

The total increase is **21%**, not 20%.

This is why we use logarithms: log changes **do** add up, making them convenient for analyzing growth over time.

B.1.4 Logarithms

Logarithms appear constantly in econometrics. We use them to model percentage changes, interpret coefficients as elasticities, and transform skewed variables like income and wages. Understanding log rules is essential for working with regression models.

B.1.4.1 What is a Logarithm?

The natural logarithm, denoted $\ln(x)$, is the inverse of the exponential function e^x . If $y = \ln(x)$, then $e^y = x$.

Some key values to remember:

- $\ln(1) = 0$ because $e^0 = 1$
- $\ln(e) = 1$ because $e^1 = e$
- $\ln(x)$ is only defined for $x > 0$

B.1.4.2 Essential Log Rules

Three rules form the foundation for working with logarithms:

Rule 1: Product Rule. The log of a product is the sum of the logs:

$$\ln(ab) = \ln(a) + \ln(b)$$

Rule 2: Quotient Rule. The log of a ratio is the difference of the logs:

$$\ln\left(\frac{a}{b}\right) = \ln(a) - \ln(b)$$

Rule 3: Power Rule. The log of a power brings the exponent down:

$$\ln(a^k) = k \ln(a)$$

These rules are used constantly when manipulating regression equations and interpreting coefficients.

B.1.4.3 Logs and Percentage Changes

Here is the most important property of logs for econometrics:

! The Key Insight

For small changes, the **difference in logs approximately equals the percentage change**:

$$\ln(Y_2) - \ln(Y_1) \approx \frac{Y_2 - Y_1}{Y_1}$$

In other words: $\Delta \ln(Y) \approx \% \text{ change in } Y$

This approximation works well for changes up to about 20%. It follows from the quotient rule:

$$\ln(Y_2) - \ln(Y_1) = \ln\left(\frac{Y_2}{Y_1}\right)$$

When Y_2 is close to Y_1 , the ratio Y_2/Y_1 is close to 1, and $\ln(1+x) \approx x$ for small x .


Example: If $\ln(\text{wage})$ increases from 2.50 to 2.58, the change is 0.08. This means wages increased by approximately **8%**.

B.1.4.4 Other Uses of Logs

Logarithms are useful in economics for several reasons:

1. **Percentage interpretation:** Coefficients in log models have natural percentage interpretations (more on this in Chapter 7).
2. **Elasticities:** In a log-log model like $\ln(Y) = \beta_0 + \beta_1 \ln(X)$, the coefficient β_1 is the **elasticity**—the percentage change in Y for a 1% change in X .
3. **Reducing skewness:** Variables like income, wages, and prices are often right-skewed. Taking logs makes them more symmetric and closer to normal.
4. **Multiplicative relationships:** Logs transform multiplicative relationships into additive ones, which are easier to estimate with linear regression.

Figure B.1: Left: The distribution of wages is right-skewed. Right: The distribution of log wages is more symmetric.



B.1.5 Partial Derivatives

When y is a function of multiple variables, say $y = f(x_1, x_2)$, we use **partial derivatives** to measure how y changes with respect to one variable while holding the others constant.

The partial derivative of y with respect to x_1 , holding x_2 constant, is denoted $\frac{\partial y}{\partial x_1}$. Similarly, the partial derivative with respect to x_2 , holding x_1 constant, is $\frac{\partial y}{\partial x_2}$.

For example, if $y = \beta_0 + \beta_1 x_1 + \beta_2 x_2$, then:

$$\frac{\partial y}{\partial x_1} = \beta_1, \quad \frac{\partial y}{\partial x_2} = \beta_2$$

The concept of “holding other variables constant” is central to econometrics. When we estimate a regression coefficient, we are trying to measure the partial effect of one variable on the outcome, controlling for other factors.

B.1.6 Optimization: Maxima and Minima

In econometrics, we often need to find the value of a variable that maximizes or minimizes a function. For example, Ordinary Least Squares (OLS) regression finds coefficient estimates by minimizing the sum of squared residuals. Understanding optimization is essential for understanding how these estimates are derived.

B.1.6.1 Finding Critical Points

To find the maximum or minimum of a function $f(x)$, we take the derivative, set it equal to zero, and solve for x . The solutions are called **critical points**.

For example, consider the quadratic function:

$$f(x) = ax^2 + bx + c$$

Taking the derivative and setting it equal to zero:

$$\frac{df}{dx} = 2ax + b = 0$$

Solving for x :

$$x^* = -\frac{b}{2a}$$

This critical point x^* is where the function reaches its maximum or minimum.

B.1.6.2 Determining Maximum vs. Minimum

How do we know if a critical point is a maximum or a minimum? We use the **second derivative test**. The second derivative tells us about the curvature of the function:

$$\frac{d^2f}{dx^2} = 2a$$


The rule is:

- If $\frac{d^2f}{dx^2} > 0$ at the critical point, the function is **concave up** (bowl-shaped), and we have a **minimum**
- If $\frac{d^2f}{dx^2} < 0$ at the critical point, the function is **concave down** (hill-shaped), and we have a **maximum**

For our quadratic function $f(x) = ax^2 + bx + c$:

- If $a > 0$: The parabola opens upward, and $x^* = -\frac{b}{2a}$ is a **minimum**
- If $a < 0$: The parabola opens downward, and $x^* = -\frac{b}{2a}$ is a **maximum**

Figure B.2: Left: When $a > 0$, the parabola opens upward and the critical point is a minimum. Right: When $a < 0$, the parabola opens downward and the critical point is a maximum.



💡 Optimization in OLS

In OLS regression, we minimize the sum of squared residuals, which is a quadratic function of the coefficients. The second derivative is positive (the function is convex), confirming that the solution is indeed a minimum. This is why setting the first derivative equal to zero gives us the best-fitting line.

B.1.6.3 Optimization with Multiple Variables

When a function depends on multiple variables, we find critical points by setting all partial derivatives equal to zero simultaneously. For a function $f(x_1, x_2)$:

$$\frac{\partial f}{\partial x_1} = 0 \quad \text{and} \quad \frac{\partial f}{\partial x_2} = 0$$

Determining whether a critical point is a maximum, minimum, or neither (a saddle point) requires examining the second partial derivatives, which we will encounter when deriving the OLS estimator.

B.2 Population vs. Sample

Before diving into probability, it's important to understand the distinction between a **population** and a **sample**, as this distinction underlies all of statistical inference.

The **population** is the entire group of individuals, objects, or observations that we are interested in studying. For example, if we want to understand the wages of all U.S. workers, the population is every person employed in the United States. If we want to know the effect of a drug on blood pressure, the population might be all adults with hypertension.

In practice, we almost never observe the entire population. Instead, we collect data on a **sample**—a subset of the population. We might survey 5,000 workers rather than all 160 million, or run a clinical trial with 500 patients rather than everyone with hypertension.

! The Goal of Statistical Inference

The fundamental goal of statistics is to use information from a **sample** to draw conclusions about the **population**. We observe sample statistics (like the sample mean $\bar{x}$) and use them to make inferences about population parameters (like the population mean μ).

This distinction affects our notation:

Concept	Population Parameter	Sample Statistic
Mean	μ	$\bar{x}$
Variance	σ^2	s^2
Standard deviation	σ	s
Correlation	ρ	r

Population parameters are typically denoted with Greek letters and are usually unknown. Sample statistics are computed from our data and are used to estimate the population parameters.

For example, if we want to know the average wage of all U.S. workers (μ), we collect a sample and compute the sample mean ($\bar{x}$). Under certain conditions, $\bar{x}$ is a good estimate of μ . Much of econometrics is about understanding when and how well sample statistics estimate population parameters, and quantifying the uncertainty in those estimates.

B.3 Probability Basics

B.3.1 Random Variables

A **random variable** is a variable whose value is determined by a random phenomenon. It represents a number, but the exact value is determined by chance. Examples include the outcome of a dice roll, whether a coin lands heads or tails, or the percentage of flights that arrive on time.

Most variables we work with in econometrics can be thought of as random variables. A person's wage, years of education, or health status all have elements of randomness—we cannot perfectly predict them in advance.

B.3.2 Probability

Probability quantifies uncertainty. If X is a random variable that takes on k possible values $\{x_1, x_2, \dots, x_k\}$, then the probability that X takes a particular value x_j is denoted:

$$p_j = P(X = x_j)$$

Each probability p_j must be between 0 and 1, and the probabilities must sum to 1: $p_1 + p_2 + \dots + p_k = 1$.

B.3.3 Probability Distributions

The **probability density function (pdf)**, denoted $f(x)$, gives the probability that a random variable takes on a particular value x . For discrete random variables, $f(x) = P(X = x)$.

The **cumulative distribution function (cdf)**, denoted $F(x)$, gives the probability that the random variable is less than or equal to x :

$$F(x) = P(X \leq x)$$

The cdf is useful for answering questions like “what is the probability that X is below some threshold?”

B.3.4 Expected Values

The **expected value** (or **mathematical expectation**) of a random variable X , denoted $E[X]$, represents the “average” outcome we would expect if we could repeat the random process many times. It is our population “best guess” for the value of X . We often denote it with μ or μ_X .

For a discrete random variable with pdf $f(x)$, the expected value is the weighted average of all possible values, where the weights are the probabilities:

$$E[X] = \sum_{j=1}^k x_j f(x_j) = \mu$$

Example: Let X be the outcome of rolling a fair six-sided die. Each value (1 through 6) has probability $1/6$. The expected value is:

$$E[X] = \frac{1}{6}(1) + \frac{1}{6}(2) + \frac{1}{6}(3) + \frac{1}{6}(4) + \frac{1}{6}(5) + \frac{1}{6}(6) = 3.5$$

Notice that 3.5 is not a possible outcome of any single roll, but it is the average we would expect over many rolls.

B.3.4.1 Properties of Expected Values

The properties of expected values mirror those of the summation operator:

Property 1: For any constant c , $E[c] = c$.

Property 2: For any constants a and b , $E[aX + b] = aE[X] + b$.

Property 3: If $\{a_1, a_2, \dots, a_n\}$ are constants and $\{X_1, X_2, \dots, X_n\}$ are random variables, then:

$$E \left[\sum_{i=1}^n a_i X_i \right] = \sum_{i=1}^n a_i E[X_i]$$

B.4 Measures of Variability

Knowing the central tendency of a variable is useful, but it doesn't tell us everything. Two distributions can have the same mean but look very different if one is tightly clustered around the mean while the other is spread out. We need measures of **variability** or **dispersion**.

B.4.1 Variance

Let X be a random variable with mean $\mu = E[X]$. The **variance** of X , denoted σ^2 or $\text{Var}(X)$, measures how spread out the distribution is:

$$\text{Var}(X) = E[(X - \mu)^2] = \sigma^2$$

The variance is the expected squared deviation from the mean. Squaring serves two purposes: it ensures all deviations are positive (so positive and negative deviations don't cancel out), and it gives more weight to observations far from the mean.

B.4.2 Standard Deviation

The **standard deviation** of X , denoted $\text{sd}(X)$ or σ , is the positive square root of the variance:

$$\text{sd}(X) = \sqrt{\text{Var}(X)} = \sigma$$

The standard deviation is often more interpretable than the variance because it is measured in the same units as X itself. If X is measured in dollars, the standard deviation is also in dollars, whereas the variance would be in “dollars squared.”

When computing the variance and standard deviation from a sample of n observations, we use the **sample variance** $s^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2$ and **sample standard deviation** $s = \sqrt{s^2}$. The $n - 1$ in the denominator (rather than n) is called Bessel’s correction and ensures the sample variance is an unbiased estimator of the population variance.

B.4.3 Covariance

A central concern in econometrics is understanding how variables move together. When X is large, is Y also large? Or is it small? Or is there no relationship at all?

The **covariance** quantifies the linear relationship between two random variables. Let X and Y be random variables with means $\mu_X = E[X]$ and $\mu_Y = E[Y]$. The covariance is defined as:

$$\text{Cov}(X, Y) = E[(X - \mu_X)(Y - \mu_Y)]$$

The covariance measures how X and Y move together:

- **Positive covariance:** When X is above its mean, Y tends to be above its mean (and vice versa). The variables move in the same direction.
- **Negative covariance:** When X is above its mean, Y tends to be below its mean. The variables move in opposite directions.
- **Zero covariance:** There is no linear relationship between X and Y .

B.4.4 Correlation Coefficient

The covariance tells us the direction of the relationship, but its magnitude depends on the scales of X and Y , making it hard to interpret. The **correlation coefficient** standardizes the covariance to produce a unitless measure:

$$\text{Corr}(X, Y) = \frac{\text{Cov}(X, Y)}{\text{sd}(X) \cdot \text{sd}(Y)} = \rho_{XY}$$

The correlation coefficient is always between -1 and 1 :

- $\rho_{XY} = 1$: Perfect positive linear relationship
- $\rho_{XY} = -1$: Perfect negative linear relationship
- $\rho_{XY} = 0$: No linear relationship

Perfect correlations rarely occur in practice. Values of ρ_{XY} closer to 1 or -1 indicate stronger linear relationships.

Correlation vs. Causation

Remember from Chapter 1: correlation does not imply causation! A strong correlation between X and Y tells us the variables move together, but it does not tell us whether X causes Y , Y causes X , or some third factor causes both.

B.4.5 Conditional Expectations

While correlation measures the overall linear relationship between two variables, we often want to describe how the expected value of one variable changes with another. This is where **conditional expectations** come in.

Let X and Y be random variables. The conditional expectation of Y given that X takes a specific value x is denoted:

$$E[Y|X = x]$$

This asks: what is our “best guess” for Y when we know that X equals x ?

Example: What is the expected wage (Y) for people with a college degree ($X = 1$) versus those without ($X = 0$)? $E[Y|X = 1]$ and $E[Y|X = 0]$ answer these questions.

Conditional expectations are fundamental to regression analysis. When we estimate a regression of Y on X , we are essentially modeling how $E[Y|X]$ changes with X .

B.5 Distributions

B.5.1 The Normal Distribution


The **normal distribution** (also called the Gaussian distribution) is the most important distribution in statistics and econometrics. Many naturally occurring phenomena are approximately normally distributed, and many statistical procedures assume normality.

A random variable X with mean μ and variance σ^2 that follows a normal distribution is denoted:

$$X \sim N(\mu, \sigma^2)$$

The normal distribution has a characteristic “bell curve” shape: symmetric around the mean, with most observations clustered near the center and fewer observations in the tails.

Figure B.3: A histogram of 10,000 draws from a normal distribution with mean 20 and standard deviation 10.



Two key facts about the normal distribution:

- Approximately **68%** of observations lie within one standard deviation of the mean (between $\mu - \sigma$ and $\mu + \sigma$)
- Approximately **95%** of observations lie within two standard deviations of the mean (between $\mu - 2\sigma$ and $\mu + 2\sigma$)

B.5.2 The Standard Normal Distribution

The **standard normal distribution** is a special case of the normal distribution with mean 0 and variance 1:


$$Z \sim N(0, 1)$$

The standard normal is important because any normal random variable can be converted to a standard normal through **standardization**. If $X \sim N(\mu, \sigma^2)$, then:

$$Z = \frac{X - \mu}{\sigma} \sim N(0, 1)$$

Standardization subtracts the mean (centering the distribution at zero) and divides by the standard deviation (scaling it to have unit variance).

Figure B.4: A histogram of 10,000 draws from a standard normal distribution (mean 0, standard deviation 1).



The standard normal is what z-tables (or statistical software) use to compute probabilities. Once you standardize a variable, you can look up the probability of observing values in any range.

B.6 Summary

This appendix reviewed the mathematical and statistical foundations needed for econometrics. We covered the summation operator and its properties, which are essential for understanding formulas throughout the course. We introduced derivatives and partial derivatives, which help us interpret how changes in one variable affect another, and learned how to find and classify maxima and minima of functions. We distinguished between populations and samples—a fundamental concept for statistical inference. We then turned to probability, defining random variables, expected values, variance, covariance, and correlation. Finally, we introduced the normal distribution, which plays a central role in statistical inference.

B.6.1 Key Formulas

Concept	Formula
Sample mean	$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$
Percent change	$\% \Delta Y = \frac{Y_2 - Y_1}{Y_1} \times 100\%$
Value after % increase	$Y_{new} = Y_{old} \times (1 + p/100)$
Log product rule	$\ln(ab) = \ln(a) + \ln(b)$
Log quotient rule	$\ln(a/b) = \ln(a) - \ln(b)$
Log power rule	$\ln(a^k) = k \ln(a)$
Logs and % changes	$\ln(Y_2) - \ln(Y_1) \approx \% \Delta Y$
Derivative of $\ln(x)$	$\frac{d}{dx} \ln(x) = \frac{1}{x}$
Critical point (quadratic)	$x^* = -\frac{b}{2a}$ for $f(x) = ax^2 + bx + c$
Second derivative test	Max if $f''(x^*) < 0$; Min if $f''(x^*) > 0$
Expected value	$E[X] = \sum_{j=1}^k x_j f(x_j) = \mu$
Variance	$\text{Var}(X) = E[(X - \mu)^2] = \sigma^2$
Standard deviation	$\text{sd}(X) = \sqrt{\text{Var}(X)} = \sigma$
Covariance	$\text{Cov}(X, Y) = E[(X - \mu_X)(Y - \mu_Y)]$
Correlation	$\rho_{XY} = \frac{\text{Cov}(X, Y)}{\text{sd}(X) \cdot \text{sd}(Y)}$
Standardization	$Z = \frac{X - \mu}{\sigma}$

B.7 Check Your Understanding

Note: This section contains interactive content only available in the HTML version.

C R Functions and Packages Reference

This appendix provides a quick reference for all the R functions and packages used throughout this book. Use this as a handy guide when you need to remember the syntax or arguments for a particular function.

C.1 Packages Used in This Book

C.1.1 tidyverse

The `tidyverse` is a collection of R packages designed for data science. When you load `tidyverse`, it automatically loads several packages including `ggplot2`, `dplyr`, `tidyr`, `readr`, and others.

What it's used for:

- Data manipulation and transformation (`dplyr`)
- Data visualization (`ggplot2`)
- Reshaping data (`tidyr`)
- Reading data files (`readr`)

Loading:

```
library(tidyverse)
```

C.1.2 modelsummary

The `modelsummary` package creates publication-quality tables for regression results and descriptive statistics.

What it's used for:

- Creating formatted regression tables comparing multiple models
- Generating descriptive statistics tables

- Exporting tables to Word, LaTeX, or HTML

Loading:

```
library(modelsummary)
```

C.1.3 palmerpenguins

A dataset package containing measurements of penguins from Palmer Station, Antarctica. Great for learning data visualization and exploration.

What it's used for:

- Practice dataset for learning R
- Data visualization examples
- Regression examples

Loading:

```
library(palmerpenguins)  
data(penguins)
```

C.1.4 wooldridge

Contains all datasets from Wooldridge's *Introductory Econometrics* textbook.

What it's used for:

- Real-world econometric datasets
- Regression examples (wages, crime, etc.)

Loading:

```
library(wooldridge)  
data(wage1) # Example: load the wage1 dataset
```

C.1.5 fixest

A fast and powerful package for fixed effects and panel data regression.

What it's used for:

- Panel data regression with fixed effects
- Clustered standard errors
- Very large datasets (faster than `lm()`)

Loading:

```
library(fixest)
```

C.1.6 lmtest

Provides diagnostic tests for linear regression models.

What it's used for:

- Breusch-Pagan test for heteroskedasticity
- Other specification tests

Loading:

```
library(lmtest)
```

C.1.7 sandwich

Provides robust covariance matrix estimators (heteroskedasticity-consistent standard errors).

What it's used for:

- Robust standard errors
- Heteroskedasticity-consistent (HC) standard errors

Loading:

```
library(sandwich)
```

C.1.8 patchwork

Easily combine multiple ggplot2 plots into a single figure.

What it's used for:

- Arranging multiple plots side by side
- Creating multi-panel figures

Loading:

```
library(patchwork)
```

C.2 Data Inspection Functions

C.2.1 head()

Display the first few rows of a data frame.

Arguments:

- x: A data frame or vector
- n: Number of rows to display (default: 6)

Example:

```
head(mtcars, n = 3)
```

	mpg	cyl	disp	hp	drat	wt	qsec	vs	am	gear	carb
Mazda RX4	21.0	6	160	110	3.90	2.620	16.46	0	1	4	4
Mazda RX4 Wag	21.0	6	160	110	3.90	2.875	17.02	0	1	4	4
Datsun 710	22.8	4	108	93	3.85	2.320	18.61	1	1	4	1

C.2.2 glimpse()

Get a compact overview of a data frame's structure and column types.

Arguments:

- x: A data frame

Example:

```
glimpse(mtcars)
```

```
Rows: 32
```

```
Columns: 11
```

```
$ mpg <dbl> 21.0, 21.0, 22.8, 21.4, 18.7, 18.1, 14.3, 24.4, 22.8, 19.2, 17.8,~  
$ cyl <dbl> 6, 6, 4, 6, 8, 6, 8, 4, 4, 6, 6, 8, 8, 8, 8, 8, 8, 4, 4, 4, 4, 8,~  
$ disp <dbl> 160.0, 160.0, 108.0, 258.0, 360.0, 225.0, 360.0, 146.7, 140.8, 16~  
$ hp <dbl> 110, 110, 93, 110, 175, 105, 245, 62, 95, 123, 123, 180, 180, 180~  
$ drat <dbl> 3.90, 3.90, 3.85, 3.08, 3.15, 2.76, 3.21, 3.69, 3.92, 3.92, 3.92,~  
$ wt <dbl> 2.620, 2.875, 2.320, 3.215, 3.440, 3.460, 3.570, 3.190, 3.150, 3.~  
$ qsec <dbl> 16.46, 17.02, 18.61, 19.44, 17.02, 20.22, 15.84, 20.00, 22.90, 18~  
$ vs <dbl> 0, 0, 1, 1, 0, 1, 0, 1, 1, 1, 1, 0, 0, 0, 0, 0, 1, 1, 1, 1, 0,~  
$ am <dbl> 1, 1, 1, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 1, 1, 1, 0, 0,~  
$ gear <dbl> 4, 4, 4, 3, 3, 3, 3, 4, 4, 4, 4, 3, 3, 3, 3, 3, 4, 4, 4, 3, 3,~  
$ carb <dbl> 4, 4, 1, 1, 2, 1, 4, 2, 2, 4, 4, 3, 3, 3, 4, 4, 4, 1, 2, 1, 1, 2,~
```

C.2.3 summary()

Generate summary statistics for each variable.

Arguments:

- object: A data frame, vector, or model object

Example:

```
summary(mtcars$mpg)
```

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
10.40	15.43	19.20	20.09	22.80	33.90

C.2.4 `nrow()` and `ncol()`

Return the number of rows or columns in a data frame.

Arguments:

- `x`: A data frame or matrix

Example:

```
nrow(mtcars)
```

```
[1] 32
```

```
ncol(mtcars)
```

```
[1] 11
```

C.2.5 `colnames()`

Return or set the column names of a data frame.

Arguments:

- `x`: A data frame or matrix

Example:

```
colnames(mtcars)
```

```
[1] "mpg"  "cyl"  "disp" "hp"    "drat" "wt"    "qsec" "vs"    "am"    "gear"  
[11] "carb"
```

C.2.6 class() and typeof()

Return the class or underlying type of an object.

Arguments:

- `x`: Any R object

Example:

```
class(mtcars$mpg)
```

```
[1] "numeric"
```

```
typeof(mtcars$mpg)
```

```
[1] "double"
```

C.3 Descriptive Statistics Functions

C.3.1 mean()

Calculate the arithmetic mean.

Arguments:

- `x`: A numeric vector
- `na.rm`: Remove missing values? (default: FALSE)

Example:

```
x <- c(10, 20, 30, NA, 50)
mean(x, na.rm = TRUE)
```

```
[1] 27.5
```

C.3.2 sd()

Calculate the standard deviation.

Arguments:

- **x**: A numeric vector
- **na.rm**: Remove missing values? (default: FALSE)

Example:

```
sd(mtcars$mpg)
```

```
[1] 6.026948
```

C.3.3 median()

Calculate the median (middle value).

Arguments:

- **x**: A numeric vector
- **na.rm**: Remove missing values? (default: FALSE)

Example:

```
median(mtcars$mpg)
```

```
[1] 19.2
```

C.3.4 min() and max()

Find the minimum or maximum value.

Arguments:

- ...: Numeric vectors
- na.rm: Remove missing values? (default: FALSE)

Example:

```
min(mtcars$mpg)
```

```
[1] 10.4
```

```
max(mtcars$mpg)
```

```
[1] 33.9
```

C.3.5 sum()

Calculate the sum of all values.

Arguments:

- ...: Numeric vectors
- na.rm: Remove missing values? (default: FALSE)

Example:

```
sum(mtcars$mpg)
```

```
[1] 642.9
```

C.3.6 var()

Calculate the variance.

Arguments:

- **x**: A numeric vector
- **na.rm**: Remove missing values? (default: FALSE)

Example:

```
var(mtcars$mpg)
```

```
[1] 36.3241
```

C.3.7 cor()

Calculate the correlation between two variables.

Arguments:

- **x, y**: Numeric vectors
- **use**: How to handle missing values (e.g., "complete.obs")

Example:

```
cor(mtcars$mpg, mtcars$hp)
```

```
[1] -0.7761684
```

C.4 Data Manipulation Functions (dplyr)

C.4.1 `select()`

Choose which columns to keep.

Arguments:

- `.data`: A data frame
- `...`: Column names or selection helpers

Example:

```
mtcars |>
  select(mpg, cyl, hp) |>
  head(3)
```

	mpg	cyl	hp
Mazda RX4	21.0	6	110
Mazda RX4 Wag	21.0	6	110
Datsun 710	22.8	4	93

C.4.2 `filter()`

Keep rows that meet a condition.

Arguments:

- `.data`: A data frame
- `...`: Logical conditions

Example:

```
mtcars |>
  filter(mpg > 25) |>
  head(3)
```

	mpg	cyl	disp	hp	drat	wt	qsec	vs	am	gear	carb
Fiat 128	32.4	4	78.7	66	4.08	2.200	19.47	1	1	4	1
Honda Civic	30.4	4	75.7	52	4.93	1.615	18.52	1	1	4	2
Toyota Corolla	33.9	4	71.1	65	4.22	1.835	19.90	1	1	4	1

C.4.3 mutate()

Create new variables or modify existing ones.

Arguments:

- `.data`: A data frame
- `...`: Name-value pairs of expressions

Example:

```
mtcars |>
  mutate(kpl = mpg * 0.425) |> # Convert to km per liter
  select(mpg, kpl) |>
  head(3)
```

	mpg	kpl
Mazda RX4	21.0	8.925
Mazda RX4 Wag	21.0	8.925
Datsun 710	22.8	9.690

C.4.4 group_by()

Group data by one or more variables (usually followed by `summarize()`).

Arguments:

- `.data`: A data frame
- `...`: Variables to group by

Example:

```
mtcars |>
  group_by(cyl) |>
  summarize(avg_mpg = mean(mpg))
```



```
# A tibble: 3 x 2
  cyl avg_mpg
<dbl> <dbl>
1     4    26.7
2     6    19.7
3     8    15.1
```

C.4.5 summarize() / summarise()

Calculate summary statistics for each group.

Arguments:

- `.data`: A data frame (usually grouped)
- `...`: Name-value pairs of summary functions

Example:

```
mtcars |>
  group_by(cyl) |>
  summarize(
    avg_mpg = mean(mpg),
    sd_mpg = sd(mpg),
    n = n()
  )
```

```
# A tibble: 3 x 4
  cyl avg_mpg sd_mpg    n
<dbl> <dbl> <dbl> <int>
1     4    26.7   4.51    11
2     6    19.7   1.45     7
3     8    15.1   2.56    14
```

C.4.6 count()

Count observations by group.

Arguments:

- **x**: A data frame
- **...**: Variables to count by
- **sort**: Sort by count? (default: FALSE)

Example:

```
mtcars |>
  count(cyl)
```

```
   cyl  n
1    4 11
2    6  7
3    8 14
```

C.4.7 arrange()

Sort rows by one or more variables.

Arguments:

- **.data**: A data frame
- **...**: Variables to sort by (use **desc()** for descending)

Example:

```
mtcars |>
  arrange(desc(mpg)) |>
  head(3)
```

```
      mpg cyl disp hp drat   wt  qsec vs am gear carb
Toyota Corolla 33.9   4  71.1 65 4.22 1.835 19.90 1  1    4    1
Fiat 128       32.4   4  78.7 66 4.08 2.200 19.47 1  1    4    1
Honda Civic    30.4   4  75.7 52 4.93 1.615 18.52 1  1    4    2
```

C.4.8 case_when()

Vectorized if-else for creating categorical variables.

Arguments:

- ...: A sequence of two-sided formulas: `condition ~ value`

Example:

```
mtcars |>
  mutate(
    mpg_category = case_when(
      mpg < 15 ~ "Low",
      mpg < 25 ~ "Medium",
      TRUE ~ "High"
    )
  ) |>
  count(mpg_category)
```

	mpg_category	n
1	High	6
2	Low	5
3	Medium	21

C.4.9 ifelse()

Simple conditional: if condition is TRUE, return one value; otherwise, another.

Arguments:

- `test`: A logical condition
- `yes`: Value if TRUE
- `no`: Value if FALSE

Example:

```
mtcars |>
  mutate(efficient = ifelse(mpg > 20, 1, 0)) |>
  count(efficient)
```

	efficient	n
1	0	18
2	1	14

C.4.10 `n()`

Count the number of observations in the current group (used inside `summarize()`).

Arguments: None

Example:

```
mtcars |>
  group_by(cyl) |>
  summarize(count = n())
```

```
# A tibble: 3 x 2
  cyl count
<dbl> <int>
1     4    11
2     6     7
3     8    14
```

C.5 Regression Functions

C.5.1 `lm()`

Fit a linear model using Ordinary Least Squares (OLS).

Arguments:

- **formula:** A formula like `y ~ x1 + x2`
- **data:** A data frame

Example:

```
reg <- lm(mpg ~ hp + wt, data = mtcars)
summary(reg)
```

Call:

```
lm(formula = mpg ~ hp + wt, data = mtcars)
```

Residuals:

	Min	1Q	Median	3Q	Max
	-3.941	-1.600	-0.182	1.050	5.854

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	37.22727	1.59879	23.285	< 2e-16 ***
hp	-0.03177	0.00903	-3.519	0.00145 **
wt	-3.87783	0.63273	-6.129	1.12e-06 ***

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 2.593 on 29 degrees of freedom

Multiple R-squared: 0.8268, Adjusted R-squared: 0.8148

F-statistic: 69.21 on 2 and 29 DF, p-value: 9.109e-12

C.5.2 summary() (for regression)

Display detailed regression results including coefficients, standard errors, t-statistics, and p-values.

Arguments:

- object: A fitted model object

Example:

```
reg <- lm(mpg ~ hp, data = mtcars)
summary(reg)
```

C.5.3 `coef()`

Extract the estimated coefficients from a model.

Arguments:

- `object`: A fitted model object

Example:

```
reg <- lm(mpg ~ hp, data = mtcars)
coef(reg)
```

```
(Intercept)          hp
30.09886054 -0.06822828
```

C.5.4 `confint()`

Calculate confidence intervals for model coefficients.

Arguments:

- `object`: A fitted model object
- `level`: Confidence level (default: 0.95)

Example:

```
reg <- lm(mpg ~ hp, data = mtcars)
confint(reg, level = 0.95)
```

```
                2.5 %    97.5 %
(Intercept) 26.76194879 33.4357723
hp          -0.08889465 -0.0475619
```

C.5.5 predict()

Generate predicted values from a fitted model.

Arguments:

- **object**: A fitted model object
- **newdata**: Optional data frame with new predictor values

Example:

```
reg <- lm(mpg ~ hp, data = mtcars)
# Predict for existing data
head(predict(reg))
```

	Mazda RX4	Mazda RX4 Wag	Datsun 710	Hornet 4 Drive
	22.59375	22.59375	23.75363	22.59375
Hornet Sportabout		Valiant		
	18.15891	22.93489		

```
# Predict for new data
predict(reg, newdata = data.frame(hp = c(100, 150, 200)))
```

1	2	3
23.27603	19.86462	16.45320

C.5.6 residuals()

Extract residuals (prediction errors) from a fitted model.

Arguments:

- **object**: A fitted model object

Example:

```
reg <- lm(mpg ~ hp, data = mtcars)
head(residuals(reg))
```

	Mazda RX4	Mazda RX4 Wag	Datsun 710	Hornet 4 Drive
	-1.5937500	-1.5937500	-0.9536307	-1.1937500
Hornet Sportabout		Valiant		
	0.5410881	-4.8348913		

C.5.7 feols() (fixest package)

Fit linear models with fixed effects and robust standard errors.

Arguments:

- `fml`: A formula (use `|` for fixed effects)
- `data`: A data frame
- `vcov`: Type of standard errors (e.g., "HC1" for robust)

Example:

```
library(fixest)
reg <- feols(mpg ~ hp + wt, data = mtcars, vcov = "HC1")
summary(reg)
```

C.5.8 modelsummary() (modelsummary package)

Create publication-quality regression tables.

Arguments:

- `models`: A model or list of models
- `stars`: Show significance stars?
- `gof_map`: Which goodness-of-fit statistics to show

Example:

```
library(modelsummary)
reg1 <- lm(mpg ~ hp, data = mtcars)
reg2 <- lm(mpg ~ hp + wt, data = mtcars)
modelsummary(list(reg1, reg2), stars = TRUE)
```

C.6 Visualization Functions (ggplot2)

C.6.1 ggplot()


Initialize a ggplot object.

Arguments:

- **data:** A data frame
- **mapping:** Aesthetic mappings created by `aes()`

Example:

```
ggplot(mtcars, aes(x = hp, y = mpg)) +  
  geom_point()
```



C.6.2 aes()

Define aesthetic mappings (which variables map to x, y, color, etc.).


Arguments:

- **x, y:** Variables for axes
- **color, fill:** Variables for color

- **size, shape:** Variables for size and shape

Example:

```
ggplot(mtcars, aes(x = hp, y = mpg, color = factor(cyl))) +  
  geom_point()
```



C.6.3 geom_point()


Add points to a plot (scatterplot).

Arguments:

- **size:** Point size
- **color:** Point color
- **alpha:** Transparency (0 to 1)

Example:

```
ggplot(mtcars, aes(x = hp, y = mpg)) +  
  geom_point(size = 3, color = "steelblue", alpha = 0.7)
```



C.6.4 `geom_line()`


Add lines connecting points.

Arguments:

- `linewidth`: Line thickness
- `color`: Line color
- `linetype`: Line type (e.g., “dashed”)

Example:

```
# Line plot (useful for time series)
df <- data.frame(x = 1:10, y = cumsum(rnorm(10)))
ggplot(df, aes(x = x, y = y)) +
  geom_line(color = "steelblue", linewidth = 1)
```



C.6.5 geom_histogram()


Create a histogram.

Arguments:

- **binwidth**: Width of each bin
- **bins**: Number of bins
- **fill**: Bar fill color
- **color**: Bar outline color

Example:

```
ggplot(mtcars, aes(x = mpg)) +  
  geom_histogram(binwidth = 3, fill = "steelblue", color = "white")
```



C.6.6 geom_boxplot()


Create a boxplot.

Arguments:

- `fill`: Box fill color
- `outlier.color`: Color of outlier points

Example:

```
ggplot(mtcars, aes(x = factor(cyl), y = mpg)) +  
  geom_boxplot(fill = "lightblue")
```



C.6.7 geom_bar() and geom_col()


Create bar charts. `geom_bar()` counts observations; `geom_col()` uses values directly.

Arguments:

- `fill`: Bar fill color
- `stat`: For `geom_bar()`, use "identity" to plot values directly

Example:

```
# geom_bar counts automatically
ggplot(mtcars, aes(x = factor(cyl))) +
  geom_bar(fill = "steelblue")
```



C.6.8 geom_smooth()

Add a smoothed conditional mean (often a regression line).


Arguments:

- **method**: Smoothing method (e.g., "lm" for linear)
- **se**: Show confidence interval? (default: TRUE)
- **color**: Line color

Example:

```
ggplot(mtcars, aes(x = hp, y = mpg)) +  
  geom_point() +  
  geom_smooth(method = "lm", se = TRUE, color = "red")
```

``geom_smooth()`` using `formula = 'y ~ x'`



C.6.9 geom_hline() and geom_vline()


Add horizontal or vertical reference lines.

Arguments:

- `yintercept` / `xintercept`: Where to draw the line
- `linetype`: Line type (e.g., "dashed")
- `color`: Line color

Example:

```
ggplot(mtcars, aes(x = hp, y = mpg)) +  
  geom_point() +  
  geom_hline(yintercept = mean(mtcars$mpg), linetype = "dashed", color = "red")
```

C.6.10 labs()


Add labels to the plot (title, axis labels, etc.).

Arguments:

- title: Plot title
- x, y: Axis labels
- color, fill: Legend titles

Example:

```
ggplot(mtcars, aes(x = hp, y = mpg)) +  
  geom_point() +  
  labs(  
    title = "Fuel Efficiency vs. Horsepower",  
    x = "Horsepower",  
    y = "Miles per Gallon"  
  )
```



C.6.11 facet_wrap()


Create small multiples (separate panels for each level of a variable).

Arguments:

- **facets:** A formula like `~ variable`
- **nrow, ncol:** Number of rows or columns
- **scales:** Should scales be fixed or free?

Example:

```
ggplot(mtcars, aes(x = hp, y = mpg)) +  
  geom_point() +  
  facet_wrap(~ cyl, nrow = 1)
```




C.6.12 `theme_minimal()`

Apply a clean, minimal theme to the plot.

Arguments: None (or see `theme()` for customization)

Example:

```
ggplot(mtcars, aes(x = hp, y = mpg)) +  
  geom_point() +  
  theme_minimal()
```



C.7 Utility Functions

C.7.1 c()

Combine values into a vector.

Arguments:

- ...: Values to combine

Example:

```
x <- c(1, 2, 3, 4, 5)
x
```

```
[1] 1 2 3 4 5
```

C.7.2 seq()

Generate a sequence of numbers.

Arguments:

- **from**: Starting value
- **to**: Ending value
- **by**: Increment
- **length.out**: Desired length of sequence

Example:

```
seq(from = 0, to = 10, by = 2)
```

```
[1] 0 2 4 6 8 10
```

```
seq(from = 0, to = 1, length.out = 5)
```

```
[1] 0.00 0.25 0.50 0.75 1.00
```

C.7.3 rep()

Repeat values.

Arguments:

- **x**: Value(s) to repeat
- **times**: Number of times to repeat

Example:

```
rep(c("A", "B"), times = 3)
```

```
[1] "A" "B" "A" "B" "A" "B"
```

C.7.4 sample()

Take a random sample.

Arguments:

- **x**: Vector to sample from
- **size**: Number of items to sample
- **replace**: Sample with replacement?

Example:

```
set.seed(123)
sample(1:10, size = 5, replace = FALSE)
```

```
[1]  3 10  2  8  6
```

C.7.5 set.seed()

Set the random number seed for reproducibility.

Arguments:

- **seed**: An integer

Example:

```
set.seed(42)
rnorm(3) # Will always produce the same values
```

```
[1]  1.3709584 -0.5646982  0.3631284
```

C.7.6 rnorm()

Generate random numbers from a normal distribution.

Arguments:

- **n**: Number of values to generate
- **mean**: Mean of the distribution (default: 0)
- **sd**: Standard deviation (default: 1)

Example:

```
set.seed(123)
rnorm(5, mean = 100, sd = 15)
```

```
[1] 91.59287 96.54734 123.38062 101.05763 101.93932
```

C.7.7 factor()

Create a factor (categorical variable).

Arguments:

- **x**: A vector
- **levels**: The allowed levels (in order)
- **labels**: Labels for the levels

Example:

```
education <- c("HS", "College", "HS", "Graduate")
factor(education, levels = c("HS", "College", "Graduate"))
```

```
[1] HS      College HS      Graduate
Levels: HS College Graduate
```

C.7.8 `as.factor()`, `as.numeric()`, `as.character()`

Convert objects to a different type.

Arguments:

- `x`: Object to convert

Example:

```
x <- c("1", "2", "3")
as.numeric(x)
```

```
[1] 1 2 3
```

C.7.9 `is.na()`

Check for missing values.

Arguments:

- `x`: A vector or data frame

Example:

```
x <- c(1, 2, NA, 4)
is.na(x)
```

```
[1] FALSE FALSE  TRUE FALSE
```

```
sum(is.na(x)) # Count missing values
```

```
[1] 1
```

C.7.10 library()

Load an installed package.

Arguments:

- **package:** Name of the package (unquoted)

Example:

```
library(tidyverse)
```

C.7.11 install.packages()

Install a package from CRAN (run once, not in scripts).

Arguments:

- **pkgs:** Package name (quoted)

Example:

```
install.packages("tidyverse")
```

C.8 The Pipe Operator: |>

The pipe operator takes the output from the left side and passes it as the first argument to the function on the right. This allows you to chain operations together in a readable way.

Example without pipe:

```
# Nested functions are hard to read  
summary(select(filter(mtcars, mpg > 20), mpg, hp))
```

Example with pipe:

```
# Piped version is much clearer
mtcars |>
  filter(mpg > 20) |>
  select(mpg, hp) |>
  summary()
```

```
      mpg      hp
Min.   :21.00  Min.   : 52.0
1st Qu.:21.43  1st Qu.: 66.0
Median :23.60  Median : 94.0
Mean   :25.48  Mean    : 88.5
3rd Qu.:29.62  3rd Qu.:109.8
Max.   :33.90  Max.    :113.0
```

💡 Keyboard Shortcut

In RStudio, type **Cmd/Ctrl + Shift + M** to insert the pipe operator.

C.9 Quick Reference Tables

C.9.1 Descriptive Statistics Functions

Table C.1: Descriptive statistics functions

Function	Purpose	Key Argument
<code>mean()</code>	Average	<code>na.rm = TRUE</code>
<code>sd()</code>	Standard deviation	<code>na.rm = TRUE</code>
<code>median()</code>	Middle value	<code>na.rm = TRUE</code>
<code>min(), max()</code>	Extremes	<code>na.rm = TRUE</code>
<code>sum()</code>	Total	<code>na.rm = TRUE</code>
<code>var()</code>	Variance	<code>na.rm = TRUE</code>
<code>cor()</code>	Correlation	<code>use = "complete.obs"</code>
<code>summary()</code>	Multiple stats	—

C.9.2 Data Manipulation Functions (dplyr)

Table C.2: Data manipulation functions

Function	Purpose	Example
<code>select()</code>	Choose columns	<code>select(data, col1, col2)</code>
<code>filter()</code>	Choose rows	<code>filter(data, x > 5)</code>
<code>mutate()</code>	Create/modify variables	<code>mutate(data, new = x * 2)</code>
<code>group_by()</code>	Group data	<code>group_by(data, category)</code>
<code>summarize()</code>	Aggregate	<code>summarize(data, avg = mean(x))</code>
<code>count()</code>	Count rows	<code>count(data, category)</code>
<code>arrange()</code>	Sort rows	<code>arrange(data, desc(x))</code>

C.9.3 Regression Functions

Table C.3: Regression functions

Function	Purpose	Package
<code>lm()</code>	OLS regression	base R
<code>summary()</code>	Model results	base R
<code>coef()</code>	Coefficients	base R
<code>confint()</code>	Confidence intervals	base R
<code>predict()</code>	Fitted values	base R
<code>residuals()</code>	Residuals	base R
<code>feols()</code>	Fixed effects + robust SE	fixest
<code>modelsummary()</code>	Formatted tables	modelsummary

C.9.4 ggplot2 Geometries

Table C.4: ggplot2 geometries

Function	Plot Type
<code>geom_point()</code>	Scatterplot
<code>geom_line()</code>	Line plot
<code>geom_histogram()</code>	Histogram
<code>geom_boxplot()</code>	Boxplot
<code>geom_bar()</code>	Bar chart (counts)
<code>geom_col()</code>	Bar chart (values)

Function	Plot Type
<code>geom_smooth()</code>	Smoothed line/regression
<code>geom_hline()</code>	Horizontal line
<code>geom_vline()</code>	Vertical line