

Computability

Rasmus Guldberg Pedersen

January 2015

Deterministic and non-deterministic finite automata

Define the two types of finite automata. Are the classes of languages they define the same? Explain and justify your answer.

Finite Automata

$(Q, \Sigma, q_0, A, \delta)$

Q is a finite set of *states*;

Σ is a finite *input alphabet*;

$q_0 \in Q$ is the *initial* state;

$A \subseteq Q$ is the set of *accepting* states;

δ is the *transition* function.

Deterministic & Nondeterministic FA

DFA

$$\delta : Q \times \Sigma \rightarrow Q$$

For $q \in Q$ and $\sigma \in \Sigma$ then $\delta(q, \sigma)$ denotes the state transition from q on input σ .

NFA

$$\delta : Q \times (\Sigma \cup \{\Lambda\}) \rightarrow 2^Q$$

For $q \in Q$ and $\sigma \in (\Sigma \cup \{\Lambda\})$ then $\delta(q, \sigma)$ denotes the set of states the NFA can move to from q on input σ .

Eliminating Λ -transitions

$$M = (Q, \Sigma, q_0, A, \delta)$$

$$M_1 = (Q, \Sigma, q_0, A_1, \delta_1)$$

Eliminating Λ -transitions

$$M = (Q, \Sigma, q_0, A, \delta)$$

$$M_1 = (Q, \Sigma, q_0, A_1, \delta_1)$$

$$A_1 = \begin{cases} A \cup \{q_0\} & \text{if } \Lambda \in L \\ A & \text{if not} \end{cases}$$

Eliminating Λ -transitions

$$M = (Q, \Sigma, q_0, A, \delta)$$

$$M_1 = (Q, \Sigma, q_0, A_1, \delta_1)$$

$$A_1 = \begin{cases} A \cup \{q_0\} & \text{if } \Lambda \in L \\ A & \text{if not} \end{cases}$$

For every $q \in Q$ and every $\sigma \in \Sigma$ then $\delta_1(q, \sigma) = \delta^*(q, \sigma)$

DFA accepting the same language as an NFA

NFA $M = (Q, \Sigma, q_0, A, \delta)$

FA $M_1 = (Q_1, \Sigma, q_1, A_1, \delta_1)$

DFA accepting the same language as an NFA

NFA $M = (Q, \Sigma, q_0, A, \delta)$

FA $M_1 = (Q_1, \Sigma, q_1, A_1, \delta_1)$

$Q_1 = 2^Q$

$q_1 = q_0$

DFA accepting the same language as an NFA

NFA $M = (Q, \Sigma, q_0, A, \delta)$

FA $M_1 = (Q_1, \Sigma, q_1, A_1, \delta_1)$

$Q_1 = 2^Q$

$q_1 = q_0$

$A_1 = \{q \in Q_1 \mid q \cap A \neq \emptyset\}$

DFA accepting the same language as an NFA

NFA $M = (Q, \Sigma, q_0, A, \delta)$

FA $M_1 = (Q_1, \Sigma, q_1, A_1, \delta_1)$

$$Q_1 = 2^Q$$

$$q_1 = q_0$$

$$A_1 = \{q \in Q_1 \mid q \cap A \neq \emptyset\}$$

$$\delta_1(q, \sigma) = \bigcup \{\delta(p, \sigma) \mid p \in q\}$$

The End

The End