# Overview of optimal observables for EFT DM sensitivity

Rebecca Pickles, Darren Price

February 23, 2016



The University of Manchester

### Status

- ullet Added phase space VBFDM100: VBFDM with Jet 1 pT  $> 100 {
  m GeV}$
- Scaled the different dimensions with the effective field theory constraints:
  - D5c :  $\Lambda = 3.3 \text{ TeV}$
  - D5d : Λ = 6.6 TeV
  - D6a : Λ = 230 GeV
  - D6b : Λ = 330 GeV
  - This has meant that some dimensions would not be seen by any parameters
- Added ratio plots of  $\frac{Z \rightarrow \nu \nu + DM}{Z \rightarrow \nu \nu}$
- Started to look at which set of distributions will give us the broadest sensitivity to dark matter models
- No jet veto is applied for QCD suppression right now, so the background QCD+EWK Z is pessimistic


$$\frac{(Z \rightarrow \nu \nu) + DM}{Z \rightarrow \nu \nu}$$
:  $Y > 10 > M > 5 > N$ 


## VBFDM Phase space:

Parameter	D5a	D5b	D5c	D5d	D6a	D6b	D7a	D7b	D7c	D7d	
Etmiss	N	М	N	N	Υ	Υ	Υ	Υ	Υ	Υ	Г
Mjj	Y	Υ	N	N	N	N	M	M	M	M	
Jet1pT	М	M	N	N	Ν	N	M	M	M	M	
Jet2pT	N	N	N	N	Ν	N	M	M	M	M	
Jet1Eta	Y	Υ	N	N	Ν	Ν	N	Ν	N	Ν	
Jet2Eta	М	M	N	N	Ν	Ν	N	Ν	N	Ν	
DeltaEta	Y	Υ	N	N	Ν	Ν	N	Ν	N	Ν	
DeltaPhi	М	М	N	N	N	N	М	М	N	N	

• The parameters that have good sensitivity to the most dimensions are Etmiss and Mjj for the VBFDM phase space.

#### These are for a DM mass of 100GeV






$$\frac{(Z \to \nu \nu) + DM}{Z \to \nu \nu}$$
:  $Y > 10 > M > 5 > N$ 


## VBFDM100 Phase space:

Parameter	D5a	D5b	D5c	D5d	D6a	D6b	D7a	D7b	D7c	D7d	ı
Etmiss	N	М	N	N	Υ	Υ	Υ	Υ	Υ	Υ	П
Mjj	Y	Υ	Ν	N	N	Ν	М	М	M	M	
Jet1pT	M	М	Ν	N	N	Ν	М	М	M	M	
Jet2pT	N	Ν	Ν	N	N	Ν	М	М	M	M	
Jet1Eta	Y	Υ	Ν	N	Ν	N	Ν	Ν	Ν	Ν	
Jet2Eta	М	М	N	N	N	N	N	N	N	N	
DeltaEta	Y	Υ	Ν	N	N	Ν	N	Ν	Ν	Ν	
DeltaPhi	М	М	N	N	Ν	N	М	М	Ν	Ν	

• The parameters that have good sensitivity to the most dimensions are Etmiss and Mjj for the VBFDM100 phase space.

#### These are for a DM mass of 100GeV






$$\frac{(Z\to\nu\nu)+DM}{Z\to\nu\nu}:Y>10>M>5>N$$

## VBFDM OR Monojet Phase space:

Parameter	D5a	D5b	D5c	D5d	D6a	D6b	D7a	D7b	D7c	D7d	
Etmiss	М	М	М	М	Υ	Υ	Υ	Y	Y	Υ	Г
Mjj	Y	Υ	N	Ν	Ν	N	Ν	Ν	M	M	
Jet1pT	М	M	N	N	N	Ν	М	М	N	N	
Jet2pT	N	N	N	N	N	Ν	M	М	M	M	
Jet1Eta	Y	Υ	N	N	N	Ν	N	N	N	N	
Jet2Eta	М	M	N	N	N	Ν	N	N	N	N	
DeltaEta	Y	Υ	N	N	N	Ν	N	N	N	N	
DeltaPhi	N	Ν	Ν	Ν	Ν	Ν	Ν	N	Ν	Ν	

 The parameter that has the best sensitivity to the most dimensions is Etmiss, for the VBFDM OR Monojet phase space.

#### This is for a DM mass of 100GeV



 None of the dimensions scaled by the EFT constraints can bee seen here


$$\frac{(Z\to\nu\nu)+DM}{Z\to\nu\nu}:Y>10>M>5>N$$

## VBFDM OR Monojet High pT Phase space:

Parameter	D5a	D5b	D5c	D5d	D6a	D6b	D7a	D7b	D7c	D7d	ı
Etmiss	М	М	N	N	Υ	Υ	Υ	Υ	Υ	Υ	Г
Mjj	Y	Υ	Ν	N	Ν	N	Ν	Ν	M	M	
Jet1pT	М	М	N	N	N	N	М	М	N	N	
Jet2pT	N	Ν	N	N	N	N	М	М	M	M	
Jet1Eta	Y	Υ	N	N	N	Ν	N	N	N	N	
Jet2Eta	М	М	N	N	N	Ν	N	N	N	N	
DeltaEta	Y	Υ	N	N	N	N	N	N	N	N	
DeltaPhi	М	М	N	N	N	N	N	N	N	N	


• The parameter that has the best sensitivity to the most dimensions is Etmiss for the VBFDM OR Monojet High pT phase space.

### Most broadly sensitive parameter plots: VBFDM OR Monojet High pT Phase space




## Best parameter?

- Due to the scaling with the EFT constraints, some dimensions would not be seen by any parameters. ( D5c, D5d, D6a, D6b )
- Etmiss seems to be the parameter that is sensitive to the most dimensions.
- This doesn't change for the different masses that have been looked at:




# Backup slides

### Phase space VBFDM : Other Parameters



### Phase space VBFDM or Monojet: Other Parameters



### Phase space VBFDM or Monojet High pT: Other Parameters

