

Update on observable phase spaces

Rebecca Pickles, Darren Price

April 26, 2016



Update on observable phase space plots


Done:

- Fixed bug causing EFT scale to not make a difference.
- Found issue with the SM being normalised to 1: Though still not as big as the corrected data from Emily. This could be due to not having the jet veto as only two jets are currently being produced.

To-do:


- Add more masses and scales (iterate though a large number of scales).
- Change calculation of ratio to prevent double counting of uncertainties.
- Make more presentable (Swap the colour scheme around, Change the range of the y axis on the ratio plots, Fix the y axis on the phase space plots, etc.).

Observable phase space plots : Mjj : D7a




- Large range of p-values (Some so small they don't show up on plot).
- Exclude region at high EFT scale and high DM mass.

Observable phase space plots : M_{jj} : D5d




- All p-values here are 1 as they are so close to the SM due to the EFT scale constraint.

Observable phase space plots : $\Delta\eta$: D7a




- Again a large range of p-values.
- Exclude region at high EFT scale and high DM mass.
- Gives better exclusion than M_{jj} , but still in same area.

Observable phase space plots : E_T : D7a



- Again a large range of p-values.
- Exclude region at high EFT scale and high DM mass.
- Gives more varied exclusion than M_{jj} or $\Delta\eta$, but still in same area.

Observable phase space plots : Jet 1 pT : D7a



- Again a large range of p-values.
- Exclude region at high EFT scale and high DM mass.
- Very similar exclusion to Mjj.