



# The arches and spandrels of maize domestication, adaptation, and improvement

María Jazmín Abraham-Juárez<sup>1,a</sup>, Allison C. Barnes<sup>2,a</sup>,  
 Alejandro Aragón-Raygoza<sup>2,3,a</sup>, Destiny Tyson<sup>2,4</sup>, Andi Kur<sup>2</sup>,  
 Josh Strable<sup>2</sup> and Rubén Rellán-Álvarez<sup>2</sup>

## Abstract

People living in the Balsas River basin in southwest México domesticated maize from the bushy grass teosinte. Nine thousand years later, in 2021, Ms. Deb Haaland — a member of the Pueblo of Laguna tribe of New Mexico — wore a dress adorned with a cornstalk when she was sworn in as the Secretary of Interior of the United States of America. This choice of garment highlights the importance of the coevolution of maize and the farmers who, through careful selection over thousands of years, domesticated maize and adapted the physiology and shoot architecture of maize to fit local environments and growth habits. Some traits such as tillering were directly selected on (arches), and others such as tassel size are the by-products (spandrels) of maize evolution. Here, we review current knowledge of the underlying cellular, developmental, physiological, and metabolic processes that were selected by farmers and breeders, which have positioned maize as a top global staple crop.

## Addresses

<sup>1</sup> Laboratorio Nacional de Genómica para la Biodiversidad (LANGEBIO), Unidad de Genómica Avanzada, Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional (CINVESTAV), Irapuato, 36821, México

<sup>2</sup> Department of Molecular and Structural Biochemistry, North Carolina State University, Raleigh, NC, 27695, USA

<sup>3</sup> Unidad de Genómica Avanzada, Cinvestav Sede Irapuato, Km. 9.6 Libramiento Norte Carretera Irapuato-León, Guanajuato, México

<sup>4</sup> Department of Crop and Soil Sciences, North Carolina State University, Raleigh, NC, 27695, USA

Corresponding authors: Rellán-Álvarez, Rubén ([rrellan@ncsu.edu](mailto:rrellan@ncsu.edu)); Strable, Josh ([jjstrabl@ncsu.edu](mailto:jjstrabl@ncsu.edu))

<sup>a</sup> These authors contributed equally.

**Current Opinion in Plant Biology** 2021, **64**:102124

This review comes from a themed issue on **Cell Biology**

Edited by **Siobhan Braybrook** and **David Mendoza-Cózatl**

For a complete overview see the [Issue](#) and the [Editorial](#)

Available online xxx

<https://doi.org/10.1016/j.pbi.2021.102124>

1369-5266/© 2021 Elsevier Ltd. All rights reserved.


## Keywords

Maize adaptation, Maize domestication, Maize improvement.

The domestication of maize from teosinte ([Figure 1](#)) [1,2], as well as its expansion to new environments, local adaptation [3–5], and genetic improvement [6,7], reduced overall genetic diversity through selection and bottlenecks. Teosinte is a key resource for identifying loci that regulate architectural traits that have been lost during domestication, because maize shoot morphology has a considerable impact on yield-related traits, such as plant density and inflorescence and kernel production [8–10]. Recently, several quantitative trait loci (QTL) and candidate alleles related to the evolution of maize shoot architecture were mapped using maize-teosinte recombinant inbred line populations [11–17]. Using this approach, Tian et al. identified *UPA1* (*Upright Plant Architecture 1*) and *UPA2*, two major QTL for upright plant architecture. *UPA2* is a *cis*-regulatory variant, whereas *UPA1* encodes a biosynthesis enzyme for the phytohormone brassinosteroid [16]. Introgression of the teosinte allele of *UPA2*, which promotes more upright blades, into maize hybrid lines permitted denser planting and enhanced grain yield [16]. Although this study demonstrates the importance of wild relative alleles on shoot architecture and yield, the average selection intensity for reproductive traits was found to be double that for vegetative traits, highlighting the importance of altering ear morphology [17].

Reduced axillary branching is a common trait in many domesticated crop plants compared with their wild ancestors [18], and modifications of these developmental patterns are fine-tuned at the regulatory level by dynamic interactions between proteins, transcription factors, and noncoding elements [19]. In maize, increased apical dominance was key for domestication, and it was achieved by the gain of function of the transcription factor *tb1* (*teosinte branched1*). *Tb1* has been proposed as a crucial negative regulator of cell growth that modulates several domestication genes, including *gt1* (*grassy tillers1*), *tru1* (*tassels replace uppers ear1*), and *tga1* (*teosinte glume architecture1*) [18,20]. Indeed, some groups of genes related to crop domestication and improvement have been reported, with maize TCP, bHLH, and MADS families highly represented [21]. As per *tb1* function, mutant *tb1*, *gt1*, and *tru1* plants overproduce tillers and aerial branches owing to altered bud dormancy [22–24].

Figure 1



**Axes of maize selection.** Three axes on which human selection has acted on during maize domestication, adaptation, and improvement and some of the traits that have been the results of this selection. The Y-axis represents the environmental and geographic changes that maize had to adapt after domestication in current Mexico. Coloring in the map of America indicates day length. The X-axis indicates how selection by farmers and breeders has acted to change whole plant architecture such as tillering, ear height, tassel size, and so on. The z-axis shows how changes at the metabolic, physiological, genomic, and cellular levels underlie whole plant architecture changes. References indicate some of the key studies and genes involved in processes in each of the three axes.

Furthermore, TB1 targets phytohormones and sugars by positively regulating abscisic acid and jasmonic acid and negatively regulating gibberellin and the sucrose transporter *SWEET15b* [18].

RNA-seq and ChIP-seq experiments have contributed to a better understanding of the regulatory network controlled by TB1 and GT1, explaining how allelic variation in this transcriptional hub produced drastic architectural changes that were agronomically beneficial. Among TB1 targets were genes related to light perception and response to red and far-red light [18]. This observation fits the previously described function of *tb1*, where *PHYTOCHROME B* perceives shading as a low red/far-red light ratio and initiates a signaling cascade promoting *TB1* and *GT1* suppression of lateral bud outgrowth in the shade [18,23]. In domesticated maize, the *TB1-GT1* module conditions constitutive repression on axillary bud growth, which is perceived as insensitivity to shade avoidance response, a distinctive trait compared with teosinte [23,25]. Selection for bud

growth repression together with selection for upright leaf angles has contributed to the increase of plant densities in modern breeding [6], a fundamental factor of increased yields per surface area [26].

Plant architecture depends on the activity of shoot apical meristems (SAMs) (Figure 1) [27]. The CLV-WUS (CLAVATA-WUSCHEL) module is a key feedback pathway that regulates communication between cells within the SAM in different plant species [28]; in maize, CLV orthologs, FEA (FASCIATED EAR) receptor-like proteins, and WUS proteins have been studied primarily in the inflorescences [29,30]. In this same pathway, ZmCRN (CORYNE), a signaling protein, functions downstream of FEA2/CLV2, which transmits signals from CLE (CLV3/ESR-RELATED) peptides through interactions with CT2 (COMPACT PLANT2) and ZmCRN [31]. Interestingly, other FEA proteins have been identified functioning in parallel pathways to CLV-WUS, for example, FEA4 promotes differentiation in the SAM periphery in opposition to

KN1 (KNOTTED1) and WUS [32]. These pathways illustrate how cell–cell communication through signal transduction pathways fine-tunes inflorescence meristem activity, a target in maize domestication in search of larger ear meristems. Recently, it was reported that use of CRISPR-Cas9 genome editing to make weak promoter alleles of maize *CLE* genes increased multiple grain yield-related traits, two *ZmCLE7* fragment promoter deletion alleles showed a significant increase in most yield-related traits including weight and grain yield per ear, and some others showed compensation and redundancy of *ZmCLE7* null alleles [33]. Showing that in the genome editing era, rapid domestication can be achieved more efficiently. Furthermore, recreation of ancestral proteins in maize into novel versions that harbor small changes in sequences, either by genome editing or transgenesis, highlights the potential for inducing gradual evolution in plant morphology [34,35].

Advances in plant single-cell and single nucleus ‘omics’ are forging new opportunities to overcome organ and tissue heterogeneity [36]. Single-cell transcriptomic studies in maize highlight previously unappreciated developmental and physiological changes in germinal cells [37]. Vegetative and reproductive meristems historically have been recalcitrant to isolating viable protoplasts (plant cells without their cell wall). Recent reports in the maize shoot apex and developing ear have cleared technical hurdles for obtaining thousands of protoplasts from shoot meristems for single-cell transcriptomics [38,39]. In more differentiated maize leaf tissues, scRNA: single cell RNA sequencing (scRNA-seq) uncovered novel transcript accumulation of *SWEET13* paralogs in abaxial bundle sheath cells that surround rank-2 veins [40]. Furthermore, individualizing nuclei and assaying accessible chromatin by sequencing identified cell-specific lineages by enrichment of accessible chromatin regions (ACRs), which often have transcription factor DNA-binding motifs in proximity to genes with cell-specific expression [41]. Moreover, physical interactions between distal ACRs and genes, especially for agronomically important loci, that were found previously in whole tissues [42] were also detected in single nuclei [41]. ACRs are highly conserved across plant species [43], and single-cell/nucleus applications have the potential to revolutionize comparative studies. For example, by deciphering cell-type-specific transcriptomes or chromatin status between teosinte and domesticated maize and/or by incorporating plants grown in modeled domestication conditions [44], such strategies could lead to understanding how environmental changes affected gene expression in certain cell lineages during domestication.

Humans have selected secondary metabolites that impact color, sweetness, and other kernel traits that are related to flavor and general organoleptic properties

[45]. Indeed, genes such as *ZmSWEET4c* involved in sugar transport into the seed show signals of selection in domesticated maize (and rice) when compared with teosinte [46]. Vallebueno et al. [47] suggest that the timing of selection on *TB1* may point out that domestication could first be targeted to increase stalk biomass for direct consumption. Later on, sweet corn was developed by indigenous people, with mutant *SU1* alleles having been fixed and cultivated three separate times in North America [48]. Selection for a reduction in grain bitterness may have led to a reduction of alkaloids in maize when compared with teosinte [49]. Many of the metabolic changes taking place during domestication were unintended, so to compare them is informative in assessing the nutritional composition and differences between crops and their progenitors. Fang et al. [50] found that there is similar genetic architecture in maize and teosinte oil and carotenoid variation and that maize traits underwent a strong and recent selection during and after domestication. Carotenoids in particular are an area where work is being performed to harness natural variation to increase levels and regain nutritional content for locations where maize is a nutritional staple [51]. A recent study compared the metabolites of a BC2S7 population created by crossing a highland teosinte (*Zea mays* ssp. *Mexicana*) with Mo17, a modern maize inbred line, and showed that long-term domestication and breeding reshaped amino acid metabolism, likely to meet demands in high-yielding modern varieties [52]. The origin of many of the most favorable alleles resulting in altered sugars, amino acids, or tricarboxylic acid cycle intermediates came from teosinte [52]. Other favorable metabolic traits introduced from highland teosinte may have helped maize adapt to higher altitudes and lower temperatures. A teosinte introgression into some highland maize landraces carries a favorable allele that alters membrane lipid quantities by increasing the amount of phosphatidylcholine [53]. Several other genes in the phospholipid pathways of maize have shown accelerated evolution to cold, as well [54].

Humans did not just adapt teosinte into modern maize, but they have also adapted and relocated tropical maize into more temperate climates that required additional metabolic changes. Deng et al. [55] found that 39.8% of metabolites analyzed were significantly altered between tropical and temperate maize kernels. The genetic architecture of genes involved in metabolomic differences between temperate maize and tropical maize is simpler (fewer changed genes and larger effect sizes) than maize and its wild relatives, probably reflecting the shorter evolutionary divergence between temperate and tropical maize. Alkaloids, terpenoids, and lipids were targeted when tropical and temperate maize diverged. For example, 2,4-dihydroxy-7-methoxy-1,4-benzoxazin-3-one (DIMBOA) compound concentrations are higher in temperate maize than tropical maize and at least one

such compound, 2,4-dihydroxy-7-methoxy-1,4-benzoxazin-3-one glucoside (DIMBOA-Glc), may be important for local adaptation of temperate maize to resist new insect pressure [49].

Photoperiod adaptation was a key component that enabled maize to expand across the Americas after domestication [4,56] (Fig. 1). This was particularly crucial as it moved northward to the United States where day lengths are longer during the growing season, delaying flowering time in unadapted tropical maize. Photoperiod adaptation was not an issue as maize moved southward, and this may have facilitated the rapid expansion of maize to Central and South America after domestication [57,58]. The nested association panel was used to identify the major QTL involved in photoperiod sensitivity [59]. Major genes that increase photoperiod sensitivity include *ZmCCT* (*CO-like/COL and TOC1*) 9 [60], *ZmCCT10* [61], and *ZmRap2.7* [62]. On the other hand, *ZCN8* (*CENTRORADIALIS8*), the most commonly known/studied maize florigen gene, decreases photoperiod sensitivity [63]. Adaptation to temperate latitudes resulted in additive selection of genetic variants that increase the activity of *ZCN8* [63] and decrease the activity of *ZmCCT9* [60] and *ZmCCT10* [64] leading to reduced photoperiod sensitivity. Artificial evolution experiments that selected for shorter flowering of tropical maize populations recapitulated the evolutionary process of tropical maize adaptation and identified the same major key regulators of photoperiod sensitivity [65]. Other genes, such as *zagl1* (*zea agamous-like1*), that are not necessarily involved in photoperiod sensitivity but can reduce flowering time, were also under strong selection during maize domestication [66]. Genes involved in maize flowering time can also exhibit pleiotropic effects, as observed with *zagl1*'s effect on increased ear size [66] and the effect of a *ZmCCT* gene on resistance to *Gibberella* stalk rot [67].

Before maize was brought to northern latitudes, it first was adapted to the highlands of México, Central America, and South América. In highland environments, low temperatures that lead to slow heat unit accumulation have imposed a strong selection for early flowering genetic variants and/or against late-flowering variants. Genomic scans of highland adapted materials have identified that flowering time genes are under selection in highland maize [68,69]. Highland teosinte (*Z. mays* ssp. *Mexicana*) is a source of early flowering alleles in highland maize [53]. Introgressed early flowering alleles have been conserved in temperate maize [53,63], pointing to a relevant role of highland teosinte *mexicana* introgression [70] in maize adaptation to environments where the ability to photosynthesize in low-temperature conditions [71] and flower early is advantageous. Teosinte *mexicana* is found in the Mesomerica highlands but not in South America.

However, evidence shows significant gene flow between highland populations of Mesomerica and South America [69], so the role of teosinte *mexicana* derived alleles in South America highland maize adaptation cannot be ruled out. Strong selection for early flowering affects several developmental and genomic traits that enable faster development, growth, and flowering, which contributed to local adaptation. SAM size and flowering time are inversely correlated such that maize with large meristems flowers earlier than maize with small meristems [72,73]. Larger cells and faster leaf cell production rates are also correlated with smaller genome size in maize and teosintes, and in turn, maize and teosintes with smaller genome sizes are more prevalent at high elevations [74,75]. In theory, smaller genome sizes should enable faster cell replication rates and facilitate faster development; indeed, artificial selection experiments have shown that selection for early flowering does indeed reduce genome size [76].

## Perspective and outlook

Comparative morphology and genomics will continue to be foundational to future studies on maize domestication and adaptation. The current genomics era is providing a more complete understanding of maize evolution as more high-quality genome assemblies for maize inbred lines [77], landraces [78], and wild relatives [70] become available, in addition to the construction of a pangenome graphical representation of haplotypes [79]. This collective wealth of genomic information, together with the development of new gene-editing technologies in an ever-growing number of genotypes [80,81], is enabling CRISPR/Cas9 high-throughput mutagenesis [82], trait stacking in precise genome locations [83], and even targeted chromosomal inversions [84]. Technologies that leverage this wealth of genomics, such as single-cell/single-nucleus 'omics' [37,38,40,41,85], assays that infer genome-wide DNA binding landscapes and associations between transposons, regulatory regions, and target genes [86–90] and evolutionary-guided machine learning methods to predict levels of gene products [91] will continue to foster and inform innovative strategies to study and modify maize genetic variation, conceivably at precise cell-type or pathway resolution.

The development of more diverse mapping populations with open-pollinated maize landraces [53,65] and wild relatives [92–94] together with genomic scans of selection has proven to be a powerful tool to uncover loci contributing to domestication and local adaptation, and the important roles that teosinte *parviglumis* [16] and the highland teosinte *mexicana* have in modern maize [53,63]. The ability to identify domestication- and adaptive-QTL/QTN and QTL by environment interactions will gain further momentum as more maize-teosinte mapping populations are generated and large-scale common garden experiments [68,95] with an ever-growing number of genotypes are implemented.



Finally, we hope that our highlighting the decades of attention to understanding the genetic basis of maize domestication and adaptation is accompanied by an appreciation (Figure 1) of the fundamental roles that smallholder farmers and campesinos have played — and continue to play — in the development and maintenance of maize diversity and, ultimately, in ensuring present and future food security [96].

## Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

## Acknowledgements

The authors are grateful to the editors for the opportunity to contribute to this review and the two reviewers who have provided valuable suggestions. To those colleagues whose work was not referenced in the review, we apologize for the omission owing to strict space limitations. Our work is supported by NSF-PGRP PRFB grant 2010703 to A.C.B., North Carolina State University startup funds and USDA Hatch projects 1026392 and 1020249 to J.S. and R.R-A respectively.

## References

Papers of particular interest, published within the period of review, have been highlighted as:

- \* of special interest
- \*\* of outstanding interest

1. Wright SI, Bi IV, Schroeder SG, Yamasaki M, Doebley JF, McMullen MD, Gaut BS: **The effects of artificial selection on the maize genome.** *Science* 2005, **308**:1310–1314.
2. Hufford MB, Xu X, van Heerwaarden J, Pyhäjärvi T, Chia J-M, Cartwright RA, Elshire RJ, Glaubitz JC, Guill KE, Kaeppler SM, *et al.*: **Comparative population genomics of maize domestication and improvement.** *Nat Genet* 2012, **44**:808–811.
3. Liu H, Wang X, Warburton ML, Wen W, Jin M, Deng M, Liu J, Tong H, Pan Q, Yang X, *et al.*: **Genomic, transcriptomic, and phenomic variation reveals the complex adaptation of modern maize breeding.** *Mol Plant* 2015, **8**:871–884.
4. Swarts K, Gutaker RM, Benz B, Blake M, Bukowski R, Holland J, Kruse-Peebles M, Lepak N, Prim L, Romay MC, *et al.*: **Genomic estimation of complex traits reveals ancient maize adaptation to temperate North America.** *Science* 2017, **357**:512–515.
5. Wang L, Beissinger TM, Loran A, Ross-Ibarra C, Ross-Ibarra J, Hufford MB: **The interplay of demography and selection during maize domestication and expansion.** *Genome Biol* 2017, **18**:215.
6. Wang B, Lin Z, Li X, Zhao Y, Zhao B, Wu G, Ma X, Wang H, Xie Y, Li Q, *et al.*: **Genome-wide selection and genetic improvement during modern maize breeding.** *Nat Genet* 2020, **52**:565–571.
- Genomic and phenotypic analysis of a chronological sampling of Chinese and USA inbreds revealed convergent phenotypic and genomic selection.
7. Jiao Y, Zhao H, Ren L, Song W, Zeng B, Guo J, Wang B, Liu Z, Chen J, Li W, *et al.*: **Genome-wide genetic changes during modern breeding of maize.** *Nat Genet* 2012, **44**:812–815.
8. Dong Z, Alexander M, Chuck G: **Understanding grass domestication through maize mutants.** *Trends Genet* 2019, **35**:118–128.
9. Strable J: **Developmental genetics of maize vegetative shoot architecture.** *Mol Breed* 2021, **41**:19.
10. Chen Z, Gallavotti A: **Improving architectural traits of maize inflorescences.** *Mol Breed* 2021, **41**:21.
11. Wills DM, Whipple CJ, Takuno S, Kursel LE, Shannon LM, Ross-Ibarra J, Doebley JF: **From many, one: genetic control of prolificacy during maize domestication.** *PLoS Genet* 2013, **9**:e1003604.
12. Li D, Wang X, Zhang X, Chen Q, Xu G, Xu D, Wang C, Liang Y, Wu L, Huang C, *et al.*: **The genetic architecture of leaf number and its genetic relationship to flowering time in maize.** *New Phytol* 2016, **210**:256–268.
13. Calderón CI, Yandell BS, Doebley JF: **Fine mapping of a QTL associated with kernel row number on chromosome 1 of maize.** *PLoS One* 2016, **11**:e0150276.
14. Xu G, Wang X, Huang C, Xu D, Li D, Tian J, Chen Q, Wang C, Liang Y, Wu Y, *et al.*: **Complex genetic architecture underlies maize tassel domestication.** *New Phytol* 2017, **214**:852–864.
15. Liu L, Huang J, He L, Liu N, Du Y, Hou R, Du H, Qiu F, Zhang Z: **Dissecting the genetic architecture of important traits that enhance wild germplasm resource usage in modern maize breeding.** *Mol Breed* 2019, **39**:157.
16. Tian J, Wang C, Xia J, Wu L, Xu G, Wu W, Li D, Qin W, Han X, Chen Q, *et al.*: **Teosinte ligule allele narrows plant architecture and enhances high-density maize yields.** *Science* 2019, **365**:658–664.
- Using a maize-teosinte RIL population, the authors identify QTL associations for upright plant architecture. Cloning the QTL uncovered a *cis*-regulatory variant that contributes to brassinosteroid levels and blade angle. Introgression of the teosinte allele for upright blade angle enhanced grain yield in high-density maize hybrids.
17. Yang CJ, Samayoa LF, Bradbury PJ, Olukolu BA, Xue W, York AM, Tuholski MR, Wang W, Daskalska LL, Neumeyer MA, *et al.*: **The genetic architecture of teosinte catalyzed and constrained maize domestication.** *Proc Natl Acad Sci U S A* 2019, **116**:5643–5652.
18. Dong Z, Xiao Y, Govindarajulu R, Feil R, Siddoway ML, Nielsen T, Lunn JE, Hawkins J, Whipple C, Chuck G: **The regulatory landscape of a core maize domestication module controlling bud dormancy and growth repression.** *Nat Commun* 2019, **10**:1–15.
- The genetic pathways regulated by TB1 were identified through ChIP seq, RNAseq and hormone and sugar measurements in maize axillary bud tissue. Demonstrating the key role of TB1 on top of domestication hierarchy.
19. Parvathaneni RK, Bertolini E, Shamimuzzaman M, Vera DL, Lung P-Y, Rice BR, Zhang J, Brown PJ, Lipka AE, Bass HW, *et al.*: **The regulatory landscape of early maize inflorescence development.** *Genome Biol* 2020, **21**:165.
20. Doebley J: **The genetics of maize evolution.** *Annu Rev Genet* 2004, **38**:37–59.
21. Doebley JF, Gaut BS, Smith BD: **The molecular genetics of crop domestication.** *Cell* 2006, **127**:1309–1321.
22. Doebley J, Stec A, Hubbard L: **The evolution of apical dominance in maize.** *Nature* 1997, **386**:485–488.
23. Whipple CJ, Kebrom TH, Weber AL, Yang F, Hall D, Meeley R, Schmidt R, Doebley J, Brutnell TP, Jackson DP: **Grassy tillers1 promotes apical dominance in maize and responds to shade signals in the grasses.** *Proc Natl Acad Sci U S A* 2011, **108**:E506–E512.
24. Dong Z, Li W, Unger-Wallace E, Yang J, Vollbrecht E, Chuck G: **Ideal crop plant architecture is mediated by tassels replace upper ears1, a BTB/POZ ankyrin repeat gene directly targeted by TEOSINTE BRANCHED1.** *Proc Natl Acad Sci U S A* 2017, **114**:E8656–E8664.
25. Wang H, Wu G, Zhao B, Wang B, Lang Z, Zhang C, Wang H: **Regulatory modules controlling early shade avoidance response in maize seedlings.** *BMC Genom* 2016, **17**:269.
26. Mansfield BD, Mumm RH: **Survey of plant density tolerance in U.S. maize germplasm.** *Crop Sci* 2014, **54**:157–173.
27. Wu Q, Xu F, Jackson D: **All together now, a magical mystery tour of the maize shoot meristem.** *Curr Opin Plant Biol* 2018, **45**:26–35.

28. Somssich M, Je BI, Simon R, Jackson D: **CLAVATA-WUSCHEL signaling in the shoot meristem.** *Development* 2016, **143**: 3238–3248.
  29. Bommert P, Je BI, Goldshmidt A, Jackson D: **The maize G $\alpha$  gene COMPACT PLANT2 functions in CLAVATA signalling to control shoot meristem size.** *Nature* 2013, **502**:555–558.
  30. Je BI, Gruel J, Lee YK, Bommert P, Arevalo ED, Eveland AL, Wu Q, Goldshmidt A, Meeley R, Bartlett M, *et al.*: **Signaling from maize organ primordia via FASCIATED EAR3 regulates stem cell proliferation and yield traits.** *Nat Genet* 2016, **48**:785–791.
  31. Je BI, Xu F, Wu Q, Liu L, Meeley R, Gallagher JP, Corcilius L, Payne RJ, Bartlett ME, Jackson D: **The CLAVATA receptor FASCIATED EAR4 responds to distinct CLE peptides by signaling through two downstream effectors.** *Elife* 2018, **7**.
  32. Pautler M, Eveland AL, LaRue T, Yang F, Weeks R, Lunde C, Je BI, Meeley R, Komatsu M, Vollbrecht E, *et al.*: **FASCIATED EAR4 encodes a bZIP transcription factor that regulates shoot meristem size in maize.** *Plant Cell* 2015, **27**:104–120.
  33. Liu L, Gallagher J, Arevalo ED, Chen R, Skopelitis T, Wu Q, Bartlett M, Jackson D: **Enhancing grain-yield-related traits by CRISPR-Cas9 promoter editing of maize CLE genes.** *Nat Plants* 2021, **7**:287–294.
  34. Bartlett M, Thompson B, Brabazon H, Del Gizzi R, Zhang T, Whipple C: **Evolutionary dynamics of floral homeotic transcription factor protein–protein interactions.** *Mol Biol Evol* 2016, **33**:1486–1501.
  35. Abraham-Juárez MJ, Schrager-Lavelle A, Man J, Whipple C, Handakumbura P, Babbitt C, Bartlett M: **Evolutionary variation in MADS box dimerization affects floral development and protein abundance in maize.** *Plant Cell* 2020, **32**:3408–3424.
- Using the maize B-class MADS-box protein STS1 and one of its evolutionary variants, the authors evaluated experimentally the deep conservation of the floral quartet model and the consequences of change STS1 dimerization, at transcriptional and protein interaction level. Highlighting the importance of gradual evolutionary changes.
36. Seyferth C, Renema J, Wendrich JR, Eekhout T, Seurinck R, Vandamme N, Blob B, Saeys Y, Helariutta Y, Birnbaum KD, *et al.*: **Advances and opportunities of single-cell transcriptomics for plant research.** *Annu Rev Plant Biol* 2021, **72**:847–866, <https://doi.org/10.1146/annurev-arplant-081720-010120>.
  37. Nelms B, Walbot V: **Defining the developmental program leading to meiosis in maize.** *Science* 2019, **364**:52–56.
  38. Satterlee JW, Strable J, Scanlon MJ: **Plant stem-cell organization and differentiation at single-cell resolution.** *Proc Natl Acad Sci U S A* 2020, **117**:33689–33699.
  39. Xu X, Crow M, Rice BR, Li F, Harris B, Liu L, Demesa-Arevalo E, Lu Z, Wang L, Fox N, *et al.*: **Single-cell RNA sequencing of developing maize ears facilitates functional analysis and trait candidate gene discovery.** *Dev Cell* 2021, **56**:557–568.e6.
  40. Bezruczyk M, Zöllner NR, Kruse CPS, Hartwig T, Lautwein T, Köhrer K, Frommer WB, Kim J-Y: **Evidence for Phloem loading via the abaxial bundle sheath cells in maize leaves.** *Plant Cell* 2021, <https://doi.org/10.1093/plcell/koaa055>.
- Generated the first single-cell transcriptome from differentiated leaf tissue and were able to identify an unknown developmental process for sugar loading in maize leaves where the bundle sheath is involved.
41. Marand AP, Chen Z, Gallavotti A, Schmitz RJ: **A cis-regulatory atlas in maize at single-cell resolution.** *Cell* 2021 May 27, **184**(11):3041–3055.e21, <https://doi.org/10.1016/j.cell.2021.04.014>. Epub 2021 May 7.
  42. Ricci WA, Lu Z, Ji L, Marand AP, Ethridge CL, Murphy NG, Noshay JM, Galli M, Mejía-Guerra MK, Colomé-Tatché M, *et al.*: **Widespread long-range cis-regulatory elements in the maize genome.** *Nat Plants* 2019, **5**:1237–1249.
  43. Lu Z, Marand AP, Ricci WA, Ethridge CL, Zhang X, Schmitz RJ: **The prevalence, evolution and chromatin signatures of plant regulatory elements.** *Nat Plants* 2019, **5**:1250–1259.
  44. Lorant A, Pedersen S, Holst I, Hufford MB, Winter K, Piperno D, Ross-Ibarra J: **The potential role of genetic assimilation during maize domestication.** *PLoS One* 2017, **12**, e0184202.
  45. Alseekh S, Scossa F, Wen W, Luo J, Yan J, Beleggia R, Klee HJ, Huang S, Papa R, Fernie AR: **Domestication of crop metabolomes: desired and unintended consequences.** *Trends Plant Sci* 2021, <https://doi.org/10.1016/j.tplants.2021.02.005>.
  46. Sosso D, Luo D, Li Q-B, Sasse J, Yang J, Gendrot G, Suzuki M, Koch KE, McCarty DR, Chourey PS, *et al.*: **Seed filling in domesticated maize and rice depends on SWEET-mediated hexose transport.** *Nat Genet* 2015.
  47. Vallebuena-Estrada M, Rodríguez-Arévalo I, Rougon-Cardoso A, Martínez González J, García Cook A, Montiel R, Vielle-Calzada J-P: **The earliest maize from San Marcos Tehuacán is a partial domesticate with genomic evidence of inbreeding.** *Proc Natl Acad Sci U S A* 2016, **113**:14151–14156.
  48. Hu Y, Colantonio V, Müller BSF, Leach KA, Nanni A, Finegan C, Wang B, Baseggio M, Newton CJ, Juhl EM, *et al.*: **Genome assembly and population genomic analysis provide insights into the evolution of modern sweet corn.** *Nat Commun* 2021, **12**:1227.
  49. Xu G, Cao J, Wang X, Chen Q, Jin W, Li Z, Tian F: **Evolutionary metabolomics identifies substantial metabolic divergence between maize and its wild ancestor, teosinte.** *Plant Cell* 2019, **31**:1990–2009.
- The authors compared the metabolic changes during domestication from teosinte to maize, and adaptation from tropical maize to temperate maize. Identified several differentially expressed alkaloids, terpenoids, and lipids, and contextualized how their changes helped domesticate or adapt maize.
50. Fang H, Fu X, Wang Y, Xu J, Feng H, Li W, Xu J, Jittham O, Zhang X, Zhang L, *et al.*: **Genetic basis of kernel nutritional traits during maize domestication and improvement.** *Plant J* 2020, **101**:278–292.
  51. Diepenbrock CH, Ilut DC, Magallanes-Lundback M, Kandianis CB, Lipka AE, Bradbury PJ, Holland JB, Hamilton JP, Wooldridge E, Vaillancourt B, *et al.*: **Eleven biosynthetic genes explain the majority of natural variation in carotenoid levels in maize grain.** *Plant Cell* 2020, <https://doi.org/10.1093/plcell/koab032>.
- Measured 7 carotenoid compounds in maize grain, analyzed the underlying genetics, and resolved 11 carotenoid-related QTL to their genes, elucidating the key precursor and biosynthetic pathway genes.
52. Li K, Wen W, Alseekh S, Yang X, Guo H, Li W, Wang L, Pan Q, Zhan W, Liu J, *et al.*: **Large-scale metabolite quantitative trait locus analysis provides new insights for high-quality maize improvement.** *Plant J* 2019, **99**:216–230.
  53. Rodríguez-Zapata F, Barnes AC, Blöcher-Juárez KA, Gates D, Kur A, Wang L, Janzen GM, Jensen S, Estévez-Palmas JM, Crow T, *et al.*: **Teosinte introgression modulates phosphatidylcholine levels and induces early maize flowering time.** *Cold Spring Harbor Laboratory* 2021, <https://doi.org/10.1101/2021.01.25.426574>.
- Using a combination of genomic selection scans, linkage mapping, lipidomics the authors identified HPC1 a teosinte mexicana introgression gene encoding a phospholipase that leads to high levels of phosphatidylcholine and reduces flowering time in highland adapted landraces and Northern US and EU Flint inbreds.
54. Yan L, Kenchanmane Raju SK, Lai X, Zhang Y, Dai X, Rodríguez O, Mahboub S, Roston RL, Schnable JC: **Parallels between natural selection in the cold-adapted crop-wild relative *Tripsacum dactyloides* and artificial selection in temperate adapted maize.** *Plant J* 2019, **99**: 965–977.
  55. Deng M, Zhang X, Luo J, Liu H, Wen W, Luo H, Yan J, Xiao Y: **Metabolomics analysis reveals differences in evolution between maize and rice.** *Plant J* 2020, **103**:1710–1722.
  56. da Fonseca RR, Smith BD, Wales N, Cappellini E, Skoglund P, Fumagalli M, Samaniego JA, Carøe C, Ávila-Arcos MC, Hufnagel DE, *et al.*: **The origin and evolution of maize in the Southwestern United States.** *Nat Plants* 2015, **1**:14003.
  57. Kistler L, Maezumi SY, Gregorio de Souza J, Przelomska NAS, Malaquias Costa F, Smith O, Loiselle H, Ramos-Madrigr J, Wales N, Ribeiro ER, *et al.*: **Multiproxy evidence highlights a complex evolutionary legacy of maize in South America.** *Science* 2018, **362**:1309–1313.

58. Stephen Athens J, Ward JV, Pearsall DM, Chandler-Ezell K, Blinn DW, Morrison AE: **Early prehistoric maize in northern highland Ecuador.** *Lat Am Antiq* 2016, **27**:3–21.
  59. Hung H-Y, Shannon LM, Tian F, Bradbury PJ, Chen C, Flint-Garcia SA, McMullen MD, Ware D, Buckler ES, Doebley JF, *et al.*: **ZmCCT and the genetic basis of day-length adaptation underlying the postdomestication spread of maize.** *Proc Natl Acad Sci Unit States Am* 2012, **109**:E1913–E1921.
- The authors dissect the genetic architecture of photoperiod sensitivity by evaluating the Nested Association Mapping population in neutral and long day environments. They identify key components of the photoperiod response including ZmCCT
60. Huang C, Sun H, Xu D, Chen Q, Liang Y, Wang X, Xu G, Tian J, Wang C, Li D, *et al.*: **ZmCCT9 enhances maize adaptation to higher latitudes.** *Proc Natl Acad Sci U S A* 2018, **115**:E334–E341.
  61. Stephenson E, Estrada S, Meng X, Ourada J, Muszynski MG, Habben JE, Danilevskaya ON: **Over-expression of the photoperiod response regulator ZmCCT10 modifies plant architecture, flowering time and inflorescence morphology in maize.** *PLoS One* 2019, **14**, e0203728.
  62. Liang Y, Liu Q, Wang X, Huang C, Xu G, Hey S, Lin H-Y, Li C, Xu D, Wu L, *et al.*: **ZmMADS69 functions as a flowering activator through the ZmRap2.7-ZCN8 regulatory module and contributes to maize flowering time adaptation.** *New Phytol* 2019, **221**:2335–2347.
  63. Guo L, Wang X, Zhao M, Huang C, Li C, Li D, Yang CJ, York AM, Xue W, Xu G, *et al.*: **Stepwise cis-regulatory changes in ZCN8 contribute to maize flowering-time adaptation.** *Curr Biol* 2018, **28**:3005–3015.e4.
- The authors identified two cis-variants of ZCN8, SNP-1245 and Indel-2339, that arose independently and contributed to maize flowering time adaptation. SNP-1245 was shown to contribute to flowering time adaptation by activating ZCN8 via the ZmMADS1 flowering activator, while Indel-2339 was shown to most-likely play a role in maize's ability to adapt to higher latitudes.
64. Yang Q, Li Z, Li W, Ku L, Wang C, Ye J, Li K, Yang N, Li Y, Zhong T, *et al.*: **CACTA-like transposable element in ZmCCT attenuated photoperiod sensitivity and accelerated the postdomestication spread of maize.** *Proc Natl Acad Sci U S A* 2013, **110**:16969–16974.
  65. Wisser RJ, Fang Z, Holland JB, Teixeira JEC, Dougherty J, Weldekidan T, de Leon N, Flint-Garcia S, Lauter N, Murray SC, *et al.*: **The genomic basis for short-term evolution of environmental adaptation in maize.** *Genetics* 2019, <https://doi.org/10.1534/genetics.119.302780>.
- Using a tropical maize landrace that was adapted to temperate environments over the course of 10 years, the authors were able to explain the mechanisms of adaptive evolution to long days and showed that artificial selection experiments can recapitulate evolutionary history of maize adaptation.
66. Wills DM, Fang Z, York AM, Holland JB, Doebley JF: **Defining the role of the MADS-box gene, *Zea agamos-like1*, a target of selection during maize domestication.** *J Hered* 2018, **109**:333–338.
- The authors suggest that *Zag1*, a transcription factor that acts as a flowering time gene in maize was a target in early maize domestication, resulting in larger ear sizes in maize versus teosinte. *Zag1* was also shown to have a strong association with flowering time and plant height. One of the first cloned genes first identified as genomic selection scan candidate.
67. Wang C, Yang Q, Wang W, Li Y, Guo Y, Zhang D, Ma X, Song W, Zhao J, Xu M: **A transposon-directed epigenetic change in ZmCCT underlies quantitative resistance to Gibberella stalk rot in maize.** *New Phytol* 2017, **215**:1503–1515.
  68. Gates DJ, Runcie D, Janzen GM, Navarro AR: **Single-gene resolution of locally adaptive genetic variation in Mexican maize.** *BioRxiv* 2019.
  69. Wang L, Josephs EB, Lee KM, Roberts LM: **Molecular parallelism underlies convergent highland adaptation of maize landraces.** *bioRxiv* 2020.
  70. Yang N, Xu X-W, Wang R-R, Peng W-L, Cai L, Song J-M, Li W, Luo X, Niu L, Wang Y, *et al.*: **Contributions of *Zea mays* subspecies mexicana haplotypes to modern maize.** *Nat Commun* 2017, **8**:1874.
  71. Crow T, Ta J, Nojoomi S, Aguilar-Rangel MR, Torres Rodríguez JV, Gates D, Rellán-Álvarez R, Sawers R, Runcie D: **Gene regulatory effects of a large chromosomal inversion in highland maize.** *PLoS Genet* 2020, **16**, e1009213.
  72. Thompson AM, Yu J, Timmermans MCP, Schnable P, Crants JC, Scanlon MJ, Muehlbauer GJ: **Diversity of maize shoot apical meristem architecture and its relationship to plant morphology.** *G3* 2015, **5**:819–827.
  73. Leiboff S, Li X, Hu H-C, Todt N, Yang J, Li X, Yu X, Muehlbauer GJ, Timmermans MCP, Yu J, *et al.*: **Genetic control of morphometric diversity in the maize shoot apical meristem.** *Nat Commun* 2015, **6**:8974.
  74. Bilinski P, Albert PS, Berg JJ, Birchler JA, Grote MN, Lorant A, Quezada J, Swarts K, Yang J, Ross-Ibarra J: **Parallel altitudinal clines reveal trends in adaptive evolution of genome size in *Zea mays*.** *PLoS Genet* 2018, **14**, e1007162.
- The authors reveal striking altitudinal dependent gradients of genome size variation in maize and its wild relative and the possible role of genome size variation in local adaptation.
75. Díez CM, Gaut BS, Meca E, Scheinvar E, Montes-Hernandez S, Eguarte LE, Tenaillon MI: **Genome size variation in wild and cultivated maize along altitudinal gradients.** *New Phytol* 2013, **199**:264–276.
  76. Rayburn AL, Dudley JW, Biradar DP: **Selection for early flowering results in simultaneous selection for reduced nuclear DNA content in maize.** *Plant Breed* 1994, **112**:318–322.
  77. Hufford MB, Seetharam AS, Woodhouse MR, Chougule KM, Ou S, Liu J, Ricci WA, Guo T, Olson A, Qiu Y, *et al.*: **De novo assembly, annotation, and comparative analysis of 26 diverse maize genomes.** *bioRxiv* 2021, <https://doi.org/10.1101/2021.01.14.426684>.
  78. Yang N, Liu J, Gao Q, Gui S, Chen L, Yang L, Huang J, Deng T, Luo J, He L, *et al.*: **Genome assembly of a tropical maize inbred line provides insights into structural variation and crop improvement.** *Nat Genet* 2019, **51**:1052–1059.
  79. Valdes Franco JA, Gage JL, Bradbury PJ, Johnson LC, Miller ZR, Buckler ES, Cinta Romay M: **A maize practical haplotype Graph leverages diverse NAM assemblies.** *bioRxiv* 2020, <https://doi.org/10.1101/2020.08.31.268425>.
- The authors leverage whole genome assemblies of the NAM parents to construct a pangenome representation (Practical Haplotype Graph) that captures haplotype diversity and sets the basis for future haplotype imputation of non sequenced varieties using shallow or sparse genotyping.
80. Kelliher T, Starr D, Su X, Tang G, Chen Z, Carter J, Wittich PE, Dong S, Green J, Burch E, *et al.*: **One-step genome editing of elite crop germplasm during haploid induction.** *Nat Biotechnol* 2019, **37**:287–292.
  81. Lowe K, La Rota M, Hoerster G, Hastings C, Wang N, Chamberlin M, Wu E, Jones T, Gordon-Kamm W: **Rapid genotype “independent” *Zea mays* L. (maize) transformation via direct somatic embryogenesis.** *In Vitro Cell Dev Biol Plant* 2018, **54**:240–252.
  82. Liu H-J, Jian L, Xu J, Zhang Q, Zhang M, Jin M, Peng Y, Yan J, Han B, Liu J, *et al.*: **High-throughput CRISPR/Cas9 mutagenesis streamlines trait gene identification in maize.** *Plant Cell* 2020, **32**:1397–1413.
  83. Gao H, Mutti J, Young JK, Yang M, Schroder M, Lenderts B, Wang L, Peterson D, St Clair G, Jones S, *et al.*: **Complex trait loci in maize enabled by CRISPR-Cas9 mediated gene insertion.** *Front Plant Sci* 2020, **11**:535.
  84. Schwartz C, Lenderts B, Feigenbutz L, Barone P, Liaca V, Fengler K, Svitashv S: **CRISPR–Cas9-mediated 75.5-Mb inversion in maize.** *Nature Plants* 2020, **6**:1427–1431.



85. Xu G, Lyu J, Li Q, Liu H, Wang D, Zhang M, Springer NM, Ross-Ibarra J, Yang J: **Evolutionary and functional genomics of DNA methylation in maize domestication and improvement.** *Nat Commun* 2020, **11**:5539.
86. Peng Y, Xiong D, Zhao L, Ouyang W, Wang S, Sun J, Zhang Q, Guan P, Xie L, Li W, *et al.*: **Chromatin interaction maps reveal genetic regulation for quantitative traits in maize.** *Nat Commun* 2019, **10**:2632.
87. Tu X, Mejía-Guerra MK, Valdes Franco JA, Tzeng D, Chu P-Y, Shen W, Wei Y, Dai X, Li P, Buckler ES, *et al.*: **Reconstructing the maize leaf regulatory network using ChIP-seq data of 104 transcription factors.** *Nat Commun* 2020, **11**:5089.
88. Crisp PA, Marand AP, Noshay JM, Zhou P, Lu Z, Schmitz RJ, Springer NM: **Stable unmethylated DNA demarcates expressed genes and their cis-regulatory space in plant genomes.** *Proc Natl Acad Sci U S A* 2020, **117**:23991–24000.
89. Noshay JM, Marand AP, Anderson SN, Zhou P, Mejia Guerra MK, Lu Z, O'Connor CH, Crisp PA, Hirsch CN, Schmitz RJ, *et al.*: **Assessing the regulatory potential of transposable elements using chromatin accessibility profiles of maize transposons.** *Genetics* 2021, **217**:1–13.
90. Sun Y, Dong L, Zhang Y, Lin D, Xu W, Ke C, Han L, Deng L, Li G, Jackson D, *et al.*: **3D genome architecture coordinates trans and cis regulation of differentially expressed ear and tassel genes in maize.** *Genome Biol* 2020, **21**:143.
91. Washburn JD, Mejia-Guerra MK, Ramstein G, Kremling KA, Valluru R, Buckler ES, Wang H: **Evolutionarily informed deep learning methods for predicting relative transcript abundance from DNA sequence.** *Proc Natl Acad Sci U S A* 2019, **116**:5542–5549.
92. Liu Z, Cook J, Melia-Hancock S, Guill K, Bottoms C, Garcia A, Ott O, Nelson R, Recker J, Balint-Kurti P, *et al.*: **Expanding maize genetic resources with predomestication alleles: maize-teosinte introgression populations.** *Plant Genome* 2016, **9**.
93. Chen Q, Yang CJ, York AM, Xue W, Daskalska LL, DeValk CA, Krueger KW, Lawton SB, Spiegelberg BG, Schnell JM, *et al.*: **TeoNAM: a nested association mapping population for domestication and agronomic trait analysis in maize.** *Genetics* 2019, **213**:1065–1078.
94. Chen Q, Samayoa LF, Yang CJ, Bradbury PJ, Olukolu BA, Neumeyer MA, Romay MC, Sun Q, Lorant A, Buckler ES, *et al.*: **The genetic architecture of the maize progenitor, teosinte, and how it was altered during maize domestication.** *PLoS Genet* 2020, **16**, e1008791.
95. Rogers AR, Dunne JC, Romay C, Bohn M, Buckler ES, Ciampitti IA, Edwards J, Ertl D, Flint-Garcia S, Gore MA, *et al.*: **The importance of dominance and genotype-by-environment interactions on grain yield variation in a large-scale public cooperative maize experiment.** *G3* 2021:11.
96. Bellon MR, Mastretta-Yanes A, Ponce-Mendoza A, Ortiz-Santamaría D, Oliveros-Galindo O, Perales H, Acevedo F, Sarukhán J: **Evolutionary and food supply implications of ongoing maize domestication by Mexican campesinos.** *Proc Biol Sci* 2018, **285**.