

Bibliography

- Broman KW** (2015) R/qtlcharts: interactive graphics for quantitative trait locus mapping. *Genetics* **199**: 359–361
- Cruz-Ramirez A, López-Bucio J, Ramírez-Pimentel G, Zurita-Silva A, Sanchez-Calderon L, Ramírez-Chávez E, González-Ortega E, Herrera-Estrella L** (2004) The xip1 mutant of *Arabidopsis* reveals a critical role for phospholipid metabolism in root system development and epidermal cell integrity. *The Plant Cell* **16**: 2020–2034
- Degenkolbe T, Giavalisco P, Zuther E, Seiwert B, Hinch DK, Willmitzer L** (2012) Differential remodeling of the lipidome during cold acclimation in natural accessions of *Arabidopsis thaliana*. *The Plant Journal* **72**: 972–982
- Hufford MB, Lubinsky P, Pyhäjärvi T, Devengenzo MT, Ellstrand NC, Ross-Ibarra J** (2013) The genomic signature of crop-wild introgression in maize. *PLoS Genetics* **9**: e1003477
- Kind T, Liu K-H, Lee DY, Defelice B, Meissen JK, Fiehn O** (2013) LipidBlast in silico tandem mass spectrometry database for lipid identification. *Nature methods* **10**: 755–758
- Krasilnikov P, Carmen Gutiérrez-Castorena M del** (2013) The soils of Mexico.
- Lin Y-C, Liu Y-c, Nakamura Y** (2015) The Choline/Ethanolamine Kinase Family in *Arabidopsis*: Essential Role of CEK4 in Phospholipid Biosynthesis and Embryo Development. *The Plant Cell* **27**: 1497–1511
- Matsuoka Y, Vigouroux Y, Goodman MM, Sanchez G J, Buckler E, Doebley J** (2002) A single domestication for maize shown by multilocus microsatellite genotyping. *Proceedings of the National Academy of Sciences of the United States of America* **99**: 6080–6084
- Matyash V, Liebisch G, Kurzchalia TV, Shevchenko A, Schwudke D** (2008) Lipid extraction by methyl-tert-butyl ether for high-throughput lipidomics. *Journal of lipid research* **49**: 1137–1146
- Nakamura Y, Andrés F, Kanehara K, Liu Y-c, Dörmann P, Coupland G** (2014) *Arabidopsis* florigen FT binds to diurnally oscillating phospholipids that accelerate flowering. *Nature Communications* **5**: 3553
- Okazaki Y, Otsuki H, Narisawa T, Kobayashi M, Sawai S, Kamide Y, Kusano M, Aoki T, Hirai MY, Saito K** (2013) A new class of plant lipid is essential for protection against phosphorus depletion. *Nature Communications* **4**: 1510
- Petersen A-K, Krumsiek J, Wägele B, Theis FJ, Wichmann H-E, Gieger C, Suhre K** (2012) On the hypothesis-free testing of metabolite ratios in genome-wide and metabolome-wide association studies. *BMC Bioinformatics* **13**: 120
- Riedelsheimer C, Lisec J, Czedik-Eysenberg A, Sulpice R, Flis A, Grieder C, Altmann T, Stitt M, Willmitzer L, Melchinger AE** (2012) Genome-wide association mapping of leaf metabolic profiles for dissecting complex traits in maize. *Proceedings of the National Academy of Sciences* **109**: 8872–8877
- Takuno S, Ralph P, Swarts K, Elshire RJ, Glaubitz JC, Buckler ES, Hufford MB, Ross-Ibarra J** (2015) Independent Molecular Basis of Convergent Highland Adaptation in Maize. *Genetics* **195**: 115.178327
- Veneklaas EJ, Lambers H, Bragg J, Finnegan PM, Lovelock CE, Plaxton WC, Price CA, Scheible W-R, Shane MW, White PJ et al** (2012) Opportunities for improving phosphorus-use efficiency in crop plants. *New Phytologist* **195**: 306–320
- Wen W, Li D, Li X, Gao Y, Li W, Li H, Liu J, Liu H, Chen W, Luo J et al** (2014) Metabolome-based genome-wide association study of maize kernel leads to novel biochemical insights.

Nature Communications 5:

Xue H-W, Chen X, Mei Y (2009) Function and regulation of phospholipid signalling in plants. *Biochemical Journal* **421**: 145–156

Yong-Villalobos L, González-Morales SI, Wrobel K, Gutiérrez-Alanis D, Cervantes-Peréz SA, Hayano-Kanashiro C, Oropeza-Aburto A, Cruz-Ramirez A, Martínez O, Herrera-Estrella L (2015) Methylome analysis reveals an important role for epigenetic changes in the regulation of the Arabidopsis response to phosphate starvation. *Proceedings of the National Academy of Sciences* **112**: E7293–302