

Chapter 2: 5.4, 5.5, 6.4

Chapter 3: 1.7, 2.3, 2.4, 2.6, 2.8, 2.11, 2.12, 2.16, 2.20

**Exercise 5.4.** Let  $a$  and  $b$  be real numbers. Show that the following three equalities hold:

$$\bigcap_{x>b} (a, x) = (a, b], \quad \bigcup_{n=1}^{\infty} \left[a + \frac{1}{n}, b - \frac{1}{n}\right] = (a, b), \quad \bigcap_{n=1}^{\infty} (a + n, +\infty) = \emptyset.$$

■

**Exercise 5.5.** Let  $a_1, a_2, \dots$  be any enumeration of the negative rational numbers; let  $b_1, b_2, \dots$  be any enumeration of the positive rational numbers. Show that the following two equalities hold:

$$\bigcap_{j=1}^{\infty} (a_j, b_j) = \{0\}, \quad \bigcup_{j=1}^{\infty} (a_j, b_j) = \mathbb{R},$$

.

■

**Exercise 6.4.** Prove there exists no order  $\leq$  that makes  $(\mathbb{C}, +, \cdot, \leq)$  into an ordered field.

Let us argue by contradiction. Assume that  $\mathbb{C}$  is a ordered field. Let us analyze the case  $(0, 1) = i$ .

- If  $i > 0$ ,  $x, y \in \mathbb{C}$  and  $x = i, y = i, i \cdot i > 0 \implies -1 > 0$ , which is a contradiction.
- If  $i < 0$ ,  $x, y \in \mathbb{C}$  and  $x = -i, y = -i$  (since  $x, y > 0$  for the second condition to hold),  $-i \cdot -i < 0 \implies i^2 > 0 \implies -1 > 0$ , which is a contradiction.

■

**Exercise 1.7.** Let  $\|\cdot\|$  be a norm on a real vector space  $V$ . Prove the *reverse triangle inequality*:

$$|||x| - |y|| \leq \|x - y\|$$

■

**Exercise 2.3.** Let  $X$  be any set. Prove that the discrete metric  $d : X \times X \rightarrow \mathbb{R}$  (defined by  $d(x, y) = 1$  if  $x \neq y$  and  $d(x, x) = 0$  for  $x \in X$ ) satisfies the triangle inequality and is therefore a metric on  $X$ .

**Exercise 2.4.** Determine which of the following functions are metrics on  $\mathbb{R}$ . Prove your answer in each case.

- $d_1(x, y) = \sqrt{|x - y|}$ .
- $d_2(x, y) = |x - 2y|$ .
- $d_3(x, y) = \frac{|x-y|}{1+|x-y|}$ .

**Exercise 2.6.** Let  $\|\cdot\|$  denote the Euclidean norm  $\mathbb{R}^2$ , i.e.  $\|(x_1, x_2)\| = \sqrt{x_1^2 + x_2^2}$ . Consider the function from  $\mathbb{R}^2 \times \mathbb{R}^2 \rightarrow \mathbb{R}$ , defined by

$$d(x, y) = \|x_1 - y_1\| + \|x_2 - y_2\|, \quad (x = (x_1, x_2), y = (y_1, y_2)).$$

1. Prove that  $d$  is a metric on  $\mathbb{R}^2$ .
2. On a sheet of graph paper, draw the set  $B_d((5, 1), 3)$ . Use dotted lines to indicate the boundary, which is not included in the set you are drawing. (Hint: it may be easier to figure out what the set looks like if you first consider  $B_d((0, 0), 3)$ ).
3. On the same graph as in the previous part, draw  $B_{d_u}((-3, 2), 1)$ , where  $d_u$  denotes the square metric.

**Exercise 2.8.** Let  $(X, d)$  be a metric space, and let  $E$  be a subset of  $X$ . The *diameter* of  $E$  in  $(X, d)$  is defined by the formula

$$\text{diam}_d(E) = \sup\{d(x, y) : x, y \in E\}.$$

1. Prove that for any  $r > 0$  and  $x \in X$ , we have  $\text{diam}(B(x, r)) \leq 2r$ .
2. If  $X$  is any set and  $d$  is the discrete metric, show  $\text{diam}(B(x, r)) = 0$ .
3. If  $X = \mathbb{R}^n$  for some  $n \in \mathbb{N}$  and  $d$  is the Euclidean metric, prove that  $\text{diam}(B(x, r)) = 2r$ .

**Exercise 2.11.** As in Example 2.7, let  $X = \mathbb{R}^2$ ,  $Y = [-1, 3] \times [2, 4]$ , and let  $d$  denote the Euclidean metric on  $X = \mathbb{R}^2$ . Let  $p = (3, 4)$  and let  $q = (2, 4)$ . Arguing *directly from the definition of an interior point* (i.e., without using Exercise 2.12), show that  $q$  is an interior point  $B_Y(p, 2)$  with respect to  $Y$ , but  $q$  is not an interior point  $B_Y(p, 2)$  with respect to  $X$ . In addition, draw a picture on a piece of graph paper that illustrates the idea of your proof.

■

**Exercise 2.12.** Let  $(X, d)$  be a metric space, and let  $Y$  be a subset of  $X$ . Prove that

$$(*) \operatorname{Int}_X(U) = \operatorname{Int}_Y(U) \cap \operatorname{Int}_X(Y).$$

In the notation of Exercise 2.11, the equality  $(*)$  gives an alternate explanation of why  $q$  is not an interior point of  $B_Y(p, 2)$  with respect to  $X$ : It is because  $q \notin \operatorname{Int}_X(Y)$ , as can be seen from the picture you drew in that Exercise.

■

**Exercise 2.16.** Let  $(X, d)$  be a metric space, and let  $U$  be a subset of  $X$ . Use Proposition 2.15 to prove that  $\operatorname{Int}_X(U)$  is open in  $X$ .

■

**Exercise 2.20.** Let  $(X, d)$  be a metric space. Assume that  $U \subset Y \subset X$ , and additionally that  $Y$  is open in  $X$ . Prove that  $U$  is open in  $Y$  if and only if  $U$  is open in  $X$ . (Note: There are at least two possible solutions; one uses Theorem 2.19, the other uses Exercise 2.12.)

■