

Numerical schemes for Classical Chemotaxis Equations PC-239



Daniel Acosta Soba¹, Alba M. Navarro Izquierdo², J. Rafael Rodríguez Galván³

Departamento de Matemáticas, Universidad de Cádiz

Introduction

Chemotaxis (movement of biological cells in response to chemical signals) was modeled by Keller-Segel in 1970. Although there are several models, we focus on the classical one, given by the following equations in $\Omega \subset \mathbb{R}^n$:

$$\begin{cases} u_t = \alpha_0 \Delta u - \alpha_1 \nabla \cdot (u \nabla v), & x \in \Omega, t > 0, & (1a) \\ v_t = \alpha_2 \Delta v - \alpha_3 v + \alpha_4 u, & x \in \Omega, t > 0, & (1b) \\ \nabla u \cdot \mathbf{n} = \nabla v \cdot \mathbf{n} = 0, & x \in \partial\Omega, t > 0, & (1c) \\ u(x, 0) = u_0(x), v(x, 0) = v_0(x), & x \in \Omega, & (1d) \end{cases}$$

where u and v represent density of **cells** and **chemical-signal**, respectively.

From an analytical point of view, a lot of research has been recently done (see e.g. [2] and references therein) and interesting results about global in time existence, mass conservation, energy, blow-up and positivity of solution have been published. However, there is not a large literature on *numerical analysis* for (1), and reproducing former properties is an interesting challenge. This work is mainly focused on development of *positivity preserving numerical schemes*, related to discontinuous *Galerkin methods*, which *decouple* calculus of u and v .

Energy-Stable Semi-Discretization in Time

Given a partition of time interval $(0, T)$ into subintervals of size $k > 0$, we approximate u and v at each time step t^{m+1} by an implicit Euler scheme^a as follows:

$$\begin{cases} \delta_t u^{m+1} - \nabla u^{m+1} + \alpha_1 \nabla \cdot (u^{m+1} \nabla v^{m+1}) = 0, & (2a) \\ \delta_t v^{m+1} - \alpha_2 \nabla v^{m+1} + \alpha_3 v^{m+1} - \alpha_4 u^m = 0, & (2b) \end{cases}$$

where δ_t is the backward difference operator. For this scheme, we can show the **energy-stability**: **(COMPLETAR A PARTIR DE TFG DE ALBA?)**

MPP Space Discretization for (2a)

Let \mathcal{T}_h a mesh of Ω and let U_h be a space of \mathbb{P}_k^d (discontinuous) polynomials in elements $K \in \mathcal{T}_h$ (**mejorar esta definición, como en [1], sección 2.1**). We fix^b $k = 1$. Let V_h be an space of (conforming or not) FE. For each $m \geq 0$ let $v_h^{m+1} \in V_h$ computed from (2b), where u_h^{m+1} is replaced by $P_{V_h}(u_h^{m+1})$ (its $L^2(\Omega)$ -projection on W_h) and let $\mathbf{w}_h^{m+1} = P_{U_h^2}(\nabla v_h^{m+1})$.

Let us consider the following discrete problem: find $u_h^{m+1} \in U_h$,

$$\int_{\Omega} \delta_t u_h^{m+1} \phi + a^{sip}(u_h^{m+1}, \phi) + a^{god}(u_h^{m+1}, \phi) = 0 \quad \forall \phi \in U_h, \quad (3)$$

where

$$\begin{aligned} a^{sip}(u, \phi) &= \dots \text{completar} \dots, \\ a^{god}(u, \phi) &= - \sum_K \int_K u (\mathbf{w}_h^{m+1} \cdot \nabla \phi) + \sum_{e \in \mathcal{E}_h} \{u \mathbf{w}_h^{m+1} \cdot \mathbf{n}_e\}_* \end{aligned}$$

and

$$\{u \mathbf{w}_h^{m+1} \cdot \mathbf{n}_e\}_* = \dots \text{completar} \dots$$

is the Godunov (upwind) flux [1].

^aHowever, this work can be applied to higher order implicit methods or, following [1] and references therein, to parallel explicit high-order approximations via Strong Stability Preserving (SSP) methods

^bResults presented here might be improved to high order space approximations by using Bernstein polynomials, see[1]

References

- [1] R. Anderson, V. Dobrev, T. Kolev, D. Kuzmin, M. Quezada de Luna, R. Rieben, and V. Tomov. High-order local maximum principle preserving (MPP) discontinuous Galerkin finite element method for the transport equation. *Journal of Computational Physics*, 334:102–124, 2017.
- [2] N. Bellomo, A. Belloquid, Y. Tao, and M. Winkler. Towards a mathematical theory of keller-segel models of pattern formation in biological tissues. *Math. Models Meth. Appl. Sci.*, 25(9):1663–1763, 2015.
- [3] M. Farina, M. Marras, and G. Viglialoro. On explicit lower bounds and blow-up times in a model of chemotaxis. *Discret. Contin. Dyn. Syst. Suppl.*, pages 409–417, 2015.
- [4] F. Hecht. New development in freefem++. *J. Numer. Math.*, 20(3-4):251–265, 2012.

Test 1. Time Schemes Convergence Order

Exact solution: In order to compute error orders, we introduce the exact solution $u = v = e^{-2t} \cos x \cos y + 2$, adding the following term to the right hand side of (1.a):

$$f = -\alpha_1(4 \cos x \cos y(e^{2t} + \cos x \cos y) - \cos^2 x - \cos^2 y)e^{-4t}.$$

Experiment data: $\Omega = [0, \pi]^2 \subset \mathbb{R}^2$, $\alpha_i = 1$ for $i = 1, \dots, 4$ and $u_0 = v_0 = 2 + \cos x \cos y$. Discretization 50×50 P2-Lagrange, $k_i = 4 \cdot 10^{-1}/2^i$ for $i = 0, \dots, 5$. Here we show errors orders in norm $L^2(0, T; H^1(\Omega))$.

Test 2. Blow-up and Positivity

Acknowledgement

Second author is partially supported by the research group FQM-315 of Junta de Andalucía and by the project: New methods for predicting, manufacturing and application of shimming (FUTURE SHIMMING) -UIC AIRBUS, and the research team: Mathematics for Computational Intelligence Systems (M-CIS).

Contact

¹daniel.acostsoba@alum.uca.es

²alba.navarroiz@alum.uca.es

³rafael.rodriguez@uca.es