

# Math 104A Homework #3 \*

Instructor: Lihui Chai

**General Instructions:** Please write your homework papers neatly. You need to turn in both full printouts of your codes and the appropriate runs you made. Write your own code, individually. Do not copy codes!

1. (a) Let  $f \in C^2[x_0, x_1]$  and  $P_1$  its interpolation linear polynomial at  $x_0$  and  $x_1$ . Prove that

$$\|f - P_1\|_\infty \leq \frac{1}{8}(x_1 - x_0)^2 M_2, \quad (1)$$

where  $|f''(x)| \leq M_2$  for all  $x \in [x_0, x_1]$  and  $\|f - P_1\|_\infty = \max_{x \in [x_0, x_1]} |f(x) - P_1(x)|$ .

- (b) Let  $P_1(x)$  be the linear polynomial that interpolates  $f(x) = \sin x$  at  $x_0 = 0$  and  $x_1 = \pi/2$ . Using (a) find a bound for the maximum error  $\|f - P_1\|_\infty$  and compare this bound with the actual error at  $x = \pi/4$ .
2. (a) Equating the leading coefficient of in the Lagrange form of the interpolation polynomial  $P_n(x)$  with that of the Newton's form deduce that

$$f[x_0, x_1, \dots, x_n] = \sum_{j=0}^n \frac{f(x_j)}{\prod_{\substack{k=0 \\ k \neq j}}^n (x_j - x_k)}. \quad (2)$$

- (b) Use (a) to conclude that divided differences are symmetric functions of their arguments, i.e. any permutation of  $x_0, x_1, \dots, x_n$  leaves the corresponding divided difference unchanged.
3. In Newton's form of the interpolation polynomial we need to compute the coefficients,  $c_0 = f[x_0]$ ,  $c_1 = f[x_0, x_1]$ , ...,  $c_n = f[x_0, x_1, \dots, x_n]$ . In the table of divided differences we proceed column by column and the needed coefficients are in the uppermost diagonal. A simple 1D array,  $c$  of size  $n + 1$ , can be used to store and compute these values. We just have to compute them from bottom to top to avoid losing values we have already computed. The following pseudocode does precisely this:

**for**  $j = 0, 1, \dots, n$  **do**

---

\*All course materials (class lectures and discussions, handouts, homework assignments, examinations, web materials) and the intellectual content of the course itself are protected by United States Federal Copyright Law, the California Civil Code. The UC Policy 102.23 expressly prohibits students (and all other persons) from recording lectures or discussions and from distributing or selling lectures notes and all other course materials without the prior written permission of Prof. Hector D. Ceniceros.

```

     $c_j = f_j$ 
end for
for  $k = 1, \dots, n$  do
    for  $j = n, n-1, \dots, k$  do
         $c_j = (c_j - c_{j-1}) / (x_j - x_{j-k})$ 
    end for
end for

```

The evaluation of the interpolation polynomial in Newton's form can be then done with the Horner-like scheme:

```

 $p = c_n$ 
for  $j = n-1, n-2, \dots, 0$  do
     $p = c_j + (x - x_j)p$ 
end for

```

- (a) Write computer codes to compute the coefficients  $c_0, c_1, \dots, c_n$  and to evaluate the corresponding interpolation polynomial at an arbitrary point  $x$ . Test your codes and turn in a run of your test.
  - (b) Consider the function  $f(x) = e^{-x^2}$  for  $x \in [-1, 1]$  and the nodes  $x_j = -1 + j(2/10)$ ,  $j = 0, 1, \dots, 10$ . Use your code in (a) to evaluate  $P_{10}(x)$  at the points  $\bar{x}_j = -1 + j(2/100)$ ,  $j = 0, 1, \dots, 100$  and plot the error  $f(x) - P_{10}(x)$ .
4. Obtain the Hermite interpolation polynomial corresponding to the data  $f(0) = 0$ ,  $f'(0) = 0$ ,  $f(1) = 2$ ,  $f'(1) = 3$ .