

Math 104A Homework #2 *

Instructor: Lihui Chai

General Instructions: Please write your homework papers neatly. You need to turn in both full printouts of your codes and the appropriate runs you made. Write your own code, individually. Do not copy codes!

1. (a) Write the Lagrangian form of the interpolating polynomial $P_2(x)$ corresponding to the data in the table below:

x_j	$f(x_j)$
0	1
1	1
3	-5

- (b) Use $P_2(x)$ you obtained in (a) to approximate $f(2)$.

2. **(optional)** We proved in class that

$$\|f - P_n\|_\infty \leq (1 + \Lambda_n) \|f - P_n^*\|_\infty, \quad (1)$$

where P_n is the interpolating polynomial of f at the nodes x_0, \dots, x_n , P_n^* is the best approximation of f , in the maximum (infinity) norm, by a polynomial of degree at most n , and

$$\Lambda_n = \left\| \sum_{j=0}^n \left| l_j^{(n)} \right| \right\|_\infty, \quad (2)$$

is the Lebesgue constant (here the $l_j^{(n)}$ are the elementary Lagrange polynomials).

- (a) Write a computer code to evaluate the Lebesgue function

$$L^{(n)}(x) = \sum_{j=0}^n \left| l_j^{(n)}(x) \right|, \quad (3)$$

associated to a given set of pairwise distinct nodes x_0, \dots, x_n .

*All course materials (class lectures and discussions, handouts, homework assignments, examinations, web materials) and the intellectual content of the course itself are protected by United States Federal Copyright Law, the California Civil Code. The UC Policy 102.23 expressly prohibits students (and all other persons) from recording lectures or discussions and from distributing or selling lectures notes and all other course materials without the prior written permission of Prof. Hector D. Ceniceros.

- (b) Consider the equidistributed points $x_j = -1 + j(2/n)$ for $j = 0, \dots, n$. Write a computer code that uses (a) to evaluate and plot $L^{(n)}(x)$ (evaluate $L^{(n)}(x)$ at a large number of points \bar{x} to have a good plotting resolution, e.g. $\bar{x}_k = -1 + k(2/n_e)$, $k = 0, \dots, n_e$ with $n_e = 1000$) for $n = 4, 10$, and 20 . Estimate Λ_n for these three values of n .
- (c) Repeat (b) with the nodes given by $x_j = \cos(\frac{j\pi}{n})$, $j = 0, \dots, n$. Contrast the behavior of $L^{(n)}(x)$ and Λ_n with those corresponding to the equidistributed points in (b).
3. (a) Implement the Barycentric Formula for evaluating the interpolating polynomial for arbitrarily distributed nodes x_0, \dots, x_n ; you need to write a function or script that computes the barycentric weights $\lambda_j^{(n)} = 1/\Pi_{k \neq j}(x_j - x_k)$ first and another code to use these values in the Barycentric Formula. Make sure to test your implementation.
- (b) Consider the following table of data

x_j	$f(x_j)$
0.00	0.0000
0.25	0.7071
0.50	1.0000
0.75	0.7071
1.25	-0.7071
1.50	-1.0000

Use your code in (a) to find $P_5(2)$ as an approximation of $f(2)$.

4. *The Runge Example.* Let

$$f(x) = \frac{1}{1+x^2}, \quad x \in [-5, 5]. \quad (4)$$

Using your Barycentric Formula code (Prob. 3) and (5) and (6) below, evaluate and plot the interpolating polynomial of $f(x)$ corresponding to

- (a) the equidistributed nodes $x_j = -5 + j(10/n)$, $j = 0, \dots, n$ for $n = 4, 8$, and 12 .
- (b) the nodes $x_j = 5 \cos(\frac{j\pi}{n})$, $j = 0, \dots, n$ for $n = 4, 8, 12$, and 100 .
- (c) Repeat (a) for $f(x) = e^{-x^2/5}$ for $x \in [-5, 5]$ and comment on the result.

Remark 1. *It can be shown that for equidistributed nodes one can use the barycentric weights*

$$\lambda_j^{(n)} = (-1)^j \binom{n}{j}, \quad j = 0, \dots, n, \quad (5)$$

where $\binom{n}{j}$ is the binomial coefficient (`nchoosek(n,j)` in Matlab). It can be shown that for the nodes $x_j = \frac{a+b}{2} + \frac{b-a}{2} \cos(\frac{j\pi}{n})$, $j = 0, \dots, n$, in $[a, b]$, one can use

$$\lambda_j^{(n)} = \begin{cases} \frac{1}{2}(-1)^j & \text{for } j = 0 \text{ or } j = n \\ (-1)^j & \text{for } j = 1, \dots, n-1. \end{cases} \quad (6)$$

Make sure to employ (5) and (6) in your Barycentric Formula code for this problem. To plot the corresponding $P_n(x)$ evaluate $P_n(x)$ at a large number of points \bar{x} to have a good plotting

resolution, e.g. $\bar{x}_k = -5 + k(10/n_e)$, $k = 0, \dots, n_e$ with $n_e = 5000$. Note that your Barycentric Formula cannot be used to evaluate $P_n(x)$ when x coincides with an interpolating node! Plot also f for comparison. Compare (a) and (b) and comment on the result in view of what you observed in Prob. 2.