

Cannabinoids Disrupt Memory Encoding by Functionally Isolating Hippocampal CA1 from CA3

Roman A. Sandler ^{*1}, Dustin Fetterhoff ², Robert E. Hampson ², Sam A. Deadwyler ² & Vasilis Z. Marmarelis¹

¹Department of Biomedical Engineering, University of Southern California, Los Angeles, CA, USA

²Department of Physiology & Pharmacology, Wake Forest University, Winston-Salem, NC, USA

March 29, 2017

Abstract

Much of the research on cannabinoids (CBs) has focused on their effects at the molecular and synaptic level. However, the effects of CBs on the dynamics of neural circuits remains poorly understood. This study aims to disentangle the effects of CBs on the functional dynamics of the hippocampal Schaffer collateral synapse by using data-driven nonparametric modeling. Multi-unit activity was recorded from rats doing an episodic memory task in control sessions and under the influence of exogenously administered tetrahydrocannabinol (THC), the primary CB found in marijuana. It was found that THC left firing rate unaltered and only slightly reduced theta oscillations. Multivariate autoregressive models, estimated from spontaneous spiking activity, were then used to describe the dynamical transformation from CA3 to CA1. They revealed that THC served to functionally isolate CA1 from CA3 by reducing feedforward excitation and theta information flow. The functional isolation was compensated by increased feedback excitation within CA1, thus leading to unaltered firing rates. Finally, both of these effects were shown to be correlated with memory impairments in the episodic memory task. By elucidating the circuit mechanisms of CBs, these results help close the gap in knowledge between the cellular and behavioral effects of CBs.

Author Summary

Research into cannabinoids (CBs) over the last several decades has found that they induce a large variety of oftentimes opposing effects on various neuronal receptors and processes. Due to this plethora of effects, disentangling how CBs influence neuronal circuits has proven challenging. This paper contributes to our understanding of the circuit level effects of CBs by using data driven modeling to examine how THC affects the input-output relationship in the Schaffer collateral synapse in the hippocampus. It was found that THC functionally isolated CA1 from CA3 by reducing feedforward excitation and theta information flow while simultaneously increasing feedback excitation within CA1. By elucidating the circuit mechanisms of CBs, these results help close the gap in knowledge between the cellular and behavioral effects of CBs.

*Corresponding Author: rsandler00@gmail.com

1 Introduction

Recent years have seen a resurgence of interest in the therapeutic role of cannabinoids (CBs) for several diseases and neuropsychiatric disorders such as psychosis, anxiety disorders, PTSD, and multiple sclerosis [1, 2]. In particular, CB agonists have shown promising but mixed results in the treatment of epilepsy, as various types of agonists at various doses have been shown to be both pro- and anticonvulsant [3, 4, 5, 6, 7, 8, 7]. Parallel to increasing therapeutic research, much work has been done on the chemical structure of various cannabinoids and cannabinoid receptors, along with their cellular interactions and pharmacology [9].

Nonetheless, between the large bodies of literature on cannabinoids from chemical, disease, and behavioral perspectives, much less work has been done to explore the effects of cannabinoids on the neural circuit level. This is particularly important since a wide range of complex and often opposing effects have been attributed to cannabinoids on a molecular and cellular level. For example, cannabinoid activation of CB1 receptors, which are found on both pyramidal cells and interneurons, reduces the quantity of neurotransmitter released during an action potential; consequently, increased extracellular cannabinoid levels reduce both excitatory (glutamatergic) and inhibitory (GABAergic) transmission [10]. Furthermore, cannabinoids have been shown to interact with astrocytes [11], mitochondria [12], glycine receptors [13], vanilloid receptors [14], potassium ion channels [15], and to reduce GABA and glutamate reuptake [16, 17]. Consequently, it is very difficult to extrapolate the emergent network level changes simply from a catalogue of effects cannabinoids have at a cellular/molecular level.

Here, we studied the effects of Δ^9 -tetrahydrocannabinol (THC) on hippocampal networks during memory encoding using spiking activity recorded in rodents in-vivo performing the Delayed-NonMatch-to-Sample (DNMS) episodic memory task. Multivariate autoregressive (MVAR) models [18], were used in both control and THC sessions to estimate feedforward and feedback dynamical filters, which are akin to the waveform shapes of the CA3→CA1 EPSP and CA1 afterhyperpolarization, respectively. MVAR models, which are a type of linear nonparametric model, are 'data-driven' in the sense that they estimate model parameters directly from recorded neural spiketrains and, unlike more biologically realistic models, make very few *a priori* assumptions on the nature of the neural dynamics [19, 20]. This characteristic makes them particularly well suited for this study, since as previously mentioned the emergent effects of THC on neural circuits are highly complex and unclear. Overall our results suggest that cannabinoids impair memory encoding by functionally isolating CA1 from CA3 via reduced theta information flow and altered excitatory-inhibitory balance across the Schaffer collateral synapse.

2 Results

2.1 Changes in rate and temporal coding under Cannabinoids

To evaluate the effects of exogenous cannabinoids on the hippocampal network 1 mg/kg THC was injected intraperitoneally into $N = 6$ rodents during certain sessions while they were performing a DNMS task (Fig. S1). All data was previously used in a study on the effects of cannabinoids on hippocampal multifractality [21, 22]. Briefly, in the sample phase, the rats were presented one of two levers. After a variable length delay, both levers were presented in the match phase and the rat had to choose the opposite lever to receive a reward. On the behavioral level, it was found that THC impaired rodent-performance on the DNMS task by about $12.2 \pm .6\%$ (Fig. 1a, [23]).

While performing the DNMS task, single-unit activity was recorded from the hippocampal CA3 and CA1 regions using a multi-electrode array. No mean firing rate (MFR) changes were seen during THC sessions in either CA3 or CA1 cells ($P=.502$, Fig. 1b). This includes activity around the DNMS sample phase, during the entire session, and in both general cells and sample-presentation cells (see below). The lack of any cannabinoid-induced changes in firing rates at this dosage has been observed in previous studies [24, 25].

Two types of temporal coding were identified in the recorded spiketrains. First, on slower timescales, several neurons fired preferentially in response to lever presentation in the sample phase of the DNMS task [26]. It was found that THC reduced the proportion of sample-presentation cells in both CA3 and CA1 by roughly equal amounts ($\Delta = 13 \pm 4\%$, $P < .001$; Fig. 1c). Interestingly, some sample-presentation cells lost all of their preferential firing in THC sessions (Fig.1d); this contrasts with place cells whose receptive field stays largely intact under cannabinoids [27]. There was an interesting but insignificant trend connecting sample-presentation cell reduction with behavioral deficits ($R^2 = .27$, $P = .052$, Fig. S3a).

On faster timescales, it was found that several CA3 and CA1 neurons had theta band rhythmicity (4-7 Hz). Hippocampal theta oscillations are known to be intimately related to cognitive function [28, 29, 30] and have previously been linked to performance in the DNMS task [31]; furthermore, theta oscillations are known to be reduced by systemic injections of cannabinoids on both the single unit [24] and network level [32]. It was found that CA1 theta power was slightly but significantly reduced in THC sessions ($\Delta = 2.52\%$, $CI : [.61, 4.4]\%$ $P = .004$; Fig. 1e). A similar, albeit slightly weaker, theta power reduction was seen in CA3 cells ($\Delta = 1.94\%$, $P = .045$; Fig. S2). Unlike previous results in a different task [24], the reduction in CA1 theta power was not found to be correlated with behavioral deficits in the DNMS task ($P = .674$, Fig. S3b).

Overall, these results show that THC has minor effects on the actual neuronal spiketimes: quantity of spikes (MFR) was not affected and spike rhythmicity (theta oscillations) were only slightly affected. Furthermore behavioral deficits induced by cannabinoids could not be explained by any of these factors, which are the traditional markers of rate and temporal coding in the hippocampus.




Figure 1: (A) Behavioral performance on Delayed-NonMatch-to-Sample (DNMS) task in both control and THC sessions. Dashed lines show individual animal performance, while solid lines show mean performance over all animals. Bars indicate SEM. Dashed red line indicates performance at chance level. (B) Individual neuron mean firing rate (MFR). (C) Sample-presentation cell proportion in CA3 and CA1 cells in control & THC sessions (D) Example of a sample-presentation cell in a control session (top) which lost its firing specificity under THC (bottom). X-axis shows MFR (Hz) (E) Average CA1 spiketrain spectra (top). Bottom shows mean difference in individual cell spectra (thus it is not simply the difference between the signals in above which are averaged over whole population). Gray error bounds indicate 99% confidence bounds. In (B) and (E), only neurons recorded in at least one control & THC session were included. Results for neurons recorded in several control or THC sessions were averaged over those sessions.

2.2 Systems Analysis

The remainder of the study will focus on systems analysis of the Schaffer collateral synapse connecting CA3 to CA1, and how this synapse is affected by THC. Systems analysis aims to identify the input-output "blackbox" by which the input spiketrains are transformed into the output spiketrain. On a more abstract level, it aims to identify how the information encoded in CA3 is propagated into CA1. This is distinct from the *signal* analysis done in the previous section which only looks at features of individual spiketrains rather than the causal relationship between multiple spiketrains as done in systems analysis.

The relationship between an arbitrary number of input CA3 spiketrains and the output CA1 spiketrain was modeled using a multivariate autoregressive model described by Eq. 1 and an example of which is pictured in Fig. 2a. Each system consists of N input CA3 neurons and N feedforward filters describing the dynamical input-output relationship between the given CA3 and CA1 neurons. Intuitively, these filters can be thought of as the EPSP elicited in the output CA1 neuron in response to an action potential (AP) in the input CA3 neuron. However, unlike EPSPs which traditionally only encapsulate ion-conductances from neurotransmitter-gated ion channels, the "blackbox" nature of the feedforward filters means they also include more complex dynamical effects such as dendritic integration, spike generation, active membrane conductances, and feedforward interneuronal inhibition (thereby allowing the filters between two pyramidal cells to be inhibitory). Each model also includes a feedback (autoregressive) filter which describes the effects of past output spikes onto the output present. This filter, which can be intuitively thought of as the afterhyperpotential (AHP) [33] includes intracellular processes such as the absolute and relative refractory periods, slow potassium conductances, and I_h conductances. It also includes more complex intercellular processes such as the recurrent connections between CA1 pyramidal cells and interneurons [34]. Neuronal connectivity was estimated using a stepwise input selection procedure. Filter parameters were estimated with Laguerre basis regression using neuronal activity around the sample phase. Model significance was verified using ROC plots and shuffling methods (see supplementary methods).

A representative connectivity grid from a recorded THC session with 10 recorded neurons (4 CA3, 6 CA1) is shown in Fig. 2a. Fig. 2b shows a sample system from this session between 3 CA3 pyramidal cells and 1 CA1 pyramidal cell. Note that two of the feedforward filters are excitatory (above the x-axis) while the third has both excitatory and inhibitory components, presumably arising through feedforward inhibition involving interneurons [35, 36]. The system also involves a feedback filter which shows a relatively long refractory period (~ 40 ms) followed oscillatory bursting activity. Oscillations in the CA1 pyramidal cell AHP are a well known phenomena caused by slow K^+ and I_h conductances, and these oscillations are known to lead to theta resonances [37, 38, 18]. In order to study the filter oscillations more closely, the filter frequency spectra were plotted in Fig. 2c. Both feedforward excitatory filters were found to have peaks in the high theta range (8-9 Hz). Intuitively, this can be understood

171 to mean that information encoded in the theta range in these input neurons is
 172 preferentially transmitted to the output CA1 neuron. Furthermore, the feedback
 173 filter has a low theta resonance of 3.5 Hz. Significance metrics for the displayed
 174 system is shown in Fig. S4, and additional systems are shown in Fig. S5. All
 175 together 66% (707/1068) of all systems were found to be significant and 2139
 176 feedforward and 707 feedback filters were obtained. THC was found to reduce
 177 the amount of significant models per session ($\Delta = -7.4\%$, $P = .011$), but the
 178 predictive power of significant models, as measured by AUC (see supplementary
 179 methods), was unaltered ($P = .24$).




Figure 2: (A) Example connectivity grid of 4 CA3 neurons (orange) and 6 CA1 neurons (blue) recorded during a single session. Note that 1 CA1 neuron has no significant granger-causal inputs. Each line represents a causal connection between those neurons, as encapsulated by a feedforward filter. (B) Example system of CA1 neuron enclosed by the red box in (A). Diagram shows 3 input CA3 spiketrains followed by their respective feedforward filters which are summed with the feedback filter to generate the output CA1 spiketrain. All feedforward filter are plotted with the same y-axis scale. Dashed red line in filter boxes indicates x-axis. (C) Normalized filter spectra computed of feedforward and feedback filters from (B).

180 To study how THC affects system dynamics on a population level, we ex-
 181 amined how features change in the entire sample of control and THC filters.
 182 The average filter frequency profile for both control and THC sessions is shown
 183 in Fig. 3a,b (top). Both feedforward and feedback spectra are found to have
 184 clear theta band peaks, thus generalizing the trend seen in the example system
 185 of Fig. 2. This is consistent previous reports which show that CA3 propagates
 186 strong theta rhythms to CA1 [39, 40] and also that CA1 is capable to generating
 187 endogenous theta rhythms [41]. A significant decline was found in the theta
 188 power of the feedback filters ($\Delta = 20.8\%$, $P < .001$; Fig. 3b). Note that the
 189 feedback filter theta reduction is about 10x stronger than the theta reduction
 190 found in the CA1 spiketrain signals (Fig. 1e. No reduction in theta power was
 191 found in the feedforward filters ($P = .61$, Fig. 3a). This surprising result sug-
 192 gests that cannabinoid-induced theta desynchronization results primarily from
 193 altered feedback properties rather than changes in CA3→CA1 dynamics.

194 Cannabinoids have been reported to affect network excitation-inhibition bal-
 195 ance (EIB) [10, 42]. Particularly, there is much debate whether cannabinoids are
 196 pro- or anticonvulsants [8, 43, 44, 4, 6]. In order to examine the effects cannabi-

197 noids have on network EIB, we quantified the excitation of the estimated filters
 198 using a metric called the excitatory index (EI), which is the ratio between posi-
 199 tive filter area and total filter area. It was found that THC had no significant
 200 effect on feedforward EI ($P = .14$); however, there was an interesting but in-
 201 significant trend showing that THC-induced decreases in feedforward EI were
 202 correlated with behavioral deficits ($R^2 = .27, P = .063$, Fig. 3c). Addition-
 203 ally, THC reduced the amount of casually connected CA3-CA1 neuronal pairs
 204 ($\Delta = -8.9\%$, $P < .001$). These findings, together with the THC-induced de-
 205 crease of CA3→CA1 significant models, suggest that THC reduces the causal
 206 influence CA3 neurons have on CA1 spiketimes. In other words, THC can be
 207 said to functionally isolate CA1 from CA3. It was also found that THC sig-
 208 nificantly increased feedback EI ($\Delta = 3.5\%$, $P = .022$) and that the increased
 209 feedback EI was correlated with behavioral deficits ($R^2 = .38, P = .007$, Fig.
 210 3d).




Figure 3: Average feedforward (A) and feedback (B) filter spectra in control and THC sessions (top), and their differences (bottom). Same format and analysis as Fig. 1a. (C) Correlation between feedforward filter excitatory index (EI) reduction and behavioral deficits. Each point represents a specific THC session, with points of the same color coming from the same animal. X-axis shows reduction in feedforward EI, while y-axis shows reduction in behavioral performance. Both reductions were taken relative to control sessions (see supplemental methods). (D) Same as (C) but for feedback EI increase.

2.3 PDM Analysis

The large quantity (>2800) and variability of the obtained filters describing the CA3→CA1 dynamic transformation presents a challenge of interpretation. Namely, how could one identify features from the entire filter population which are representative of the CA3→CA1 transformation rather than just the input-output relationship found in this or that particular pair of neurons. In essence this is an unsupervised learning problem which aims to identify hidden structure within the filter population for the purpose of knowledge discovery. Our group has developed the concept of the global principal dynamic modes (gPDMs) towards this effort [19, 45, 46]. The gPDMs are a system-specific and efficient basis set which contain the essential dynamic components of the filter population and are meant to be amenable to biological interpretation. One set of gPDMs were estimated from all (control and THC) obtained filters with the hypothesis that THC would primarily change the expression strength of the gPDMs rather than their specific shapes.

Fig. 4a,b shows the obtained feedforward and feedback gPDMs in both the time and frequency domain. Once again, the feedforward and feedback gPDMs represent the dominant independent components of feedforward and feedback kernels, respectively. The first, and thus most significant, feedforward gPDM was found to have almost all its energy in the 1st time bin, with an immediate decline thereafter. This gPDM represents near concurrent firing between CA3 and CA1 neurons and presumably results from both direct CA3→CA1 connections via the Schaffer collateral synapse [47, 48] and common inputs from the entorhinal cortex [49, 50]. The third feedforward gPDM, which is characterized by an initial inhibitory phase, presumably represents feedforward interneuronal inhibition which is prevalent in the CA3→CA1 connection [35, 36]. THC was not found to influence the strength of either of these gPDMs ($P = .76$, $P = .60$; Fig. S6). The second feedforward gPDM which is characterized by sustained and oscillatory excitation was found to have a strong theta peak in the frequency domain. Furthermore, it was found that THC-induced declines in the strength of this gPDM were correlated with behavioral deficits ($R^2 = .30$, $P = .032$; Fig. 4c).

The three obtained feedback gPDMs are shown in Fig. 4b. These gPDMs express the essential feedback dynamics found in CA1 neurons. As previously mentioned, these dynamics arise through the combination of intracellular processes such as the AHP and extracellular processes such as recurrent connections between CA1 pyramidal cells and interneurons. It was found that THC-induced increases in the third feedback gPDM were correlated with behavioral deficits ($R^2 = .39$, $P = .005$; Fig. 4d). This correlation was not seen in either of the first two feedback gPDMs ($P = .32$, $P = .75$; Fig. S6). Notably, the 3rd feedback gPDM was seen to be "theta-blocking" in the frequency domain - namely, it serves to counteract the 1st "theta-promoting" feedback gPDM and disrupts theta oscillations in the CA1 neuron. The THC-induced changes in the feedforward and feedback theta gPDMs paint a more complete picture of the CA1 theta reductions seen in Fig. 1e. Namely, they attribute the theta losses to spe-




Figure 4: Feedforward (A) and feedback (B) global principal dynamic modes (gPDMs) in both the time (left) and frequency domain (right). Reductions in 2nd feedforward gPDM (C) and increases in 3rd feedback gPDM (D) were found to be correlated with behavioral deficits. Same format as Fig. 3.

cific feedforward and feedback dynamical filters which may potentially be traced to specific biophysical mechanisms. Furthermore, changes in these dynamical filters have been specifically correlated with behavioral deficits, which could not be done with theta reductions in the CA1 signal (Fig. S3).

3 Discussion

The current study uses 'data-driven' nonparametric system dynamics modeling tools to study the effects of THC on the Schaffer Collateral synapse in rodents. The chief findings of the study can be summarized as: (1) THC induced little or no change in traditional rate and temporal coding metrics such as MFR and theta power, (2) THC altered the CA1 excitatory-inhibitory balance by reducing feedforward influence from CA3 while increasing feedback excitation from CA1, (3) THC reduced theta information flow through the Schaffer collateral synapse, and (4) The magnitudes of both of the previous effects were directly correlated with the severity of behavioral deficits induced by THC. Overall these results suggest the conclusion that THC impairs memory encoding by functionally isolating CA1 from CA3.

From a computational perspective, the nonparametric modeling methods used in this study proved successful in studying the network level effects of cannabinoids since, unlike biophysical models, all model parameters were es-

275 timated directly from recorded data and very few *a priori* assumptions were
 276 made about the effects of THC [19, 20, 51]. The global principal dynamic modes
 277 (gPDMs), which were derived from MVAR filters of the entire population of neu-
 278 rons, further extracted hidden dynamical structure from 'noisy' neuron-neuron
 279 variability. Importantly, THC-induced changes in the gPDMs were directly cor-
 280 related with behavioral impairments, thus justifying their utility. Furthermore,
 281 while most in-vivo studies on THC analyze macro level signals such as ECoG
 282 and EEG, this work adds to a relatively small body of literature which ana-
 283 lyzes the effects of THC on neuronal population spiking activity. Finally, to our
 284 knowledge, this is the first work which examines the effect of THC on neuronal
 285 systems dynamics, or the causal interactions between signals, rather than on
 286 neuronal signals themselves.

287 It was found that THC increased feedback excitatory index in CA1 and that
 288 the magnitude of this effect was correlated with behavioral deficits. We hypoth-
 289 esize that this is due to reduced feedback inhibition from CA1 cholecystokinin
 290 (CCK)-containing basket cells. While CCK cells only make up only 13.9% of
 291 interneurons [52], they express significantly more CB1 receptors than any other
 292 cell in the hippocampus [53], and their primary output is to CA1 pyramidal cells
 293 [52]. Increased THC concentrations would reduce CCK interneuron output by
 294 (1) reducing the amount of GABA they release per action potential (2) reducing
 295 their MFR due to reduced glutamatergic input from principal cells in both CA3
 296 and CA1 [54, 55].

297 It was also found that THC reduces the number of casually connected CA3-
 298 CA1 neuronal pairs; furthermore there was an interesting but insignificant trend
 299 for THC-induced deficits in feedforward excitation to lead to behavioral deficits.
 300 This trend may prove to be significant given a higher sample size. We hypoth-
 301 esize that this reduced feedforward influence is caused by decreased glutamate
 302 release from CA3 pyramidal cells due to CB1 receptor activation by THC [56].
 303 Even though pyramidal cells have much lower densities of CB1 receptors than
 304 interneurons [53, 57], there is evidence that CB induced reduction of excita-
 305 tion is larger than these relative densities suggest. Principal cells outnumber
 306 interneurons 20:1 in CA1 [50] and their CB1 receptors were found to be several
 307 fold more efficacious than those of interneurons [58]. Further, lower baseline ac-
 308 tivation levels of CB1 receptors on principal cells than on interneurons suggest
 309 they would be disproportionately activated by CB agonists [59]. Altogether, the
 310 decreased feedback inhibition and feedforward excitation amount to a functional
 311 isolation, or breakdown in information flow from CA3 to CA1. We suggest that
 312 this functional isolation is responsible for the behavioral impairments seen in the
 313 DNMS task.

314 The 'functional isolation' hypothesis is further supported by previous work
 315 which showed that the behavioral impairments caused by cannabinoids in the
 316 DNMS task were similar to those seen with a full pharmacological lesion of the
 317 hippocampus [60] Given the centrality of CA3→CA1 information flow to hip-
 318 pocampal function, a functional isolation of these areas could indeed presumably
 319 lead to impairments similar to that of a full lesion. Also relevant is a study by
 320 Goonawardena et al. [25] where THC was injected intraperitoneally at low 1

321 mg/kg doses as in this study and in higher doses of 3 mg/kg. It was found
322 that while both doses disrupted hippocampal synchrony, only the higher dose
323 resulted in a reduction in pyramidal cell MFR. This suggests that at the lower
324 doses both previously described phenomena are at a net balance, while at the
325 higher dose, the decrease in feedforward excitation overpowers the increase in
326 feedback excitation and results in lower MFR. Finally, the hypothesis predicts a
327 breakdown in the normal spiketime coordination between pyramidal cells and in-
328 terneurons in CA1 circuits. The breakdown of this coordination, which has been
329 extensively implicated in hippocampal oscillations [61, 62], could be responsible
330 for the observed decrease in theta oscillations and information flow.

331 Although the current results only suggest this hypothesis, several experi-
332 ments could be done to further substantiate it. Feedforward and feedback ker-
333 nels and gPDMs could be estimated at different doses of THC; the hypothesis
334 would predict that different doses would effect the two processes independently,
335 with one of the two processes potentially being more dominant at different THC
336 levels. Also significant developments in in-vivo synaptic patch clamping [63] and
337 calcium imaging in recent years could be used to directly measure the drive of
338 CCK cells and CA3 pyramidal cells onto CA1 pyramidal cells under THC.

339 Much research has been done investigating the effects THC and other cannabi-
340 noids have on seizures and epilepsy. Results so far have been mixed, with various
341 studies showing that THC is both pro- and anticonvulsant [3, 4, 5, 6, 7, 8, 7].
342 The results from this study and the presented hypothesis suggest that THC in-
343 herently is not pro- or anti-convulsant but that its effects will depend on the
344 dosage and the unique circuitry of every epileptic focus. Interestingly, a study
345 by Rudenko et al. [6] has shown that indeed the effects of a CB1 agonist were
346 dose dependant, with *lower* doses being anticonvulsant and higher doses being
347 proconvulsant. Finally, this study suggests that in order to truly understand the
348 effects of THC on epileptic circuits, one must study the systems level changes
349 in circuit dynamics rather than taking a reductionist approach and studying the
350 effects of THC on any particular receptor or cell type.

351 The present study analyzed the effects of THC from both a signals and sys-
352 tems perspective - and found that systems analysis yielded much richer results.
353 For example, while analysis of CA1 spiketrain signals showed a slight (2%) re-
354 duction in theta frequency, analysis of system kernels showed that the theta loss
355 was primarily due to CA1 feedback dynamics whose kernels lost over 20% of their
356 theta power, while theta power in feedforward kernels was unaffected. Further-
357 more, only systems analysis allows one to analyze predictive power, feedforward
358 and feedback excitation, and EPSP and AHP waveform shape. Notably, the
359 finding that feedforward influence decreased while feedback excitation increased
360 could not have been observed using only signal analysis which would have only
361 detected a constant MFR.

362 The present study also employed gPDMs as a means to extract the most
363 significant information from the kernel dynamics estimated from several animals
364 over several sessions [19, 64, 18, 48]. The utility of the gPDM method was justi-
365 fied by the finding that reductions in theta related gPDMs in a given session were
366 directly correlated with behavioral deficits, showing that the gPDMs can isolate

the particular dynamics which are most affected by THC. Furthermore, THC-induced theta power losses in spiketrain signals were not found to be correlated with behavioral deficits. Although in the present study, kernels and gPDMs were restricted to being linear in order to more easily quantify their overall strength and excitation (via the EI), future work will aim to identify the effects of THC on hippocampal nonlinear dynamics [65, 51].

Ethics Statement

All animal protocols were approved by the Wake Forest University Institutional Animal Care and Use Committee, in accordance with the Association for Assessment and Accreditation of Laboratory Animal Care and the National Institute of Health Guide for the Care and Use of Laboratory Animals (NIH Publication No. 8023).

Acknowledgements

This work was supported by NIH (www.nih.gov) grant P41-EB001978 to the Biomedical Simulations Resource at the University of Southern California.

4 Methods

4.1 Experimental Procedures

N=6 Male Long-Evans rats were trained to criterion on a two lever, spatial Delayed NonMatch-to-Sample (DNMS) task (see Fig. S1). Briefly, during the sample phase the rat was presented one of two levers (left or right). After a delay phase ranging from 1-30 seconds, the rat was presented both levers and had to choose the opposite level in order to attain a reward. Each rodent underwent 16-25 sessions of the task, which were roughly evenly divided between control and THC sessions, wherein the rodent was intraperitoneally administered 1 mg/kg of body weight Δ^9 -tetrahydrocannabinol (THC), an exogenous cannabinoid found in marijuana. During the task, spike trains were recorded in-vivo with multi-electrode arrays implanted in the left and right CA3 and CA1 regions of the hippocampus. In an effort to acquire a consistent cognitive state, only spiking activity around the sample phase of the task was used. Spikes from multiple trials were sorted, time-stamped, and concatenated into a discretized binary time series using a 4ms bin. For more details on the experimental setup, see supplementary methods.

4.2 Model Configuration and Estimation

Nonparametric multiple-input linear autoregressive models were used to model the dynamical transformation between input and output spike trains (see Fig. 2,5) [18, 51]. Each model consisted of a feedforward component, reflecting the effect of the N input cells on the output cell and a feedback (autoregressive)

404 component reflecting the subthreshold and suprathreshold effects the output
 405 cell has on itself. Thus, the output $y(t)$ is calculated as:

$$y(t) = \sum_{n=1}^N \sum_{\tau=0}^M k_n(\tau) x_n(t - \tau) + \sum_{\tau=1}^{M+1} k_{AR}(\tau) y(t - \tau) \quad (1)$$

406 where k_n reflects the feedforward filter of input $x_n(t)$, and k_{AR} reflects the feed-
 407 back filter. In order to reduce the amount of model parameters and thereby
 408 increase parameter stability, we applied the Laguerre expansion technique to ex-
 409 pand the feedforward and feedback filters over L Laguerre basis functions (see
 410 supplementary methods).

411 Effective connectivity between neurons was assessed using a Granger causality-
 412 like approach. For each output CA1 neuron, input CA3 neurons were selected
 413 in a forward stepwise procedure whereby only neurons which help predict the
 414 output CA1 spike activity were included in the model. After all input neurons
 415 were selected, a Monte Carlo approach was used to assess model significance. A
 416 model was deemed significant if the CA3 inputs could predict the output CA1
 417 activity significantly better ($P < .0001$) than randomly permuted versions of the
 418 inputs. See supplementary methods for more details.




Figure 5: Model Configuration. Each model has N point-process inputs which each go through a linear filter, K_i . These inputs are then summed with the output of the feedback filter, K_{AR} to generate the final output, $y(t)$, which is a continuous signal

419 4.3 Principal Dynamic Modes

420 The global principal dynamic modes (gPDMs) were obtained in a two step pro-
 421 cess: first, all filters of each input from every animal were concatenated in a
 422 rectangular matrix. Then singular value decomposition (SVD) was performed
 423 on the rectangular matrix to obtain all the significant singular vectors, which
 424 are the gPDMs. It was found that 3 gPDMs were sufficient to describe the lin-
 425 ear dynamics both the population of feedforward and feedback filters. gPDM
 426 strength in a given filter was computed by taking the dot product between the
 427 gPDM and the filter. gPDM strength in a given session was computed by taking
 428 the average gPDM strength in every filter of that session.

429 **A Supplementary Methods**

430 All data was previously used in a study on the effects of cannabinoids on hip-
431 pocampal multifractality [21, 22])

432 **A.1 Animals**

433 Subjects were Long-Evans rats (Harlan) aged 4–6 months ($n = 6$) individually
434 housed and allowed free access to food with water regulation to maintain 85% of
435 ad libitum body weight during testing. All animal protocols were approved by
436 the Wake Forest University Institutional Animal Care and Use Committee, in
437 accordance with the Association for Assessment and Accreditation of Laboratory
438 Animal Care and the National Institute of Health Guide for the Care and Use
439 of Laboratory Animals (NIH Publication No. 8023).

440 **A.2 Apparatus**

441 The behavioral testing apparatus for the delayed nonmatch-to-sample (DNMS)
442 task is the same as reported in other studies [23] and consisted of a 43x43x50 cm
443 Plexiglas chamber with two retractable levers (left and right) positioned on either
444 side of a water trough on the front panel. A nosepoke device (photocell) was
445 mounted in the center of the wall opposite the levers with a cue light positioned
446 immediately above the nosepoke device. A video camera was mounted on the
447 ceiling and the entire chamber was housed inside a commercially built sound-
448 attenuated cubicle.

449 **A.3 DNMS Task**

450 The DNMS task consisted of three main phases: Sample, Delay and Nonmatch.
451 The sample phase initiated the trial when either the left or right lever was
452 extended (50% probability), requiring the animal to press it as the Sample Re-
453 sponse (SR). The lever was then retracted and the Delay phase of the task
454 initiated, as signaled by the illumination of a cue light over the nosepoke pho-
455 tocell device on the wall on the opposite side of the chamber. At least one
456 nosepoke (NP) was required following the delay interval which varied randomly
457 in duration (1-30 s) on each trial during the session. The Nonmatch phase began
458 when the delay timed out, the photocell cue light turned off and both the left
459 and right levers on the front panel were extended. Correct responses consisted
460 of pressing the lever in the Nonmatch phase located in the spatial position op-
461 posite the SR (nonmatch response: NR). This produced a drop of water (0.4
462 ml) reward in the trough between the two levers. After the NR the levers were
463 retracted for a 10.0 second intertrial interval (ITI) before the next Sample lever
464 was presented to begin the next trial. A lever press at the same position as the
465 SR (match response) constituted an “error” with no water delivery and turned
466 off of the chamber house lights for 5.0s and the next trial was presented 5.0 s
467 later. Individual performance was assessed as % NRs (correct responses) with
468 respect to the total number of trials (80-100) per daily (1 hr) sessions.

469 A.4 Drug Preparation & Administration

470 Δ^9 -tetrahydrocannabinol (THC) was obtained from the National Institute on
471 Drug Abuse as a 50 mg/ml solution in ethanol. Detergent vehicle was pre-
472 pared from Pluronic F68 (Sigma, St. Louis, MO), 20 mg/ml in ethanol. THC
473 was added to the detergent-ethanol solution (0.5 ml of either THC), and then 2.0
474 ml of saline (0.9%) was slowly added to the ethanol-drug solution. The solution
475 was stirred rapidly and placed under a steady stream of nitrogen gas to evapo-
476 rate the ethanol (~ 10 min). This resulted in a detergent-drug suspension (12.5
477 mg/ml THC), which was sonicated and then diluted with saline to final injec-
478 tion concentrations (0.5-2.0 mg/ml THC). On drug administration days, animals
479 were injected intraperitoneally with the drug-detergent solution (1 mg/kg) ~ 10
480 min before the start of the behavioral session. Our experience with these ex-
481 periments has shown that performance after vehicle injection is not significantly
482 different than no injection, and therefore was omitted during this series of ex-
483 periments to minimize risk of infection to the animals. At least two no injection
484 days were imposed between each drug-testing session. All drug solutions were
485 mixed fresh each day.

486 A.5 Surgery

487 All surgical procedures conformed to National Institutes of Health and Associ-
488 ation for Assessment and Accreditation of Laboratory Animal Care guidelines,
489 and were performed in a rodent surgical facility approved by the Wake Forest
490 University Institutional Animal Care and Use Committee. After being trained to
491 criterion performance level in the DNMS task animals were anesthetized with ke-
492 tamine (100 mg/kg) and xylazine (10 mg/kg) and placed in a stereotaxic frame.
493 Craniotomies (5mm-diameter) were performed bilaterally over the dorsal hip-
494 pocampus to provide for implantation of 2 identical array electrodes (Neurolinec,
495 New York, NY), each consisting of two rows of 8 stainless steel wires (diameter:
496 20 μm) positioned such that the geometric center of each electrode array was
497 centered at co-ordinates 3.4 mm posterior to Bregma and 3.0 mm lateral (right
498 or left) to midline [66]. The array was designed such that the distance between
499 two adjacent electrodes within a row was 200 μm and between rows was 400
500 μm to conform to the locations of the respective CA3 and CA1 cell layers. The
501 longitudinal axis of the array of electrodes was angled 30° to the midline during
502 implantation to conform to the orientation of the longitudinal axis of the hip-
503 pocampus, with posterior electrode sites more lateral than anterior sites. The
504 electrode array was lowered in 25-100 μm steps to a depth of 3.0 - 4.0 mm from
505 the cortical surface for the longer electrodes positioned in the CA3 cell layer,
506 leaving the shorter CA1 electrodes 1.2 mm higher with tips in the CA1 layer.
507 Extracellular neuronal spike activity was monitored from all electrodes during
508 surgery to maximize placement in the appropriate hippocampal cell layers. After
509 placement of the array the cranium was sealed with bone wax and dental cement
510 and the animals treated with buprenorphine (0.01–0.05 mg/kg) for pain relief
511 over the next 4-6 hrs. The scalp wound was treated periodically with Neosporin
512 antibiotic and systemic injections of penicillin G (300,000 U, intramuscular) were

513 given to prevent infection. Animals were allowed to recover from surgery for at
514 least 1 week before continuing behavioral testing [67].

515 **A.6 Electrophysiological Monitoring & Preprocessing**

516 Animals were connected by cable to the recording apparatus via a 32-channel
517 headstage and harness attached to a 40-channel slip-ring commutator (Crist
518 Instruments, Hagerstown, MD) to allow free movement in the behavioral test-
519 ing chamber. Single neuron action potentials (spikes) were isolated by time-
520 amplitude window discrimination and computer-identified individual waveform
521 characteristics using a multi-neuron acquisition (MAP) processor (Plexon Inc.,
522 Dallas, TX, USA). Single neuron spikes were recorded daily and identified us-
523 ing waveform and firing characteristics within the task (perievent histograms)
524 for each of the DNMS events (SR, LNP & NR). To maintain waveform shape
525 across days, all recorded data was concatenated into one file (separately for each
526 rat) and offline sorting was performed using principal component analysis, peak-
527 valley, and nonlinear energy algorithms in Offline Sorter (Plexon Inc., Dallas,
528 TX, USA). Hippocampal neuron ensembles used to distinguish recording phases
529 and drug treatment conditions consisted of 10-30 single neurons, each recorded
530 from a separate identified electrode location on either of the bilateral arrays.
531 All isolated spike trains contained no less than a 1 ms gap at the center of the
532 autocorrelogram. No effort was made to differentiate between principal cells and
533 interneurons. Previous work has shown that hippocampal neurons recorded with
534 the same waveform from the same electrodes exhibit consistent mean, baseline
535 and DNMS task modulated firing rate alterations [68, 26], and therefore indi-
536 vidual neurons were treated as the same when recorded over multiple days. A
537 total of 189 neurons recorded during 5,143 recording phases were analyzed in
538 the reported experiments.

539 **A.7 Sample-Response Cell Identification**

540 Prior studies from this laboratory have identified hippocampal neurons recorded
541 as above by “Functional Cell Types” (FCTs) described by different behavioral
542 correlates of DNMS task-related events such as lever position and/or phase of
543 the task [26, 25]. Sample-response cells, a subtype of FCTs, were identified by
544 first constructing a smoothed (51 bin) perievent histogram around the sample
545 presentation phase of the DNMS task. The neurons background firing rate mean
546 and variance were calculated from activity 3.5-5s after sample presentation. If
547 the neuron’s MFR from the 2 second window around sample presentation was
548 4 standard deviations greater than its MFR from the background period it was
549 classified as a sample-response cell. It should be noted that for the purpose of
550 this paper other FCTs such as those which respond to a specific lever (left/right)
551 or trial-type cells were not considered [69].

552 A.8 Laguerre Expansion Technique

553 In order to apply the Laguerre expansion technique [19], the input and output
554 data records were first convolved with the Laguerre functions:

$$v_{x_i}^{(l)} = \sum_{\tau=0}^M b_l(\tau) x_i(t - \tau) \quad (2)$$

555

$$v_y^{(l)} = \sum_{\tau=0}^M b_l(\tau) y(t - \tau) \quad (3)$$

556 where b_l is the l^{th} Laguerre basis function. By first convolving with the Laguerre
557 basis functions, the dynamical effects of the past input epochs are removed and
558 we are left with a simple regression of contemporaneous data. Substituting the
559 above equations into equation 1, we have:

$$y(t) = k_0 + \sum_{n=1}^N \sum_{l=1}^L c_{l,x_i}(l) v_{l,x_i}(t) + \sum_{l=1}^L c_{l,y}(l) v_{l,y}(t) \quad (4)$$

560 where c_{l,x_i} and $c_{l,y}$ are the feedforward and feedback Laguerre expansion coeffi-
561 cients. To estimate model parameters, eq. 4 was cast in matrix form:

$$\mathbf{y} = \mathbf{V}\mathbf{c} + \epsilon \quad (5)$$

562 where \mathbf{y} is the vector of all N output samples, \mathbf{V} is the design matrix consisting
563 of the convolved inputs, \mathbf{c} are the model parameters to be estimated, and ϵ is
564 the modeling error. Eq. 5 was solved using least squares regression (LSR). The
565 memory of our system was fixed at 300ms, in accordance with previous studies
566 [65, 70]. The Laguerre parameter α was fixed at 0.6 to reflect this system memory
567 [19].

568 A.9 Model Selection

569 In theory, the most predictive model would include all recorded inputs. However,
570 such a model would be susceptible to overfitting, and would not reveal which
571 neurons are causally connected to each other. To overcome this issue a forward
572 step-wise selection procedure was used to minimize overfitting and prune out
573 all inputs which are not causally related to the output [71]. Given an output
574 cell and M potential input cells recorded during the same session, the following
575 steps were used to select the N input cells which are causally connected to the
576 output cell. First, the data was divided into training (in-sample) and testing
577 (out-of-sample) sets. Then, M single-input single-output (SISO) models were
578 constructed with each of the potential inputs. The model whose predicted output
579 had the highest correlation, as measured by the Pearson correlation-coefficient,
580 ρ , with the actual output was selected. Afterwards, N-1 models were constructed
581 with two inputs: the previously selected input and one of the remaining potential
582 inputs. If any of the inputs were able to raise ρ , the input which raised ρ the

most was selected; otherwise, the procedure was ended, and only 1 input was selected. This procedure was repeated until either none of the inputs were able to raise ρ , or all M potential neurons were selected. The N selected neurons were then used as the model input.

A.10 Model Validation

To avoid overfitting, Monte Carlo style simulations were used to select those models which represent significant causal connections between input and output neurons and do not just fit noise [72]. The following procedure was used: in each run the real input was randomly permuted with respect to the output. A model was then generated between the permuted input and the real output, and the Pearson correlation coefficient, ρ_i , was obtained as a metric of performance. T=40 such simulations were conducted for each output and a set of performance metrics, $\{\rho_i\}_i^T$, was obtained. Then, using Fisher's transformation, we tested the hypothesis, H_0 , that ρ was within the population of $\{\rho_i\}$. If this hypothesis could be rejected at the 99.99% significance level, the model was deemed significant. The very conservative threshold ($P < .0001$) was used due to the large amount of comparisons being made.

A.11 Statistical Analysis

Unless otherwise noted, the unpaired Mann-Whitney U test was used to access whether significant differences exist between two samples. This test was used since it does not assume a normal distribution, and much of our data was found to be skewed/nonnormal. Shift estimates (Hodges-Lehman) and confidence intervals were estimated as prescribed by Higgins [73]. In order to estimate the scale estimate, or the ratio between two samples, the data was first log-transformed and then scale estimate was taken to be the antilog of the shift estimate. The χ^2 test was used to compare proportions.

In addition to the Pearson correlation coefficient, ρ , Receiver Operating Characteristic (ROC) curves were used to visualize model performance. ROC curves plot the true positive rate against the false positive rate over the putative range of threshold values for the continuous output, y [72]. The area under the curve (AUC) of ROC plots are used as a performance metric of the model, and have been shown to be equivalent to the Mann-Whitney two sample statistic [74]. The AUC ranges from 0 to 1, with 0.5 indicating a random predictor and higher values indicating better model performance. The ρ and AUC metrics were chosen as they measure the similarity between a continuous 'prethreshold' signal and a spike train. The continuous 'prethreshold' signal was chosen over adding a threshold trigger and comparing true output spike train with an output 'post-threshold' spike train for two reasons. First, this allows us to avoid specifying the threshold trigger value, which relies on the somewhat arbitrary tradeoff between true-positive and false-negative spikes [45]. Also, similarity metrics between two spike trains often require the specification of a 'binning parameter' to determine

624 the temporal resolution of the metric [75, 76].¹

625 B Supplementary Figures




Figure S1: Schematic of the DNMS task. First the rat is presented with one of two levers (sample presentation), which it presses (sample response). Then following a delay phase, the rat is presented with both levers (Nonmatch), of which it must press the opposite level from which it was presented in order to successfully complete the task.




Figure S2: CA3 spectra mean frequency and differences. Same format as Fig. 1e. A weak but significant trend was found for declining CA3 theta oscillations ($\Delta = 1.94\%$, $P = .045$).

¹I should probably add sections on how behavioral correlation analysis & FFT was done...




Figure S3: (A) A suggestive but insignificant relationship was found between the THC-induced decrease in the mean number of sample-presentation cells and behavioral performance ($R^2 = .265$, $P = .052$). (B) No relationship was found between reductions in CA1 theta power and behavioral impairment ($P = .67$). Format is same as Fig. 3.




Figure S4: (A) ROC plot (see supplementary methods) for model shown in Fig. 2 showing model predictive power. The light blue line ($TPR=FPR$) indicates a model with no predictive power. (B,C) Examples of Monte Carlo simulations: For each model, 40 surrogate models, with shuffled inputs were generated. The Fisher z-scores of these models, which are derived from ρ , were plotted as a histogram, while the true ρ value is the plotted dashed red line. The P value for the hypothesis that the true ρ value is greater than the simulated ρ values is printed above the graphs. Models were deemed significant if $P < .0001$. (B) shows the results for the model in Fig. 2, which was deemed significant. (C) shows an insignificant model




Figure S5: 4 additional systems are presented. Left column shows all system filters, including feedback filter (dashed line) in the time domain. Middle column shows the filters in the frequency domain and right column shows the ROC plots of the models. All these models were found to have significant predictive power in Monte Carlo tests.




Figure S6: Top Row: neither the first (middle column) nor third feedforward gPDM were found to be significantly correlated with THC induced behavioral deficits. Bottom Row: neither the first (middle column) nor second feedback gPDM were found to be significantly correlated with THC induced behavioral deficits. Format is same as Fig. 3.

References

- [1] Barbara S Koppel, John CM Brust, Terry Fife, Jeff Bronstein, Sarah Youssof, Gary Gronseth, and David Gloss. Systematic review: Efficacy and safety of medical marijuana in selected neurologic disorders report of the guideline development subcommittee of the american academy of neurology. *Neurology*, 82(17):1556–1563, 2014.
- [2] Liana Fattore. *Cannabinoids in Neurologic and Mental Disease*. Academic Press, 2015.
- [3] Melisa J Wallace, Jenny L Wiley, Billy R Martin, and Robert J DeLorenzo. Assessment of the role of cb 1 receptors in cannabinoid anticonvulsant effects. *European Journal of Pharmacology*, 428(1):51–57, 2001.
- [4] Robert E Blair, Laxmikant S Deshpande, Sompong Sombati, Katherine W Falenski, Billy R Martin, and Robert J DeLorenzo. Activation of the cannabinoid type-1 receptor mediates the anticonvulsant properties of cannabinoids in the hippocampal neuronal culture models of acquired epilepsy and status epilepticus. *Journal of Pharmacology and Experimental Therapeutics*, 317(3):1072–1078, 2006.
- [5] Laxmikant S Deshpande, Sompong Sombati, Robert E Blair, Dawn S Carter, Billy R Martin, and Robert J DeLorenzo. Cannabinoid cb1 receptor antagonists cause status epilepticus-like activity in the hippocampal neuronal culture model of acquired epilepsy. *Neuroscience letters*, 411(1):11–16, 2007.
- [6] V Rudenko, A Rafiuddin, JR Lehesté, and LK Friedman. Inverse relationship of cannabimimetic (r+) win 55, 212 on behavior and seizure threshold during the juvenile period. *Pharmacology Biochemistry and Behavior*, 100(3):474–484, 2012.
- [7] István Katona. Cannabis and endocannabinoid signaling in epilepsy. In *Endocannabinoids*, pages 285–316. Springer, 2015.
- [8] Andrew J Hill, TD Hill, and B Whalley. The development of cannabinoid based therapies for epilepsy. *Endocannabinoids: molecular, pharmacological, behavioral and clinical features*. bentham science publishers, Oak Park, IL, pages 164–204, 2013.
- [9] Raphael Mechoulam, Lumír O Hanuš, Roger Pertwee, and Allyn C Howlett. Early phytocannabinoid chemistry to endocannabinoids and beyond. *Nature Reviews Neuroscience*, 2014.
- [10] Emma Puighermanal, Arnau Busquets-Garcia, Rafael Maldonado, and Andrés Ozaita. Cellular and intracellular mechanisms involved in the cognitive impairment of cannabinoids. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 367(1607):3254–3263, 2012.

- 665 [11] Mathilde Metna-Laurent and Giovanni Marsicano. Rising stars: Modulation
666 of brain functions by astroglial type-1 cannabinoid receptors. *Glia*, 63(3):
667 353–364, 2015.
- 668 [12] Giovanni Bénard, Federico Massa, Nagore Puente, Joana Lourenço, Luigi
669 Bellocchio, Edgar Soria-Gómez, Isabel Matias, Anna Delamarre, Mathilde
670 Metna-Laurent, Astrid Cannich, et al. Mitochondrial cb1 receptors regulate
671 neuronal energy metabolism. *Nature Neuroscience*, 15(4):558–564, 2012.
- 672 [13] Wei Xiong, KeJun Cheng, Tanxing Cui, Grzegorz Godlewski, Kenner C
673 Rice, Yan Xu, and Li Zhang. Cannabinoid potentiation of glycine receptors
674 contributes to cannabis-induced analgesia. *Nature Chemical Biology*, 7(5):
675 296–303, 2011.
- 676 [14] Jessica A Fawley, Mackenzie E Hofmann, and Michael C Andresen.
677 Cannabinoid 1 and transient receptor potential vanilloid 1 receptors dis-
678 cretely modulate evoked glutamate separately from spontaneous glutamate
679 transmission. *The Journal of Neuroscience*, 34(24):8324–8332, 2014.
- 680 [15] Stephanie C Gantz and Bruce P Bean. Cell-autonomous excitation of
681 midbrain dopamine neurons by endocannabinoid-dependent lipid signaling.
682 *Neuron*, 2017.
- 683 [16] Moyra A Coull, Andrew T Johnston, Roger G Pertwee, and Stephen N
684 Davies. Action of δ -9-tetrahydrocannabinol on gaba a receptor-mediated
685 responses in a grease-gap recording preparation of the rat hippocampal slice.
686 *Neuropharmacology*, 36(10):1387–1392, 1997.
- 687 [17] Timothy M Brown, Jonathan M Brotchie, and Stephen M Fitzjohn.
688 Cannabinoids decrease corticostriatal synaptic transmission via an effect
689 on glutamate uptake. *The Journal of Neuroscience*, 23(35):11073–11077,
690 2003.
- 691 [18] Roman A Sandler, Dong Song, Robert E Hampson, Sam A Deadwyler,
692 Theodore W Berger, and Vasilis Z Marmarelis. Model-based asesment of an
693 in-vivo predictive relationship from ca1 to ca3 in the rodent hippocampus.
694 *Journal of Computational Neuroscience*, pages 1–15, 2014.
- 695 [19] Vasilis Z Marmarelis. *Nonlinear dynamic modeling of physiological systems*.
696 Wiley-Interscience, 2004.
- 697 [20] Dong Song, Vasilis Z Marmarelis, and Theodore W Berger. Parametric and
698 non-parametric modeling of short-term synaptic plasticity. part i: Compu-
699 tational study. *Journal of Computational Neuroscience*, 26(1):1–19, 2009.
- 700 [21] Dustin Fetterhoff, Ioan Opris, Sean L Simpson, Sam A Deadwyler, Robert E
701 Hampson, and Robert A Kraft. Multifractal analysis of information pro-
702 cessing in hippocampal neural ensembles during working memory under δ
703 9-tetrahydrocannabinol administration. *Journal of Neuroscience Methods*,
704 2014.

- [22] Dustin Fetterhoff, Robert A Kraft, Roman A Sandler, Ioan Opris, Cheryl A Sexton, Vasilis Z Marmarelis, Robert E Hampson, and Sam A Deadwyler. Distinguishing cognitive state with multifractal complexity of hippocampal interspike interval sequences. *Frontiers in Systems Neuroscience*, 9, 2015.
- [23] Robert E Hampson and Sam A Deadwyler. Cannabinoids reveal the necessity of hippocampal neural encoding for short-term memory in rats. *The Journal of Neuroscience*, 20(23):8932–8942, 2000.
- [24] David Robbe, Sean M Montgomery, Alexander Thome, Pavel E Rueda-Orozco, Bruce L McNaughton, and György Buzsáki. Cannabinoids reveal importance of spike timing coordination in hippocampal function. *Nature Neuroscience*, 9(12):1526–1533, 2006.
- [25] Anushka V Goonawardena, Lianne Robinson, Robert E Hampson, and Gernot Riedel. Cannabinoid and cholinergic systems interact during performance of a short-term memory task in the rat. *Learning & Memory*, 17(10):502–511, 2010.
- [26] Robert E Hampson, John D Simeral, and Sam A Deadwyler. Distribution of spatial and nonspatial information in dorsal hippocampus. *Nature*, 402(6762):610–614, 1999.
- [27] David Robbe and György Buzsáki. Alteration of theta timescale dynamics of hippocampal place cells by a cannabinoid is associated with memory impairment. *The Journal of Neuroscience*, 29(40):12597–12605, 2009.
- [28] Gyorgy Buzsaki. *Rhythms of the Brain*. Oxford University Press, 2006.
- [29] Laura Lee Colgin. Mechanisms and functions of theta rhythms. *Annual Review of Neuroscience*, 36:295–312, 2013.
- [30] György Buzsáki and Edvard I Moser. Memory, navigation and theta rhythm in the hippocampal-entorhinal system. *Nature Neuroscience*, 16(2):130–138, 2013.
- [31] James M Hyman, Eric A Zilli, Amanda M Paley, and Michael E Hasselmo. Working memory performance correlates with prefrontal-hippocampal theta interactions but not with prefrontal neuron firing rates. *Frontiers in Integrative Neuroscience*, 4, 2010.
- [32] Mihály Hajós, William E Hoffmann, and Bernát Kocsis. Activation of cannabinoid-1 receptors disrupts sensory gating and neuronal oscillation: relevance to schizophrenia. *Biological psychiatry*, 63(11):1075–1083, 2008.
- [33] N Spruston and C McBain. Structural and functional properties of hippocampal neurons. *The Hippocampus Book*, pages 133–201, 2007.
- [34] Thomas Klausberger and Peter Somogyi. Neuronal diversity and temporal dynamics: the unity of hippocampal circuit operations. *Science*, 321(5885):53–57, 2008.

- [35] Frédéric Pouille and Massimo Scanziani. Enforcement of temporal fidelity in pyramidal cells by somatic feed-forward inhibition. *Science*, 293(5532):1159–1163, 2001.
- [36] Rita Zemankovics, Judit M Veres, Iris Oren, and Norbert Hájos. Feedforward inhibition underlies the propagation of cholinergically induced gamma oscillations from hippocampal ca3 to ca1. *The Journal of Neuroscience*, 33(30):12337–12351, 2013.
- [37] L Stan Leung and Hui-Wen Yu. Theta-frequency resonance in hippocampal ca1 neurons in vitro demonstrated by sinusoidal current injection. *Journal of neurophysiology*, 79(3):1592–1596, 1998.
- [38] Bruce Hutcheon and Yosef Yarom. Resonance, oscillation and the intrinsic frequency preferences of neurons. *Trends in Neurosciences*, 23(5):216–222, 2000.
- [39] Bernat Kocsis, Anatol Bragin, and György Buzsáki. Interdependence of multiple theta generators in the hippocampus: a partial coherence analysis. *The Journal of neuroscience*, 19(14):6200–6212, 1999.
- [40] György Buzsáki. Theta oscillations in the hippocampus. *Neuron*, 33(3):325–340, 2002.
- [41] Romain Goutagny, Jesse Jackson, and Sylvain Williams. Self-generated theta oscillations in the hippocampus. *Nature Neuroscience*, 12(12), 2009.
- [42] Krisztina Monory, Martin Polack, Anita Remus, Beat Lutz, and Martin Korte. Cannabinoid cb1 receptor calibrates excitatory synaptic balance in the mouse hippocampus. *The Journal of Neuroscience*, 35(9):3842–3850, 2015.
- [43] SA Turkanis and R Karler. Central excitatory properties of δ 9-tetrahydrocannabinol and its metabolites in iron-induced epileptic rats. *Neuropharmacology*, 21(1):7–13, 1982.
- [44] Angela B Clement, E Gregory Hawkins, Aron H Lichtman, and Benjamin F Cravatt. Increased seizure susceptibility and proconvulsant activity of anandamide in mice lacking fatty acid amide hydrolase. *The Journal of Neuroscience*, 23(9):3916–3923, 2003.
- [45] Vasilis Z Marmarelis, Dae C Shin, Dong Song, Robert E Hampson, Sam A Deadwyler, and Theodore W Berger. Nonlinear modeling of dynamic interactions within neuronal ensembles using principal dynamic modes. *Journal of computational neuroscience*, 34(1):73–87, 2013.
- [46] Roman A Sandler and Vasilis Z. Marmarelis. Understanding spike triggered covariance using wiener theory for receptive field identification. *Journal of Vision*, in press.

- 782 [47] Sam A Deadwyler, James R West, Carl W Cotman, and Gary Lynch. Phys-
783 iological studies of the reciprocal connections between the hippocampus and
784 entorhinal cortex. *Experimental Neurology*, 49(1):35–57, 1975.
- 785 [48] Roman A Sandler, Dong Song, Robert E Hampson, Sam A Deadwyler,
786 Theodore W Berger, and Vasilis Z Marmarelis. Closed-loop hippocam-
787 pal modeling and the design of neurostimulation patterns for suppressing
788 seizures. *Journal of Neural Engineering*, Under Review.
- 789 [49] Roland SG Jones. Entorhinal-hippocampal connections: a speculative view
790 of their function. *Trends in Neurosciences*, 16(2):58–64, 1993.
- 791 [50] Omar J Ahmed and Mayank R Mehta. The hippocampal rate code:
792 anatomy, physiology and theory. *Trends in neurosciences*, 32(6):329–338,
793 2009.
- 794 [51] Roman A Sandler, Samuel A Deadwyler, Robert E Hampson, Dong Song,
795 Theodore W Berger, and Vasilis Z Marmarelis. System identification of
796 point-process neural systems using probability based volterra kernels. *Jour-
797 nal of Neuroscience Methods*, 240:179–192, 2015.
- 798 [52] Marianne J Bezaire and Ivan Soltesz. Quantitative assessment of cal lo-
799 cal circuits: Knowledge base for interneuron-pyramidal cell connectivity.
800 *Hippocampus*, 23(9):751–785, 2013.
- 801 [53] I Katona, B Sperlagh, Zs Maglóczy, E Santha, A Köfalvi, S Czirjak,
802 K Mackie, ES Vizi, and TF Freund. Gabaergic interneurons are the targets
803 of cannabinoid actions in the human hippocampus. *Neuroscience*, 100(4):
804 797–804, 2000.
- 805 [54] Ferenc Mátyás, Tamás F Freund, and Attila I Gulyás. Convergence of
806 excitatory and inhibitory inputs onto cck-containing basket cells in the cal
807 area of the rat hippocampus. *European Journal of Neuroscience*, 19(5):
808 1243–1256, 2004.
- 809 [55] Sang-Hun Lee, Csaba Földy, and Ivan Soltesz. Distinct endocannabinoid
810 control of gaba release at perisomatic and dendritic synapses in the hip-
811 pocampus. *Journal of Neuroscience*, 30(23):7993–8000, 2010.
- 812 [56] Maoxing Shen, Timothy M Piser, Virginia S Seybold, and Stanley A Thayer.
813 Cannabinoid receptor agonists inhibit glutamatergic synaptic transmission
814 in rat hippocampal cultures. *The Journal of Neuroscience*, 16(14):4322–
815 4334, 1996.
- 816 [57] Takako Ohno-Shosaku, Hiroshi Tsubokawa, Ichiro Mizushima, Norihide
817 Yoneda, Andreas Zimmer, and Masanobu Kano. Presynaptic cannabinoid
818 sensitivity is a major determinant of depolarization-induced retrograde sup-
819 pression at hippocampal synapses. *Journal of Neuroscience*, 22(10):3864–
820 3872, 2002.

- [58] Frauke Steindel, Raissa Lerner, Martin Häring, Sabine Ruehle, Giovanni Marsicano, Beat Lutz, and Krisztina Monory. Neuron-type specific cannabinoid-mediated g protein signalling in mouse hippocampus. *Journal of neurochemistry*, 124(6):795–807, 2013.
- [59] Sabine Ruehle, A Aparisi Rey, Floortje Remmers, and Beat Lutz. The endocannabinoid system in anxiety, fear memory and habituation. *Journal of Psychopharmacology*, 26(1):23–39, 2012.
- [60] Robert E Hampson and Sam A Deadwyler. Role of cannabinoid receptors in memory storage. *Neurobiology of disease*, 5(6):474–482, 1998.
- [61] Miles A Whittington and Roger D Traub. Interneuron diversity series: Inhibitory interneurons and network oscillations in vitro. *Trends in neurosciences*, 26(12):676–682, 2003.
- [62] Horacio G Rotstein, Dmitri D Pervouchine, Corey D Acker, Martin J Gillies, John A White, Eberhardt H Buhl, Miles A Whittington, and Nancy Kopell. Slow and fast inhibition and an h-current interact to create a theta rhythm in a model of ca1 interneuron network. *J Neurophysiol*, 94:1509–1518, 2005.
- [63] Can Tao, Guangwei Zhang, Ying Xiong, and Yi Zhou. Functional dissection of synaptic circuits: in vivo patch-clamp recording in neuroscience. *Frontiers in Neural Circuits*, 9:23, 2015.
- [64] Vasilis Z Marmarelis, Dae C Shin, Dong Song, Robert E Hampson, Sam A Deadwyler, and Theodore W Berger. On parsing the neural code in the prefrontal cortex of primates using principal dynamic modes. *Journal of Computational Neuroscience*, 36(3):321–337, 2014.
- [65] Dong Song, Rosa HM Chan, Vasilis Z Marmarelis, Robert E Hampson, Sam A Deadwyler, and Theodore W Berger. Nonlinear dynamic modeling of spike train transformations for hippocampal-cortical prostheses. *Biomedical Engineering, IEEE Transactions on*, 54(6):1053–1066, 2007.
- [66] George Paxinos and CH Watson. The rat brain in stereotaxic coordinates 2nd edn. *Academic Press, New York.*, 11(4):237–243, 1986.
- [67] Theodore W Berger, Dong Song, Rosa HM Chan, Vasilis Z Marmarelis, Jeff LaCoss, Jack Wills, Robert E Hampson, Sam A Deadwyler, and John J Granacki. A hippocampal cognitive prosthesis: multi-input, multi-output nonlinear modeling and vlsi implementation. *Neural Systems and Rehabilitation Engineering, IEEE Transactions on*, 20(2):198–211, 2012.
- [68] Sam A Deadwyler, Terence Bunn, and Robert E Hampson. Hippocampal ensemble activity during spatial delayed-nonmatch-to-sample performance in rats. *Journal of Neuroscience*, 16(1):354–372, 1996.
- [69] Robert E Hampson, Dong Song, Rosa HM Chan, Andrew J Sweatt, Mitchell R Riley, Anushka V Goonawardena, Vasilis Z Marmarelis, Greg A

- 860 Gerhardt, Theodore W Berger, and Sam A Deadwyler. Closing the loop
861 for memory prosthesis: Detecting the role of hippocampal neural ensembles
862 using nonlinear models. *Neural Systems and Rehabilitation Engineering*,
863 *IEEE Transactions on*, 20(4):510–525, 2012.
- 864 [70] Ude Lu, Dong Song, and Theodore W Berger. Nonlinear dynamic model-
865 ing of synaptically driven single hippocampal neuron intracellular activity.
866 *Biomedical Engineering, IEEE Transactions on*, 58(5):1303–1313, 2011.
- 867 [71] Dong Song, Rosa HM Chan, Vasilis Z Marmarelis, Robert E Hampson,
868 Sam A Deadwyler, and Theodore W Berger. Nonlinear modeling of neural
869 population dynamics for hippocampal prostheses. *Neural Networks*, 22(9):
870 1340–1351, 2009.
- 871 [72] Theodoros P Zanos, Spiros H Courellis, Theodore W Berger, Robert E
872 Hampson, Sam A Deadwyler, and Vasilis Z Marmarelis. Nonlinear model-
873 ing of causal interrelationships in neuronal ensembles. *Neural Systems and*
874 *Rehabilitation Engineering, IEEE Transactions on*, 16(4):336–352, 2008.
- 875 [73] James J. Higgins. *Intoruction to Modern Nonparametric Statistics*. 2003.
- 876 [74] James A Hanley and Barbara J McNeil. The meaning and use of the area
877 under a receiver operating characteristic (roc) curve. *Radiology*, 143(1):
878 29–36, 1982.
- 879 [75] Mark CW van Rossum. A novel spike distance. *Neural Computation*, 13
880 (4):751–763, 2001.
- 881 [76] Jonathan D Victor and Keith P Purpura. Metric-space analysis of spike
882 trains: theory, algorithms and application. *Network: computation in neural*
883 *systems*, 8(2):127–164, 1997.