


DIFFUSION DEPENDENT OVERDAMPED LANGEVIN DYNAMICS

Using a position dependent diffusion coefficient¹ D : favoring exploration in anisotropic or metastable potential landscapes \Rightarrow faster convergence to steady-state.

$$dq_t = \left(-D(q_t) \nabla V(q_t) + \frac{\text{div } D(q_t)}{\beta} \right) dt + \sqrt{2\beta^{-1}D(q_t)} dW_t \quad (1)$$

- **Aim:** Unbiased estimation of $\mathbb{E}(\varphi) = \int_{\Omega} \varphi(x) \pi(dx)$, $\pi = e^{-\beta V}$, $\beta^{-1} = k_B T$.



Error between empirical angle distribution and uniform distribution on $[0, 2\pi]$ after 10^5 iterations.


$$V(x, y) = 100(x^2 + y^2 - 1)^2$$

Anisotropic diffusion coefficient

$$D_{\text{Tangent}}(q) = \varepsilon I_2 + \tilde{q} \tilde{q}^T / \|q\|^2, \quad \tilde{q} = (-y \ x)^T$$

Isotropic diffusion coefficient

$$D_{\text{One}} \equiv (1 + \varepsilon) I_2, \quad \varepsilon = 0.1$$



Mean error for optimal time steps in HMC case.

- **Issue:** Rejection rates scales as $\mathcal{O}(\Delta t^{1/2})$ for Euler–Maruyama + Metropolis–Hastings.

BETTER UNBIASED NUMERICAL SAMPLING ?

- **Solution:** Riemann Manifold (Generalized) Hamiltonian Monte Carlo^{2,3} scheme based on Langevin dynamics integration.


$$\begin{cases} dq_t = \nabla_p H(q_t, p_t) dt \\ dp_t = -\nabla_q H(q_t, p_t) dt - \gamma \nabla_p H(q_t, p_t) dt + \sqrt{\frac{2\gamma}{\beta}} dW_t \end{cases}$$

- **Rejection rates scale as $\mathcal{O}(\Delta t^3)$.** But need to have a time-reversible and volume-preserving numerical integrator⁵.
- **Generalized Störmer–Verlet [GSV]: implicit** symplectic time-reversible integrator⁴

$$\begin{cases} p^{n+\frac{1}{2}} = p^n - \frac{\Delta t}{2} \nabla_q H(q^n, p^{n+\frac{1}{2}}) \\ q^{n+1} = q^n + \frac{\Delta t}{2} \left(\nabla_p H(q^n, p^{n+\frac{1}{2}}) + \nabla_p H(q^{n+1}, p^{n+\frac{1}{2}}) \right) \\ p^{n+1} = p^{n+\frac{1}{2}} - \frac{\Delta t}{2} \nabla_q H(q^{n+1}, p^{n+\frac{1}{2}}) \end{cases}$$

Proposition. GSV is first order weakly consistent with the O.L. dynamics with multiplicative noise (1) when using the Hamiltonian $H(q, p) = V(q) - \frac{1}{2} \ln(\det(D(q))) + \frac{1}{2} p^T D(q) p$.

Sampling results for a double-well confining potential $V(q) = q^2 - 1 + K e^{-q^2/(2\sigma)}$, oscillating diffusion coefficient $D(q) = \left(\frac{1 + \cos(\pi q)}{2} \right)^2$.



- **Unbiased sampling with Metropolis–Hastings scheme:** naive approach is **biased** !

IMPLICIT INTEGRATION

- Need to check for forward/backward convergence of implicit method and reversibility check when using Newton's/fixed-point iteration methods^{6,7}

- Starting from q_0 , given a time step Δt ,


1. Simulate $p_0 \sim \mathcal{N}(0, D(q_0)^{-1})$,
2. Integrate the Hamiltonian dynamics using GSV during Δt
 - if this forward integration does not converge, stay in place: return $(q_1, p_1) = (q_0, p_0)$
 - if it converges to (q_*, p_*) , integrate the Hamiltonian dynamics starting from $(q_*, -p_*)$:
 - if this backward integration does not converge, stay in place
 - else if the result differs from $(q_0, -p_0)$, stay in place

3. Apply the M–H procedure between (q_0, p_0) and (q_*, p_*) and return the position component.

GHMC can be recovered by integrating an Ornstein–Uhlenbeck process for the first step.

- **There are 4 ways to reject the proposal:** no forward/backward convergence, no numerical reversibility, M–H ratio computation.

- **Even for large time steps, unbiased sampling of the configuration space.**



REFERENCES

- [1] N. Bou-Rabee, A. Donev, and E. Vanden-Eijnden. "Metropolis integration schemes for self-adjoint diffusions". In: *Multiscale Modeling & Simulation* (2014).
- [2] S. Duane, A. D. Kennedy, B. J. Pendleton, and D. Roweth. "Hybrid Monte Carlo". In: *Physics Letters B* (1987).
- [3] M. Girolami and B. Calderhead. "Riemann Manifold Langevin and Hamiltonian Monte Carlo methods". In: *Journal of the Royal Statistical Society: Series B (Statistical Methodology)* (2011).
- [4] E. Hairer, G. Wanner, and C. Lubich. "Symplectic integration of hamiltonian systems". In: *Geometric Numerical Integration: Structure-Preserving Algorithms for Ordinary Differential Equations*. 2006.
- [5] T. Lelièvre, M. Rousset, and G. Stoltz. *Free Energy Computations: A Mathematical Perspective*. Imperial College Press, 2010.
- [6] T. Lelièvre, M. Rousset, and G. Stoltz. "Hybrid Monte Carlo methods for sampling probability measures on submanifolds". In: *Numerische Mathematik* (2019).
- [7] E. Zappa, M. Holmes-Cerfon, and J. Goodman. "Monte Carlo on manifolds: sampling densities and integrating functions". In: *Communications on Pure and Applied Mathematics* (2018).