

STAT432 Assignment 2

Rory Sarten 301005654

21 August 2020

Question 1

a)

These are independent binomial processes (despite happening in sequence), so the three trapping occasions can be denoted:

$$n_1 \sim \text{Binomial}(N, \theta) = \binom{N}{n_1} \theta^{n_1} (1 - \theta)^{N - n_1}$$

$$n_2 \sim \text{Binomial}(N - n_1, \theta) = \binom{N - n_1}{n_2} \theta^{n_2} (1 - \theta)^{N - n_1 - n_2}$$

$$n_3 \sim \text{Binomial}(N - n_1 - n_2, \theta) = \binom{N - n_1 - n_2}{n_3} \theta^{n_3} (1 - \theta)^{N - n_1 - n_2 - n_3}$$

So the probability of catching $n_1 + n_2 + n_3$ rats is given by

$$\text{Binomial}(N, \theta) \times \text{Binomial}(N - n_1, \theta) \times \text{Binomial}(N - n_1 - n_2, \theta)$$

b)

$$\begin{aligned} \ell(N, \theta) = & \ln \left[\frac{N!}{n_1(N - n_1)!} \theta^{n_1} (1 - \theta)^{N - n_1} \right. \\ & \times \frac{(N - n_1)!}{n_2(N - n_1 - n_2)!} \theta^{n_2} (1 - \theta)^{N - n_1 - n_2} \\ & \left. \times \frac{(N - n_1 - n_2)!}{n_3(N - n_1 - n_2 - n_3)!} \theta^{n_3} (1 - \theta)^{N - n_1 - n_2 - n_3} \right] \end{aligned}$$

$$\begin{aligned} \ell(N, \theta) = & \ln N! - [\ln n_1 + \ln(N - n_1)] + n_1 \ln \theta + (N - n_1) \ln(1 - \theta) + \\ & \ln(N - n_1)! - [\ln n_2 + \ln(N - n_1 - n_2)] + n_2 \ln \theta + (N - n_1 - n_2) \ln(1 - \theta) + \\ & \ln(N - n_1 - n_2)! - [\ln n_3 + \ln(N - n_1 - n_2 - n_3)] + n_3 \ln \theta + (N - n_1 - n_2 - n_3) \ln(1 - \theta) \end{aligned}$$

We can reduce the components

$$n_1 \ln \theta + n_2 \ln \theta + n_3 \ln \theta = (n_1 + n_2 + n_3) \ln \theta$$

$$(N - n_1) \ln(1 - \theta) + (N - n_1 - n_2) \ln(1 - \theta) + (N - n_1 - n_2 - n_3) \ln(1 - \theta) = (3N - 3n_1 - 2n_2 - n_3) \ln(1 - \theta)$$

$$\ln N! + \ln(N - n_1)! + \dots$$

Gives

$$\ln N! - \ln(N - n_1 - n_2 - n_3)! + (n_1 + n_2 + n_3) \ln \theta + (3N - 3n_1 - 2n_2 - n_3) \ln(1 - \theta)$$

finish that bit of algebra somehow

c)

```
llfunc <- function(par, n1, n2, n3) {  
  N <- par[1]  
  theta <- par[2]  
  lfactorial(N) - lfactorial(N - n1 - n2 - n3) +  
    sum(n1, n2, n3)*log(theta) +  
    (3*N - 3*n1 - 2*n2 - n3)*log(1 - theta)  
}
```

```

n1 <- 82L
n2 <- 54L
n3 <- 23L
## since N >= n1 + n2 + n3
N_0 <- n1 + n2 + n3
par_start <- c(N_0, 0.5)

optim_fit <- optim(par = par_start,
                  fn = llfunc,
                  n1 = n1, n2 = n2, n3 = n3,
                  lower = c(N_0, 1e-4),
                  upper = c(Inf, 1),
                  method = "L-BFGS-B",
                  control = list(fnscale = -1),
                  hessian = TRUE)

MLE <- optim_fit$par
SE <- sqrt(diag(solve(-optim_fit$hessian)))
LowerBound <- MLE - qnorm(0.975) * SE
UpperBound <- MLE + qnorm(0.975) * SE
results <- cbind(MLE, SE, LowerBound, UpperBound) %>%
  round(2)
rownames(results) <- c("N", "theta")
results

```

```

##           MLE      SE LowerBound UpperBound
## N       189.43 13.23      163.49      215.36
## theta    0.45  0.06        0.34        0.57

```

The maximum likelihood point $\hat{N} = 189.43$ with confidence interval (163.49, 215.36)

The maximum likelihood point $\hat{\theta} = 0.45$ with confidence interval (0.34, 0.57)

d)

Contour map of (N, θ) surface

```

n_vals <- seq(from = n1+n2+n3, to = 260, length = 100)
theta_vals <- seq(from = 0.2, to = 0.7, length = 100)

combos <- expand.grid(n_vals, theta_vals)
surface <- apply(expand.grid(n_vals, theta_vals), 1, llfunc, n1 = n1, n2 = n2, n3 = n3) %>%
  round(2)
outcome <- cbind(combos, surface)
names(outcome) <- c("N", "theta", "value")


ggplot(data = outcome, mapping = aes(N, theta)) +
  geom_contour(aes(z = value),
               breaks = seq(from = min(outcome$value),
                             to = max(outcome$value), by = 12)) +
  geom_contour(aes(z = value,
                  colour = factor(..level.. == max(surface) - 1.92,
                                levels = c(F, T),
                                labels = c("Single 95% CI "))),
               breaks = max(surface) - 1.92) +
  geom_contour(aes(z = value,

```

```

    colour = factor(..level.. == max(surface) - 3,
                    levels = c(F, T),
                    labels = c("Joint 95% CI ")),
    breaks = max(surface) - 3) +
geom_point(mapping = aes(x = MLE[1], y = MLE[2]), colour = "darkblue") +
labs(colour = "Of interest:") +
theme_minimal()

```



e)

$$\begin{aligned}\frac{\partial \ell}{\partial \theta} &= \frac{n_1 + n_2 + n_3}{\theta} - \frac{3N - 3n_1 - 2n_2 - n_3}{1 - \theta} \\ &= \frac{(n_1 + n_2 + n_3) - (n_1 + n_2 + n_3)\theta - (3N - 3n_1 - 2n_2 - n_3)\theta}{\theta(1 - \theta)}\end{aligned}$$

Setting this to 0

$$(n_1 + n_2 + n_3) - [(n_1 + n_2 + n_3) + (3N - 3n_1 - 2n_2 - n_3)]\theta = 0$$

$$[(n_1 + n_2 + n_3) + (3N - 3n_1 - 2n_2 - n_3)]\theta = (n_1 + n_2 + n_3)$$

$$\theta = \frac{(n_1 + n_2 + n_3)}{(n_1 + n_2 + n_3) + (3N - 3n_1 - 2n_2 - n_3)} = \frac{(n_1 + n_2 + n_3)}{3N - 3n_1 + n_1 - 2n_2 + n_2 - n_3 + n_3}$$

$$\hat{\theta} = \frac{n_1 + n_2 + n_3}{3N - 2n_1 - n_2}$$

f)

We have

$$\hat{\theta} = \frac{n_1 + n_2 + n_3}{3N - 2n_1 - n_2}$$

and

$$1 - \hat{\theta} = \frac{3N - 2n_1 - n_2}{3N - 2n_1 - n_2} - \frac{n_1 + n_2 + n_3}{3N - 2n_1 - n_2} = \frac{3N - 3n_1 - 2n_2 - n_3}{3N - 2n_1 - n_2}$$

So we can rewrite the original kernel of the log likelihood in terms of N as:

$$\begin{aligned}\ell(N) &= \ln N! - \ln(N - n_1 - n_2 - n_3)! + (n_1 + n_2 + n_3) \ln \left(\frac{n_1 + n_2 + n_3}{3N - 2n_1 - n_2} \right) + \\ &\quad (3N - 3n_1 - 2n_2 - n_3) \ln \left(\frac{3N - 3n_1 - 2n_2 - n_3}{3N - 2n_1 - n_2} \right)\end{aligned}$$

$$\begin{aligned}\ell(N) &= \ln N! - \ln N! - \ln(N - n_1 - n_2 - n_3)! + \\ &\quad (n_1 + n_2 + n_3) \ln(n_1 + n_2 + n_3) - (n_1 + n_2 + n_3) \ln(3N - 2n_1 - n_2) + \\ &\quad (3N - 3n_1 - 2n_2 - n_3) \ln(3N - 3n_1 - 2n_2 - n_3) - (3N - 3n_1 - 2n_2 - n_3) \ln(3N - 2n_1 - n_2)\end{aligned}$$

$$\begin{aligned}\ell(N) &= \ln N! - \ln N! - \ln(N - n_1 - n_2 - n_3)! + \\ &\quad (3N - 3n_1 - 2n_2 - n_3) \ln(3N - 3n_1 - 2n_2 - n_3) - \\ &\quad (n_1 + n_2 + n_3 + 3N - 3n_1 - 2n_2 - n_3) \ln(3N - 2n_1 - n_2)\end{aligned}$$

Therefore

$$\begin{aligned}\ell(N) &= \ln N! - \ln N! - \ln(N - n_1 - n_2 - n_3)! + \\ &\quad (3N - 3n_1 - 2n_2 - n_3) \ln(3N - 3n_1 - 2n_2 - n_3) - \\ &\quad (3N - 2n_1 - n_2) \ln(3N - 2n_1 - n_2)\end{aligned}$$

g)

```
llfunc_N <- function(N, n1, n2, n3) {  
  lfactorial(N) - lfactorial(N - n1 - n2 - n3) +  
    (3*N - 3*n1 - 2*n2 - n3)*log(3*N - 3*n1 - 2*n2 - n3) -  
    (3*N - 2*n1 - n2)*log(3*N - 2*n1 - n2)  
}  
  
n1 <- 82L  
n2 <- 54L  
n3 <- 23L  
## since N >= n1 + n2 + n3  
N_0 <- n1 + n2 + n3  
N_start <- c(N_0)  
  
optim_fit <- optim(par = N_start,  
  fn = llfunc_N,  
  n1 = n1, n2 = n2, n3 = n3,  
  lower = c(N_0, 1e-4),  
  upper = c(Inf, 1),  
  method = "L-BFGS-B",  
  control = list(fnscale = -1),  
  hessian = TRUE)  
  
N_hat <- optim_fit$par  
log_max <- optim_fit$value  
  
## calculate 95% CI  
crit_point <- log_max - 0.5 * qchisq(0.95, df = 1)  
  
ci_func <- function(critical_value, ...) {  
  llfunc_N(...) - critical_value  
}  
  
lower_ci <- uniroot(ci_func, interval = c(N_0, N_hat),  
  n1 = n1, n2 = n2, n3 = n3,  
  critical_value = crit_point)$root  
upper_ci <- uniroot(ci_func, interval = c(N_hat, 300),  
  n1 = n1, n2 = n2, n3 = n3,  
  critical_value = crit_point)$root  
  
cbind(N_hat, lower_ci, upper_ci)
```

```
##           N_hat lower_ci upper_ci  
## [1,] 189.4255 170.918 229.9337
```

The univariate log-likelihood estimate for $\hat{N} = 189.4254905$


with 95% confidence interval (170.92, 229.93).

Compared with the confidence interval in (c), this interval is similar in size, but is no longer symmetric around the MLE.

```
n_vals <- seq(from = 163, to = 260, length = 100)  
data <- data.frame(x = n_vals,  
  y = llfunc_N(n_vals, n1 = n1, n2 = n2, n3 = n3))
```

```
ggplot(data = data, mapping = aes(x = x, y = y)) +
  geom_line() +
  geom_point(aes(x = N_hat, y = log_max, color = "red")) +
  geom_text(aes(x = N_hat + 0.4, y = log_max + 0.4, label = "MLE")) +
  geom_hline(yintercept=crit_point, linetype="dashed", color = "red") +
  geom_vline(xintercept=lower_ci, linetype="dashed", color = "red") +
  geom_vline(xintercept=upper_ci, linetype="dashed", color = "red") +
  geom_point(aes(x = lower_ci, y = crit_point, color = "red")) +
  geom_point(aes(x = upper_ci, y = crit_point, color = "red")) +
  geom_text(aes(x = lower_ci + 5, y = crit_point - 0.4, label = "Lower")) +
  geom_text(aes(x = upper_ci - 5, y = crit_point - 0.4, label = "Upper")) +
  ggtitle("Confidence interval boundaries for MLE of Univariate log-likelihood for N") +
  xlab("N") + ylab("Log-likelihood") +
  theme(legend.position="none")
```

Confidence interval boundaries for MLE of Univariate log-likelihood for N



h)

Given the 95% profile likelihood-ratio for N calculated above, and that we know a total of 159 rats were caught over the three occasions, we can our confidence interval for R to be $(lower - 159, upper - 159) = (171 - 159, 230 - 159) = (12, 71)$ (with rounding to the nearest whole rat).

Question 2

a)

$$\begin{aligned}
\ell(\lambda, x) &= \log\left(\prod_{i=1}^n \left(\frac{e^{-\lambda} \lambda^{x_i}}{x_i!}\right)\right) \\
&= \sum_{i=1}^n (-\lambda + x_i \ln \lambda - \ln(x_i!)) \\
&= -n\lambda + \left(\sum_{i=1}^n x_i\right) \ln \lambda - \sum_{i=1}^n \ln(x_i!)
\end{aligned}$$

The last term can be dropped, giving the kernel of the log likelihood

$$-n\lambda + \left(\sum_{i=1}^n x_i\right) \ln \lambda$$

Differentiating gives

$$\frac{d\ell}{d\lambda} = -n + \frac{\sum_{i=1}^n x_i}{\lambda}$$

Setting this to 0 gives

$$\hat{\lambda} = \frac{\sum_{i=1}^n x_i}{n}$$

We can calculate the standard error by taking the second derivative of the log-likelihood at the MLE and calculating the negative inverse

$$\frac{d^2\ell}{d\lambda^2} = -\frac{\sum_{i=1}^n x_i}{\lambda^2}$$

$$\left.\frac{d^2\ell}{d\lambda^2}\right|_{\hat{\lambda}} = -\left[\frac{n}{\sum_{i=1}^n x_i}\right]^2 \times \sum_{i=1}^n x_i = -\frac{n^2}{\sum_{i=1}^n x_i}$$

The negative inverse is

$$\frac{\sum_{i=1}^n x_i}{n^2}$$

So we can calculate the standard error as

$$SE[\hat{\lambda}] = \frac{\sqrt{\sum_{i=1}^n x_i}}{n}$$

```
counts <- readr::read_rds("counts.rds")
MLE <- sum(counts)/length(counts) # or mean(counts)
SE <- (sqrt(sum(counts))/length(counts)) %>% round(digits = 2)
LOWER <- (MLE - SE * 1.96) %>% round(digits = 2)
UPPER <- (MLE + SE * 1.96) %>% round(digits = 2)
cbind(MLE, SE, LOWER, UPPER)
```

```
##      MLE    SE LOWER UPPER
## [1,] 4.66 0.31  4.05  5.27
```

```

poisson_llfunc <- function(lambda, counts) {
  -(length(counts) * lambda) + sum(counts) * log(lambda)
}

optim_fit <- optim(par = 1,
  fn = poisson_llfunc,
  counts = counts,
  lower = 1e-3,
  upper = Inf,
  method = "L-BFGS-B",
  control = list(fnscale = -1),
  hessian = TRUE)
MLE <- optim_fit$par

n_vals <- seq(from = MLE - 1, to = MLE + 1, length = 100)
data <- data.frame(x = n_vals,
  y = poisson_llfunc(n_vals, counts))

log_max <- optim_fit$value
crit_point <- optim_fit$value - 0.5 * qchisq(0.95, df = 1)
ci_func <- function(critical_point, ...) poisson_llfunc(...) - critical_point

lower_ci <- uniroot(ci_func,
  interval = c(0, MLE),
  counts = counts,
  critical_point = crit_point)$root
upper_ci <- uniroot(ci_func,
  interval = c(MLE, MLE*5),
  counts = counts,
  critical_point = crit_point)$root
cbind(MLE, lower_ci, upper_ci)

```

```


##      MLE lower_ci upper_ci
## [1,] 4.66 4.086982 5.284223

```

```

ggplot(data = data, mapping = aes(x = x, y = y)) +
  geom_line() +
  geom_point(aes(x = MLE, y = log_max, color = "red")) +
  geom_text(aes(x = MLE, y = log_max + 0.4, label = "MLE")) +
  geom_hline(yintercept=crit_point, linetype="dashed", color = "red") +
  geom_vline(xintercept=lower_ci, linetype="dashed", color = "red") +
  geom_vline(xintercept=upper_ci, linetype="dashed", color = "red") +
  geom_point(aes(x = lower_ci, y = crit_point, color = "red")) +
  geom_point(aes(x = upper_ci, y = crit_point, color = "red")) +
  geom_text(aes(x = lower_ci + 0.13, y = crit_point - 0.4, label = "Lower")) +
  geom_text(aes(x = upper_ci - 0.13, y = crit_point - 0.4, label = "Upper")) +
  ggtitle(expression(paste("Likelihood ratio confidence interval boundaries for MLE of ", lambda))) +
  xlab(expression(lambda)) + ylab("Log-likelihood") +
  theme(legend.position="none")

```

c)

d)

$$\ell(\boldsymbol{\lambda}, \boldsymbol{\pi} | \mathbf{x}) = \sum_{i=1}^n \ln \left(\sum_{g=1}^G \pi_g \frac{e^{-\lambda} \lambda^{x_i}}{x_i!} \right)$$

$$\ell_C(\boldsymbol{\lambda}, \boldsymbol{\pi} | \mathbf{x}, Z) = \sum_{i=1}^n \ln \left(\sum_{g=1}^G z_{ig} \pi_g \frac{e^{-\lambda} \lambda^{x_i}}{x_i!} \right) \text{ where } z_{ig} \text{ has value 0 or 1}$$

ii)

If we are looking at each $g = 1, \dots, G$ then we can ignore the π_g term, as we are not concerned with the proportion in the group g . We can also exploit that z_{ig} is 0 or 1, by treating it as an exponent.

$$\begin{aligned} \ell_C(\boldsymbol{\lambda}, \boldsymbol{\pi} | \mathbf{x}, Z) &= \sum_{i=1}^n \ln \left(z_i \frac{e^{-\lambda} \lambda^{x_i}}{x_i!} \right) \\ &= \sum_{i=1}^n \ln \left(\left[\frac{e^{-\lambda} \lambda^{x_i}}{x_i!} \right]^{z_i} \right) \\ &= \sum_{i=1}^n z_i \ln \left(\frac{e^{-\lambda} \lambda^{x_i}}{x_i!} \right) \\ &= \sum_{i=1}^n z_i (-\lambda + x_i \ln \lambda - \ln x_i!) \\ &= -\lambda \sum_{i=1}^n z_i + \sum_{i=1}^n z_i x_i \ln \lambda - \sum_{i=1}^n z_i \ln x_i! \end{aligned}$$

$$\frac{\partial \ell_C}{\partial \lambda} = - \sum_{i=1}^n z_i + \frac{\sum_{i=1}^n z_i x_i}{\lambda}$$

Setting equal to 0 and solving

$$\frac{\sum_{i=1}^n z_i x_i}{\lambda} = \sum_{i=1}^n z_i$$

therefore for each group g

$$\hat{\lambda}_g = \frac{\sum_{i=1}^n z_{ig} x_i}{\sum_{i=1}^n z_{ig}}$$

iv)

```
parameters <- list(g1 = c(lambda_est = 3, pi_est = 0.5),
                  g2 = c(lambda_est = 1, pi_est = 0.5))

## start with high max distance from parameters
previous_parameters <- list(g1 = c(lambda_est = 30, pi_est = 0.5),
                           g2 = c(lambda_est = 1, pi_est = 0.5))

z_col <- function(parameters, x)
  parameters["pi_est"] * exp(-parameters["lambda_est"]) * parameters["lambda_est"]^x

z_matrix <- function(parameters, x) {
  Z <- lapply(parameters, z_col, x) %>%
    data.frame() %>%
    as.matrix()
  Z/apply(Z, 1, sum)
}

calc_lambda <- function(Z, x)
  as.data.frame(Z) %>% `*`(x) %>% apply(2, sum) %>% `/(`(apply(Z, 2, sum))

not_converged <- function(current_para, previous_para, threshold = 0.01)
  any(max(abs(unlist(current_para) - unlist(previous_para))) > threshold)

while (not_converged(parameters, previous_parameters)) {

  previous_parameters <- parameters
  ## E-step - estimate Z from current parameters
  Z <- z_matrix(parameters, counts)

  ## M-step - update estimates for parameters
  new_pi <- apply(Z, 2, mean)
  new_lambda <- calc_lambda(Z, counts)
  parameters <- purrr::map2(parameters, new_pi, function(x, y) {x["pi_est"] <- y; x})
  parameters <- purrr::map2(parameters, new_lambda, function(x, y) {x["lambda_est"] <- y; x})
}

parameters

## $g1
## lambda_est      pi_est
## 7.0142441 0.4809512
##
```

```
## $g2
## lambda_est      pi_est
## 2.4785544 0.5190488
```

```
as.data.frame(Z) %>% lapply(function(x) as.integer(x >= 0.5)) %>% as.data.frame() %>% as.matrix()
```

```
##      g1 g2
## [1,]  1  0
## [2,]  1  0
## [3,]  0  1
## [4,]  1  0
## [5,]  0  1
## [6,]  1  0
## [7,]  0  1
## [8,]  1  0
## [9,]  1  0
## [10,] 0  1
## [11,] 0  1
## [12,]  1  0
## [13,]  1  0
## [14,]  0  1
## [15,]  1  0
## [16,]  0  1
## [17,]  0  1
## [18,]  1  0
## [19,]  0  1
## [20,]  1  0
## [21,]  0  1
## [22,]  1  0
## [23,]  0  1
## [24,]  0  1
## [25,]  0  1
## [26,]  0  1
## [27,]  0  1
## [28,]  1  0
## [29,]  0  1
## [30,]  1  0
## [31,]  1  0
## [32,]  1  0
## [33,]  0  1
## [34,]  1  0
## [35,]  0  1
## [36,]  0  1
## [37,]  0  1
## [38,]  0  1
## [39,]  1  0
## [40,]  0  1
## [41,]  0  1
## [42,]  0  1
## [43,]  1  0
## [44,]  1  0
## [45,]  0  1
## [46,]  1  0
## [47,]  0  1
## [48,]  1  0
```

```
## [49,] 0 1
## [50,] 1 0
```

```
pandoc -s assignment2.Rmd -t latex -o assignment.pdf --highlight-style=tango -S "pandoc -s", paste0(filen, "-out.md"), " -t latex -o ", paste0(filen, ".pdf"), " --highlight-style=tango -S"
```