

# On the Trajectories of the $3x + 1$ Problem

Roy Burson

July 27, 2019

## Abstract

This paper studies certain trajectories of the Collatz function. I show that if for each odd number  $n$ ,  $n \sim 3n + 2$  then every positive integer  $n \in \mathbb{N} \setminus 2^{\mathbb{N}}$  has the representation

$$n = \left( 2^{a_{k+1}} - \sum_{i=0}^k 2^{a_i} 3^{k-i} \right) / 3^{k+1}$$

where  $0 \leq a_0 \leq a_1 \leq \dots \leq a_{k+1}$ . As a consequence, in order to prove Collatz Conjecture I illustrate that it is sufficient to prove  $n \sim 3n + 2$  for any odd  $n \in \mathbb{N} \setminus 2^{\mathbb{N}}$ . This is the main result of the paper.

## 1. Introduction

Some problems in mathematics are easy to state but take very complex tools to prove and often it takes new tools to be developed. The *Collatz Conjecture* is either one of this type or else it might be a conspiracy theory to slow down the field of mathematics (a joke that spread across Yale University). The *Collatz Conjecture* is a well known unsolved mathematical problem that concerns the recursive behavior of the function

$$T(n) = \begin{cases} \frac{n}{2} & \text{if } n \equiv 0(\text{mod}2) \\ 3n + 1 & \text{if } n \equiv 1(\text{mod}2) \end{cases}$$

over the set of integers  $\mathbb{Z}$ . In this paper I focus on the Collatz function over the positive whole numbers  $\mathbb{N}$ . The problem is most commonly referred to as the " $3x + 1$ " *problem*. The history and exact origin of the problem is somewhat vague. Some early history on the problem is discussed by Jeffrey C. Lagarias at <http://www.cec.m.sfu.ca/organics/papers/lagarias/paper/html/node1.html>. Lagarias gives 197 different documentations on the topic in his annotated bibliographies [10] and [11]. Dr. Lothar Collatz is credited for the discovery of the problem during his career as a student. Dr. Lothar Collatz even asserts himself that he was the first to study this problem in his letter <http://www.cec.m.sfu.ca/organics/papers/lagarias/paper/html/letter.html>.

The problem states that  $\forall n \in \mathbb{N}$  there is a value  $k \in \mathbb{N}$  such that  $T^k(n) = 1$  were  $T^k$  is the  $k$ th application of the map  $T$ . For example, for the value  $n = 11$  we can view the iterations with arrows (to indicate direction) as followed:

$$11 \rightarrow 34 \rightarrow 17 \rightarrow 52 \rightarrow 26 \rightarrow 13 \rightarrow 40 \rightarrow 20 \rightarrow 10 \rightarrow 5 \rightarrow 16 \rightarrow \dots \rightarrow 1$$

We can also visualize this backwards by reversing the operations of the map and traces our steps in the reverse direction. Doing so for this example we have the following:

$$\begin{aligned} 1 &\rightarrow \dots \rightarrow \frac{2^4 - 1}{3} \rightarrow \frac{2^5 - 2}{3} \rightarrow \frac{2^6 - 2^2}{3} \rightarrow \frac{2^7 - 2^3}{3} \rightarrow \frac{2^7 - 2^3 - 3}{3^2} \\ &\rightarrow \frac{2^8 - 2^4 - 2 \cdot 3}{3^2} \rightarrow \frac{2^9 - 2^5 - 2^2 \cdot 3}{3^2} \rightarrow \frac{2^9 - 2^5 - 2^2 \cdot 3 - 3^2}{3^3} \\ &\rightarrow \frac{2^{10} - 2^6 - 2^3 \cdot 3 - 2 \cdot 3^2}{3^3} \rightarrow \frac{2^{10} - 2^6 - 2^3 \cdot 3 - 2 \cdot 3^2 - 3^3}{3^4} = 11 \end{aligned}$$

it has been verified that all natural numbers  $n < 87 \cdot 2^{60}$  iterate to the value 1 under Collatz function. A neat discussion about the empirical results and the record-holders are discussed by Tomás Oliveria e Silvia at his home page <http://sweet.ua.pt/tos/3x+1.html>, and by Eric Roosendaal <http://www.ericr.nl/wondrous/index.html>. A complete list of the record

holders can be found here <http://www.ericr.nl/wondrous/pathrecs.html> which was accomplished by the yoyo@home project in 2017. In [10] and [11] Lagarias also mentioned the empirical evidence of the problem given by Oliveira e Silva. An interesting study on properties of divergent trajectories of the Collatz function (if one exist) can be found here <http://www.csun.edu/~vcmth02i/Collatz.pdf>. A similar result of this paper is given by Charles C. Cadogan in his works [5] and [4], which can also be found in Lagarias's bibliography [11]. Some more interesting discussions and findings that relate to this work can be found in [6, 9, 12, 13, 14].

## 2. Terminology

In order to prove the main result of the paper the following definitions are needed.

**Definition 1.** *let  $m \in \mathbb{N}$ . Write  $m \equiv \mathcal{R}$  if and only if the number  $m$  has the representation*

$$m = \left( 2^{a_{k+1}} - \sum_{i=0}^k 2^{a_i} 3^{k-i} \right) / 3^{k+1}$$

*where  $(a_i)$  is a monotonically increasing sequence of positive integers for  $i \leq k+1$ .*

**Definition 2.** *The Trajectory or Forward Orbit of a positive integer  $n$  is the set*

$$O^+(n) = \{n, T(n), T^2(n), \dots\}$$

*where  $T : \mathbb{N} \rightarrow \mathbb{N}$  is the Collatz function defined by  $n \rightarrow \frac{n}{2}$  if  $n$  is even and  $n \rightarrow 3n + 1$  if  $n$  is odd, and  $T^k$  is the function  $T : \mathbb{N} \times \mathbb{N} \rightarrow \mathbb{N}$  defined by*

$$T^k(n) = \underbrace{(T \circ T \circ \dots \circ T)}_{k\text{-times}}$$

**Definition 3.** *Given two integers  $n_1$  and  $n_2$  define the relation  $n_1 \sim n_2$  if and only if the two trajectories  $O^+(n_1)$  and  $O^+(n_2)$  coalesce, i.e.  $O^+(n_1) \cap O^+(n_2) \neq \emptyset$ .*

### 3. Backwards Iterations and Integer Representations

As shown by [1, 2, 3, 7, 15], and discussed in [8] we have that

$$\forall m \in \mathbb{N} \setminus 2^{\mathbb{N}}, 1 \in O^+(m) \iff m = \left( 2^{a_{k+1}} - \sum_{i=0}^k 2^{a_i} 3^{k-i} \right) / 3^{k+1}$$

for some sequence  $(a_j)$  where  $0 \leq a_0 \leq a_1 \leq a_2 \leq \dots \leq a_{k+1}$ . If  $m$  has this representation given above then  $m$  cannot be a power of 2 because any element of the set  $2^{\mathbb{N}} = \{2, 2^2, 2^3, \dots\}$  cannot be represented in this form. This can easily be verified by observing that the numbers that can be written in the form  $\left( 2^{a_{k+1}} - \sum_{i=0}^k 2^{a_i} 3^{k-i} \right) / 3^{k+1}$  are exactly those numbers obtained by iterating the functions  $n \rightarrow 2n$  and  $n \rightarrow \frac{n-1}{3}$  in succession, beginning at the number  $n = 1$ , where as each function is applied at least once and the function  $n \rightarrow \frac{n-1}{3}$  is never applied twice in succession. The numbers in the set  $2^{\mathbb{N}}$  can only be obtained by iterating the function  $n \rightarrow 2n$ , beginning at  $n = 1$ , so that the function  $n \rightarrow \frac{n-1}{3}$  is never applied. This shows that no integer in the set  $2^{\mathbb{N}}$  has the representation

$$\left( 2^{a_{k+1}} - \sum_{i=0}^k 2^{a_i} 3^{k-i} \right) / 3^{k+1}$$

The next lemma shows that if a number  $n$  has the representation  $n = \left( 2^{a_{k+1}} - \sum_{i=0}^k 2^{a_i} 3^{k-i} \right) / 3^{k+1}$  then the number  $2n$  and  $\frac{2n-1}{3}$  must also have the same representation.

**Lemma 1.** (*Closure of  $\mathcal{R}$* ) If  $n \equiv \mathcal{R}$  then  $2n \equiv \mathcal{R}$  and if  $n \equiv \mathcal{R}$  then  $\frac{2n-1}{3} \equiv \mathcal{R}$ .

*Proof.* (Direct proof) First suppose  $n \equiv \mathcal{R}$ . Then there is a sequence  $(a_j)$  so that

$$n = \left( 2^{a_{k+1}} - \sum_{i=0}^k 2^{a_i} 3^{k-i} \right) / 3^{k+1}$$

where  $0 \leq a_0 \leq a_1 \leq a_2 \leq \dots \leq a_{k+1}$ . Write

$$2n = 2 \left( \left( 2^{a_{k+1}} - \sum_{i=0}^k 2^{a_i} 3^{k-i} \right) / 3^{k+1} \right) = \left( 2^{a_{k+1}+1} - \sum_{i=0}^k 2^{a_i+1} 3^{k-i} \right) / 3^{k+1}$$

Define the sequence  $(b_j)$  by  $b_j = a_{j+1} + 1$  for each  $0 \leq j \leq k$  so that  $(b_j)$  is also positive and increasing. Then we have

$$2n = \left( 2^{b_j} - \sum_{i=0}^k 2^{b_i} 3^{k-i} \right) / 3^{k+1} \equiv \mathcal{R}$$

Therefore  $2n \equiv \mathcal{R}$ . Now suppose that  $n \equiv \mathcal{R}$  then by the above  $2n \equiv \mathcal{R}$ . Write

$$2n = \left( 2^{a_{k+1}} - \sum_{i=0}^k 2^{a_i} 3^{k-i} \right) / 3^{k+1}$$

were as  $1 \leq a_0 \leq a_1 \leq \dots \leq a_{k+1}$ . Then

$$\frac{2n-1}{3} = \frac{\left( 2^{a_{k+1}} - \sum_{i=0}^k 2^{a_i} 3^{k-i} \right) / 3^{k+1} - 1}{3}$$

Define the sequence  $(b_j)$  by

$$b_j = \begin{cases} 0 & \text{if } j = 0 \\ a_{j-1} & \text{if } 1 \leq j \leq k+1 \end{cases}$$

Then

$$\frac{2n-1}{3} = \left( 2^{b_{k+1}} - \sum_{i=0}^{k+1} 2^{b_i} 3^{k+1-i} \right) / 3^{k+2}$$

and therefore  $\frac{2n-1}{3} \equiv \mathcal{R}$

□

**Lemma 2.**

$$3^{\frac{a}{2}+1} + 2 < 2^a + 1$$

for all  $a \geq 8$ .

*Proof.* (proof by induction) The proof follows by induction. Let

$$S = \{1, 2, 3, \dots, 7\} \cup \{a \in \mathbb{N} : 3^{\frac{a}{2}+1} + 2 \leq 2^a + 1\}$$

The value  $a = 8$  is in  $S$  since

$$245 = 3^5 + 2 < 2^8 + 1 = 257$$

Now if  $a \geq 8 \in S$  then

$$\begin{aligned} 3^{\frac{a}{2}+1} + 2 &\leq 2^a + 1 \Rightarrow 3^{\frac{1}{2}}(3^{\frac{a}{2}+1} + 2) \leq 2(2^a + 1) \\ \therefore 3^{\frac{1}{2}}(3^{\frac{1}{2}(a+2)} + 2) &\leq 2(2^a + 1) \\ \therefore 3^{\frac{1}{2}(a+3)} + 2 \cdot 3^{\frac{1}{2}} &\leq 2^{a+1} + 2 \\ \Rightarrow 3^{\frac{1}{2}(a+3)} + 3 &\leq 2^{a+1} + 2 \\ \therefore 3^{\frac{1}{2}(a+3)} + 2 &\leq 2^{a+1} + 1 \\ \therefore 3^{\frac{a+1}{2}+1} + 2 &\leq 2^{a+1} + 1 \end{aligned}$$

Therefore  $a + 1 \in S$  so that  $S = \mathbb{N}$ .  $\square$

**Lemma 3.** Assume that  $a \in \mathbb{N}$ . If  $a$  is even, then there exist a value  $k$  such that  $T^k(2^a + 1) = 3^{\frac{a}{2}} + 1$ . If  $a$  is a odd, then there exist a value  $k$  such that  $T^k(2^a + 1) = 3^{\lfloor \frac{a}{2} \rfloor + 1} + 2$ .

*Proof.* (Direct proof) First suppose  $a$  is even. For any  $a$  apply the map  $T$  and note that  $T(2^a + 1) = 3 \cdot 2^a + 2^2$ . Then since  $a$  is even apply  $T$  twice in succession to obtain  $T^3(2^a + 1) = 32^{a-2} + 1$ . Now notice that every 3 successions of the map  $T$  takes away a value of 2 from the power  $a$  and increasing the coefficient of  $2^a$  by a multiple of 3. Noticing this pattern simply take  $k = 3\frac{a}{2}$  then we obtain  $T^k(2^a + 1) = 3^{\frac{a}{2}} + 1$ .

The second part of the proof is similar. If  $a$  is odd then we cannot apply the map  $T$   $3\frac{a}{2}$  times by the simply fact that  $3\frac{a}{2}$  is is not an integer and the map  $T^k$  only takes in positive integers for  $k$ . Whence, we can only apply the function  $T$   $3\lfloor \frac{a}{2} \rfloor$  times and then once more in addition to be rid of all the powers of 2 in the expression  $2^a + 1$ . So simply take  $k = 3\lfloor \frac{a}{2} \rfloor + 1$  then  $T^k(2^a + 1) = 3^{\lfloor \frac{a}{2} \rfloor + 1} + 2$   $\square$

**Theorem 1.** If for all  $n \in \mathbb{N} \setminus 2^{\mathbb{N}}$   $n \sim 3n + 2$ , then for all  $n \in \mathbb{N} \setminus 2^{\mathbb{N}}$  it follows that  $n \equiv \mathcal{R}$

*Proof.* (Proof by induction) First suppose for all  $n \in \mathbb{N} \setminus 2^{\mathbb{N}}$  that  $n \sim 3n + 2$ . Let  $X = \{x_1, x_2, \dots\} = \mathbb{N} \setminus 2^{\mathbb{N}}$  were as  $x_1 < x_2 < x_3 < \dots$  and let  $S$  be the

set defined by  $S = \{n \in \mathbb{N} : x_n \equiv \mathcal{R}\}$ . Now  $1 \in S$  since  $x_1 = \inf(\mathbb{N} \setminus 2^{\mathbb{N}}) = 3$  and  $3 = \frac{2^5 - 2^2 - 1}{3^2} \equiv \mathcal{R}$ . Now Suppose  $k \in S$  for  $1 \leq k \leq n$ . The proof that  $n + 1 \in S$  is broken into three cases.

In the first case suppose  $x_n$  is not one less than a power of 2 and  $x_n$  is even. Then it follows that  $x_{n+1}$  is odd. Since  $x_{n+1}$  is odd there exist a value  $t \geq 2 \in \mathbb{N}$  such that  $x_{n+1} = 2t + 1$ . Write

$$x_{n+1} = 2t + 1 = \frac{2(3t + 2) - 1}{3}$$

By the inductive hypothesis we know  $n \equiv \mathcal{R}$ . Also under the assumption  $t \sim 3t + 2$  for each value  $t$  it follows that  $3t + 2 \equiv \mathcal{R}$  or else a power of 2. If  $3t + 2 \equiv \mathcal{R}$  then by direct application of Lemma 3 it follows that  $\frac{2(3t+2)-1}{3} \equiv \mathcal{R}$ . Otherwise, if  $3t + 2 = 2^a$  for some positive integer  $a$  then  $\frac{2(3t+2)-1}{3} = \frac{2^{a+1}-1}{3} \equiv \mathcal{R}$ . Therefore in either case  $x_{n+1} \equiv \mathcal{R}$ .

In the second case, suppose  $x_n$  is not one less than a power of 2 and that  $x_n$  is odd. Then it follows that  $x_{n+1}$  is even. Since  $x_{n+1}$  is even there is a value  $t$  such that  $x_{n+1} = 2t$ . Notice that  $t$  cannot be a power of 2 since  $x_{n+1} = 2t = x_n + 1 \neq (2^a - 1) + 1 = 2^a$ . From this it follows that  $t$  cannot be a power of 2. Therefore, since  $t = \frac{x_{n+1}}{2} = \frac{x_n+1}{2} < x_n$  it follows that  $t$  is an element of the inductive set  $S$ , and hence  $t \equiv \mathcal{R}$ . By Lemma 1 it follows that  $2t \equiv \mathcal{R}$ . Therefore,  $x_{n+1} \equiv \mathcal{R}$ .

In the third and final case, suppose that  $x_n$  is exactly one less than a power of 2. That is, suppose  $x_n = 2^a - 1$  for some positive integer  $a \geq 8$ . Then it follows that  $x_{n+1} = 2^a + 1$ . Now if  $a$  is even then by direct application of Lemma 3 there exist a value  $k$  such that  $T^k(2^a + 1) = 3^{\frac{a}{2}} + 1$ . However by Lemma 2 we have the inequality

$$3^{\frac{a}{2}} + 1 < 3^{\frac{a}{2}+1} + 2 < 2^a + 1 = x_{n+1}$$

whenever  $a \geq 8$ . Therefore  $x_{n+1} = 2^a + 1$  iterates to the number  $3^{\frac{a}{2}} + 1$  and this number is either a power of 2 or it is an elemental of the inductive set  $S$ , in either case we have  $x_{n+1} \equiv \mathcal{R}$ .

Now if  $a$  is odd then by direct application of Lemma 3 there exist a value  $k$  such that  $T^k(2^a + 1) = 3^{\lfloor \frac{a}{2} \rfloor} + 2$ . By Lemma 2 we have the inequality

$$3^{\lfloor \frac{a}{2} \rfloor + 1} + 2 < 3^{\frac{a}{2}+1} + 2 < 2^a + 1 = x_{n+1}$$

whenever  $a \geq 8$ . Therefore  $x_{n+1} = 2^a + 1$  iterates to the number  $3^{\lfloor \frac{a}{2} \rfloor} + 2$  and this number is either a power of 2 or it is an elemental of the inductive set  $S$ , in either case we have  $x_{n+1} \equiv \mathcal{R}$ . The separate cases  $a < 8$  can be checked and verified by strait forward computation.

Now in all three we found that  $x_{n+1} \equiv \mathcal{R}$ . Since there are no more cases it follows that  $S = \mathbb{N}$ . This completes the proof.  $\square$



## References

- [1] J. Amigo, Representing the integers with powers of 2 and 3, *Acta Inform.*, 43 (2006) 293–306.
- [2] S., Andrei, M. Kudlek, and R. S., Niculescu, Some results on the Collatz Problem, *Acta Inform.*, 37 (2000) 145–160
- [3] C. Bohm and G. Sontacchi, On the existence of cycles of given length in integer sequences like  $x_n + 1 = x_n/2$  if  $x_n$  even, and  $x_n + 1 = 3x_n + 1$  otherwise, *Atti della Accademia Nazionale dei Lincei. Rendiconti. Classe di Scienze Fisiche, Matematiche e Naturali*, 64(8) (1978) 260–264
- [4] Charles C. Cadogan (2000), The  $3x+1$  problem: towards a solution, *Caribbean J. Math. Comput. Sci.* 10 (2000), paper 2, 11pp. (MR 2005g:11032)
- [5] Charles C. Cadogan (2003), Trajectories in the  $3x+1$  problem, *J. of Combinatorial Mathematics and Combinatorial Computing*, 44 (2003), 177–187. (MR 2004a:11017)
- [6] L. Collatz, On the motivation and origin of the  $(3n + 1)$ –Problem, *J. Qufu Normal University, Natural Science Edition*, 12(3) (1986) 9–11
- [7] R. E. Crandall, On the “ $3x+1$ ” Problem, *Math. Comp.*, 32(144) (1978) 1281–1292.
- [8] Goodwin, J. (2015). The  $3x + 1$  Problem and Integer Representations.
- [9] M. S. Klamkin, Problem 63–12, *SIAM Review*, 5 (1963) 275–276. States the problem: Consider the permutation defined by , , . Does  $n = 8$  have an infinite trajectory? How many finite cycles are there for  $f$ ? This function was the original one proposed by L. Collatz in 1932
- [10] J. C. Lagarias, The  $3x + 1$  problem: An annotated bibliography (1963–1999), web document [arXiv:math/0309224v13](http://arxiv.org/abs/math/0309224v13) [math.NT], available at arXiv.org: <http://arxiv.org/abs/math/0309224v13>
- [11] J. C. Lagarias, The  $3x + 1$  Problem: An Annotated Bibliography, II (2000–2009), web document [arXiv:math/0608208v6](http://arxiv.org/abs/math/0608208v6) [math.NT], available at arXiv.org: <http://arxiv.org/abs/math/0608208v6>

- [12] lagarius, J. C "The  $3x+1$  problem and its generalization " Amer Math. Monthly 3-23, 1985.
- [13] lagarius, J. C " The  $3x+1$  problem: An Annotated Bibliography" Amer Math. Monthly 3-23, 2012.
- [14] J. C. Lagarias, The Ultimate Challenge: The  $3x+1$  Problem, American Mathematical Society, 2010
- [15] G. Wirsching, On the combinatorial structure of  $3n+1$  predecessor sets, Discrete Math., 148 (1996) 265-286.
- [16] <http://www.cecm.sfu.ca/organics/papers/lagarias/paper/html/node1.html>
- [17] <http://www.cecm.sfu.ca/organics/papers/lagarias/paper/html/letter.html>
- [18] <http://sweet.ua.pt/tos/3x+1.html>
- [19] <http://www.ericr.nl/wondrous/index.html>
- [20] <http://www.ericr.nl/wondrous/pathrecs.html>
- [21] <http://www.csun.edu/~vcmth02i/Collatz.pdf>