

Dewlap color variation in *Anolis sagrei* is maintained among habitats within islands of the West Indies

RAPHAËL SCHERRER^{1,3} *, COLIN M. DONIHUE^{1,4},
R. GRAHAM REYNOLDS², JONATHAN B. LOSOS^{1,4} AND ANTHONY J. GENEVA^{1,5}

¹ Department of Organismic and Evolutionary Biology and Museum of Comparative Zoology
Harvard University, Cambridge, MA, USA

² Department of Biology, University of North Carolina Asheville, Asheville, NC, USA

³ Current address: Groningen Institute for Evolutionary Life Sciences,
University of Groningen, Groningen, The Netherlands

⁴ Current address: Department of Biology, Washington University, St. Louis, MO, USA

⁵ Current address: Department of Biology, Center for Computational and Integrative Biology,
Rutgers University–Camden, Camden, NJ, USA

Abstract

Animal signals evolve in an ecological context. Locally adapting animal sexual signals can be especially important for initiating or reinforcing reproductive isolation during the early stages of speciation. Previous studies have demonstrated that dewlap color in *Anolis* lizards can be highly variable between populations in relation to both biotic and abiotic adaptive drivers at relatively large geographical scales. Here, we investigated differentiation of dewlap coloration among habitat types at a small spatial scale, within multiple islands of the West Indies, to test the hypothesis that similar local adaptive processes occur over smaller spatial scales. We explored variation in dewlap coloration in the most widespread species of anole, *Anolis sagrei*, across three characteristic habitats spanning the Bahamas and the Cayman Islands, namely beach scrub, primary coppice forest and mangrove forest. Using reflectance spectrometry paired with supervised machine learning, we found significant differences in spectral properties of the dewlap between habitats within small islands, sometimes over very short distances. Passive divergence in dewlap phenotype associated with isolation-by-distance did not explain our results. On the other hand, these habitat-specific dewlap differences varied in magnitude and direction across islands, and thus our primary test for adaptation – parallel responses across islands – was not supported. We suggest that neutral processes or selection could be involved in several ways, including sexual selection. Our results shed new light on the scale at which signal color polymorphism can be maintained in the presence of gene flow, and the relative role of local adaptation and other processes in driving these patterns.

Keywords — reflectance, adaptation, sexual signal, machine learning, polymorphism

Introduction

The staggering diversity of animal communication signals has long been of interest to evolutionary biologists. Animals use chemical, mechanical, electromagnetic, and visual signals to communicate in a wide variety of contexts, including, competition for mates, species recognition, aposematism, and cooperation (Bradbury and Vehrenkamp, 2011). A primary evolutionary factor shaping communication signals is the sensory system and behavior of recipients (the sensory drive hypothesis; Endler and McLellan 1988; Endler 1992, 1998). Over the past decades, scientists have established that signals evolve in an ecological context and are dependent on environmental conditions (Endler, 1992, 1993a,b). Just as different habitats may favor different combinations of ecomorphological traits to maximize performance and fitness (Arnold, 1983), they may also shape different forms of a signal, so as to maximize its transmission and detection (e.g. Seehausen 1997), or reduce its detection by unintended recipients such as predators (Endler, 1984, 1990, 1991; Halfwerk et al.,

*Corresponding author: r.scherrer@rug.nl

37 This selective pressure may drive the local adaptation of communication signals.

38
39 One potential barrier to the maintenance of localized signal divergence is the homogenizing effect
40 of gene flow. Population genetics theory suggests that gene flow may counteract local adaptation
41 between localities and prevent divergence altogether, especially at small spatial scales, because
42 of the inflow of maladapted alleles or because of the breaking of linkage between coevolving loci
43 (Felsenstein, 1976; García-Ramos and Kirkpatrick, 1997; Dieckmann and Doebeli, 1999; Lenor-
44 mand, 2002; Hendry et al., 2007a). This genetic homogenization has been confirmed empirically
45 in systems such as stick insects (Nosil and Crespi, 2004) and stickleback (Hendry et al., 2007b).
46 Yet, examples of microgeographic adaptation, i.e. adaptation at smaller scales than the range of
47 dispersal, exist, highlighting the potential of some organisms to respond to selection in the face of
48 gene flow (see Richardson et al. 2014 and references therein). Examples include small scale adapta-
49 tion in fragmented areas in Australian fruit flies (Willi and Hoffmann, 2012), and local adaptation
50 to predation pressure in North American salamanders (Richardson and Urban, 2013). Therefore,
51 despite evidence that local adaptation may be particularly difficult at small spatial scales where
52 gene flow tends to cause adjoining populations to remain genetically homogeneous, the potential
53 adaptive response of species traits, in particular communication signals, to localized differences in
54 habitats remains relatively unknown (Richardson et al., 2014). Lizards of the neotropical genus
55 *Anolis* are an excellent group for studying the eco-evolutionary dynamics of local adaptation and
56 natural selection (Losos, 2009). A particularly conspicuous trait of anoles is their dewlap, an
57 extensible flap of skin that is typically sexually dimorphic and used as a communication signal
58 in courtship (Sigmund, 1983; Driessens et al., 2014, 2015) and territorial displays (Losos, 1985;
59 Macedonia and Stamps, 1994; Macedonia et al., 2013) as well as in predator deterrence (Leal
60 and Rodríguez-Robles, 1995, 1997; Leal and Rodriguez-Robles, 1997). Dewlap characteristics vary
61 widely among the approximately 400 species of the genus (Nicholson et al., 2007). Interspecific
62 variation in dewlap coloration is implicated in species recognition (Rand and Williams, 1970;
63 Williams, 1969; Williams and Rand, 1977; Losos, 1985; Macedonia and Stamps, 1994; Fleishman,
64 2000; Macedonia et al., 2013), and this function could have had a role in initiating or reinforcing
65 reproductive isolation during speciation (Lambert et al., 2013; Geneva et al., 2015; Ng et al., 2017).

66
67 Within species, studies have shown a link between variation in dewlap coloration and differ-
68 ences in habitats or climatic conditions (Macedonia, 2001; Leal and Fleishman, 2002; Thorpe and
69 Stenson, 2002; Thorpe, 2002; Leal and Fleishman, 2004; Vanhooydonck et al., 2009; Ng et al., 2012,
70 2013, 2016; Vanhooydonck et al., 2009; Driessens et al., 2017). Some studies suggest that those
71 differences may be adaptive and that dewlaps may have evolved to maximize detectability given
72 local light conditions (Fleishman and Persons, 2001; Leal and Fleishman, 2002, 2004). Although
73 this claim is further supported by recent findings that dewlap colors are perceived differently un-
74 der different levels of shading (Fleishman et al., 2020), other studies found conflicting patterns
75 of between-habitat variation that did not support the sensory drive hypothesis (Fleishman et al.,
76 2009; Ng et al., 2012; Macedonia et al., 2014).

77
78 Previous studies investigating variation in anole dewlaps compared populations at relatively
79 large geographical scales, e.g. between islands (Vanhooydonck et al., 2009; Driessens et al., 2017)
80 or within large islands such as Puerto Rico (Leal and Fleishman, 2004) or Hispaniola (Ng et al.,
81 2012, 2016). These large scales and marine barriers should reduce gene flow (Ng and Glor, 2011;
82 Lambert et al., 2013; Richardson et al., 2014; Ng et al., 2017). That said, examples do exist of
83 divergence in dewlap coloration at smaller scales or between populations with high degrees of gene
84 flow (Thorpe and Stenson, 2002; Thorpe, 2002; Stapley et al., 2011; Ng et al., 2016).

85
86 The species *Anolis sagrei* is widespread across islands of the West Indies (Reynolds et al., 2020).
87 It has been the subject of numerous studies concerning local adaptation (Losos et al., 1994, 1997,
88 2001; Kolbe et al., 2012), biological invasion (Kolbe et al., 2008), and sexual selection (Tokarz,
89 2002; Tokarz et al., 2005; Tokarz, 2006; Driessens et al., 2014; Steffen and Guyer, 2014; Driessens
90 et al., 2015) among many other topics. Between-island variation in the mainly orange-red color of
91 its dewlap was shown to be better explained by climatic variables (Driessens et al., 2017) than by
92 proxies for biotic factors such as sexual selection or predation pressure (Vanhooydonck et al., 2009;
93 Baeckens et al., 2018). How intra-island differences in habitat may contribute to the diversity of
94 dewlap coloration, however, remains unexplored, and may reveal new insights into the scale of local

differentiation despite gene flow.

Here, we analyzed the color characteristics of *A. sagrei* dewlaps within nine islands in the Bahamas and Cayman Islands. These island systems presently, if not historically, comprise relatively small islands, with no major geographic barriers within islands limiting dispersal for this species (Kamath and Losos, 2018). These islands all share three characteristic native West Indian small-island habitat-types – beach scrub bush, closed-canopy primary coppice forest, and mangrove forest – that are often spatially intermingled. These habitats contrast in environmental parameters including vegetation community, light irradiance, humidity, and temperature (Howard, 1950; Schoener, 1968). The Cayman Islands and the Bahamas have been colonized independently by *A. sagrei* from Cuba (van de Schoot 2016 unpublished thesis; Reynolds et al. 2020), such that these archipelagos constitute an ideal suite of natural replicates to explore within-island dewlap diversity across multiple islands.

Our sampling design included sites in close proximity; the median distance between two sites within an island was 11.2km. Combining reflectance spectrometry and supervised machine learning, we tested for divergence in dewlap phenotype between habitats within islands and between islands across part of the range of *A. sagrei*. We predicted that if light conditions in the environment indeed drive color evolution, dewlaps should be most similar between beach scrub and mangrove forest, which both have high levels of light irradiance, compared to the darker, closed-canopy coppice forest. Similar, if detectability is maximized given the local conditions, we expected darker and more contrasting dewlaps in high irradiance habitats. Finally, if habitat characteristics are strong determinants of dewlap color variation, similar patterns should be observed across multiple islands (Harvey and Pagel, 1991; Losos, 2011).

119 **Methods**

120 **Data collection**

121 We sampled 466 male *A. sagrei* from seven islands in the Bahamas Archipelago – Abaco, North
122 Andros, South Andros, South Bimini, Eleuthera, Long Island, and Ragged Island – and two in the
123 Cayman Islands – Cayman Brac and Little Cayman (Figure S1A). These islands were chosen to
124 span the breadth of the West Indian range of *A. sagrei*, because they have highly similar habitat
125 types, and because the *A. sagrei* on each island group are derived from ancient and distinct
126 colonization events from Cuba (i.e. relatively evolutionarily independent, Reynolds et al. 2020).
127 Three habitats were sampled on each island based on characterizations by Howard (1950) and
128 Schoener (1968). Each habitat is clearly distinguishable by its dominant vegetation type — xeric
129 beach scrub (open, relatively dry habitat consisting of low vegetation or isolated trees), primary
130 coppice forest (closed-canopy forest) and mangrove forest (wet coastal habitat with trees growing
131 in brackish water and high light penetration, although lizards were sampled in dry soil areas).
132 Sample sizes are given in Table S1. Our sampling design enabled us to test for differences between
133 habitats at a coarse and fine geographical scale. The median distance between two localities within
134 an island was ~ 11km (Figure S1B), and 80.3% of all pairwise distances within islands were less
135 than 50km. Additionally, there are no major barriers to dispersal (such as mountains or grassland)
136 on any of the islands that we sampled.

137 **Reflectance measurements**

138 We measured reflectance between 300nm and 700nm wavelength, a range from ultraviolet to red
139 that encompasses the colors visible to most lizards and vertebrates in general (Lazareva et al.,
140 2012). Measurements were taken with an Ocean Optics USB4000 spectrometer, a pulsed Xenon
141 light source (PX-2, Ocean Optics, Largo, FL, USA) and a reflectance probe protected by a black
142 anodized aluminum sheath. Measurements were taken with a 45-degree inclination to prevent
143 specular reflection (Endler, 1990). The device was regularly standardized with a Spectralon white
144 standard (Labsphere, North Sutton, NH, USA). Reflectance was measured at the center of the
145 dewlap. Reflectance curves were smoothed using the R package pavo (Maia et al., 2013) as well
146 as with custom R functions, down to one reflectance value at each nanometer in wavelength from
147 300 to 700nm.

148 **Analysis**

149 We tested for detectable differences in dewlap coloration between populations from different habi-
150 tats across islands by following an analytic pipeline in several steps. First, we used multivariate
151 analyses of variance to assess the relative contributions of islands, habitats and habitat-by-island
152 interactions on the partitioning of variation in color space. Second, and provided that habitat-by-
153 island interactions were found, we investigated habitat-differences in dewlap color for each island
154 separately using machine learning classification. Third, for each island where multivariate differ-
155 ences were detected using our machine learning pipeline, we used univariate analyses of variance
156 to decompose the signal among the different dimensions of color space. Fourth, for each signifi-
157 cant between-habitat variation found in univariate analyses, we used post-hoc tests to determine
158 which habitats were responsible for the differences. Last, to get insights into the spatial scale of
159 phenotypic variation, for each significant contrast between two habitats detected along a given
160 dimension on a given island, we performed multiple pairwise Wilcoxon tests to assess differences
161 in dewlap coloration among the sites involved in that significant contrast, and recorded the geo-
162 graphical distance between sites that were found significant. In parallel, we tested a possible effect
163 of isolation-by-distance, an alternative cause of phenotypic divergence between localities, based on
164 diffusion approximation and dispersal distance, irrespective of habitat types. We did so using a
165 permutation test to assess the significance of the correlation between geographical distances and
166 phenotypic distances among sites within each island.

167
168 All analyses in this study were performed in R 3.6.1 (R Core Team, 2019) and the source code
169 can be found at <https://github.com/rscherrer/dewlap>, presently private.

Dimensionality reduction

170

Because neighboring wavelengths are highly collinear and redundant in reflectance, we reduced the dimensionality of the data using principal component analysis (PCA), as per Cuthill et al. (1999) and Leal and Fleishman (2002). We performed PCA on data from all islands combined, as well as on each island separately and systematically retained the first four principal components (PC), which together always explained more than 88.8% of the variance across islands (Table S2). PCs need not represent the same wavelengths across islands because they are fitted on different datasets. Nevertheless, PC1 was highly collinear with brightness for all islands (Figure S11), while the other PCs captured the chromatic variation (i.e. irrespective of brightness) in dewlap color.

Among-island variance partitioning

179

We performed a two-way nonparametric multivariate analysis of variance (PERMANOVA, Anderson 2001, R package vegan, Oksanen et al. 2019) to identify differences in coloration between islands, habitats and habitats within islands, using principal components fitted on data from all islands together. We used a nonparametric test because although no multivariate outliers were detected based on the Mahalanobis distance, the assumption of multivariate normality was violated in several habitats on several islands (Henze-Zirkler's test, Henze and Zirkler 1990, R package MVN, Korkmaz et al. 2014, $P < 0.05$, Table S3).

Within-island machine learning

187

We performed a machine learning classification analysis on the first four principal components within each island separately, using random forests (Breiman, 2001). Random forests are a versatile, intuitive, and powerful algorithm commonly used in machine learning, using decision trees to predict the labels of particular observations based on their multivariate coordinates. These coordinates, or variables, are passed through a series of successive decision nodes, each examining a given variable of any given observation (James et al., 2013). The prediction for each observation is an aggregate over a large number of decision trees, each tree being trained on a subset of observations sampled with replacement from the dataset, and each tree being allowed to examine only a subset of the variables. This allows the random forest to overcome the individual errors of all trees in the predictions it makes.

To detect differences in dewlap coloration between habitats, we measured the success of random forests in reassigning individual lizards to their correct habitat of origin, based solely on their principal component scores. In machine learning, this so-called cross-validation procedure is typically done in two steps (James et al., 2013). First, a random forest is trained in recognizing features of dewlap coloration most associated with the different habitats, by being presented with multiple observations, making predictions about them, and updating its own decision rules based on whether the prediction deviates from the truth. Then, once trained, the patterns that the random forest has learned to recognize are tested by presenting new, previously unseen observations to the random forest, and measuring the proportion of correct predictions. This proportion, or success score, can then be statistically assessed against random guessing using a binomial test.

The cross-validation procedure requires that the data be split into a training set and a testing set. To remove any bias due to the set that is being sampled for training, it is common practice to use k-fold cross-validation (James et al., 2013), where the data are split into k random bins and k independent machines are trained, each taking one of the bins as a testing set and the rest for training, and where classification success is measured by summing all correct classifications from the k machines.

Here, we used a k-fold cross-validation procedure with $k = 5$, where each training set consisted of 80% of the data and the machine was tested on the remaining 20%. Each training set was conditioned on containing at least five lizards from each of the three habitats. We also down-sampled the training set to the sample size of the least represented habitat, to ensure that the different habitats were equally represented. To further remove any bias due to the specific random split into the different bins, we replicated each k-fold cross-validation five times. We then averaged the five resulting confusion matrices across replicates, where each confusion matrix shows the number

of lizards from each habitat reassigned into each habitat. For each island, we then used the average proportion of correctly reassigned lizards (i.e. the proportion of observations on the diagonal of the average confusion matrix) as an estimate of classification success. This score was tested against random guessing by comparing it to a binomial distribution with number of trials being the number of lizards on that island and success probability 1/3, representing the rate of successful classification by chance when three habitats are involved.

We used the machine learning fitting functions in the R package rminer (Cortez, 2020), which calls random forest routines from the randomForest package (Liaw and Wiener 2002, implementation from the original random forest algorithm Breiman, 2001). For each random forest, we optimized the number of trees in the forest and the number of variables examined by each tree using the grid hyperparameter search procedure implemented in rminer, to choose between two numbers of trees (500 or 1,000) and four numbers of principal components examined per tree (1 to 4), using rminer's ordered holdout validation method with 2/3 of the data used for training.

We validated the results of our analysis by using two other widely used machine learning classification methods: linear discriminant analysis and support vector machines (Cristianini and Shawe-Taylor, 2000; James et al., 2013), both accessible in rminer (Cortez, 2020).

To know which wavelengths were most used to assign data points to each habitat, we trained another set of random forests, this time directly on reflectance data (taken every 5nm from 300 to 700nm) instead of principal components. We recorded the relative importance of each wavelength for each habitat, as measured by the mean decrease in accuracy during wavelength permutation, implemented in the randomForest package (Liaw and Wiener, 2002).

Univariate analyses

For each island where significant differences in dewlap coloration were detected between habitats, we used multiple univariate analyses of variance (ANOVA) to identify possible principal components underlying the observed differences. We constructed our ANOVA models in two steps, as per Zuur (2009). In a first step, we accounted for heterogeneity of variances across groups by systematically comparing the goodness-of-fit of an ANOVA model estimated with ordinary least squares (OLS) with that of a model estimated with generalized least squares (GLS), which allowed one estimate of residual variance per habitat (using the R package nlme, Pinheiro and Bates 2000; Pinheiro et al. 2020). Both models were fitted with restricted maximum likelihood (REML). Goodness-of-fit was estimated using Akaike's Information Criterion corrected for small sample sizes (AICc, R package MuMIn, Bartoń 2019), and the estimation method yielding the lowest AICc was retained. In a second step, we re-fitted the retained model with maximum likelihood (ML) to test for the effect of habitat type using likelihood ratio tests (LRT) between a model including a habitat-term and a null model lacking the habitat-term.

We evaluated the normality of the standardized residuals (residuals divided by their standard error, which can differ among habitats in a GLS model) of each fitted ANOVA model using Shapiro-Wilk's test, with P-values adjusted for multiple testing using the Benjamini-Hochberg correction (Benjamini and Hochberg, 1995). In cases where significant deviations from normality were detected ($P_{adj} < 0.05$, Table S4) we performed Kruskal-Wallis's nonparametric test to back up the ANOVA results.

To know which habitat-populations were different from which in dewlap coloration, we performed different post-hoc multiple comparison tests (all implemented in the PMCMRplus package, Pohlert, 2020), depending on which assumptions were met. In cases where normality and homoscedasticity were met (i.e. OLS-ANOVA was the best fit), we used Tukey's honest significant difference test. When normality was met but not homoscedasticity (i.e. GLS-ANOVA was the best fit), we used Dunnett's T3 test. Finally, whenever we used Kruskal-Wallis's test because the ANOVA residuals were not normally distributed, we used Nemenyi's test for post-hoc comparisons.

Spatial autocorrelation

277

We tested for within-island spatial autocorrelation between the geographical distances among sampling sites and their Euclidean distances in multivariate color space (mean PC1 to PC4 per site, Table S5), regardless of habitat type. For this, we performed Mantel's test (Legendre and Legendre 2012, R package vegan; Oksanen et al. 2019) on each island, using 999 permutations and geographical distances computed as geodesic distances from latitude and longitude data (R package geosphere, Hijmans 2019).

278

279

280

281

282

283

284

Site differences

285

In this study, we were interested in the minimum spatial scale at which significant differences between habitats could be detected within islands. We performed multiple pairwise nonparametric Wilcoxon-Mann-Whitney tests (Hollander et al., 2013) to compare dewlap coloration between sites with different habitat types, for each pair of habitats and each variable where significant differences were detected with our analyses of variance. The P-values were adjusted using a Benjamini-Hochberg correction for multiple testing (Benjamini and Hochberg, 1995).

286

287

288

289

290

291

292 **Results**

293 We tested for variation in *A. sagrei* dewlap coloration between populations living in three charac-
294 teristic habitat types across nine islands that span the West Indian range of the brown anole (beach
295 scrub, primary coppice and mangroves). We found that most of the variation in coloration was
296 partitioned between islands (two-way PERMANOVA, $F(df = 8) = 43.7$, $P = 0.0001$, explained
297 variance $R^2 = 40.9\%$). Nonetheless, we did find evidence for differences in dewlap coloration
298 between habitat types, and those were mostly island-specific (habitat-by-island interaction term,
299 $F(16) = 3.53$, $P = 0.0001$, $R^2 = 6.6\%$), with a significant portion of the variation explained by
300 an ~~an~~ habitat effect across all islands, but this effect was relatively small ($F(2) = 4.7$, $P = 0.0001$,
301 $R^2 = 1.1\%$).

302

303 We subsequently tested for differences in dewlap coloration between habitat-populations within
304 each island, using within-island principal component scores (to maximize the variation captured
305 for each island, see Methods). Our within-island random forest classification analyses revealed
306 detectable differences in dewlap coloration on eight out of the nine islands in our sample: Abaco,
307 Bimini, Cayman Brac, Eleuthera, Little Cayman, Long Island, North Andros and South Andros.
308 The accuracy of random forest classification exceeded random expectation more often than ex-
309 pected by chance for all these islands (Table 1). Accuracy was as high as 73% for Cayman Brac.
310 We obtained similar results using other machine learning approaches such as support vector ma-
311 chines (Table S7) and linear discriminant analysis (Table S8), except that these methods did not
312 detect significant differences on Eleuthera and North Andros. We describe in details the specific
313 differences detected on each island in the Appendix, and ~~instead~~ focus here on the general patterns
314 emerging from our data.

315

316 Overall, we found significant differences in dewlap coloration between populations that were
317 often in close geographical proximity. On Bimini, notably, we found a significant difference be-
318 tween dewlaps from beach scrub and primary coppice forest, at a distance of a few hundred meters,
319 making this contrast the smallest geographical scale at which differences in coloration were found
320 in our study (Fig. S3). We also detected significant differences in dewlap coloration at distances
321 below one kilometer on Abaco (Fig. S2G), and at distances between one and ten kilometers on
322 Bimini (Fig. S3G), Cayman Brac (Fig. S4G), Little Cayman (Fig. S6G), Long Island (Fig. S7G)
323 and South Andros (Fig. S10G).

324

325 We found evidence of spatial autocorrelation in dewlap coloration between the sites within is-
326 lands for Abaco (Table 2), suggesting that populations from closer sites tend to have more similar
327 dewlaps on this island than expected by chance. Abaco was the island we sampled at the largest
328 scale, with some sites nearly a hundred kilometers away from each other (Fig. 2A). That said, some
329 sites were also in close proximity, and significant differences in coloration were detected between
330 habitats sometimes less than a kilometer away (Fig. S2G), suggesting that differences in dewlap
331 coloration between distant sites may be partly attributable to isolation-by-distance, but this may
332 not necessarily be the case for sites in close proximity. We did not find evidence for spatial auto-
333 correlation on other islands than Abaco (Table 2).

334

335 A striking feature of our data was inconsistencies in between-habitat differences among islands,
336 in terms of which habitats differ from which, which dimensions of coloration were involved, and
337 in which direction. For example, while on Cayman Brac the random forests could well distinguish
338 between all three habitats (Fig. S4D), on Abaco dewlaps from beach scrub and primary coppice
339 were often mistaken, and on Bimini beach scrub dewlaps were more often classified into primary
340 coppice or mangrove than into beach scrub (Fig. S3D). In terms of variable importance, for multi-
341 ple islands the random forests used information in the UV range to discriminate between at least
342 some habitats, particularly on Abaco (Fig. S2F), Bimini (Fig. S3F), Cayman Brac (Fig. S4F),
343 Little Cayman (Fig. S6F) and Long Island (Fig. S7F), but differences in UV reflectance involved
344 different habitats and were in different directions among these islands.

345

Discussion

346

Two main insights follow from our results. First, we detected significant differences in dewlap coloration between habitats within seven out of the nine islands investigated (excluding North Andros where the follow-up univariate analyses were not significant), suggesting a putatively high potential for local differentiation of dewlap coloration in *Anolis sagrei*. Second, we found differences in coloration along different dimensions of color space, and in different directions, on different islands.

347

348

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

402

Detectable differences in dewlap color between populations are surprising, as habitats were often in close geographical proximity to each other (as close as a few hundred meters on Bimini and most of the time within ten kilometers). Indeed, given that (1) the populations were continuously distributed between the habitats, (2) populations from different habitats were not monophyletic with respect to mitochondrial haplotypes (van de Schoot 2016 unpublished thesis), and (3) *A. sagrei* has been shown to be a highly mobile species (Kamath and Losos, 2018), we would have expected more homogeneous distributions of color phenotypes within islands due to extensive gene flow, with fewer differences between populations, especially those in close proximity.

Several scenarios could account for these findings. One explanation is an adaptive one: populations living in different habitats could be phenotypically adapted to their local environmental conditions. Given that the brightly colored dewlap of *A. sagrei* is used as a communication signal, its color may be a target for selection if the transmission or perception of the signal differs from one habitat to another, for example because of differences in ambient light, according to the sensory-drive hypothesis (Endler and McLellan, 1988; Endler, 1992, 1998). The sensory-drive hypothesis has been tested multiple times for dewlap coloration in *Anolis* lizards, with mixed results. Some authors found support for it (Leal and Fleishman, 2002, 2004), while others did find differences in dewlap coloration between habitats, but those were inconsistent with the sensory-drive hypothesis (Fleishman et al., 2009; Ng et al., 2012).

If our results were an example of sensory drive, we would have expected to see consistent differences between populations from different habitats across islands, given the apparent environmental consistency each of the three habitat types across the islands we sampled. In particular, we would have expected divergence in line with increased detectability given local light conditions, such as the high contrasts with background vegetation found in the UV range in Leal and Fleishman (2002) and Leal and Fleishman (2004). We might also have expected mangrove and beach scrub lizards, both inhabiting areas with high light penetration, to harbor more similar dewlaps, and to differ significantly from lizards from the coppice habitat, where irradiance is low. Instead, we found inconsistencies in the way dewlap color differed between habitats across islands. While short-wavelengths (UV reflectance) were often involved in color differences, they were not involved on all islands where significant differences were detected. On some islands, other or additional variables differed, such as brightness, red reflectance or the reflectance at the ends of the spectrum visible to *Anolis* lizards (UV and red, Lazareva et al. 2012) relative to intermediate wavelengths (blue-to-yellow). Similar portions of the spectrum were sometimes involved in opposite directions on different islands, such as on Abaco and Cayman Brac, where mangrove lizards had a higher UV-reflectance than beach scrub lizards on the former, but a lower UV-reflectance on the latter. Overall, the observed heterogeneity of divergence patterns across islands provides no support to a sensory-drive explanation.

It is presently not known if the reported differences in coloration have a genetic basis. Yet, we find it unlikely that these differences arose through phenotypic plasticity, as although the carotenoids that partly make up the red and orange colors of anole dewlaps must be found in the diet (Goodwin, 1984; Hill et al., 2002; Hill and McGraw, 2006), studies testing the effect of carotenoid deprivation (Steffen et al., 2010; Ng et al., 2013) and heritability (Cox et al., 2017) of dewlap coloration in *A. sagrei* and *A. distichus* (another species with a carotenoid-based dewlap), found little support for phenotypic and developmental plasticity in dewlap coloration. One exception is a study demonstrating that lizards heavily parasitized by skin mites had duller dewlaps (Cook et al., 2013), but we found no sign of that in our study.

We found no evidence for a role of genetic drift in explaining the observed patterns either.

403 First, *A. sagrei* was distributed across the islands continuously, usually at relatively high popula-
404 tion densities, ~~and not~~ⁱⁿ small and isolated populations where drift might be expected to have a
405 strong effect. Second, we found no evidence of isolation-by-distance except on Abaco, which was
406 sampled at the largest geographical scale, with sites nearly a hundred kilometers apart from each
407 other. Hence, while isolation-by-distance may explain long-range differences on this island, most
408 of the differences among habitats across ~~islands~~, which were often in close proximity (less than
409 10km), are unlikely to be explained by genetic drift.

410
411 In this study, we found larger differences among than within islands, a pattern already reported
412 and linked to climatic conditions (Vanhooydonck et al., 2009) and to densities of predators and
413 of anole congeners (Baeckens et al., 2018). Differences among habitats within islands, however,
414 are still difficult to account for. Remaining hypotheses may include, for example, runaway sex-
415 ual selection arbitrarily operating in different directions across islands (Andersson, 1994), but no
416 evidence so far suggests that dewlap is a target of mate choice in anoles (Tokarz, 2002; Tokarz
417 et al., 2005; Nicholson et al., 2007). Another hypothesis is that the different genetic constitutions
418 of different islands, perhaps resulting from founder effects (the islands have been colonized inde- 
419 pendently from Cuba, van de Schoot 2016 unpubl.; Driessens et al. 2017; Reynolds et al. 2020),
420 may have predisposed populations to adapt differently to similarly selective circumstances. Either
421 way, new data would be needed to test these hypotheses.

422
423 Altogether, our results show that dewlap color of *A. sagrei* commonly varies between habitat
424 types, even in close geographical proximity, within islands of the West Indies. ~~Besides~~, coloration
425 differs in different ways across similar habitats from one island to another. We discussed sev-
426 eral non-mutually exclusive mechanisms that could explain these observations, ~~but more data are~~
427 ~~needed to investigate these scenarios.~~ Nevertheless, heterogeneous patterns of divergence across is-
428 lands do not support an adaptive sensory-drive scenario, and we propose that within-island dewlap
429 color variation may be underlain by a more subtle mosaic of factors.

Acknowledgements

430

Collection permission was granted by the Bahamas Environment, Science and Technology Commission, the Bahamas National Trust, the Bahamas Ministry of Agriculture, and the Cayman Islands Department of the Environment. The authors thank Sofia Prado-Irwin, Pavitra Muralidhar, Nicholas Herrmann, Richard E. Glor, Alberto R. Puente-Rolón, Kevin Aviles-Rodriguez, Kristin Winchell, Jason Fredette, Quyhn Quach, Wendy Jesse, Inbar Maayan, Alexis Harrison and Melissa Kemp for assistance in the field and Pratik Gupte, Max Lambert and James Stroud for helpful discussions. This project was made possible through the support of a grant from the John Templeton Foundation (to JBL). The opinions expressed in this publication are those of the author(s) and do not necessarily reflect the views of the John Templeton Foundation. Additional funding for this work was provided by NSF DEB #1927194 (to JBL and AJG), NSF DEB #1500761 (to AJG), NSF DBI #1609284 (to CMD), and a Harvard Museum of Comparative Zoology Putnam Expedition Grant (to RGR).

431

432

433

434

435

436

437

438

439

440

441

442

443 Figures




Figure 1: Overview of our study design, including a map of the Bahamas and the Cayman Islands, on which are indicated the nine islands we sampled, two representatives of our study species *Anolis sagrei* with their dewlaps deployed, and the three types of habitats we considered on each island.




Figure 2: Comparison of dewlap coloration across habitats on Abaco. (A) Map of the island with the sampling sites colored by habitat. (B) Reflectance profiles of all the dewlaps on the island. (C) How reflectance profiles map onto the within-island principal components. (D) Confusion matrix showing the proportion of lizards from each (true) habitat reassigned to each (predicted) habitat by the random forests, based on the first four within-island principal components and averaged across replicates. Each column sums to one. (E) Within-island principal component scores across habitats. Bars indicate significant contrasts. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$.

Tables

444

Table 1: Random forest classification results. For each island are shown the sample size (N) and the proportion of correctly reassigned observations (or success score). P-values were computed using a binomial test and assess the significance of the observed success score relative to the score expected under random guessing. *, $P < 0.05$; **, $P < 0.01$, ***, $P < 0.001$.

Island	N	Score	P
Abaco	86	0.612	< 0.0001 ***
Bimini	67	0.510	0.0011 **
Cayman Brac	50	0.728	< 0.0001 ***
Eleuthera	56	0.493	0.0072 **
Little Cayman	45	0.649	< 0.0001 ***
Long Island	53	0.634	< 0.0001 ***
North Andros	28	0.507	0.0216 *
Ragged Island	50	0.368	0.2874
South Andros	31	0.484	0.0270 *

Table 2: Mantel's test of spatial autocorrelation. For each island are shown the correlation (Pearson's ρ) between the matrix of phenotypic distances between populations from each site and the matrix of geographic distances between sites, where phenotypic distances are Euclidean distances between the mean phenotypes of each site in the multivariate space consisting of the first four within-island principal components. P-values assess the significance of the observed correlation against the correlation expected if population means were randomly permuted among sites (999 permutations). *, $P < 0.05$.

Island	ρ	P
Abaco	0.448	0.033 *
Bimini	0.810	0.250
Cayman Brac	-0.737	0.833
Eleuthera	0.844	0.058
Little Cayman	-0.042	0.667
Long Island	0.367	0.333
North Andros	0.051	0.667
Ragged Island	-0.363	0.708
South Andros	-0.979	1.000

Table 3: Significance of habitat differences in dewlap coloration, using ANOVA for all islands where significant multivariate differences in dewlap coloration were detected by random forests. Model, best-fitting model (either OLS or GLS). AICc, corrected AIC score of the best-fitting model. ΔAICc , difference in AICc between the best-fitting model and the OLS-model. AICcw, AICc weight. Log-lik., log-likelihood. χ^2 , likelihood ratio. df, degrees of freedom. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$.

Island	Variable	AICc	ΔAICc	AICcw	Model	Log-lik.	χ^2	df	P
Abaco	PC1	255.81	2.16	0.746	OLS	-121.45	0.14	2	0.9308
Abaco	PC2	225.32	4.02	0.882	OLS	-105.66	31.74	2	< 0.0001 ***
Abaco	PC3	229.53	2.01	0.732	OLS	-107.84	27.37	2	< 0.0001 ***
Abaco	PC4	254.64	0.78	0.596	OLS	-120.85	1.36	2	0.5070
Bimini	PC1	194.16	0.77	0.595	OLS	-90.87	7.40	2	0.0248 *
Bimini	PC2	193.49	1.29	0.656	OLS	-90.52	8.09	2	0.0175 *
Bimini	PC3	184.22	-0.23	0.529	GLS	-83.46	10.39	2	0.0056 **
Bimini	PC4	200.91	3.54	0.854	OLS	-94.40	0.33	2	0.8499
Cayman Brac	PC1	136.64	-4.05	0.884	GLS	-59.29	13.81	2	0.0010 **
Cayman Brac	PC2	144.75	3.51	0.853	OLS	-66.24	8.41	2	0.0149 *
Cayman Brac	PC3	127.13	2.77	0.800	OLS	-56.86	27.16	2	< 0.0001 ***
Cayman Brac	PC4	147.37	4.33	0.897	OLS	-67.63	5.63	2	0.0600
Eleuthera	PC1	168.72	2.42	0.770	OLS	-78.46	1.00	2	0.6074 **
Eleuthera	PC2	160.03	-2.20	0.750	GLS	-70.89	11.34	2	0.0034 **
Eleuthera	PC3	163.49	-0.20	0.525	GLS	-72.69	5.57	2	0.0617
Eleuthera	PC4	164.08	3.49	0.852	OLS	-76.01	5.89	2	0.0525
Little Cayman	PC1	130.60	2.50	0.777	OLS	-59.26	8.18	2	0.0167 *
Little Cayman	PC2	112.66	-3.61	0.859	GLS	-46.74	29.76	2	< 0.0001 ***
Little Cayman	PC3	118.32	1.41	0.669	OLS	-52.68	21.34	2	< 0.0001 ***
Little Cayman	PC4	135.58	2.53	0.780	OLS	-61.92	2.85	2	0.2410
Long Island	PC1	154.54	-2.09	0.740	GLS	-68.62	2.91	2	0.2331
Long Island	PC2	155.80	-3.08	0.823	GLS	-68.92	4.52	2	0.1043
Long Island	PC3	150.54	3.67	0.862	OLS	-69.08	11.24	2	0.0036 **
Long Island	PC4	155.05	2.38	0.767	OLS	-71.47	6.46	2	0.0395 *
North Andros	PC1	88.64	0.27	0.534	OLS	-38.84	0.75	2	0.6864
North Andros	PC2	85.36	2.17	0.748	OLS	-37.01	4.42	2	0.1100
North Andros	PC3	85.31	5.82	0.948	OLS	-36.98	4.48	2	0.1065
North Andros	PC4	88.45	4.83	0.918	OLS	-38.74	0.96	2	0.6194
South Andros	PC1	95.12	0.44	0.554	OLS	-41.93	3.10	2	0.2125 *
South Andros	PC2	89.93	-0.05	0.506	GLS	-35.84	7.76	2	0.0206 **
South Andros	PC3	87.21	-6.14	0.956	GLS	-34.05	10.35	2	0.0056 **
South Andros	PC4	83.01	2.94	0.813	OLS	-35.23	16.51	2	0.0003 ***

References

- Anderson, M. J. (2001). A new method for non-parametric multivariate analysis of variance. *Austral Ecology*, 26(1):32–46. 445
446
447
- Andersson, M. B. (1994). *Sexual Selection*. Monographs in Behavior and Ecology. Princeton University Press, Princeton, N.J. 448
449
- Arnold, S. J. (1983). Morphology, Performance and Fitness. *American Zoologist*, 23(2):347–361. 450
- Baeckens, S., Driessens, T., and Van Damme, R. (2018). The brown anole dewlap revisited: Do predation pressure, sexual selection, and species recognition shape among-population signal diversity? *PeerJ*, 6:e4722. 451
452
453
- Bartoń, K. (2019). MuMIn: Multi-Model Inference. 454
- Benjamini, Y. and Hochberg, Y. (1995). Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society Series B*, 57:289–300. 455
456
457
- Bradbury, J. W. and Vehrenkamp, S. L. (2011). *Principles of Animal Communication*. Sinauer Associates, Sunderland, Mass, 2nd ed edition. 458
459
- Breiman, L. (2001). Random forests. *Machine Learning*, 45(1):5–32. 460
- Cook, E. G., Murphy, T. G., and Johnson, M. A. (2013). Colorful displays signal male quality in a tropical anole lizard. *Naturwissenschaften*, 100(10):993–996. 461
462
- Cortez, P. (2020). Rminer: Data Mining Classification and Regression Methods. 463
- Cox, R. M., Costello, R. A., Camber, B. E., and McGlothlin, J. W. (2017). Multivariate genetic architecture of the *Anolis* dewlap reveals both shared and sex-specific features of a sexually dimorphic ornament. *Journal of Evolutionary Biology*, 30(7):1262–1275. 464
465
466
- Cristianini, N. and Shawe-Taylor, J. (2000). *An Introduction to Support Vector Machines and Other Kernel-Based Learning Methods*. Cambridge University Press, first edition. 467
468
- Cuthill, I. C., Bennett, A. T. D., Partridge, J. C., and Maier, E. J. (1999). Plumage Reflectance and the Objective Assessment of Avian Sexual Dichromatism. *The American Naturalist*, 153(2):183–200. 469
470
471
- Dieckmann, U. and Doebeli, M. (1999). On the origin of species by sympatric speciation. *Nature*, 400(6742):354–357. 472
473
- Driessens, T., Baeckens, S., Balzarolo, M., Vanhooydonck, B., Huyghe, K., and Van Damme, R. (2017). Climate-related environmental variation in a visual signalling device: The male and female dewlap in *Anolis sagrei* lizards. *Journal of Evolutionary Biology*, 30(10):1846–1861. 474
475
476
- Driessens, T., Huyghe, K., Vanhooydonck, B., and Van Damme, R. (2015). Messages conveyed by assorted facets of the dewlap, in both sexes of *Anolis sagrei*. *Behavioral Ecology and Sociobiology*, 69(8):1251–1264. 477
478
479
- Driessens, T., Vanhooydonck, B., and Van Damme, R. (2014). Deterring predators, daunting opponents or drawing partners? Signaling rates across diverse contexts in the lizard *Anolis sagrei*. *Behavioral Ecology and Sociobiology*, 68(2):173–184. 480
481
482
- Endler, J. (1998). Sensory ecology, receiver biases and sexual selection. *Trends in Ecology & Evolution*, 13(10):415–420. 483
484
- Endler, J. A. (1984). Natural and sexual selection on color patterns in poeciliid fishes. In Balon, E. K. and Zaret, T. M., editors, *Evolutionary Ecology of Neotropical Freshwater Fishes*, volume 3, pages 95–111. Springer Netherlands, Dordrecht. 485
486
487
- Endler, J. A. (1990). On the measurement and classification of colour in studies of animal colour patterns. *Biological Journal of the Linnean Society*, 41(4):315–352. 488
489

- 490 Endler, J. A. (1991). Variation in the appearance of guppy color patterns to guppies and their
491 predators under different visual conditions. *Vision Research*, 31(3):587–608.
- 492 Endler, J. A. (1992). Signals, Signal Conditions, and the Direction of Evolution. *The American
493 Naturalist*, 139:S125–S153.
- 494 Endler, J. A. (1993a). The Color of Light in Forests and Its Implications. *Ecological Monographs*,
495 63(1):1–27.
- 496 Endler, J. A. (1993b). Some general comments on the evolution and design of animal communi-
497 cation systems. *Philosophical Transactions of the Royal Society of London. Series B: Biological
498 Sciences*, 340(1292):215–225.
- 499 Endler, J. A. and McLellan, T. (1988). The Processes of Evolution: Toward a Newer Synthesis.
500 *Annual Review of Ecology and Systematics*, 19(1):395–421.
- 501 Felsenstein, J. (1976). The Theoretical Population Genetics of Variable Selection and Migration.
502 *Annual Review of Genetics*, 10(1):253–280.
- 503 Fleishman, L. J. (2000). *Signal Function, Signal Efficiency and the Evolution of Anoline Lizard
504 Dewlap Color*, pages 209–236. Tapir Academic, Trondheim.
- 505 Fleishman, L. J., Leal, M., and Persons, M. H. (2009). Habitat light and dewlap color diversity in
506 four species of Puerto Rican anoline lizards. *Journal of Comparative Physiology A*, 195(11):1043–
507 1060.
- 508 Fleishman, L. J. and Persons, M. (2001). The influence of stimulus and background colour on
509 signal visibility in the lizard *Anolis cristatellus*. *The Journal of Experimental Biology*, 204(Pt
510 9):1559–1575.
- 511 Fleishman, L. J., Wadman, C. S., and Maximov, K. J. (2020). The interacting effects of total
512 light intensity and chromatic contrast on visual signal visibility in an *Anolis* lizard. *Animal
513 Behaviour*, page S0003347220302037.
- 514 García-Ramos, G. and Kirkpatrick, M. (1997). Genetic Models of Adaptation and Gene Flow in
515 Peripheral Populations. *Evolution*, 51(1):21–28.
- 516 Geneva, A. J., Hilton, J., Noll, S., and Glor, R. E. (2015). Multilocus phylogenetic analyses of
517 Hispaniolan and Bahamian trunk anoles (*distichus* species group). *Molecular Phylogenetics and
518 Evolution*, 87:105–117.
- 519 Goodwin, T. W. (1984). *The Biochemistry of the Carotenoids*. Springer Netherlands, Dordrecht.
- 520 Halfwerk, W., Jones, P. L., Taylor, R. C., Ryan, M. J., and Page, R. A. (2014). Risky Ripples Allow
521 Bats and Frogs to Eavesdrop on a Multisensory Sexual Display. *Science*, 343(6169):413–416.
- 522 Harvey, P. H. and Pagel, M. D. (1991). *The Comparative Method in Evolutionary Biology*. Oxford
523 Series in Ecology and Evolution. Oxford University Press, Oxford ; New York.
- 524 Hendry, A. P., Day, T., and Taylor, E. B. (2007a). Population Mixing and the Adaptive Divergence
525 of Quantitative Traits in Discrete Populations: A Theoretical Framework for Empirical Tests.
526 *Evolution*, 55(3):459–466.
- 527 Hendry, A. P., Taylor, E. B., and McPhail, J. D. (2007b). Adaptive Divergence and the Balance
528 Between Selection and Gene Flow: Lake and Stream Stickleback in the Misty System. *Evolution*,
529 56(6):1199–1216.
- 530 Henze, N. and Zirkler, B. (1990). A class of invariant consistent tests for multivariate normality.
531 *Communications in Statistics - Theory and Methods*, 19(10):3595–3617.
- 532 Hijmans, R. J. (2019). Geosphere: Spherical Trigonometry.
- 533 Hill, G. E., Inouye, C. Y., and Montgomerie, R. (2002). Dietary carotenoids predict plumage
534 coloration in wild house finches. *Proceedings of the Royal Society of London. Series B: Biological
535 Sciences*, 269(1496):1119–1124.

Hill, G. E. and McGraw, K. J., editors (2006). <i>Bird Coloration</i> . Harvard University Press, Cambridge, Mass.	536 537
Hollander, M., Wolfe, D. A., and Chicken, E. (2013). <i>Nonparametric Statistical Methods</i> . Wiley Series in Probability and Statistics. John Wiley & Sons, Inc, Hoboken, New Jersey, third edition edition.	538 539 540
Howard, R. A. (1950). Vegetation of the Bimini Island Group: Bahamas, B. W. I. <i>Ecological Monographs</i> , 20(4):317–349.	541 542
James, G., Witten, D., Hastie, T., and Tibshirani, R. (2013). <i>An Introduction to Statistical Learning</i> , volume 103 of <i>Springer Texts in Statistics</i> . Springer New York, New York, NY.	543 544
Kamath, A. and Losos, J. B. (2018). Estimating encounter rates as the first step of sexual selection in the lizard <i>Anolis sagrei</i> . <i>Proceedings of the Royal Society B: Biological Sciences</i> , 285(1873):20172244.	545 546 547
Kolbe, J. J., Larson, A., Losos, J. B., and de Queiroz, K. (2008). Admixture determines genetic diversity and population differentiation in the biological invasion of a lizard species. <i>Biology Letters</i> , 4(4):434–437.	548 549 550
Kolbe, J. J., Leal, M., Schoener, T. W., Spiller, D. A., and Losos, J. B. (2012). Founder Effects Persist Despite Adaptive Differentiation: A Field Experiment with Lizards. <i>Science</i> , 335(6072):1086–1089.	551 552 553
Korkmaz, S., Goksuluk, D., and Zararsiz, G. (2014). MVN: An R Package for Assessing Multivariate Normality. <i>The R Journal</i> , 6(2):151–162.	554 555
Lambert, S. M., Geneva, A. J., Luke Mahler, D., and Glor, R. E. (2013). Using genomic data to revisit an early example of reproductive character displacement in Haitian <i>Anolis</i> lizards. <i>Molecular Ecology</i> , 22(15):3981–3995.	556 557 558
Lazareva, O. F., Shimizu, T., and Wasserman, E. A. (2012). <i>How Animals See the WorldComparative Behavior, Biology, and Evolution of Vision</i> . Oxford University Press.	559 560
Leal, M. and Fleishman, L. J. (2002). Evidence for habitat partitioning based on adaptation to environmental light in a pair of sympatric lizard species. <i>Proceedings of the Royal Society of London. Series B: Biological Sciences</i> , 269(1489):351–359.	561 562 563
Leal, M. and Fleishman, L. J. (2004). Differences in Visual Signal Design and Detectability between Allopatric Populations of <i>Anolis</i> Lizards. <i>The American Naturalist</i> , 163(1):26–39.	564 565
Leal, M. and Rodríguez-Robles, J. A. (1995). Antipredator Responses of <i>Anolis cristatellus</i> (Sauria: Polychrotidae). <i>Copeia</i> , 1995(1):155–161.	566 567
Leal, M. and Rodriguez-Robles, J. A. (1997). Antipredator Responses of the Puerto Rican Giant Anole, <i>Anolis cuvieri</i> (Squamata: Polychrotidae). <i>Biotropica</i> , 29(3):372–375.	568 569
Leal, M. and Rodríguez-Robles, J. A. (1997). Signalling displays during predator–prey interactions in a Puerto Rican anole, <i>Anolis cristatellus</i> . <i>Animal Behaviour</i> , 54(5):1147–1154.	570 571
Legendre, P. and Legendre, L. (2012). <i>Numerical Ecology</i> . Number 24 in Developments in Environmental Modelling. Elsevier, Amsterdam, third english edition edition.	572 573
Lenormand, T. (2002). Gene flow and the limits to natural selection. <i>Trends in Ecology & Evolution</i> , 17(4):183–189.	574 575
Liaw, A. and Wiener, M. (2002). Classification and Regression by randomForest. <i>R News</i> , 2(3):18–22.	576 577
Losos, J. B. (1985). An Experimental Demonstration of the Species-Recognition Role of <i>Anolis</i> Dewlap Color. <i>Copeia</i> , 1985(4):905–910.	578 579
Losos, J. B. (2009). <i>Lizards in an Evolutionary Tree: Ecology and Adaptive Radiation of Anoles</i> . University of California Press.	580 581

- 582 Losos, J. B. (2011). Convergence, Adaptation, and Constraint. *Evolution*, 65(7):1827–1840.
- 583 Losos, J. B., Irschick, D. J., and Schoener, T. W. (1994). Adaptation and Constraint in the
584 Evolution of Specialization of Bahamian *Anolis* Lizards. *Evolution*, 48(6):1786–1798.
- 585 Losos, J. B., Schoener, T. W., Warheit, K. I., and Creer, D. (2001). Experimental studies of
586 adaptive differentiation in Bahamian *Anolis* lizards. *Genetica*, 112-113:399–415.
- 587 Losos, J. B., Warheit, K. I., and Schoener, T. W. (1997). Adaptive differentiation following
588 experimental island colonization in *Anolis* lizards. *Nature*, 387(6628):70–73.
- 589 Macedonia, J. M. (2001). Habitat light, colour variation, and ultraviolet reflectance in the Grand
590 Cayman anole, *Anolis conspersus*. *Biological Journal of the Linnean Society*, 73(3):299–320.
- 591 Macedonia, J. M., Clark, D. L., Riley, R. G., and Kemp, D. J. (2013). Species recognition of color
592 and motion signals in *Anolis grahami*: Evidence from responses to lizard robots. *Behavioral
593 Ecology*, 24(4):846–852.
- 594 Macedonia, J. M., Clark, D. L., and Tamasi, A. L. (2014). Does Selection Favor Dewlap Colors that
595 Maximize Detectability? A Test with Five Species of Jamaican *Anolis* Lizards. *Herpetologica*,
596 70(2):157–170.
- 597 Macedonia, J. M. and Stamps, J. A. (1994). Species Recognition in *Anolis grahami* (Sauria,
598 Iguanidae): Evidence from Responses to Video Playbacks of Conspecific and Heterospecific
599 Displays. *Ethology*, 98(3-4):246–264.
- 600 Maia, R., Eliason, C. M., Bitton, P.-P., Doucet, S. M., and Shawkey, M. D. (2013). Pavo: An
601 R package for the analysis, visualization and organization of spectral data. *Methods in Ecology
602 and Evolution*, pages n/a–n/a.
- 603 Ng, J., Geneva, A. J., Noll, S., and Glor, R. E. (2017). Signals and Speciation: *Anolis* Dewlap
604 Color as a Reproductive Barrier. *Journal of Herpetology*, 51(3):437–447.
- 605 Ng, J. and Glor, R. E. (2011). Genetic differentiation among populations of a Hispaniolan trunk
606 anole that exhibit geographical variation in dewlap colour. *Molecular Ecology*, 20(20):4302–4317.
- 607 Ng, J., Kelly, A. L., MacGuigan, D. J., and Glor, R. E. (2013). The Role of Heritable and Dietary
608 Factors in the Sexual Signal of a Hispaniolan *Anolis* Lizard, *Anolis distichus*. *Journal of Heredity*,
609 104(6):862–873.
- 610 Ng, J., Landeen, E. L., Logsdon, R. M., and Glor, R. E. (2012). Correlation Between *Anolis*
611 Lizard Dewlap Phenotype and Environmental Variation Indicates Adaptive Divergence of a
612 Signal Important to Sexual Selection and Species Recognition. *Evolution*, 67(2):573–582.
- 613 Ng, J., Ossip-Klein, A. G., and Glor, R. E. (2016). Adaptive signal coloration maintained in the
614 face of gene flow in a Hispaniolan *Anolis* Lizard. *BMC Evolutionary Biology*, 16(1):193.
- 615 Nicholson, K. E., Harmon, L. J., and Losos, J. B. (2007). Evolution of *Anolis* Lizard Dewlap
616 Diversity. *PLoS ONE*, 2(3):e274.
- 617 Nosil, P. and Crespi, B. J. (2004). Does Gene Flow Constrain Adaptive Divergence or Vice Versa? A
618 Test Using Ecomorphology and Sexual Isolation in *Timema cristinae* Walking-Sticks. *Evolution*,
619 58(1):102–112.
- 620 Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R.,
621 O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., and Wagner, H.
622 (2019). Vegan: Community Ecology Package. R package version 2.5-6.
- 623 Pinheiro, J. and Bates, D. (2000). *Mixed-Effects Models in S and S-PLUS*. Statistics and Com-
624 puting. Springer-Verlag, New York.
- 625 Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., and Team, R. C. (2020). Nlme: Linear and
626 Nonlinear Mixed Effects Models.

Pohlert, T. (2020). PMCMRplus: Calculate Pairwise Multiple Comparisons of Mean Rank Sums Extended. R package version 1.4.4.	627 628
R Core Team (2019). R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing.	629 630
Rand, A. S. and Williams, E. E. (1970). An Estimation of Redundancy and Information Content of Anole Dewlaps. <i>The American Naturalist</i> , 104(935):99–103.	631 632
Reynolds, R. G., Kolbe, J. J., Glor, R. E., López-Darias, M., Gómez Pourroy, C. V., Harrison, A. S., Queiroz, K., Revell, L. J., and Losos, J. B. (2020). Phylogeographic and phenotypic outcomes of brown anole colonization across the Caribbean provide insight into the beginning stages of an adaptive radiation. <i>Journal of Evolutionary Biology</i> , 33(4):468–494.	633 634 635 636
Richardson, J. L. and Urban, M. C. (2013). Strong Selection Barriers Explain Microgeographic Adaptation in Wild Salamander Populations. <i>Evolution</i> , 67(6):1729–1740.	637 638
Richardson, J. L., Urban, M. C., Bolnick, D. I., and Skelly, D. K. (2014). Microgeographic adaptation and the spatial scale of evolution. <i>Trends in Ecology & Evolution</i> , 29(3):165–176.	639 640
Schoener, T. W. (1968). The <i>Anolis</i> Lizards of Bimini: Resource Partitioning in a Complex Fauna. <i>Ecology</i> , 49(4):704–726.	641 642
Seehausen, O. (1997). Cichlid Fish Diversity Threatened by Eutrophication That Curbs Sexual Selection. <i>Science</i> , 277(5333):1808–1811.	643 644
Sigmund, W. R. (1983). Female Preference for <i>Anolis carolinensis</i> Males as a Function of Dewlap Color and Background Coloration. <i>Journal of Herpetology</i> , 17(2):137–143.	645 646
Stapley, J., Wordley, C., and Slate, J. (2011). No Evidence of Genetic Differentiation Between Anoles With Different Dewlap Color Patterns. <i>Journal of Heredity</i> , 102(1):118–124.	647 648
Steffen, J. E. and Guyer, C. C. (2014). Display behaviour and dewlap colour as predictors of contest success in brown anoles: Dewlap Colour and Behaviour in Contests. <i>Biological Journal of the Linnean Society</i> , 111(3):646–655.	649 650 651
Steffen, J. E., Hill, G. E., and Guyer, C. (2010). Carotenoid Access, Nutritional Stress, and the Dewlap Color of Male Brown Anoles. <i>Copeia</i> , 2010(2):239–246.	652 653
Thorpe, R. S. (2002). Analysis of Color Spectra in Comparative Evolutionary Studies: Molecular Phylogeny and Habitat Adaptation in the St. Vincent Anole (<i>Anolis trinitatis</i>). <i>Systematic Biology</i> , 51(4):554–569.	654 655 656
Thorpe, R. S. and Stenson, A. G. (2002). Phylogeny, Paraphyly and Ecological Adaptation of the Colour and Pattern in the Anolis Roquet Complex on Martinique: Interaction Between Phylogeny and Adaptation. <i>Molecular Ecology</i> , 12(1):117–132.	657 658 659
Tokarz, R. R. (2002). An Experimental Test of the Importance of the Dewlap in Male Mating Success in the Lizard <i>Anolis sagrei</i> . <i>Herpetologica</i> , 58(1):87–94.	660 661
Tokarz, R. R. (2006). Importance of Prior Physical Contact with Familiar Females in the Development of a Male Courtship and Mating Preference for Unfamiliar Females in the Lizard <i>Anolis Sagrei</i> . <i>Herpetologica</i> , 62(2):115–124.	662 663 664
Tokarz, R. R., Paterson, A. V., and McMann, S. (2005). Importance of Dewlap Display in Male Mating Success in Free-Ranging Brown Anoles (<i>Anolis sagrei</i>). <i>Journal of Herpetology</i> , 39(1):174–177.	665 666 667
van de Schoot, M. (2016). <i>Within and between Island Radiation and Genetic Variation in Anolis Sagrei</i> . PhD thesis, Wageningen University, Wageningen, The Netherlands.	668 669
Vanhooydonck, B., Herrel, A., Meyers, J. J., and Irschick, D. J. (2009). What determines dewlap diversity in <i>Anolis</i> lizards? An among-island comparison. <i>Journal of Evolutionary Biology</i> , 22(2):293–305.	670 671 672

- 673 Willi, Y. and Hoffmann, A. A. (2012). Microgeographic adaptation linked to forest fragmentation
674 and habitat quality in the tropical fruit fly *Drosophila birchii*. *Oikos*, 121(10):1627–1637.
- 675 Williams, E. E. (1969). The Ecology of Colonization as Seen in the Zoogeography of Anoline
676 Lizards on Small Islands. *The Quarterly Review of Biology*, 44(4):345–389.
- 677 Williams, E. E. and Rand, A. S. (1977). Species Recognition, Dewlap Function and Faunal Size.
678 *American Zoologist*, 17(1):261–270.
- 679 Zuur, A. F., editor (2009). *Mixed Effects Models and Extensions in Ecology with R*. Statistics for
680 Biology and Health. Springer, New York, NY.

Appendix

681

Here we describe more precisely the patterns identified on each island.

682

On Abaco, dewlaps from the mangrove habitat were the best discriminated, while dewlaps from the beach scrub habitat were often mistaken for dewlaps from the coppice habitat (Fig. 2D). Importance analysis revealed that beach scrub and mangrove lizards mostly differed in reflectance in the ultraviolet (UV) end of the spectrum (below 400nm, Fig. S2F), where mangrove dewlaps had higher UV reflectance relative to beach scrub lizards, and coppice lizards had an intermediate UV reflectance between the two other habitats (Fig. 2B). Consistent with this, our analyses of variance detected significantly lower PC2 scores in mangrove lizards than in the two other habitats (Fig. 2E, Table 3), representing a higher UV-reflectance relative to red (Fig. 2C). Beach scrub lizards also scored lower on PC3 (Fig. 2E, Table 3), indicating less curvature of the reflectance profile and relatively higher reflectance at intermediate wavelengths (blue-to-yellow) than at the ends of the range (Fig. 2C). Differences were detected between sites both at large ($\sim 100\text{km}$) and short ($< 1\text{km}$) distances (Fig. S2G). Abaco was the only island where we detected significant spatial autocorrelation (Table 2), that is, sites that were closer geographically tended to have populations of lizards with more similar dewlap colors.

683

684

685

686

687

688

689

690

691

692

693

694

695

696

697

698

699

700

701

702

703

704

705

706

707

708

709

710

711

712

713

714

715

716

717

718

719

720

721

722

723

724

725

726

727

728

729

730

731

732

733

734

735

736

737

On Bimini, the random forests mostly correctly classified lizards from the coppice and mangrove habitats while often misclassifying lizards from the beach scrub habitat (Fig. S3D). Relatively flat importance profiles for beach scrub lizards suggested that brightness was used instead of a particular wavelength to identify some of the beach scrub dewlaps (Fig. S3F). Indeed, some beach scrub dewlaps were substantially brighter than the rest (Fig. S3B, C), a pattern that was captured by our analysis of variance along PC1 (i.e. brightness, Fig. S3C, E, Table 3). The random forests also used UV reflectance to discriminate between coppice and mangrove dewlaps (Fig. S3F), which could reflect the significant difference we detected along PC3 between these two habitats (Fig. S3C, E, Table 3). Beach scrub lizards were characterized by elevated red reflectance relative to UV (as represented by PC2, (Fig. S3C, E)), and beach scrub and mangrove lizards were characterized by a more even distribution of the reflectance along the spectrum (as represented by PC3, (Fig. S3C, E)), in contrast to coppice lizards which harbored a stronger curvature at intermediate wavelengths (Fig. S3C, E). On this island, the beach scrub and coppice habitats were separated by a few hundred meters, making this contrast the smallest geographical scale at which differences in coloration were found in our study (Fig. S3G).

On Cayman Brac, all three habitats could be well discriminated against each other (Fig. S4D), with UV reflectance appearing to be an important variable differentiating beach scrub and mangrove dewlaps (Fig. S4F). In contrast, coppice dewlaps had a relatively flat importance profile, suggesting that brightness made them more distinct rather than any particular wavelength (Fig. S4F). Consistent with this, coppice dewlaps were significantly different from all other dewlaps along PC1 (Fig. S4E, Table 3). At a distance between 2 and 3km (Fig. S4G), dewlaps in the beach scrub habitat reflected more red light (as represented by PC2, Fig. S4C, E) and more UV (as represented by PC3, Fig. S4C, E) than in the mangrove habitat. Coppice lizards were also characterized by a higher UV reflectance than mangrove dewlaps (PC3, Fig. S4C, E, Table 3).

On Eleuthera, although random forests detected between-habitat differences in dewlap color, other approaches did not (Tables S8 and S7), suggesting that the differences may be small. Consistent with this, the only significant univariate difference detected was for PC2 between beach scrub and mangrove lizards, where beach scrub lizards had higher levels of red reflectance and mangrove lizards higher levels of UV reflectance (Fig. S5C, E, Table 3).

Little Cayman was characterized by a better discrimination of mangrove lizards from the rest than between beach scrub and coppice lizards (Fig. S6D). Mangrove dewlaps were most distinct with respect to their reflectance in short wavelengths (Fig. S6F), with significantly lower UV reflectance (as represented by PC2, Fig. S6C, E, Table 3). Beach scrub lizards were characterized by brighter dewlaps than coppice lizards (PC1), and also more convex curves, i.e. slightly higher UV and red reflectance (as represented by higher PC3 scores), than lizards from the other two habitats (Fig. S6C, E, Table 3).

738
739 On Long Island the three habitats were relatively well discriminated (Fig. S7D). Importance
740 profiles indicated that short wavelengths were used to discriminate between beach scrub and man-
741 grove lizards (Fig. S7F). Beach scrub lizards had more curved reflectance profiles than in either of
742 the two other habitats, with higher levels of UV and red reflectance relative to intermediate wave-
743 lengths (PC3, Fig. S7C, E, Table 3). Beach scrub lizards also differed from mangrove lizards along
744 PC4 (Fig. S7E), which represented a rather small portion of the variance not already explained
745 by the first three principal components, and is therefore difficult to interpret (Fig. S7C). Coppice
746 lizards were significantly darker than mangrove and beach scrub lizards (PC1, Fig. S7C, E, Table
747 3).

748
749 On North Andros, although the random forest classification was significant ($P = 0.0216$, Table
750 1) and the average confusion matrix indicated that lizards from beach scrub were particularly well
751 predicted (Fig. S8D), no significant univariate differences were detected along any of the four PCs
752 (Fig. S8E, Table 3). Importance analysis of full-spectrum random forests showed higher impor-
753 tance scores near the UV-end of the spectrum in discriminating beach scrub dewlaps from the rest
754 (Fig. S8F). Besides, reflectance curves of beach scrub dewlaps appeared more similar to each other
755 in the UV range than dewlaps from other habitats (Fig. S8B), suggesting that the machines may
756 have used this low within-habitat variance, as opposed to between-habitat differences in means,
757 to correctly classify beach scrub lizards. A small sample size on this island may also have con-
758 tributed to a lack of power in detecting univariate differences using analyses of variance (Table S1).

759
760 On South Andros beach scrub and coppice dewlaps could be discriminated better against
761 each other than with mangrove dewlaps (Fig. S10D), with importance profiles supporting UV-
762 reflectance as a predictor of coppice lizards (Fig. S10F). Coppice lizards had more curved re-
763 flectance profiles than beach scrub lizards (PC3), and lizards from both habitats differed along
764 PC4, which is again more difficult to interpret (Fig. S10C, E, Table 3). Beach scrub lizards also
765 differed from mangrove lizards in PC4 (Fig. S10C, E, Table 3).

766
767 Classification success was not significantly better than expected by chance on Ragged Island
768 (Table 1) where nearly no habitat could be differentiated from any other based on reflectance.

Supplementary Figures

769




Figure S1: Maps of the islands. (A) Map of the West Indies with sampled islands highlighted in black. (B) Sampling sites within islands colored after their respective habitat types.




Figure S2: Comparison of dewlap coloration across habitats on Abaco, with extended results. (A–E) Legend as per Figure 2. (F) One-dimensional sensitivity analysis showing the relative importance (mean decrease in accuracy) of the various wavelengths in random forest classification of the whole spectrum. (G) Geographical distance between sites where significant differences were detected in within-island principal component scores (Wilcoxon test, Benjamini-Hochberg correction, $P < 0.05$), including only pairs of sites whose habitats were involved in between-habitat dewlap differences.




Figure S3: Comparison of dewlap coloration across habitats on Bimini. Legend is as per Figure S2.




Figure S4: Comparison of dewlap coloration across habitats on Cayman Brac. Legend is as per Figure S2.




Figure S5: Comparison of dewlap coloration across habitats on Eleuthera. Legend is as per Figure S2.




Figure S6: Comparison of dewlap coloration across habitats on Little Cayman. Legend is as per Figure S2.




Figure S7: Comparison of dewlap coloration across habitats on Long Island. Legend is as per Figure S2.




Figure S8: Comparison of dewlap coloration across habitats on North Andros. Legend is as per Figure S2, but without panel G.




Figure S9: Comparison of dewlap coloration across habitats on Ragged Island. Legend is as per Figure S2, but without panel G.




Figure S10: Comparison of dewlap coloration across habitats on South Andros. Legend is as per Figure S2.




Figure S11: PC1 captures brightness across all islands. (A) Correlation between dewlap brightness (as measured by the mean reflectance from 300 to 700nm in wavelength) and PC1 score across all islands. (B) Correlation between brightness and within-island PC1, for each island. Pearson's correlation coefficients are reported. ***, $P < 0.001$.

770 Supplementary Tables

Table S1: Number of lizards sampled in each habitat on each island.

	coastal	coppice	mangrove
Abaco	41	24	21
Bimini	38	14	15
Cayman Brac	15	18	17
Eleuthera	22	25	9
Little Cayman	17	12	16
Long Island	26	14	13
North Andros	9	9	10
Ragged Island	18	15	17
South Andros	10	9	12

Table S2: Proportion of variance explained by the first four principal components on each island, as well as in the PCA performed on all islands together (last row).

Island	Total	PC1	PC2	PC3	PC4
Abaco	0.906	0.400	0.279	0.147	0.079
Bimini	0.921	0.502	0.208	0.160	0.051
Cayman Brac	0.888	0.438	0.190	0.155	0.105
Eleuthera	0.926	0.490	0.233	0.138	0.066
Little Cayman	0.907	0.441	0.212	0.176	0.078
Long Island	0.925	0.515	0.205	0.161	0.043
North Andros	0.937	0.560	0.170	0.152	0.054
Ragged Island	0.907	0.483	0.226	0.127	0.072
South Andros	0.948	0.488	0.247	0.146	0.067
All islands	0.913	0.473	0.197	0.164	0.079

Table S3: Henze-Zirkler's test of multivariate normality, performed on global principal components (i.e. fitted on data from all islands together) in each habitat and on each island. The number of outlier points detected based on the Mahalanobis distance is reported. HZ , test statistic. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$.

Island	Habitat	Outliers	HZ	P
Abaco	coastal	0	1.099	0.0027 **
Abaco	coppice	0	1.074	0.0022 **
Abaco	mangrove	0	1.063	0.0023 **
Bimini	coastal	0	1.281	< 0.0001 ***
Bimini	coppice	0	0.850	0.0482 *
Bimini	mangrove	0	1.191	0.0001 ***
Cayman Brac	coastal	0	0.647	0.5311
Cayman Brac	coppice	0	0.701	0.3940
Cayman Brac	mangrove	0	0.657	0.5357
Eleuthera	coastal	0	1.614	< 0.0001 ***
Eleuthera	coppice	0	1.481	< 0.0001 ***
Eleuthera	mangrove	0	0.729	0.1423
Little Cayman	coastal	0	0.624	0.6599
Little Cayman	coppice	0	0.638	0.4867
Little Cayman	mangrove	0	0.873	0.0413 *
Long Island	coastal	0	0.824	0.1468
Long Island	coppice	0	0.923	0.0150 *
Long Island	mangrove	0	0.773	0.1289
North Andros	coastal	0	0.658	0.3174
North Andros	coppice	0	0.763	0.0900
North Andros	mangrove	0	0.668	0.3185
Ragged Island	coastal	0	0.756	0.2268
Ragged Island	coppice	0	0.797	0.1115
Ragged Island	mangrove	0	0.542	0.9022
South Andros	coastal	0	0.660	0.3451
South Andros	coppice	0	0.659	0.3154
South Andros	mangrove	0	0.911	0.0144 *

Table S4: Shapiro-Wilk's test of univariate normality of the standardized residuals from OLS and GLS-ANOVAs performed on each island where significant differences were detected by random forest classification. W , test statistic. *, $P_{adj} < 0.05$; **, $P_{adj} < 0.01$; ***, $P_{adj} < 0.001$.

Island	Variable	W	P	P_{adj}	
Abaco	PC1	0.961	0.0109	0.0497	*
Abaco	PC2	0.960	0.0089	0.0473	*
Abaco	PC3	0.988	0.5938	0.7308	
Abaco	PC4	0.982	0.2709	0.4816	
Bimini	PC1	0.890	0.0000	0.0008	***
Bimini	PC2	0.984	0.5179	0.6906	
Bimini	PC3	0.959	0.0281	0.0997	
Bimini	PC4	0.980	0.3386	0.5418	
Cayman Brac	PC1	0.986	0.8236	0.8785	
Cayman Brac	PC2	0.989	0.9299	0.9299	
Cayman Brac	PC3	0.934	0.0079	0.0473	*
Cayman Brac	PC4	0.981	0.5926	0.7308	
Eleuthera	PC1	0.930	0.0031	0.0330	*
Eleuthera	PC2	0.975	0.3090	0.5204	
Eleuthera	PC3	0.981	0.4989	0.6906	
Eleuthera	PC4	0.970	0.1788	0.4082	
Little Cayman	PC1	0.955	0.0821	0.2539	
Little Cayman	PC2	0.982	0.6854	0.7833	
Little Cayman	PC3	0.891	0.0005	0.0081	**
Little Cayman	PC4	0.977	0.4858	0.6906	
Long Island	PC1	0.937	0.0074	0.0473	*
Long Island	PC2	0.989	0.9039	0.9299	
Long Island	PC3	0.971	0.2170	0.4341	
Long Island	PC4	0.983	0.6261	0.7420	
North Andros	PC1	0.937	0.0952	0.2539	
North Andros	PC2	0.978	0.8075	0.8785	
North Andros	PC3	0.905	0.0147	0.0587	
North Andros	PC4	0.949	0.1913	0.4082	
South Andros	PC1	0.941	0.0879	0.2539	
South Andros	PC2	0.946	0.1199	0.2952	
South Andros	PC3	0.965	0.3950	0.6019	
South Andros	PC4	0.957	0.2480	0.4668	

Table S5: Locations of the sampling sites across islands, with mean within-island principal component scores per site.

Island	Longitude	Latitude	Habitat	PC1	PC2	PC3	PC4
Abaco	-77.7	26.9	mangrove	1.006	0.117	-0.057	-1.259
Abaco	-77.6	26.9	coastal	0.155	0.532	0.079	-2.046
Abaco	-77.6	26.9	coppice	-0.084	0.033	0.636	-0.652
Abaco	-77.2	26.1	coastal	0.304	-0.093	-0.987	-0.133
Abaco	-77.0	26.3	mangrove	-0.316	-1.224	0.899	0.185
Abaco	-77.0	26.3	coppice	0.092	0.314	0.124	0.483
Abaco	-77.0	26.3	coastal	-0.337	0.750	-0.207	0.540
Bimini	-79.3	25.6	coastal	-0.263	-1.270	-0.185	0.164
Bimini	-79.3	25.7	coastal	0.468	0.083	-0.242	0.011
Bimini	-79.3	25.7	coppice	-0.270	0.171	0.926	-0.124
Bimini	-79.3	25.7	mangrove	-0.447	0.533	-0.288	-0.014
Cayman Brac	-79.9	19.7	coastal	0.483	-0.523	-0.781	0.443
Cayman Brac	-79.8	19.7	mangrove	0.219	0.453	0.810	-0.015
Cayman Brac	-79.8	19.7	coppice	-0.610	0.008	-0.114	-0.355
Eleuthera	-76.3	24.8	coppice	-0.123	-0.857	0.045	0.117
Eleuthera	-76.3	24.8	coastal	-0.520	-0.642	-0.492	-0.538
Eleuthera	-76.3	24.8	mangrove	-0.163	-0.651	0.126	0.685
Eleuthera	-76.2	24.9	coppice	0.391	0.602	0.540	-0.539
Eleuthera	-76.1	24.9	coastal	0.254	1.228	-0.315	0.402
Little Cayman	-80.1	19.7	coppice	-0.621	0.614	-0.179	-0.404
Little Cayman	-80.0	19.7	coastal	0.395	0.355	0.743	0.175
Little Cayman	-80.0	19.7	mangrove	0.047	-0.838	-0.655	0.118
Long Island	-75.2	23.5	mangrove	0.207	0.366	0.484	0.587
Long Island	-75.2	23.4	coastal	-0.097	-0.798	-0.140	-0.347
Long Island	-75.2	23.4	coppice	-0.485	-0.133	0.366	-0.148
Long Island	-75.1	23.4	coastal	0.316	0.317	-0.626	-0.130
North Andros	-77.9	24.8	coastal	-0.098	-0.516	-0.403	0.066
North Andros	-77.8	24.8	coppice	-0.139	0.090	0.519	0.192
North Andros	-77.8	24.7	mangrove	0.213	0.384	-0.104	-0.232
Ragged Island	-75.7	22.2	coppice	-0.269	0.440	-0.088	-0.360
Ragged Island	-75.7	22.2	coastal	0.049	0.076	-0.221	0.029
Ragged Island	-75.7	22.2	mangrove	0.246	-0.273	0.118	0.933
Ragged Island	-75.7	22.2	mangrove	0.099	-0.749	0.588	-0.636
South Andros	-77.6	24.2	mangrove	-0.250	-0.539	0.316	-0.211
South Andros	-77.6	24.1	coppice	0.466	0.657	-0.586	-0.687
South Andros	-77.5	24.1	coastal	-0.120	0.055	0.149	0.872

Table S6: Nonparametric Kruskal-Wallis tests performed on each variable on each island where deviations from normality were detected. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$.

Island	Variable	χ^2	df	P
Abaco	PC1	0.74	2	0.6924
Abaco	PC2	23.13	2	< 0.0001 ***
Bimini	PC1	7.38	2	0.0250 *
Cayman Brac	PC3	22.46	2	< 0.0001 ***
Eleuthera	PC1	0.29	2	0.8666
Little Cayman	PC3	19.95	2	< 0.0001 ***
Long Island	PC1	10.98	2	0.0041 **

Table S7: Support vector machine classification results. Legend is as per Table 1.

Island	<i>N</i>	Score	<i>P</i>	
Abaco	86	0.581	< 0.0001	***
Bimini	67	0.555	< 0.0001	***
Cayman Brac	50	0.708	< 0.0001	***
Eleuthera	56	0.443	0.0513	
Little Cayman	45	0.711	< 0.0001	***
Long Island	53	0.664	< 0.0001	***
North Andros	28	0.429	0.1039	
Ragged Island	50	0.424	0.0756	
South Andros	31	0.574	0.0040	**

Table S8: Linear discriminant analysis classification results. Legend is as per Table 1.

Island	<i>N</i>	Score	<i>P</i>	
Abaco	86	0.644	< 0.0001	***
Bimini	67	0.585	< 0.0001	***
Cayman Brac	50	0.780	< 0.0001	***
Eleuthera	56	0.425	0.0871	
Little Cayman	45	0.738	< 0.0001	***
Long Island	53	0.574	0.0001	***
North Andros	28	0.414	0.1911	
Ragged Island	50	0.400	0.1259	
South Andros	31	0.587	0.0013	**