

Dewlap color variation in *Anolis sagrei* is maintained among habitats within islands of the West Indies

1
2

3

Abstract

Animal signals evolve in an ecological context. Locally adapting animal sexual signals can be especially important for initiating or reinforcing reproductive isolation during the early stages of speciation. Previous studies have demonstrated that dewlap color in *Anolis* lizards can be highly variable between populations in relation to both biotic and abiotic adaptive drivers at relatively large geographical scales. Here, we investigated differentiation of dewlap coloration among habitat types at a small spatial scale, within multiple islands of the West Indies, to test the hypothesis that similar local adaptive processes occur over smaller spatial scales. We explored variation in dewlap coloration in the most widespread species of anole, *Anolis sagrei*, across three characteristic habitats spanning the Bahamas and the Cayman Islands, namely beach scrub, primary coppice forest and mangrove forest. Using reflectance spectrometry paired with supervised machine learning, we found significant differences in spectral properties of the dewlap between habitats within small islands, sometimes over very short distances. Passive divergence in dewlap phenotype associated with isolation-by-distance did not seem to explain our results. On the other hand, these habitat-specific dewlap differences varied in magnitude and direction across islands, and thus our primary test for adaptation – parallel responses across islands – was not supported. We suggest that neutral processes or selection could be involved in several ways, including sexual selection. Our results shed new light on the scale at which signal color polymorphism can be maintained in the presence of gene flow, and the relative role of local adaptation and other processes in driving these patterns of dewlap color variation across islands.

Keywords — reflectance, adaptation, sexual signal, machine learning, polymorphism

4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

26 Introduction

27 The staggering diversity of animal communication signals has long been of interest to evolutionary
28 biologists. Animals use chemical, mechanical, electromagnetic, and visual signals to communicate
29 in a wide variety of contexts, including, competition for mates, species recognition, aposematism,
30 and cooperation (Bradbury and Vehrenberg, 2011). A primary evolutionary factor shaping com-
31 munication signals is the sensory system and behavior of recipients (the sensory drive hypothesis;
32 Endler and McLellan 1988; Endler 1992, 1998). Over the past decades, scientists have established
33 that signals evolve in an ecological context and are dependent on environmental conditions (Endler,
34 1992, 1993a,b). Just as different habitats may favor different combinations of ecomorphological
35 traits to maximize performance and fitness (Arnold, 1983), they may also shape different forms
36 of a signal, so as to maximize its transmission and detection (e.g. Seehausen 1997), or reduce its
37 detection by unintended recipients such as predators (Endler, 1984, 1990, 1991; Halfwerk et al.,
38 2014). This selective pressure may drive the local adaptation of communication signals.

39 One potential barrier to the maintenance of localized signal divergence is the homogenizing effect
40 of gene flow. Population genetics theory suggests that gene flow may counteract local adaptation
41 between localities and prevent divergence altogether, especially at small spatial scales, because
42 of the inflow of maladapted alleles or because of the breaking of linkage between coevolving loci
43 (Felsenstein, 1976; García-Ramos and Kirkpatrick, 1997; Dieckmann and Doebeli, 1999; Lenor-
44 mand, 2002; Hendry et al., 2007a). This genetic homogenization has been confirmed empirically
45 in systems such as stick insects (Nosil and Crespi, 2004) and stickleback (Hendry et al., 2007b).
46 Yet, examples of microgeographic adaptation, i.e. adaptation at smaller scales than the range of
47 dispersal, exist, highlighting the potential of some organisms to respond to selection in the face of
48 gene flow (see Richardson et al. 2014 and references therein). Examples include small scale adapta-
49 tion in fragmented areas in Australian fruit flies (Willi and Hoffmann, 2012), and local adaptation
50 to predation pressure in North American salamanders (Richardson and Urban, 2013). Therefore,
51 despite evidence that local adaptation may be particularly difficult at small spatial scales where
52 gene flow tends to cause adjoining populations to remain genetically homogeneous, the potential
53 adaptive response of species traits, in particular communication signals, to localized differences in
54 habitats remains relatively unknown (Richardson et al., 2014). Lizards of the neotropical genus
55 *Anolis* (Squamata: Dactyloidae) are an excellent group for studying the eco-evolutionary dynam-
56 ics of local adaptation and natural selection (Losos, 2009). A particularly conspicuous trait of
57 anoles is their dewlap, an extensible flap of skin that is typically sexually dimorphic and used as
58 a communication signal in courtship (Sigmund, 1983; Driessens et al., 2014, 2015) and territorial
59 displays (Losos, 1985; Macedonia and Stamps, 1994; Macedonia et al., 2013) as well as in predator
60 deterrence (Leal and Rodríguez-Robles, 1995, 1997; Leal and Rodriguez-Robles, 1997). Dewlap
61 characteristics vary widely among the approximately 400 species of the genus (Nicholson et al.,
62 2007). Interspecific variation in dewlap coloration is implicated in species recognition (Rand and
63 Williams, 1970; Williams, 1969; Williams and Rand, 1977; Losos, 1985; Macedonia and Stamps,
64 1994; Fleishman, 2000; Macedonia et al., 2013), and this function could have had a role in initiating
65 or reinforcing reproductive isolation during speciation (Lambert et al., 2013; Geneva et al., 2015;
66 Ng et al., 2017).

67 Within species, studies have shown a link between variation in dewlap coloration and differ-
68 ences in habitats or climatic conditions (Macedonia, 2001; Leal and Fleishman, 2002; Thorpe and
69 Stenson, 2002; Thorpe, 2002; Leal and Fleishman, 2004; Vanhooydonck et al., 2009; Ng et al., 2012,
70 2013, 2016; Vanhooydonck et al., 2009; Driessens et al., 2017). Some studies suggest that those
71 differences may be adaptive and that dewlaps may have evolved to maximize detectability given
72 local light conditions (Fleishman and Persons, 2001; Leal and Fleishman, 2002, 2004). Although
73 this claim is further supported by recent findings that dewlap colors are perceived differently un-
74 der different levels of shading (Fleishman et al., 2020), other studies found conflicting patterns
75 of between-habitat variation that did not support the sensory drive hypothesis (Fleishman et al.,
76 2009; Ng et al., 2012; Macedonia et al., 2014).

77 Previous studies investigating variation in anole dewlaps compared populations at relatively
78 large geographical scales, e.g. between islands (Vanhooydonck et al., 2009; Driessens et al., 2017)
79 or within large islands such as Puerto Rico (Leal and Fleishman, 2004) or Hispaniola (Ng et al.,

2012, 2016). These large scales and marine barriers should reduce gene flow (Ng and Glor, 2011; Lambert et al., 2013; Richardson et al., 2014; Ng et al., 2017). That said, examples do exist of divergence in dewlap coloration at smaller scales or between populations with high degrees of gene flow (Thorpe and Stenson, 2002; Thorpe, 2002; Stapley et al., 2011; Ng et al., 2016).

Anolis sagrei is widespread across islands of the West Indies (Reynolds et al., 2020). It has been the subject of numerous studies concerning local adaptation (Losos et al., 1994, 1997, 2001; Kolbe et al., 2012), biological invasion (Kolbe et al., 2008), and sexual selection (Tokarz, 2002; Tokarz et al., 2005; Tokarz, 2006; Driessens et al., 2014; Steffen and Guyer, 2014; Driessens et al., 2015) among many other topics. Between-island variation in the mainly orange-red color of its dewlap was shown to be better explained by climatic variables such as annual precipitation and solar radiation (proposed to affect the average vegetation type on each island and among other things, its ambient light environment, Driessens et al. 2017), than by proxies for biotic factors such as sexual selection or predation pressure (Vanhooydonck et al., 2009; Baeckens et al., 2018). How intra-island differences in habitat may contribute to the diversity of dewlap coloration, however, remains unexplored, and may reveal new insights into the scale of local differentiation despite gene flow.

Here, we analyzed the color characteristics of *A. sagrei* dewlaps within nine islands in the Bahamas and Cayman Islands. These island systems presently, if not historically, comprise relatively small islands, with no major geographic barriers within islands limiting dispersal for this species. These islands all share three characteristic native West Indian small-island habitat-types – beach scrub bush, closed-canopy primary coppice forest, and mangrove forest – that are often spatially intermingled. These habitats contrast in environmental parameters including vegetation community, light irradiance, humidity, and temperature (Howard, 1950; Schoener, 1968). The Cayman Islands and the Bahamas have been colonized independently by *A. sagrei* from Cuba (van de Schoot 2016 unpublished thesis; Reynolds et al. 2020), such that these archipelagos constitute an ideal suite of natural replicates to explore within-island dewlap diversity across multiple islands.

Our sampling design included sites in close proximity; the median distance between two sites within an island was 8.45km. While this species has traditionally been considered territorial, a recent study revealed that they are polygynandrous and that gene flow is not impeded by territorial-like behaviors exhibited by some males (Kamath and Losos 2018). Combining reflectance spectrometry and supervised machine learning, we tested for divergence in dewlap phenotype between habitats within islands and between islands across part of the range of *A. sagrei*. We predicted that if light conditions in the environment indeed drive color evolution, dewlaps should be most similar between beach scrub and mangrove forest, which both have high levels of light irradiance, compared to the darker, closed-canopy coppice forest. If detectability is maximized given the local conditions, we expected darker and more contrasting dewlaps in high irradiance habitats. Finally, if habitat characteristics are strong determinants of dewlap color variation, similar patterns should be observed across multiple islands (Harvey and Pagel, 1991; Losos, 2011).

124 **Methods**

125 **Data collection**

126 We sampled 455 male *A. sagrei* from seven islands in the Bahamas Archipelago – Abaco, North
127 Andros, South Andros, South Bimini, Eleuthera, Long Island, and Ragged Island – and two in the
128 Cayman Islands – Cayman Brac and Little Cayman (Figure S1A). These islands were chosen to
129 span the breadth of the West Indian range of *A. sagrei*, because they have highly similar habitat
130 types, and because the *A. sagrei* on each island group are derived from ancient and distinct
131 colonization events from Cuba (i.e. relatively evolutionarily independent, Reynolds et al. 2020).
132 Three habitats were sampled on each island based on characterizations by Howard (1950) and
133 Schoener (1968). Each habitat is clearly distinguishable by its dominant vegetation type — xeric
134 beach scrub (open, relatively dry habitat consisting of low vegetation or isolated trees), primary
135 coppice forest (closed-canopy forest) and mangrove forest (wet coastal habitat with trees growing
136 in brackish water and high light penetration, although lizards were sampled in dry soil areas).
137 Sample sizes are given in Table S1. Our sampling design enabled us to test for differences between
138 habitats at a coarse and fine geographical scale. The median distance between two localities within
139 an island was 8.5km (Figure S1B), and 79.3% of all pairwise distances within islands were less
140 than 50km. Additionally, there are no major barriers to dispersal (such as mountains or grassland)
141 on any of the islands that we sampled.

142 **Reflectance measurements**

143 We measured reflectance between 300nm and 700nm wavelength, a range from ultraviolet to red
144 that encompasses the colors visible to most lizards and vertebrates in general (Lazareva et al.,
145 2012). Measurements were taken with an Ocean Optics USB4000 spectrometer, a pulsed Xenon
146 light source (PX-2, Ocean Optics, Largo, FL, USA) and a reflectance probe protected by a black
147 anodized aluminum sheath. Measurements were taken with a 45-degree inclination to prevent
148 specular reflection (Endler, 1990). The device was regularly standardized with a Spectralon white
149 standard (Labsphere, North Sutton, NH, USA). Reflectance was measured at the center of the
150 dewlap. Reflectance curves were smoothed using the R package pavo (Maia et al., 2013) as well
151 as with custom R functions, down to one reflectance value at each nanometer in wavelength from
152 300 to 700nm.

153 **Analysis**

154 We tested for detectable differences in dewlap coloration between populations from different habi-
155 tats across islands by following an analytic pipeline in several steps. First, we used multivariate
156 analyses of variance to assess the relative contributions of islands, habitats and habitat-by-island
157 interactions on the partitioning of variation in color space. Second, and provided that habitat-by-
158 island interactions were found, we investigated habitat-differences in dewlap color for each island
159 separately using machine learning classification. Third, for each island where multivariate differ-
160 ences were detected using our machine learning pipeline, we used univariate analyses of variance
161 to decompose the signal among the different dimensions of color space. Fourth, for each signifi-
162 cant between-habitat variation found in univariate analyses, we used post-hoc tests to determine
163 which habitats were responsible for the differences. Last, to get insights into the spatial scale of
164 phenotypic variation, for each significant contrast between two habitats detected along a given
165 dimension on a given island, we performed multiple pairwise Wilcoxon tests to assess differences
166 in dewlap coloration among the sites involved in that significant contrast, and recorded the geo-
167 graphical distance between sites that were found significant. In parallel, we tested a possible effect
168 of isolation-by-distance, an alternative cause of phenotypic divergence between localities, based on
169 diffusion approximation and dispersal distance, irrespective of habitat types. We did so using a
170 permutation test to assess the significance of the correlation between geographical distances and
171 phenotypic distances among sites within each island.

172

173 All analyses in this study were performed in R 3.6.1 (R Core Team, 2019).

Dimensionality reduction

174

Because neighboring wavelengths are highly collinear and redundant in reflectance, we reduced the dimensionality of the data using principal component analysis (PCA), as per Cuthill et al. (1999) and Leal and Fleishman (2002). We performed PCA on data from all islands combined, as well as on each island separately and systematically retained the first four principal components (PC), which together always explained more than 88.8% of the variance across islands (Table S2). PCs need not represent the same wavelengths across islands because they are fitted on different datasets. Nevertheless, PC1 was highly collinear with brightness for all islands (Figure S11), while the other PCs captured the chromatic variation (i.e. irrespective of brightness) in dewlap color.

Among-island variance partitioning

183

We performed a two-way nonparametric multivariate analysis of variance (PERMANOVA, Anderson 2001, R package vegan, Oksanen et al. 2019) to identify differences in coloration between islands, habitats and habitats within islands, using principal components fitted on data from all islands together. We used a nonparametric test because although no multivariate outliers were detected based on the Mahalanobis distance, the assumption of multivariate normality was violated in several habitats on several islands (Henze-Zirkler's test, Henze and Zirkler 1990, R package MVN, Korkmaz et al. 2014, $P < 0.05$, Table S3).

Within-island machine learning

191

We performed a machine learning classification analysis on the first four principal components within each island separately, using random forests (Breiman, 2001). Random forests are a versatile, intuitive, and powerful algorithm commonly used in machine learning, using decision trees to predict the labels of particular observations based on their multivariate coordinates. These coordinates, or variables, are passed through a series of successive decision nodes, each examining a given variable of any given observation (James et al., 2013). The prediction for each observation is an aggregate over a large number of decision trees, each tree being trained on a subset of observations sampled with replacement from the dataset, and each tree being allowed to examine only a subset of the variables. This allows the random forest to overcome the individual errors of all trees in the predictions it makes.

To detect differences in dewlap coloration between habitats, we measured the success of random forests in reassigning individual lizards to their correct habitat of origin, based solely on their principal component scores. In machine learning, this so-called cross-validation procedure is typically done in two steps (James et al., 2013). First, a random forest is trained in recognizing features of dewlap coloration most associated with the different habitats, by being presented with multiple observations, making predictions about them, and updating its own decision rules based on whether the prediction deviates from the truth. Then, once trained, the patterns that the random forest has learned to recognize are tested by presenting new, previously unseen observations to the random forest, and measuring the proportion of correct predictions. This proportion, or success score, can then be statistically assessed against random guessing using a binomial test.

The cross-validation procedure requires that the data be split into a training set and a testing set. To remove any bias due to the set that is being sampled for training, it is common practice to use k-fold cross-validation (James et al., 2013), where the data are split into k random bins and k independent machines are trained, each taking one of the bins as a testing set and the rest for training, and where classification success is measured by summing all correct classifications from the k machines.

Here, we used a k-fold cross-validation procedure with $k = 5$, where each training set consisted of 80% of the data and the machine was tested on the remaining 20%. Each training set was conditioned on containing at least five lizards from each of the three habitats. We also down-sampled the training set to the sample size of the least represented habitat, to ensure that the different habitats were equally represented. To further remove any bias due to the specific random split into the different bins, we replicated each k-fold cross-validation five times. We then averaged the five resulting confusion matrices across replicates, where each confusion matrix shows the number

of lizards from each habitat reassigned into each habitat. For each island, we then used the average proportion of correctly reassigned lizards (i.e. the proportion of observations on the diagonal of the average confusion matrix) as an estimate of classification success. This score was tested against random guessing by comparing it to a binomial distribution with number of trials being the number of lizards on that island and success probability 1/3, representing the rate of successful classification by chance when three habitats are involved.

We used the machine learning fitting functions in the R package rminer (Cortez, 2020), which calls random forest routines from the randomForest package (Liaw and Wiener 2002, implementation from the original random forest algorithm Breiman, 2001). For each random forest, we optimized the number of trees in the forest and the number of variables examined by each tree using the grid hyperparameter search procedure implemented in rminer, to choose between two numbers of trees (500 or 1,000) and four numbers of principal components examined per tree (1 to 4), using rminer's ordered holdout validation method with 2/3 of the data used for training.

We validated the results of our analysis by using two other widely used machine learning classification methods: linear discriminant analysis and support vector machines (Cristianini and Shawe-Taylor, 2000; James et al., 2013), both accessible in rminer (Cortez, 2020).

To know which wavelengths were most used to assign data points to each habitat, we trained another set of random forests, this time directly on reflectance data (taken every 5nm from 300 to 700nm) instead of principal components. We recorded the relative importance of each wavelength for each habitat, as measured by the mean decrease in accuracy during wavelength permutation, implemented in the randomForest package (Liaw and Wiener, 2002).

Univariate analyses

For each island where significant differences in dewlap coloration were detected between habitats, we used multiple univariate analyses of variance (ANOVA) to identify possible principal components underlying the observed differences. We constructed our ANOVA models in two steps, as per Zuur (2009). In a first step, we accounted for heterogeneity of variances across groups by systematically comparing the goodness-of-fit of an ANOVA model estimated with ordinary least squares (OLS) with that of a model estimated with generalized least squares (GLS), which allowed one estimate of residual variance per habitat (using the R package nlme, Pinheiro and Bates 2000; Pinheiro et al. 2020). Both models were fitted with restricted maximum likelihood (REML). Goodness-of-fit was estimated using Akaike's Information Criterion corrected for small sample sizes (AICc, R package MuMIn, Bartoń 2019), and the estimation method yielding the lowest AICc was retained. In a second step, we re-fitted the retained model with maximum likelihood (ML) to test for the effect of habitat type using likelihood ratio tests (LRT) between a model including a habitat-term and a null model lacking the habitat-term.

We evaluated the normality of the standardized residuals (residuals divided by their standard error, which can differ among habitats in a GLS model) of each fitted ANOVA model using Shapiro-Wilk's test, with P-values adjusted for multiple testing using the Benjamini-Hochberg correction (Benjamini and Hochberg, 1995). In cases where significant deviations from normality were detected ($P_{adj} < 0.05$, Table S4) we performed Kruskal-Wallis's nonparametric test to back up the ANOVA results.

To know which habitat-populations were different from which in dewlap coloration, we performed different post-hoc multiple comparison tests (all implemented in the PMCMRplus package, Pohlert, 2020), depending on which assumptions were met. In cases where normality and homoscedasticity were met (i.e. OLS-ANOVA was the best fit), we used Tukey's honest significant difference test. When normality was met but not homoscedasticity (i.e. GLS-ANOVA was the best fit), we used Dunnett's T3 test. Finally, whenever we used Kruskal-Wallis's test because the ANOVA residuals were not normally distributed, we used Nemenyi's test for post-hoc comparisons.

Spatial autocorrelation

281

We tested for within-island spatial autocorrelation between the geographical distances among sampling sites and their Euclidean distances in multivariate color space (mean PC1 to PC4 per site, Table S5), regardless of habitat type. For this, we performed Mantel's test (Legendre and Legendre 2012, R package vegan; Oksanen et al. 2019) on each island, using 999 permutations and geographical distances computed as geodesic distances from latitude and longitude data (R package geosphere, Hijmans 2019).

282

283

284

285

286

287

288

Site differences

289

In this study, we were interested in the minimum spatial scale at which significant differences between habitats could be detected within islands. We performed multiple pairwise nonparametric Wilcoxon-Mann-Whitney tests (Hollander et al., 2013) to compare dewlap coloration between sites with different habitat types, for each pair of habitats and each variable where significant differences were detected with our analyses of variance. The P-values were adjusted using a Benjamini-Hochberg correction for multiple testing (Benjamini and Hochberg, 1995).

290

291

292

293

294

295

296 **Results**

297 We tested for variation in *A. sagrei* dewlap coloration between populations living in three characteristic habitat types across nine islands that span the West Indian range of the brown anole (beach
298 scrub, primary coppice and mangroves). We found that most of the variation in coloration was
299 partitioned between islands (two-way PERMANOVA, $F(df = 8) = 45.38$, $P = 0.001$, explained
300 variance $R^2 = 41.4\%$). Nonetheless, we did find evidence for differences in dewlap coloration
301 between habitat types, and those were mostly island-specific (habitat-by-island interaction term,
302 $F(16) = 4.78$, $P = 0.001$, $R^2 = 8.7\%$), with a significant portion of the variation explained by
303 an habitat effect across all islands, but this effect was relatively small ($F(2) = 4.46$, $P = 0.001$,
304 $R^2 = 1\%$).

305
306 We subsequently tested for differences in dewlap coloration between habitat-populations within
307 each island, using within-island principal component scores (to maximize the variation captured
308 for each island, see Methods). Our within-island random forest classification analyses revealed
309 detectable differences in dewlap coloration on seven out of the nine islands in our sample: Abaco,
310 Bimini, Cayman Brac, Eleuthera, Little Cayman, Long Island, and North Andros. The accuracy of
311 random forest classification exceeded random expectation more often than expected by chance for
312 all these islands (Table 1). Accuracy was as high as 74.8% for Cayman Brac. We obtained similar
313 results using other machine learning approaches such as support vector machines (Table S7) and
314 linear discriminant analysis (Table S8). We describe in details the specific differences detected on
315 each island in the Appendix, and focus here on the general patterns emerging from our data.

316
317 Overall, we found significant differences in dewlap coloration between populations that were
318 often in close geographical proximity. On Bimini, notably, we found a significant difference be-
319 tween dewlaps from beach scrub and primary coppice forest, at a distance of a few hundred meters,
320 making this contrast the smallest geographical scale at which differences in coloration were found
321 in our study (Fig. S3). We also detected significant differences in dewlap coloration at distances
322 below one kilometer on Abaco (Fig. S2G), and at distances between one and ten kilometers on
323 Bimini (Fig. S3G), Cayman Brac (Fig. S4G), Little Cayman (Fig. S6G), Long Island (Fig. S7G)
324 and North Andros (Fig. S8G).

325
326 We found evidence of spatial autocorrelation in dewlap coloration between the sites within is-
327 lands for Abaco (Table 2), suggesting that populations from closer sites tend to have more similar
328 dewlaps on this island than expected by chance. Abaco was the island we sampled at the largest
329 scale, with some sites nearly a hundred kilometers away from each other (Fig. 2A). That said, some
330 sites were also in close proximity, and significant differences in coloration were detected between
331 habitats sometimes less than a kilometer away (Fig. S2G), suggesting that differences in dewlap
332 coloration between distant sites may be partly attributable to isolation-by-distance, but this may
333 not necessarily be the case for sites in close proximity. We did not find evidence for spatial auto-
334 correlation on other islands than Abaco (although Eleuthera was nearly significant, Table 2).

335
336 A striking feature of our data was inconsistency in between-habitat differences among islands,
337 in terms of which habitats differ from which, which dimensions of coloration were involved, and
338 in which direction. For example, while on Cayman Brac the random forests could well distinguish
339 between all three habitats (Fig. S4D), on Abaco dewlaps from beach scrub and primary coppice
340 were often mistaken, and on Bimini beach scrub dewlaps were more often classified into primary
341 coppice or mangrove than into beach scrub (Fig. S3D). In terms of variable importance, for multi-
342 ple islands the random forests used information in the UV range to discriminate between at least
343 some habitats, particularly on Abaco (Fig. S2F), Bimini (Fig. S3F), Cayman Brac (Fig. S4F),
344 Little Cayman (Fig. S6F) and Long Island (Fig. S7F), but differences in UV reflectance involved
345 different habitats and were in different directions among these islands.

346
347

Discussion

348

Two main insights follow from our results. First, we detected significant differences in dewlap coloration between habitats within seven out of the nine islands investigated (excluding North Andros where the follow-up univariate analyses were not significant), suggesting a putatively high potential for local differentiation of dewlap coloration in *Anolis sagrei*. Second, we found differences in coloration along different dimensions of color space, and in different directions, on different islands.

349

350

351

352

353

354

355

356

357

358

359

360

361

362

363

364

365

366

367

368

369

370

371

372

373

374

375

376

377

378

379

380

381

382

383

384

385

386

387

388

389

390

391

392

393

394

395

396

397

398

399

400

401

402

403

404

Detectable differences in dewlap color between populations are surprising, as habitats were often in close geographical proximity to each other (as close as a few hundred meters on Bimini and most of the time within ten kilometers), and we would have expected gene flow to cause a more homogeneous distribution of color phenotypes within islands. While little is known about the cruising range of individuals from our study populations (but see Steinberg and Leal 2017; Kamath and Losos 2018b for other systems), *A. sagrei* are polygynandrous (both males and females mate with multiple mates, Kamath and Losos 2017, 2018a,b), thus offering opportunity for gene flow, especially given that lizards were distributed continuously and at high densities within the islands we sampled. Consistent with that, while populations from different islands were monophyletic, individuals within islands were not monophyletic with respect to habitat based on mitochondrial haplotypes (van de Schoot 2016 unpublished thesis).

Several scenarios could account for these findings. One explanation is an adaptive one: populations living in different habitats could be phenotypically adapted to their local environmental conditions. Given that the brightly colored dewlap of *A. sagrei* is used as a communication signal, its color may be a target for selection if the transmission or perception of the signal differs from one habitat to another, for example because of differences in ambient light, according to the sensory-drive hypothesis (Endler and McLellan, 1988; Endler, 1992, 1998). The sensory-drive hypothesis has been tested multiple times for dewlap coloration in *Anolis* lizards, with mixed results. Some authors found support for it (Leal and Fleishman, 2002, 2004), while others found differences in dewlap coloration between habitats inconsistent with the sensory-drive hypothesis (Fleishman et al., 2009; Ng et al., 2012).

If our results were an example of sensory drive, we would have expected to see consistent differences between populations from different habitats across islands, given the apparent environmental consistency each of the three habitat types across the islands we sampled. In particular, we would have expected divergence in line with increased detectability given local light conditions, such as the high contrasts with background vegetation found in the UV range in Leal and Fleishman (2002) and Leal and Fleishman (2004). We might also have expected mangrove and beach scrub lizards, both inhabiting areas with high light penetration, to have more similar dewlaps, and to differ significantly from lizards from the coppice habitat, where irradiance is low. Instead, we found inconsistencies in the way dewlap color differed between habitats across islands. While short-wavelengths (UV reflectance) were often involved in color differences, they were not involved on all islands where significant differences were detected. On some islands, other or additional variables differed, such as brightness, red reflectance or the reflectance at the ends of the spectrum visible to *Anolis* lizards (UV and red, Lazareva et al. 2012) relative to intermediate wavelengths (blue-to-yellow). Similar portions of the spectrum were sometimes involved in opposite directions on different islands, such as on Abaco and Cayman Brac, where mangrove lizards had a higher UV-reflectance than beach scrub lizards on the former, but a lower UV-reflectance on the latter. Overall, the observed heterogeneity of divergence patterns across islands provides no support to a sensory-drive explanation.

It is presently not known if the reported differences in coloration have a genetic basis. Yet, we find it unlikely that these differences arose through phenotypic plasticity, as although the carotenoids that partly make up the red and orange colors of anole dewlaps must be found in the diet (Goodwin, 1984; Hill et al., 2002; Hill and McGraw, 2006), studies testing the effect of carotenoid deprivation (Steffen et al., 2010; Ng et al., 2013) and heritability (Cox et al., 2017) of dewlap coloration in *A. sagrei* and *A. distichus* (another species with a carotenoid-based dewlap), found little support for phenotypic and developmental plasticity in dewlap coloration. One exception is a study demonstrating that lizards heavily parasitized by skin mites had duller dewlaps

405 (Cook et al., 2013), but we found no sign of that in our study.

406

407 Genetic drift could contribute to some of the observed variation. Indeed, while only Abaco
408 showed significant patterns consistent with isolation-by-distance (which may emerge under limited
409 dispersal and drift, Wright 1943; Kimura and Weiss 1964; Slatkin 1987), there may have been
410 too few sites on most islands to conclusively detect it, and even then, the absence of detectable
411 isolation-by-distance may not necessarily constitute evidence for the absence of drift. Besides,
412 spatial autocorrelation was the strongest on islands sampled at the largest scales (e.g. some sites
413 on Abaco were nearly 100km apart, and Eleuthera – the second strongest signal, albeit nonsignif-
414 icant – had sites more than 30km apart), such that it is possible that neutral processes and/or
415 dispersal limitations might contribute to shaping variation over long distances. That said, many
416 significant differences were found between habitats in close proximity, contrary to what would be
417 expected under isolation-by-distance, including on islands where spatial autocorrelation was de-
418 tected. Moreover, *A. sagrei* was distributed across the islands continuously, usually at relatively
419 high population densities, rather than in small and isolated populations particularly prone to drift.
420 Together with the fact that isolation-by-distance may not necessarily only emerge from drift (e.g.
421 if there is a spatial environmental gradient), this indicates that genetic drift may have limited
422 potential to explain the differences observed between habitats, at least at a local scale.

423

424 In this study, we found larger differences among than within islands, a pattern already reported
425 and linked to climatic conditions (Driessens et al., 2017) and to densities of predators and of anole
426 congeners (Vanhooydonck et al., 2009; Baeckens et al., 2018). Differences among habitats within
427 islands, however, are still difficult to account for. Remaining hypotheses may include, for exam-
428 ple, runaway sexual selection (i.e. arbitrary preferences of females for some colors over others,
429 Andersson 1994) operating in different directions across islands, but no evidence so far suggests
430 that dewlap is a target of mate choice in anoles (Tokarz, 2002; Tokarz et al., 2005; Lailvaux and
431 Irschick, 2006; Nicholson et al., 2007). Another hypothesis is that the different genetic constitutu-
432 tions of different islands, perhaps resulting from founder effects (the islands have been colonized
433 independently, van de Schoot 2016 unpubl.; Driessens et al. 2017; Reynolds et al. 2020), may have
434 predisposed populations to adapt differently to similarly selective circumstances. Either way, new
435 data would be needed to test these hypotheses. Visual modeling, for instance, would be a valuable
436 follow-up analysis as it would indicate whether the differences we detected may be detectable by
437 the organisms themselves (as has been done in *Anolis* in Leal and Fleishman 2004; Fleishman
438 et al. 2020), but such approach requires irradiance profiles from the habitats in the field, which we
439 presently do not have.

440

441 Last, some patterns may arise owing to factors we did not measure. For one, signalling can be a
442 multifaceted behavior, and while we focused on dewlap color here, other potentially important el-
443 ements of behavioral interactions include dewlap size, dewlap patterning, display activity as well as
444 behaviors associated with dewlap extensions such as headbobs and pushups (Vanhooydonck et al.,
445 2005; Driessens et al., 2014, 2015; Lailvaux et al., 2015). These have been linked in various ways
446 (and not always consistently across studies) not only with environmental variables but also with
447 proxies for sexual selection, predation and species recognition (Vanhooydonck et al., 2005; Lailvaux
448 and Irschick, 2007; Vanhooydonck et al., 2009; Driessens et al., 2017; Baeckens et al., 2018), which
449 do vary among islands but may also vary within islands, possibly in interaction with dewlap color.
450 Besides, while the three habitats were clearly recognizable and consistent across islands to the hu-
451 man eye (Fig. 1), we did not precisely quantify irradiance within each habitat and did not consider
452 any more subtle environmental differences there might have been for a given habitat among islands.

453

454 Altogether, our results show that dewlap color of *A. sagrei* commonly varies between habitat
455 types, even in close geographical proximity, within islands of the West Indies. However, coloration
456 differs in different ways across similar habitats from one island to another. We discussed several
457 non-mutually exclusive mechanisms that could explain these observations. Nevertheless, heteroge-
458 neous patterns of divergence across islands do not support an adaptive sensory-drive scenario, and
459 we propose that within-island dewlap color variation may be underlain by a more subtle mosaic of
460 factors.

References

Anderson, M. J. (2001). A new method for non-parametric multivariate analysis of variance.	461
<i>Austral Ecology</i> , 26(1):32–46.	462
463	
Andersson, M. B. (1994). <i>Sexual Selection</i> . Monographs in Behavior and Ecology. Princeton University Press, Princeton, N.J.	464
465	
Arnold, S. J. (1983). Morphology, Performance and Fitness. <i>American Zoologist</i> , 23(2):347–361.	466
Baeckens, S., Driessens, T., and Van Damme, R. (2018). The brown anole dewlap revisited: Do predation pressure, sexual selection, and species recognition shape among-population signal diversity? <i>PeerJ</i> , 6:e4722.	467
468	
469	
Bartoń, K. (2019). MuMIn: Multi-Model Inference.	470
Benjamini, Y. and Hochberg, Y. (1995). Controlling the false discovery rate: A practical and powerful approach to multiple testing. <i>Journal of the Royal Statistical Society Series B</i> , 57:289–300.	471
472	
473	
Bradbury, J. W. and Vehrencamp, S. L. (2011). <i>Principles of Animal Communication</i> . Sinauer Associates, Sunderland, Mass, 2nd ed edition.	474
475	
Breiman, L. (2001). Random forests. <i>Machine Learning</i> , 45(1):5–32.	476
Cook, E. G., Murphy, T. G., and Johnson, M. A. (2013). Colorful displays signal male quality in a tropical anole lizard. <i>Naturwissenschaften</i> , 100(10):993–996.	477
478	
Cortez, P. (2020). Rminer: Data Mining Classification and Regression Methods.	479
Cox, R. M., Costello, R. A., Camber, B. E., and McGlothlin, J. W. (2017). Multivariate genetic architecture of the <i>Anolis</i> dewlap reveals both shared and sex-specific features of a sexually dimorphic ornament. <i>Journal of Evolutionary Biology</i> , 30(7):1262–1275.	480
481	
482	
Cristianini, N. and Shawe-Taylor, J. (2000). <i>An Introduction to Support Vector Machines and Other Kernel-based Learning Methods</i> . Cambridge University Press, first edition.	483
484	
Cuthill, I. C., Bennett, A. T. D., Partridge, J. C., and Maier, E. J. (1999). Plumage Reflectance and the Objective Assessment of Avian Sexual Dichromatism. <i>The American Naturalist</i> , 153(2):183–200.	485
486	
487	
Dieckmann, U. and Doebeli, M. (1999). On the origin of species by sympatric speciation. <i>Nature</i> , 400(6742):354–357.	488
489	
Driessens, T., Baeckens, S., Balzarolo, M., Vanhooydonck, B., Huyghe, K., and Van Damme, R. (2017). Climate-related environmental variation in a visual signalling device: The male and female dewlap in <i>Anolis sagrei</i> lizards. <i>Journal of Evolutionary Biology</i> , 30(10):1846–1861.	490
491	
492	
Driessens, T., Huyghe, K., Vanhooydonck, B., and Van Damme, R. (2015). Messages conveyed by assorted facets of the dewlap, in both sexes of <i>Anolis sagrei</i> . <i>Behavioral Ecology and Sociobiology</i> , 69(8):1251–1264.	493
494	
495	
Driessens, T., Vanhooydonck, B., and Van Damme, R. (2014). Deterring predators, daunting opponents or drawing partners? Signaling rates across diverse contexts in the lizard <i>Anolis sagrei</i> . <i>Behavioral Ecology and Sociobiology</i> , 68(2):173–184.	496
497	
498	
Endler, J. (1998). Sensory ecology, receiver biases and sexual selection. <i>Trends in Ecology & Evolution</i> , 13(10):415–420.	499
500	
Endler, J. A. (1984). Natural and sexual selection on color patterns in poeciliid fishes. In Balon, E. K. and Zaret, T. M., editors, <i>Evolutionary Ecology of Neotropical Freshwater Fishes</i> , volume 3, pages 95–111. Springer Netherlands, Dordrecht.	501
502	
503	
Endler, J. A. (1990). On the measurement and classification of colour in studies of animal colour patterns. <i>Biological Journal of the Linnean Society</i> , 41(4):315–352.	504
505	

- 506 Endler, J. A. (1991). Variation in the appearance of guppy color patterns to guppies and their
507 predators under different visual conditions. *Vision Research*, 31(3):587–608.
- 508 Endler, J. A. (1992). Signals, Signal Conditions, and the Direction of Evolution. *The American
509 Naturalist*, 139:S125–S153.
- 510 Endler, J. A. (1993a). The Color of Light in Forests and Its Implications. *Ecological Monographs*,
511 63(1):1–27.
- 512 Endler, J. A. (1993b). Some general comments on the evolution and design of animal communi-
513 cation systems. *Philosophical Transactions of the Royal Society of London. Series B: Biological
514 Sciences*, 340(1292):215–225.
- 515 Endler, J. A. and McLellan, T. (1988). The Processes of Evolution: Toward a Newer Synthesis.
516 *Annual Review of Ecology and Systematics*, 19(1):395–421.
- 517 Felsenstein, J. (1976). The Theoretical Population Genetics of Variable Selection and Migration.
518 *Annual Review of Genetics*, 10(1):253–280.
- 519 Fleishman, L. J. (2000). *Signal Function, Signal Efficiency and the Evolution of Anoline Lizard
520 Dewlap Color*, pages 209–236. Tapir Academic, Trondheim.
- 521 Fleishman, L. J., Leal, M., and Persons, M. H. (2009). Habitat light and dewlap color diversity in
522 four species of Puerto Rican anoline lizards. *Journal of Comparative Physiology A*, 195(11):1043–
523 1060.
- 524 Fleishman, L. J. and Persons, M. (2001). The influence of stimulus and background colour on
525 signal visibility in the lizard *Anolis cristatellus*. *The Journal of Experimental Biology*, 204(Pt
526 9):1559–1575.
- 527 Fleishman, L. J., Wadman, C. S., and Maximov, K. J. (2020). The interacting effects of total
528 light intensity and chromatic contrast on visual signal visibility in an *Anolis* lizard. *Animal
529 Behaviour*, page S0003347220302037.
- 530 García-Ramos, G. and Kirkpatrick, M. (1997). Genetic Models of Adaptation and Gene Flow in
531 Peripheral Populations. *Evolution*, 51(1):21–28.
- 532 Geneva, A. J., Hilton, J., Noll, S., and Glor, R. E. (2015). Multilocus phylogenetic analyses of
533 Hispaniolan and Bahamian trunk anoles (*distichus* species group). *Molecular Phylogenetics and
534 Evolution*, 87:105–117.
- 535 Goodwin, T. W. (1984). *The Biochemistry of the Carotenoids*. Springer Netherlands, Dordrecht.
- 536 Halfwerk, W., Jones, P. L., Taylor, R. C., Ryan, M. J., and Page, R. A. (2014). Risky Ripples Allow
537 Bats and Frogs to Eavesdrop on a Multisensory Sexual Display. *Science*, 343(6169):413–416.
- 538 Harvey, P. H. and Pagel, M. D. (1991). *The Comparative Method in Evolutionary Biology*. Oxford
539 Series in Ecology and Evolution. Oxford University Press, Oxford ; New York.
- 540 Hendry, A. P., Day, T., and Taylor, E. B. (2007a). Population Mixing and the Adaptive Divergence
541 of Quantitative Traits in Discrete Populations: A Theoretical Framework for Empirical Tests.
542 *Evolution*, 55(3):459–466.
- 543 Hendry, A. P., Taylor, E. B., and McPhail, J. D. (2007b). Adaptive Divergence and the Balance
544 Between Selection and Gene Flow: Lake and Stream Stickleback in the Misty System. *Evolution*,
545 56(6):1199–1216.
- 546 Henze, N. and Zirkler, B. (1990). A class of invariant consistent tests for multivariate normality.
547 *Communications in Statistics - Theory and Methods*, 19(10):3595–3617.
- 548 Hijmans, R. J. (2019). Geosphere: Spherical Trigonometry.
- 549 Hill, G. E., Inouye, C. Y., and Montgomerie, R. (2002). Dietary carotenoids predict plumage
550 coloration in wild house finches. *Proceedings of the Royal Society of London. Series B: Biological
551 Sciences*, 269(1496):1119–1124.

Hill, G. E. and McGraw, K. J., editors (2006). <i>Bird Coloration</i> . Harvard University Press, Cambridge, Mass.	552 553
Hollander, M., Wolfe, D. A., and Chicken, E. (2013). <i>Nonparametric Statistical Methods</i> . Wiley Series in Probability and Statistics. John Wiley & Sons, Inc, Hoboken, New Jersey, third edition edition.	554 555 556
Howard, R. A. (1950). Vegetation of the Bimini Island Group: Bahamas, B. W. I. <i>Ecological Monographs</i> , 20(4):317–349.	557 558
James, G., Witten, D., Hastie, T., and Tibshirani, R. (2013). <i>An Introduction to Statistical Learning</i> , volume 103 of <i>Springer Texts in Statistics</i> . Springer New York, New York, NY.	559 560 561
Kamath, A. and Losos, J. (2017). The erratic and contingent progression of research on territoriality: A case study. <i>Behavioral Ecology and Sociobiology</i> , 71(6):89.	562 563
Kamath, A. and Losos, J. (2018a). Reconsidering territoriality is necessary for understanding <i>Anolis</i> mating systems. <i>Behavioral Ecology and Sociobiology</i> , 72(7):106.	564 565
Kamath, A. and Losos, J. B. (2018b). Estimating encounter rates as the first step of sexual selection in the lizard <i>Anolis sagrei</i> . <i>Proceedings of the Royal Society B: Biological Sciences</i> , 285(1873):20172244.	566 567 568
Kimura, M. and Weiss, G. H. (1964). The Stepping Stone Model of Population Structure and the Decrease of Genetic Correlation with Distance. <i>Genetics</i> , 49(4):561–576.	569 570
Kolbe, J. J., Larson, A., Losos, J. B., and de Queiroz, K. (2008). Admixture determines genetic diversity and population differentiation in the biological invasion of a lizard species. <i>Biology Letters</i> , 4(4):434–437.	571 572 573
Kolbe, J. J., Leal, M., Schoener, T. W., Spiller, D. A., and Losos, J. B. (2012). Founder Effects Persist Despite Adaptive Differentiation: A Field Experiment with Lizards. <i>Science</i> , 335(6072):1086–1089.	574 575 576
Korkmaz, S., Goksuluk, D., and Zararsiz, G. (2014). MVN: An R Package for Assessing Multivariate Normality. <i>The R Journal</i> , 6(2):151–162.	577 578
Lailvaux, S. P. and Irschick, D. J. (2006). No Evidence for Female Association with High-Performance males in the Green Anole Lizard, <i>Anolis carolinensis</i> . <i>Ethology</i> , 112(7):707–715.	579 580
Lailvaux, S. P. and Irschick, D. J. (2007). The Evolution of Performance-Based Male Fighting Ability in Caribbean Anolis Lizards. <i>The American Naturalist</i> , 170(4):573–586.	581 582
Lailvaux, S. P., Leifer, J., Kircher, B. K., and Johnson, M. A. (2015). The incredible shrinking dewlap: Signal size, skin elasticity, and mechanical design in the green anole lizard (<i>Anolis carolinensis</i>). <i>Ecology and Evolution</i> , 5(19):4400–4409.	583 584 585
Lambert, S. M., Geneva, A. J., Luke Mahler, D., and Glor, R. E. (2013). Using genomic data to revisit an early example of reproductive character displacement in Haitian <i>Anolis</i> lizards. <i>Molecular Ecology</i> , 22(15):3981–3995.	586 587 588
Lazareva, O. F., Shimizu, T., and Wasserman, E. A. (2012). <i>How Animals See the World</i> <i>Comparative Behavior, Biology, and Evolution of Vision</i> . Oxford University Press.	589 590
Leal, M. and Fleishman, L. J. (2002). Evidence for habitat partitioning based on adaptation to environmental light in a pair of sympatric lizard species. <i>Proceedings of the Royal Society of London. Series B: Biological Sciences</i> , 269(1489):351–359.	591 592 593
Leal, M. and Fleishman, L. J. (2004). Differences in Visual Signal Design and Detectability between Allopatric Populations of <i>Anolis</i> Lizards. <i>The American Naturalist</i> , 163(1):26–39.	594 595
Leal, M. and Rodríguez-Robles, J. A. (1995). Antipredator Responses of <i>Anolis cristatellus</i> (Sauria: Polychrotidae). <i>Copeia</i> , 1995(1):155–161.	596 597

- 598 Leal, M. and Rodriguez-Robles, J. A. (1997). Antipredator Responses of the Puerto Rican Giant
599 Anole, *Anolis cuvieri* (Squamata: Polychrotidae). *Biotropica*, 29(3):372–375.
- 600 Leal, M. and Rodríguez-Robles, J. A. (1997). Signalling displays during predator–prey interactions
601 in a Puerto Rican anole, *Anolis cristatellus*. *Animal Behaviour*, 54(5):1147–1154.
- 602 Legendre, P. and Legendre, L. (2012). *Numerical Ecology*. Number 24 in *Developments in Environmental Modelling*. Elsevier, Amsterdam, third english edition edition.
- 604 Lenormand, T. (2002). Gene flow and the limits to natural selection. *Trends in Ecology & Evolution*, 17(4):183–189.
- 606 Liaw, A. and Wiener, M. (2002). Classification and Regression by randomForest. *R News*, 2(3):18–22.
- 608 Losos, J. B. (1985). An Experimental Demonstration of the Species-Recognition Role of *Anolis*
609 Dewlap Color. *Copeia*, 1985(4):905–910.
- 610 Losos, J. B. (2009). *Lizards in an Evolutionary Tree: Ecology and Adaptive Radiation of Anoles*.
611 University of California Press.
- 612 Losos, J. B. (2011). Convergence, Adaptation, and Constraint. *Evolution*, 65(7):1827–1840.
- 613 Losos, J. B., Irschick, D. J., and Schoener, T. W. (1994). Adaptation and Constraint in the
614 Evolution of Specialization of Bahamian *Anolis* Lizards. *Evolution*, 48(6):1786–1798.
- 615 Losos, J. B., Schoener, T. W., Warheit, K. I., and Creer, D. (2001). Experimental studies of
616 adaptive differentiation in Bahamian *Anolis* lizards. *Genetica*, 112–113:399–415.
- 617 Losos, J. B., Warheit, K. I., and Schoener, T. W. (1997). Adaptive differentiation following
618 experimental island colonization in *Anolis* lizards. *Nature*, 387(6628):70–73.
- 619 Macedonia, J. M. (2001). Habitat light, colour variation, and ultraviolet reflectance in the Grand
620 Cayman anole, *Anolis conspersus*. *Biological Journal of the Linnean Society*, 73(3):299–320.
- 621 Macedonia, J. M., Clark, D. L., Riley, R. G., and Kemp, D. J. (2013). Species recognition of color
622 and motion signals in *Anolis grahami*: Evidence from responses to lizard robots. *Behavioral Ecology*,
623 24(4):846–852.
- 624 Macedonia, J. M., Clark, D. L., and Tamasi, A. L. (2014). Does Selection Favor Dewlap Colors that
625 Maximize Detectability? A Test with Five Species of Jamaican *Anolis* Lizards. *Herpetologica*,
626 70(2):157–170.
- 627 Macedonia, J. M. and Stamps, J. A. (1994). Species Recognition in *Anolis grahami* (Sauria,
628 Iguanidae): Evidence from Responses to Video Playbacks of Conspecific and Heterospecific
629 Displays. *Ethology*, 98(3-4):246–264.
- 630 Maia, R., Eliason, C. M., Bitton, P.-P., Doucet, S. M., and Shawkey, M. D. (2013). Pavo: An R
631 package for the analysis, visualization and organization of spectral data. *Methods in Ecology and Evolution*,
632 pages n/a–n/a.
- 633 Ng, J., Geneva, A. J., Noll, S., and Glor, R. E. (2017). Signals and Speciation: *Anolis* Dewlap
634 Color as a Reproductive Barrier. *Journal of Herpetology*, 51(3):437–447.
- 635 Ng, J. and Glor, R. E. (2011). Genetic differentiation among populations of a Hispaniolan trunk
636 anole that exhibit geographical variation in dewlap colour. *Molecular Ecology*, 20(20):4302–4317.
- 637 Ng, J., Kelly, A. L., MacGuigan, D. J., and Glor, R. E. (2013). The Role of Heritable and Dietary
638 Factors in the Sexual Signal of a Hispaniolan *Anolis* Lizard, *Anolis distichus*. *Journal of Heredity*,
639 104(6):862–873.
- 640 Ng, J., Landeen, E. L., Logsdon, R. M., and Glor, R. E. (2012). Correlation Between *Anolis*
641 Lizard Dewlap Phenotype and Environmental Variation Indicates Adaptive Divergence of a
642 Signal Important to Sexual Selection and Species Recognition. *Evolution*, 67(2):573–582.

Ng, J., Ossip-Klein, A. G., and Glor, R. E. (2016). Adaptive signal coloration maintained in the face of gene flow in a Hispaniolan <i>Anolis</i> Lizard. <i>BMC Evolutionary Biology</i> , 16(1):193.	643 644
Nicholson, K. E., Harmon, L. J., and Losos, J. B. (2007). Evolution of <i>Anolis</i> Lizard Dewlap Diversity. <i>PLoS ONE</i> , 2(3):e274.	645 646
Nosil, P. and Crespi, B. J. (2004). Does Gene Flow Constrain Adaptive Divergence or Vice Versa? A Test Using Ecomorphology and Sexual Isolation in <i>Timema cristinae</i> Walking-Sticks. <i>Evolution</i> , 58(1):102–112.	647 648 649
Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P. R., O'Hara, R. B., Simpson, G. L., Solymos, P., Stevens, M. H. H., Szoecs, E., and Wagner, H. (2019). Vegan: Community Ecology Package. R package version 2.5-6.	650 651 652
Pinheiro, J. and Bates, D. (2000). <i>Mixed-Effects Models in S and S-PLUS</i> . Statistics and Computing. Springer-Verlag, New York.	653 654
Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., and Team, R. C. (2020). Nlme: Linear and Nonlinear Mixed Effects Models.	655 656
Pohlert, T. (2020). PMCMRplus: Calculate Pairwise Multiple Comparisons of Mean Rank Sums Extended. R package version 1.4.4.	657 658
R Core Team (2019). R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing.	659 660
Rand, A. S. and Williams, E. E. (1970). An Estimation of Redundancy and Information Content of Anole Dewlaps. <i>The American Naturalist</i> , 104(935):99–103.	661 662
Reynolds, R. G., Kolbe, J. J., Glor, R. E., López-Darias, M., Gómez Pourroy, C. V., Harrison, A. S., Queiroz, K., Revell, L. J., and Losos, J. B. (2020). Phylogeographic and phenotypic outcomes of brown anole colonization across the Caribbean provide insight into the beginning stages of an adaptive radiation. <i>Journal of Evolutionary Biology</i> , 33(4):468–494.	663 664 665 666
Richardson, J. L. and Urban, M. C. (2013). Strong Selection Barriers Explain Microgeographic Adaptation in Wild Salamander Populations. <i>Evolution</i> , 67(6):1729–1740.	667 668
Richardson, J. L., Urban, M. C., Bolnick, D. I., and Skelly, D. K. (2014). Microgeographic adaptation and the spatial scale of evolution. <i>Trends in Ecology & Evolution</i> , 29(3):165–176.	669 670
Schoener, T. W. (1968). The <i>Anolis</i> Lizards of Bimini: Resource Partitioning in a Complex Fauna. <i>Ecology</i> , 49(4):704–726.	671 672
Seehausen, O. (1997). Cichlid Fish Diversity Threatened by Eutrophication That Curbs Sexual Selection. <i>Science</i> , 277(5333):1808–1811.	673 674
Sigmund, W. R. (1983). Female Preference for <i>Anolis carolinensis</i> Males as a Function of Dewlap Color and Background Coloration. <i>Journal of Herpetology</i> , 17(2):137–143.	675 676
Slatkin, M. (1987). Gene Flow and the Geographic Structure of Natural Populations. <i>Science</i> , 236(4803):787–792.	677 678
Stapley, J., Wordley, C., and Slate, J. (2011). No Evidence of Genetic Differentiation Between Anoles With Different Dewlap Color Patterns. <i>Journal of Heredity</i> , 102(1):118–124.	679 680
Steffen, J. E. and Guyer, C. C. (2014). Display behaviour and dewlap colour as predictors of contest success in brown anoles: Dewlap Colour and Behaviour in Contests. <i>Biological Journal of the Linnean Society</i> , 111(3):646–655.	681 682 683
Steffen, J. E., Hill, G. E., and Guyer, C. (2010). Carotenoid Access, Nutritional Stress, and the Dewlap Color of Male Brown Anoles. <i>Copeia</i> , 2010(2):239–246.	684 685
Steinberg, D. S. and Leal, M. (2017). An Unexpected Case of Homing in a Territorial Lizard. <i>Integrative and Comparative Biology</i> , 57:E418.	686 687

- 688 Thorpe, R. S. (2002). Analysis of Color Spectra in Comparative Evolutionary Studies: Molecular
689 Phylogeny and Habitat Adaptation in the St. Vincent Anole (*Anolis trinitatis*). *Systematic*
690 *Biology*, 51(4):554–569.
- 691 Thorpe, R. S. and Stenson, A. G. (2002). Phylogeny, Paraphyly and Ecological Adaptation of
692 the Colour and Pattern in the *Anolis Roquet* Complex on Martinique: Interaction Between
693 Phylogeny and Adaptation. *Molecular Ecology*, 12(1):117–132.
- 694 Tokarz, R. R. (2002). An Experimental Test of the Importance of the Dewlap in Male Mating
695 Success in the Lizard *Anolis sagrei*. *Herpetologica*, 58(1):87–94.
- 696 Tokarz, R. R. (2006). Importance of Prior Physical Contact with Familiar Females in the Devel-
697 opment of a Male Courtship and Mating Preference for Unfamiliar Females in the Lizard *Anolis*
698 *Sagrei*. *Herpetologica*, 62(2):115–124.
- 699 Tokarz, R. R., Paterson, A. V., and McMann, S. (2005). Importance of Dewlap Display in
700 Male Mating Success in Free-Ranging Brown Anoles (*Anolis sagrei*). *Journal of Herpetology*,
701 39(1):174–177.
- 702 van de Schoot, M. (2016). *Within and between Island Radiation and Genetic Variation in Anolis*
703 *Sagrei*. PhD thesis, Wageningen University, Wageningen, The Netherlands.
- 704 Vanhooydonck, B., Herrel, A., Meyers, J. J., and Irschick, D. J. (2009). What determines dewlap
705 diversity in *Anolis* lizards? An among-island comparison. *Journal of Evolutionary Biology*,
706 22(2):293–305.
- 707 Vanhooydonck, B., Herrel, A. Y., Van Damme, R., and Irschick, D. J. (2005). Does dewlap size
708 predict male bite performance in Jamaican *Anolis* lizards? *Functional Ecology*, 19(1):38–42.
- 709 Willi, Y. and Hoffmann, A. A. (2012). Microgeographic adaptation linked to forest fragmentation
710 and habitat quality in the tropical fruit fly *Drosophila birchii*. *Oikos*, 121(10):1627–1637.
- 711 Williams, E. E. (1969). The Ecology of Colonization as Seen in the Zoogeography of Anoline
712 Lizards on Small Islands. *The Quarterly Review of Biology*, 44(4):345–389.
- 713 Williams, E. E. and Rand, A. S. (1977). Species Recognition, Dewlap Function and Faunal Size.
714 *American Zoologist*, 17(1):261–270.
- 715 Wright, S. (1943). Isolation by Distance. *Genetics*, 28(2):114–138.
- 716 Zuur, A. F., editor (2009). *Mixed Effects Models and Extensions in Ecology with R*. Statistics for
717 Biology and Health. Springer, New York, NY.

Figures

718




Figure 1: Overview of our study design, including a map of the Bahamas and the Cayman Islands, on which are indicated the nine islands we sampled, two representatives of our study species *Anolis sagrei* with their dewlaps deployed, and the three types of habitats we considered on each island.




Figure 2: Comparison of dewlap coloration across habitats on Abaco. (A) Map of the island with the sampling sites colored by habitat. (B) Reflectance profiles of all the dewlaps on the island. (C) How reflectance profiles map onto the within-island principal components. (D) Confusion matrix showing the proportion of lizards from each (true) habitat reassigned to each (predicted) habitat by the random forests, based on the first four within-island principal components and averaged across replicates. Each column sums to one. (E) Within-island principal component scores across habitats. Bars indicate significant contrasts. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$.

719 **Tables**

Table 1: Random forest classification results. For each island are shown the sample size (N) and the proportion of correctly reassigned observations (or success score). P-values were computed using a binomial test and assess the significance of the observed success score relative to the score expected under random guessing. *, $P < 0.05$; **, $P < 0.01$, ***, $P < 0.001$.

Island	N	Score	P	
Abaco	86	0.623	< 0.0001	***
Bimini	57	0.460	0.0194	*
Cayman Brac	50	0.748	< 0.0001	***
Eleuthera	55	0.520	0.0023	**
Little Cayman	45	0.676	< 0.0001	***
Long Island	53	0.611	< 0.0001	***
North Andros	28	0.693	< 0.0001	***
Ragged Island	50	0.412	0.1259	
South Andros	31	0.419	0.1152	

Table 2: Mantel's test of spatial autocorrelation. For each island are shown the correlation (Pearson's ρ) between the matrix of phenotypic distances between populations from each site and the matrix of geographic distances between sites, where phenotypic distances are Euclidean distances between the mean phenotypes of each site in the multivariate space consisting of the first four within-island principal components. P-values assess the significance of the observed correlation against the correlation expected if population means were randomly permuted among sites (999 permutations). *, $P < 0.05$.

Island	ρ	P	
Abaco	0.439	0.032	*
Bimini	-0.725	1.000	
Cayman Brac	-0.737	0.833	
Eleuthera	0.827	0.058	
Little Cayman	-0.042	0.667	
Long Island	-0.077	0.583	
North Andros	-0.968	1.000	
Ragged Island	-0.363	0.708	
South Andros	0.963	0.167	

Table 3: Significance of habitat differences in dewlap coloration, using ANOVA for all islands where significant multivariate differences in dewlap coloration were detected by random forests. Model, best-fitting model (either OLS or GLS). AIC_c, corrected AIC score of the best-fitting model. ΔAIC_c , difference in AIC_c between the best-fitting model and the OLS-model. AIC_{cw}, AIC_c weight. Log-lik., log-likelihood. χ^2 , likelihood ratio. df, degrees of freedom. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$.

Island	Variable	AIC _c	ΔAIC_c	AIC _{cw}	Model	Log-lik.	χ^2	df	P
Abaco	PC1	255.81	2.06	0.737	OLS	-121.46	0.14	2	0.9318
Abaco	PC2	225.29	3.98	0.880	OLS	-105.64	31.77	2	< 0.0001 ***
Abaco	PC3	229.85	1.44	0.673	OLS	-108.01	27.04	2	< 0.0001 ***
Abaco	PC4	254.59	0.72	0.589	OLS	-120.82	1.41	2	0.4945
Bimini	PC1	162.92	-0.32	0.540	GLS	-72.43	10.03	2	0.0066 **
Bimini	PC2	165.36	3.08	0.824	OLS	-76.52	7.70	2	0.0212 *
Bimini	PC3	163.58	3.13	0.827	OLS	-75.58	9.59	2	0.0083 **
Bimini	PC4	172.47	2.43	0.771	OLS	-80.27	0.20	2	0.9035
Cayman Brac	PC1	136.64	-4.05	0.884	GLS	-59.29	13.81	2	0.0010 **
Cayman Brac	PC2	144.75	3.51	0.853	OLS	-66.24	8.41	2	0.0149 *
Cayman Brac	PC3	127.13	2.77	0.800	OLS	-56.86	27.16	2	< 0.0001 ***
Cayman Brac	PC4	147.37	4.33	0.897	OLS	-67.63	5.63	2	0.0600
Eleuthera	PC1	166.33	2.26	0.756	OLS	-77.29	0.49	2	0.7827
Eleuthera	PC2	155.78	-2.38	0.767	GLS	-68.74	12.80	2	0.0017 ***
Eleuthera	PC3	160.47	-0.22	0.527	GLS	-71.18	5.59	2	0.0613
Eleuthera	PC4	160.61	3.85	0.873	OLS	-74.27	6.54	2	0.0380 *
Little Cayman	PC1	130.60	2.50	0.777	OLS	-59.26	8.18	2	0.0167 *
Little Cayman	PC2	112.66	-3.61	0.859	GLS	-46.74	29.76	2	< 0.0001 ***
Little Cayman	PC3	118.32	1.41	0.669	OLS	-52.68	21.34	2	< 0.0001 ***
Little Cayman	PC4	135.58	2.53	0.780	OLS	-61.92	2.85	2	0.2410
Long Island	PC1	145.51	3.73	0.866	OLS	-66.41	16.58	2	0.0003 ***
Long Island	PC2	158.82	-1.29	0.656	GLS	-70.56	1.35	2	0.5103
Long Island	PC3	154.36	3.02	0.819	OLS	-71.10	7.19	2	0.0274 *
Long Island	PC4	155.59	0.47	0.558	OLS	-71.75	5.89	2	0.0525
North Andros	PC1	89.00	2.87	0.808	OLS	-39.05	0.35	2	0.8406
North Andros	PC2	74.74	-0.37	0.547	GLS	-27.50	17.24	2	0.0002 ***
North Andros	PC3	87.62	0.25	0.531	OLS	-38.28	1.89	2	0.3893
North Andros	PC4	73.56	5.39	0.937	OLS	-30.40	17.64	2	0.0001 ***

720 **Appendix**

721 Here we describe more precisely the patterns identified on each island.

722 On Abaco, dewlaps from the mangrove habitat were the best discriminated, while dewlaps from
723 the beach scrub habitat were often mistaken for dewlaps from the coppice habitat (Fig. 2D). Im-
724 portance analysis revealed that beach scrub and mangrove lizards mostly differed in reflectance in
725 the ultraviolet (UV) end of the spectrum (below 400nm, Fig. S2F), where mangrove dewlaps had
726 higher UV reflectance relative to beach scrub lizards, and coppice lizards had an intermediate UV
727 reflectance between the two other habitats (Fig. 2B). Consistent with this, our analyses of variance
728 detected significantly higher PC2 scores in mangrove lizards than in the two other habitats (Fig.
729 2E, Table 3), representing a higher UV-reflectance relative to red (Fig. 2C). Beach scrub lizards
730 also scored higher on PC3 (Fig. 2E, Table 3), indicating less curvature of the reflectance profile
731 and relatively higher reflectance at intermediate wavelengths (blue-to-yellow) than at the ends of
732 the range (Fig. 2C). Differences were detected between sites both at large ($\sim 100\text{km}$) and short
733 ($< 1\text{km}$) distances (Fig. S2G). Abaco was the only island where we detected significant spatial
734 autocorrelation (Table 2), that is, sites that were closer geographically tended to have populations
735 of lizards with more similar dewlap colors.

736 On Bimini, the random forests mostly correctly classified lizards from the coppice and mangrove
737 habitats while often misclassifying lizards from the beach scrub habitat (Fig. S3D). Relatively flat
738 importance profiles for beach scrub lizards suggested that brightness was used instead of a par-
739 ticular wavelength to identify some of the beach scrub dewlaps (Fig. S3F). Indeed, some beach
740 scrub dewlaps were substantially brighter than the rest (Fig. S3B, C), a pattern that was captured
741 by our analysis of variance along PC1 (i.e. brightness, Fig. S3C, E, Table 3). Coppice dewlaps
742 had significantly higher PC2 scores than mangrove dewlaps (Fig. S3E), suggesting a higher curva-
743 ture (higher UV and red reflectance than more intermediate wavelengths, Fig. S3C). For these two
744 habitats the random forests were most sensitive to UV reflectance (Fig. S3F). Beach scrub dewlaps
745 had higher PC3 scores than coppice dewlaps but it was not clear what properties of spectral shape
746 this principal component mapped onto (Fig. S3C). On this island, the beach scrub and coppice
747 habitats were separated by a few hundred meters, making this contrast the smallest geographical
748 scale at which differences in coloration were found in our study (Fig. S3G).

751 On Cayman Brac, all three habitats could be well discriminated against each other (Fig. S4D),
752 with UV reflectance appearing to be an important variable differentiating beach scrub and man-
753 grove dewlaps (Fig. S4F). In contrast, coppice dewlaps had a relatively flat importance profile,
754 suggesting that brightness made them more distinct rather than any particular wavelength (Fig.
755 S4F). Consistent with this, coppice dewlaps were significantly different from all other dewlaps
756 along PC1 (Fig. S4E, Table 3). At a distance between 2 and 3km (Fig. S4G), dewlaps in the
757 beach scrub habitat reflected more red light (as represented by PC2, Fig. S4C, E) and more UV
758 (as represented by PC3, along which coppice dewlaps were intermediate, Fig. S4C, E) than in the
759 mangrove habitat.

760 On Eleuthera, although random forests detected between-habitat differences in dewlap color,
761 other approaches did not (Tables S8 and S7), suggesting that the differences may be small.
762 On Eleuthera, beach scrub and mangrove dewlaps were guessed relatively correctly, but coppice
763 dewlaps were more often mistaken for mangrove dewlaps (Fig. S5D). Mangrove dewlaps had lower
764 PC2 scores than beach scrub dewlaps (Fig. S5E, indicating higher UV relative to red, Fig. S5C),
765 and higher PC4 scores than coppice dewlaps (Fig. S5E, suggesting more reflectance profiles with
766 more curvature, Fig. S5C). Random forests did not seem to consistently capture the wavelengths
767 responsible for these differences (Fig. S5F).

770 Little Cayman was characterized by a better discrimination of mangrove lizards from the rest
771 than between beach scrub and coppice lizards even though all habitats were relatively well dis-
772 criminated (Fig. S6D). Mangrove dewlaps were most distinct with respect to their reflectance in
773 short wavelengths (Fig. S6F), with significantly lower UV reflectance (as represented by PC2,
774 Fig. S6C, E, Table 3). Beach scrub lizards were characterized by brighter dewlaps than cop-
775 pice lizards (PC1), and also more convex curves, i.e. slightly higher UV and red reflectance (as
776

represented by higher PC3 scores), than lizards from the other two habitats (Fig. S6C, E, Table 3). 777

On Long Island the three habitats were relatively well discriminated (Fig. S7D). Importance 778
profiles indicated that short wavelengths were used to discriminate between beach scrub and 779
mangrove lizards (Fig. S7F). Beach scrub lizards had more curved reflectance profiles than in the 780
mangrove, with higher levels of UV and red reflectance relative to intermediate wavelengths (PC3, 781
Fig. S7C, E, Table 3). Coppice lizards were significantly darker than mangrove and beach scrub 782
lizards (PC1, Fig. S7C, E, Table 3). 783

On North Andros beach scrub and coppice dewlaps could be discriminated better against each 784
other than with mangrove dewlaps (Fig. S8D), with importance profiles supporting UV-reflectance 785
as a predictor of coppice lizards (Fig. S8F). Coppice lizards had less curved reflectance profiles 786
than beach scrub and mangrove lizards (PC2), and beach scrub dewlaps had the lowest scores on 787
PC4, which was difficult to interpret (Fig. S8C, E, Table 3). 788

Classification success was not significantly better than expected by chance on Ragged Island 789
and South Andros (Table 1, S8, S7) where nearly no habitat could be differentiated from any other 790
based on reflectance. 791

⁷⁹⁵ Supplementary Figures




Figure S1: Maps of the islands. (A) Map of the West Indies with sampled islands highlighted in black. (B) Sampling sites within islands colored after their respective habitat types.




Figure S2: Comparison of dewlap coloration across habitats on Abaco, with extended results. (A–E) Legend as per Figure 2. (F) One-dimensional sensitivity analysis showing the relative importance (mean decrease in accuracy) of the various wavelengths in random forest classification of the whole spectrum. (G) Geographical distance between sites where significant differences were detected in within-island principal component scores (Wilcoxon test, Benjamini-Hochberg correction, $P < 0.05$), including only pairs of sites whose habitats were involved in between-habitat dewlap differences.




Figure S3: Comparison of dewlap coloration across habitats on Bimini. Legend is as per Figure S2.




Figure S4: Comparison of dewlap coloration across habitats on Cayman Brac. Legend is as per Figure S2.




Figure S5: Comparison of dewlap coloration across habitats on Eleuthera. Legend is as per Figure S2.




Figure S6: Comparison of dewlap coloration across habitats on Little Cayman. Legend is as per Figure S2.




Figure S7: Comparison of dewlap coloration across habitats on Long Island. Legend is as per Figure S2.




Figure S8: Comparison of dewlap coloration across habitats on North Andros. Legend is as per Figure S2.




Figure S9: Comparison of dewlap coloration across habitats on Ragged Island. Legend is as per Figure S2, but without panel G.




Figure S10: Comparison of dewlap coloration across habitats on South Andros. Legend is as per Figure S2, but without panel G.




Figure S11: PC1 captures brightness across all islands. (A) Correlation between dewlap brightness (as measured by the mean reflectance from 300 to 700nm in wavelength) and PC1 score across all islands. (B) Correlation between brightness and within-island PC1, for each island. Pearson's correlation coefficients are reported. ***, $P < 0.001$.

⁷⁹⁶ Supplementary Tables

Table S1: Number of lizards sampled in each habitat on each island.

	coastal	coppice	mangrove
Abaco	41	24	21
Bimini	28	14	15
Cayman Brac	15	18	17
Eleuthera	21	25	9
Little Cayman	17	12	16
Long Island	26	13	14
North Andros	9	9	10
Ragged Island	18	15	17
South Andros	10	9	12

Table S2: Proportion of variance explained by the first four principal components on each island, as well as in the PCA performed on all islands together (last row).

Island	Total	PC1	PC2	PC3	PC4
Abaco	0.906	0.400	0.277	0.149	0.079
Bimini	0.919	0.530	0.171	0.166	0.052
Cayman Brac	0.888	0.438	0.190	0.155	0.105
Eleuthera	0.926	0.486	0.235	0.138	0.066
Little Cayman	0.907	0.441	0.212	0.176	0.078
Long Island	0.916	0.501	0.215	0.152	0.048
North Andros	0.952	0.485	0.237	0.169	0.062
Ragged Island	0.907	0.483	0.226	0.127	0.072
South Andros	0.937	0.559	0.172	0.151	0.056
All islands	0.915	0.482	0.189	0.167	0.077

Table S3: Henze-Zirkler's test of multivariate normality, performed on global principal components (i.e. fitted on data from all islands together) in each habitat and on each island. The number of outlier points detected based on the Mahalanobis distance is reported. HZ , test statistic. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$.

Island	Habitat	Outliers	HZ	P
Abaco	coastal	0	1.097	0.0028 **
Abaco	coppice	0	1.091	0.0016 **
Abaco	mangrove	0	1.052	0.0028 **
Bimini	coastal	0	0.965	0.0199 *
Bimini	coppice	0	0.828	0.0667
Bimini	mangrove	0	1.211	< 0.0001 ***
Cayman Brac	coastal	0	0.648	0.5261
Cayman Brac	coppice	0	0.691	0.4287
Cayman Brac	mangrove	0	0.671	0.4838
Eleuthera	coastal	0	1.446	< 0.0001 ***
Eleuthera	coppice	0	1.479	< 0.0001 ***
Eleuthera	mangrove	0	0.592	0.5556
Little Cayman	coastal	0	0.626	0.6547
Little Cayman	coppice	0	0.637	0.4895
Little Cayman	mangrove	0	0.867	0.0454 *
Long Island	coastal	0	0.834	0.1291
Long Island	coppice	0	1.001	0.0034 **
Long Island	mangrove	0	0.907	0.0197 *
North Andros	coastal	0	0.747	0.1123
North Andros	coppice	0	0.644	0.3623
North Andros	mangrove	0	0.839	0.0345 *
Ragged Island	coastal	0	0.757	0.2233
Ragged Island	coppice	0	0.810	0.0944
Ragged Island	mangrove	0	0.524	0.9341
South Andros	coastal	0	0.722	0.1765
South Andros	coppice	0	0.628	0.4212
South Andros	mangrove	0	0.684	0.3250

Table S4: Shapiro-Wilk's test of univariate normality of the standardized residuals from OLS and GLS-ANOVAs performed on each island where significant differences were detected by random forest classification. W , test statistic. *, $P_{adj} < 0.05$; **, $P_{adj} < 0.01$; ***, $P_{adj} < 0.001$.

Island	Variable	W	P	P_{adj}
Abaco	PC1	0.962	0.0128	0.0599
Abaco	PC2	0.960	0.0093	0.0518
Abaco	PC3	0.982	0.2963	0.6455
Abaco	PC4	0.983	0.3168	0.6455
Bimini	PC1	0.918	0.0009	0.0125 *
Bimini	PC2	0.985	0.7104	0.7956
Bimini	PC3	0.976	0.2997	0.6455
Bimini	PC4	0.977	0.3529	0.6455
Cayman Brac	PC1	0.986	0.8236	0.8869
Cayman Brac	PC2	0.989	0.9299	0.9299
Cayman Brac	PC3	0.934	0.0079	0.0518
Cayman Brac	PC4	0.981	0.5926	0.7214
Eleuthera	PC1	0.936	0.0058	0.0518
Eleuthera	PC2	0.972	0.2199	0.6455
Eleuthera	PC3	0.981	0.5224	0.6965
Eleuthera	PC4	0.977	0.3689	0.6455
Little Cayman	PC1	0.955	0.0821	0.3094
Little Cayman	PC2	0.982	0.6854	0.7956
Little Cayman	PC3	0.891	0.0005	0.0125 *
Little Cayman	PC4	0.977	0.4858	0.6965
Long Island	PC1	0.978	0.4223	0.6956
Long Island	PC2	0.980	0.5161	0.6965
Long Island	PC3	0.981	0.5605	0.7134
Long Island	PC4	0.976	0.3580	0.6455
North Andros	PC1	0.936	0.0884	0.3094
North Andros	PC2	0.966	0.4735	0.6965
North Andros	PC3	0.982	0.8977	0.9299
North Andros	PC4	0.959	0.3348	0.6455

Table S5: Locations of the sampling sites across islands, with mean within-island principal component scores per site.

Island	Longitude	Latitude	Habitat	PC1	PC2	PC3	PC4
Abaco	-77.7	26.9	mangrove	1.005	-0.116	0.063	-1.255
Abaco	-77.6	26.9	coastal	0.157	-0.531	-0.086	-2.034
Abaco	-77.6	26.9	coppice	-0.081	-0.034	-0.631	-0.633
Abaco	-77.2	26.1	coastal	0.296	0.091	0.987	-0.158
Abaco	-77.0	26.3	mangrove	-0.312	1.224	-0.898	0.208
Abaco	-77.0	26.3	coppice	0.094	-0.314	-0.120	0.482
Abaco	-77.0	26.3	coastal	-0.334	-0.748	0.201	0.534
Bimini	-79.3	25.7	coastal	0.399	0.000	0.304	-0.049
Bimini	-79.3	25.7	coppice	-0.299	0.511	-0.657	0.097
Bimini	-79.3	25.7	mangrove	-0.466	-0.477	0.046	0.000
Cayman Brac	-79.9	19.7	coastal	0.483	-0.523	-0.781	0.443
Cayman Brac	-79.8	19.7	mangrove	0.219	0.453	0.810	-0.015
Cayman Brac	-79.8	19.7	coppice	-0.610	0.008	-0.114	-0.355
Eleuthera	-76.3	24.8	coppice	0.170	-0.855	-0.042	0.086
Eleuthera	-76.3	24.8	coastal	0.503	-0.653	0.564	-0.587
Eleuthera	-76.3	24.8	mangrove	0.067	-0.694	-0.119	0.718
Eleuthera	-76.2	24.9	coppice	-0.353	0.595	-0.544	-0.535
Eleuthera	-76.1	24.9	coastal	-0.215	1.220	0.297	0.396
Little Cayman	-80.1	19.7	coppice	-0.621	0.614	-0.179	-0.404
Little Cayman	-80.0	19.7	coastal	0.395	0.355	0.743	0.175
Little Cayman	-80.0	19.7	mangrove	0.047	-0.838	-0.655	0.118
Long Island	-75.2	23.5	mangrove	-0.390	0.217	0.541	0.535
Long Island	-75.2	23.4	coastal	0.038	-0.807	0.015	-0.326
Long Island	-75.2	23.4	coppice	0.894	0.018	0.035	-0.161
Long Island	-75.1	23.4	coastal	-0.408	0.299	-0.512	-0.134
North Andros	-77.9	24.8	coastal	-0.152	-0.074	-0.104	-0.957
North Andros	-77.8	24.8	coppice	0.112	0.860	-0.258	0.621
North Andros	-77.8	24.7	mangrove	0.036	-0.707	0.326	0.303
Ragged Island	-75.7	22.2	coppice	0.269	-0.440	0.088	-0.360
Ragged Island	-75.7	22.2	coastal	-0.049	-0.076	0.221	0.029
Ragged Island	-75.7	22.2	mangrove	-0.246	0.273	-0.118	0.933
Ragged Island	-75.7	22.2	mangrove	-0.099	0.749	-0.588	-0.636
South Andros	-77.6	24.2	mangrove	0.384	0.492	-0.333	0.472
South Andros	-77.6	24.1	coppice	0.067	-0.211	0.358	-0.018
South Andros	-77.5	24.1	coastal	-0.520	-0.401	0.078	-0.550

Table S6: Nonparametric Kruskal-Wallis tests performed on each variable on each island where deviations from normality were detected. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$.

Island	Variable	χ^2	df	P	
Bimini	PC1	9.31	2	0.0095	**
Little Cayman	PC3	19.95	2	< 0.0001	***

Table S7: Support vector machine classification results. Legend is as per Table 1.

Island	<i>N</i>	Score	<i>P</i>	
Abaco	86	0.614	< 0.0001	***
Bimini	57	0.474	0.0098	**
Cayman Brac	50	0.732	< 0.0001	***
Eleuthera	55	0.476	0.0112	*
Little Cayman	45	0.724	< 0.0001	***
Long Island	53	0.623	< 0.0001	***
North Andros	28	0.721	< 0.0001	***
Ragged Island	50	0.368	0.2874	
South Andros	31	0.445	0.1152	

Table S8: Linear discriminant analysis classification results. Legend is as per Table 1.

Island	<i>N</i>	Score	<i>P</i>
Abaco	86	0.670	< 0.0001 ***
Bimini	57	0.491	0.0046 **
Cayman Brac	50	0.736	< 0.0001 ***
Eleuthera	55	0.480	0.0112 *
Little Cayman	45	0.747	< 0.0001 ***
Long Island	53	0.562	0.0004 ***
North Andros	28	0.636	0.0008 ***
Ragged Island	50	0.388	0.1964
South Andros	31	0.458	0.0589