

Dewlap color variation in *Anolis sagrei* is maintained between habitats within islands of the West Indies

RAPHAËL SCHERRER^{1,3} *, COLIN M. DONIHUE^{1,4},
R. GRAHAM REYNOLDS², JONATHAN B. LOSOS^{1,4} AND ANTHONY J. GENEVA^{1,5}

¹ Department of Organismic and Evolutionary Biology and Museum of Comparative Zoology
Harvard University, Cambridge, MA, USA

² Department of Biology, University of North Carolina Asheville, Asheville, NC, USA

³ Current address: Groningen Institute for Evolutionary Life Sciences,
Groningen, The Netherlands

⁴ Current address: Department of Biology, Washington University, St. Louis, MO, USA

⁵ Current address: Department of Biology, Center for Computational and Integrative Biology,
Rutgers University–Camden, Camden, NJ, USA

Abstract

Animal signals evolve in an ecological context. Moreover, locally adapting animal sexual signals can be especially important for initiating or reinforcing reproductive isolation during the early stages of speciation. Dewlap color in *Anolis* lizards can be highly variable between populations in relation to both biotic and abiotic adaptive drivers, albeit at relatively large geographical scales. Here, we investigated local adaptation of the dewlap across habitat-types at a small spatial scale, as this may give an indication of how conditions for the early stages of speciation may be met. We explored variation in dewlap coloration in the most widespread species of anole, *Anolis sagrei*, across three characteristic habitats spanning the Bahamas and the Cayman Islands. Using reflectance spectrometry as well as supervised machine learning, we found significant differences in spectral properties of the dewlap between habitats within small islands. Passive divergence in dewlap phenotype associated with isolation-by-distance did not explain our results. Instead, the observed patterns in dewlap coloration are more consistent with an adaptive explanation in these *A. sagrei* populations, as one would otherwise expect differences within islands to be erased by gene flow at such small geographical scales. These habitat-specific dewlap differences varied in magnitude and direction across islands, and may suggest a role of sexual selection in driving these local patterns. While at present, populations from these different habitats probably experience too much gene flow to follow distinct evolutionary lineages, should additional barriers arise between habitat-specific populations, the observed disruptive selection on dewlap coloration may facilitate ecological speciation.

Keywords — *Anolis*, reflectance, local adaptation, sexual signal, supervised machine learning

Introduction

The staggering diversity of animal communication signals has long been of interest to evolutionary biologists. Animals use chemical, mechanical, electromagnetic, and visual signals to communicate in a wide variety of contexts, including, for example, competition for mates, species recognition, aposematism, and cooperation (Bradbury and Vehrencamp, 2011). A primary evolutionary factor shaping communication signals is the sensory system and behavior of their recipient(s) (the sensory drive hypothesis; Endler and McLellan 1988; Endler 1992, 1998). Over the past decades, scientists have established that signals evolve in an ecological context and are dependent on environmental conditions (Endler, 1992, 1993a,b). Just as different habitats may favor different combinations of eco-morphological traits to maximize performance and fitness (Arnold, 1983), they may also shape different forms of a signal, so as to maximize its transmission and detection (e.g. Seehausen 1997), or reduce its detection by unintended recipients such as predators (Endler, 1984, 1990, 1991;

*Corresponding author: r.scherrer@rug.nl

37 Halfwerk et al., 2014). This selective pressure may drive the local adaptation of communication
38 signals.

39 One potential barrier to the maintenance of localized signal divergence is the homogenizing
40 effect of gene flow. Population genetics theory suggests that gene flow may counteract local
41 adaptation between localities and prevent divergence altogether, especially at small spatial scales,
42 because of the inflow of maladapted alleles or because of the breaking of linkage between coevolving
43 loci (Felsenstein, 1976; García-Ramos and Kirkpatrick, 1997; Dieckmann and Doebeli, 1999;
44 Lenormand, 2002; Hendry et al., 2007a). This has been confirmed empirically in systems such as
45 stick-insects (Nosil and Crespi, 2004) and sticklebacks (Hendry et al., 2007b). Yet, examples of
46 microgeographic adaptation, i.e. adaptation at smaller scales than the range of dispersal, exist,
47 highlighting a high potential of some organisms to respond to selection in the face of gene flow
48 (see Richardson et al. 2014 and references therein). Examples include small scale adaptation in
49 fragmented areas in Australian fruit flies (Willi and Hoffmann, 2012), or local adaptation to pre-
50 dation pressure in North American salamanders (Richardson and Urban, 2013). Therefore, despite
51 evidence that local adaptation may be particularly difficult at small spatial scales where gene flow
52 tends to cause adjoining populations to remain genetically homogeneous, the potential adaptive
53 response of species traits, in particular communication signals, to localized differences in habitats
54 remains relatively unknown (Richardson et al., 2014).

55

56 Lizards of the neotropical genus *Anolis* are an excellent group for studying the eco-evolutionary
57 dynamics of local adaptation and natural selection (Losos, 2009). A particularly conspicuous trait
58 of anoles is their dewlap; an extensible flap of skin that is typically sexually dimorphic and used
59 as a communication signal in courtship (Sigmund, 1983; Driessens et al., 2014, 2015), territorial
60 displays (Losos, 1985; Macedonia and Stamps, 1994; Macedonia et al., 2013) as well as in predator
61 deterrence (Leal and Rodríguez-Robles, 1995, 1997; Leal and Rodriguez-Robles, 1997). Dewlap
62 characteristics vary widely among the approximately 400 species of the genus (Nicholson et al.,
63 2007). Interspecific variation in dewlap coloration is implicated in species recognition (Rand and
64 Williams, 1970; Williams, 1969; Williams and Rand, 1977; Losos, 1985; Macedonia and Stamps,
65 1994; Fleishman, 2000; Macedonia et al., 2013), and this function could have had a role in initiating
66 and/or reinforcing reproductive isolation during speciation (Lambert et al., 2013; Geneva et al.,
67 2015; Ng et al., 2017).

68

69 Within species, studies have shown a link between variation in dewlap coloration and differences
70 in habitats or climatic conditions (Macedonia, 2001; Leal and Fleishman, 2002; Thorpe and
71 Stenson, 2002; Thorpe, 2002; Leal and Fleishman, 2004; Vanhooydonck et al., 2009; Ng et al.,
72 2012, 2013, 2016; Vanhooydonck et al., 2009; Driessens et al., 2017). Some studies suggest that
73 those differences may be adaptive, and that dewlaps may have evolved to maximize detectability
74 given local light conditions (Fleishman and Persons, 2001; Leal and Fleishman, 2002, 2004). Other
75 studies testing this hypothesis, however, found no pattern (Fleishman et al., 2009; Ng et al., 2012;
76 Macedonia et al., 2014).

77

78 Previous studies investigating variation in anole dewlaps compared populations at relatively
79 large geographical scales, e.g. between islands (Vanhooydonck et al., 2009; Driessens et al., 2017)
80 or within large islands such as Puerto Rico (Leal and Fleishman, 2002, 2004) or Hispaniola (Ng
81 et al., 2012, 2016). These large scales and marine barriers should reduce gene flow (Ng and Glor,
82 2011; Lambert et al., 2013; Richardson et al., 2014; Ng et al., 2017). That said, examples do exist
83 of divergence in dewlap coloration at smaller scales or between populations with high degrees of
84 gene flow (Thorpe and Stenson, 2002; Thorpe, 2002; Stapley et al., 2011; Ng et al., 2016).

85

86 The species *Anolis sagrei* is widespread across islands of the West Indies (Reynolds et al.,
87 2020). It has been the subject of study concerning local adaptation (Losos et al., 1994, 1997, 2001;
88 Kolbe et al., 2012), biological invasion (Kolbe et al., 2008), sexual selection (Tokarz, 2002; Tokarz
89 et al., 2005; Tokarz, 2006; Driessens et al., 2014; Steffen and Guyer, 2014; Driessens et al., 2015)
90 and many other topics. Between-island variation in the mainly orange-red color of its dewlap was
91 shown to be better explained by climatic variables (Driessens et al., 2017) than by proxies for biotic
92 factors such as sexual selection or predation pressure (Vanhooydonck et al., 2009; Baeckens et al.,
93 2018). How intra-island differences in habitat may contribute to the diversity of dewlap coloration,

however, remains unexplored, and may reveal new insights into the scale of local differentiation
95
despite gene flow.
96
97

Here, we analyzed the color characteristics of *A. sagrei* dewlaps within nine islands in the Bahamas and Cayman Islands. These island systems presently, if not historically, comprise relatively small islands, with no major geographic barriers within islands limiting dispersal for this promiscuous species (Kamath and Losos, 2018). These islands all share three characteristic native West Indian habitat-types – beach scrub bush, closed-canopy primary coppice forest, and mangrove forest – that are often spatially intermingled. These habitats contrast in environmental parameters including vegetation community, light irradiance, humidity and temperature (Howard, 1950; Schoener, 1968). The Cayman Islands and the Bahamas have been colonized independently by *A. sagrei* from Cuba (Reynolds et al. 2020, van de Schoot et al. unpubl.), such that these archipelagos constitute an ideal suite of natural replicates to explore within-island dewlap diversity across multiple islands.

Our sampling design included sites in close proximity (the median distance between two sites within an island was 11.2km). Combining reflectance spectrometry and supervised machine learning, we tested for divergence in dewlap phenotype between habitats within islands and between islands across the range of *A. sagrei*. We predicted that if light conditions in the environment indeed drive color evolution, dewlaps should be most similar between beach scrub and mangrove forest, which both have high levels of light irradiance, compared to the darker, closed-canopy coppice forest. Similar, if detectability is maximized given the local conditions, we expected darker and more contrasting dewlaps in high irradiance habitats. Finally, if habitat characteristics are strong determinants of dewlap color variation, similar patterns should be observed across multiple islands (Losos, 2011).

Methods

Data collection

We sampled 466 male *Anolis sagrei* from seven islands in the Bahamas Archipelago – Abaco, North Andros, South Andros, South Bimini, Eleuthera, Long Island, Ragged Island – and two in the Cayman Islands – Cayman Brac and Little Cayman (Figure 1). These islands were chosen to span the breadth of the West Indian range of *Anolis sagrei*, because they have highly similar habitat types, and because the *A. sagrei* on each island group are derived from ancient and distinct colonization events from Cuba (i.e. relatively evolutionarily independent, Reynolds et al. 2020). Three habitats were sampled on each island based on characterizations by Howard (1950) and Schoener (1968). Each habitat is clearly distinguishable by their dominant vegetation type — xeric coastal scrub (open, relatively dry habitat consisting of low vegetation or isolated trees), primary coppice forest (closed-canopy forest) and mangrove forest (wet coastal habitat with trees growing in brackish water and high light penetration). Sample sizes are given in Table S1. Our sampling design enabled us to test for differences between habitats at a coarse and fine geographical scale. The median distance between two localities within an island was ~ 11km, with some islands being sampled at smaller or larger scales (Figure S1, Table S2). 80.3% of all pairwise distances within islands were less than 50km. Additionally, there are no major barriers to dispersal (such as mountains or grassland) on any of the islands that we sampled.

Reflectance measurements

We measured reflectance between 300 and 700nm wavelength, a range that encompasses the colors visible to most lizards and vertebrates in general (Lazareva et al., 2012). Measurements were taken with an Ocean Optics USB4000 spectrometer, a pulsed Xenon light source (PX-2, Ocean Optics, Largo, FL, USA) and a reflectance probe protected by a black anodized aluminum sheath. Measurements were taken with a 45-degree inclination to prevent specular reflection (Endler, 1990). The device was regularly standardized with a Spectralon white standard (Labsphere, North Sutton, NH, USA). Reflectance was measured at the center of the dewlap.

Analysis

146

All analyses in this study were performed in R 3.6.1 (R Core Team, 2019).

147

148 Dimensionality reduction

149 Reflectance curves were smoothed using the R package *pavo* (Maia et al., 2013) as well as with
150 custom R functions, down to one reflectance value at each nanometer in wavelength from 300
151 to 700nm. Because neighboring wavelengths are highly collinear in reflectance, we reduced the
152 dimensionality of the data using principal component analysis (PCA), as per Cuthill et al. (1999)
153 and Leal and Fleishman (2002). We performed PCA on each island separately and systematically
154 retained the first four principal components (PC), which together always explained more than
155 88.8% of the variance across islands (Table S3). PC1 explained between 40 and 56% of the variance
156 across islands; PC2 explained 17.4–27.9%; PC3 12.7–17.6% and PC4 4.3–10.5%. The first four PCs
157 explained similar proportions of variance when calculated for all islands together (Table S3). PCs
158 need not represent the same wavelengths across islands because they are fitted on different datasets.
159 Nevertheless, PC1 was very collinear with brightness for all islands (Figure S2, Table S4). PC2
160 correlated highly with the red and ultraviolet ends of the spectrum, which were inversely correlated
161 with each other (Fig. 3A). Higher PCs corresponded to various combinations of wavelengths.
162 Because PC1 correlated uniformly with all wavelengths across the spectrum we considered PC2
163 onwards to capture the chromatic dimensions of color space, i.e. the relative contributions of the
164 wavelengths regardless of brightness.

165 Pooled analyses

166 In addition to within-island PCA, we performed a PCA on pooled data from the whole archipelago.
167 The first four principal components explained 91.3% of the variance (Table S3). Again PC1 strongly
168 correlated with brightness (Fig. S3, Table S4). PC2 was positively correlated to short wavelengths
169 (ultraviolet to blue) and negatively correlated to long wavelengths (green to red, Fig. S4B). PC3
170 was strongly negatively correlated with UV reflectance and positively correlated with blue-green.
171 PC4 was made of a mosaic of wavelengths, correlating positively with blue and red but negatively
172 with ultraviolet and yellow.

173

174 We used this dataset to partition the variance in dewlap coloration among islands, habitats
175 and habitats within islands, using a two-way multivariate analysis of variance (MANOVA) with
176 an interaction term. However, because the assumptions of parametric MANOVA were violated
177 for all islands but Ragged Island (multivariate normality, Henze-Zirkler's test, Henze and Zirkler
178 1990, R package MVN, Korkmaz et al. 2014, Table S5; and homogeneity of covariance matrices,
179 Box's M-test, Box 1949; Morrison 1988, R package heplots, Fox et al. 2018, Table S6), we used
180 a semi-parametric MANOVA instead (R package MANOVA.RM, Friedrich et al. 2018), with P-
181 values calculated from a bootstrap procedure with 1,000 iterations. We calculated the proportion
182 of variance explained by islands, habitats and the habitat-by-island interaction using partial effect
183 sizes η^2 on a MANOVA-approximation of the analysis (R package heplots, Fox et al. 2018).

184 Machine learning

185 Because of the aforementioned violations of the MANOVA assumptions, and to reduce the chances
186 of false discovery, we conducted multivariate group comparisons using support vector machines
187 (SVMs), a model-free, powerful nonparametric supervised machine learning technique.

188

189 Machine learning for group comparison has become more common in ecology and evolution
190 in recent years (e.g. Pigot et al. 2020). In particular, SVMs are designed to find the best pos-
191 sible nonlinear boundaries between labelled groups of points in multidimensional spaces, without
192 assumptions about the distribution of the data (Cortes and Vapnik, 1995; Cristianini and Shawe-
193 Taylor, 2000; Kim and von Oertzen, 2018). This makes them well suited to field biological data,
194 which often violate the assumptions of classical linear modeling (Kim and von Oertzen, 2018) and
195 can be, as in the case of coloration, inherently highly multivariate (Cuthill et al., 1999). First,
196 a machine is trained to recognize differences between groups within a subset of the data called
197 the training set. Significance of differences is then assessed by testing the accuracy of that fitted
198 machine in predicting the group-labels of data points that were not included in the training, called

a testing set, based solely on their multivariate coordinates. This cross-validation procedure results in a proportion of correctly classified points, or generalization accuracy score, which can be compared to that expected under random guessing using a binomial test.

In this study, we performed SVM classifications on each island separately. We used a standard five-fold cross-validation procedure, where the data were randomly split into five bins of approximately equal sizes. Each bin was in turn taken as the testing set while the rest was used as a training set, thus resulting in five trained machines per cross-validation. We replicated this procedure 100 times for each island to account for stochastic outcomes. We performed binomial tests to evaluate the significance of deviations in observed mean generalization accuracy per island to null expectations under random guessing. Each training data set was downsampled to the size of its least represented habitat to ensure balanced training samples. We ensured that each habitat was represented by at least five data points in the training set.

All classification analyses were repeated using the more classical linear discriminant analysis (LDA), a supervised machine learning technique finding linear boundaries that maximize the differences between groups, albeit assuming multivariate normality and homogeneity of covariance matrices (Ripley, 1996). We used the R package rminer (Cortez, 2010, 2016) for SVMs, and MASS (Venables and Ripley, 2002) for LDAs. We used rminer's default heuristic search option to automatically tune the Gaussian kernel parameter σ and the complexity parameter C for the SVMs.

The same procedure was repeated on principal components from the whole archipelago (see Pooled analyses) to evaluate the significance of archipelago-wide differences in dewlap coloration across habitats.

All machine learning classifications performed on principal components were also repeated on the original reflectance datasets reduced to 50-nm spaced wavelengths from 300 to 700nm.

We conducted one-dimensional sensitivity analyses using rminer (Cortez and Embrechts, 2013) to determine the relative importance of the different input variables during classification where significant differences were detected, both on machines trained on principal components and machines trained on non-transformed reflectance at various wavelengths. In parallel, we conducted univariate analyses of variance to independently test the importance of different variables in between-habitat variation, on islands where the machines detected significant differences based on binomial tests (next section).

Univariate analyses

For each island where significant differences in multivariate dewlap coloration were detected between habitats, we used multiple univariate analyses of variance (ANOVA) to identify which variables were responsible for the observed differences. We constructed our ANOVA models in two steps, as per Zuur (2009). In a first step, we accounted for heterogeneity of variances across groups by systematically comparing the goodness-of-fit of an ANOVA model estimated with ordinary least squares (OLS) with that of a model estimated with generalized least squares (GLS), which allowed one estimate of residual variance per habitat (using the R package nlme, Pinheiro and Bates 2000; Pinheiro et al. 2020). Both models were fitted with restricted maximum likelihood (REML). Goodness-of-fit was estimated using Akaike's Information Criterion corrected for small sample sizes (AICc, R package MuMIn, Bartoń 2019), and the estimation method yielding the lowest AICc was retained. In a second step, we re-fitted the retained model with maximum likelihood (ML) to test for the effect of habitat-type using likelihood ratio tests (LRT) between a model including a habitat-term and a null model lacking the habitat-term.

We tested the assumptions of the parametric ANOVA for each island included in the univariate analyses. For all islands where deviations from multivariate normality were detected in at least one habitat (Table S5), we assessed univariate normality for each principal component (Shapiro-Wilk's test, Table S7). For skewed PCs that deviated significantly from normality, we repeated the analysis using a nonparametric Kruskal-Wallis test (Hollander et al., 2013). We found no multivariate outliers based on the Mahalanobis distance (package MVN, Korkmaz et al. 2014). We used the cases of better fit of the GLS model relative to the OLS model as evidence for heterogeneity of

256 variances, which were then accounted for by the GLS approach (Table 1).

257
258 Significant *post hoc* contrasts were assessed using Tukey's Honest Significant Difference (HSD)
259 test whenever the assumptions of normality and homogeneity of variances was met (Tukey, 1949),
260 Dunnett's T3 method when only homogeneity of variances was violated but not normality (Dun-
261 nett, 1980), and Nemenyi's test when normality was violated (Nemenyi, 1963). All *post hoc* tests
262 were performed with the R package PMCMRplus (Pohlert, 2020).

263
264 We used the same procedure to investigate which variables, if any, were involved in archipelago-
265 wide multivariate differences between habitats detected in our two-way MANOVA design (see
266 Pooled analyses). However, in the first step or our model comparison procedure, we added mixed-
267 effect equivalents of our OLS and GLS models, this time with island as a random effect. The
268 resulting four models were compared and the best fitting variance structure was retained as ex-
269 plained above.

270 Spatial autocorrelation

271 We tested for within-island spatial autocorrelation between the geographical distances among sam-
272 pling sites and their Euclidean distances in multivariate color space (mean PC1 to PC4 per site,
273 Table S2), regardless of habitat-type. Because often only a few sites were sampled per island, we
274 could not get meaningful results from tests that use sites as units of observation, such as Moran's
275 I test (Gittleman and Kot, 1990). Instead, we designed a permutation test where we randomly
276 reshuffled individual lizards across sites within islands 1,000 times each, and systematically re-
277 calculated Pearson's correlation coefficient between geographic distances (computed as geodesic
278 distances in the R package geosphere; Hijmans 2019) and phenotypic distances. We used the re-
279 sulting null distributions of correlation coefficients to assess the significance of the observed spatial
280 autocorrelation for each island.

281 Site differences

282 In this study, we were interested in the minimum spatial scale at which significant differences
283 between habitats could be detected within islands. We performed multiple pairwise nonparametric
284 Wilcoxon-Mann-Whitney tests (Hollander et al., 2013) to compare dewlap coloration between
285 sites with different habitat-types, for each pair of habitats and each variable where significant
286 differences were detected with our analyses of variances. The P-values were adjusted using a
287 Benjamini-Hochberg correction for multiple testing (Benjamini and Hochberg, 1995).

288 Results

289 We tested for variation in *A. sagrei* dewlap coloration between populations living in three charac-
290 teristic habitat types across nine islands that span the West Indian range of the brown anole (Fig
291 1, S1). We found that most of the variation in coloration is partitioned between islands (two-way
292 semi-parametric MANOVA, modified ANOVA-type statistic (MATS) = 2009.6, $P < 0.001$, Fig.
293 S5, explained variance $\eta^2 = 44.3\%$, MANOVA approximation). Nonetheless, we did find evidence
294 for differences in dewlap coloration between habitat-types, and those were mostly island-specific
295 (habitat-by-island interaction term, MATS = 384.4, $P < 0.001$, explained variance $\eta^2 = 11.4\%$),
296 with a small but significant portion of the variation explained by an archipelago-wide habitat effect
297 (MATS = 42.5, $P = 0.001$, $\eta^2 = 4.8\%$).

298
299 The small archipelago-wide effect of habitat-type was detected for PC1, PC2 and PC3 (mixed-
300 effect ANOVA with island as a random effect, Table S8), but this effect was too small for *post hoc*
301 tests to find which habitats differed. Archipelago-wide differences in dewlap coloration between
302 habitats were also detected by SVMs trained on pooled data regardless of island identity, both for
303 PCA data and reflectance scores (Fig. S6, S7). This pattern seemed to be driven by mangrove
304 lizards being correctly reassigned more often than predicted by chance. Sensitivity analyses on
305 these machines suggest a relatively small role of long wavelengths (red reflectance) in driving this
306 pattern (Fig. S9), but did not reveal strong differences between the PCs in relative importance

307 (Fig. S8). Archipelago-wide differences were not detected by LDA classifiers at all (Fig. S10, S11).

Within islands, SVM classifiers correctly assigned individuals to their habitat of origin based solely upon dewlap coloration on five islands: Abaco, Bimini, Cayman Brac, Little Cayman, and Long Island (Fig. 2). An LDA approach yielded similar success rates (Fig. S12), suggesting robust differences between these populations. Of the five islands, Little Cayman was the best discriminated with a mean SVM generalization success of 73.4% (Table S9). The results of the classification analyses on PCA data were very similar to results from SVMs and LDAs trained on reflectance values at 50nm-spaced wavelengths from 300 to 700nm (Fig. S13 and S14).

Differentiation in dewlap coloration occurred in multiple dimensions of color space. Moreover, the differences in dewlaps between habitats were not always consistent among islands, thus, we will discuss the habitat-specific variation in dewlap coloration for each island where significant differences were detected in turn (Fig. 3, Tables 1, S10). Figure 3A provides a key to map principal component scores to the underlying wavelengths.

On Abaco, dewlaps did not differ in PC1, which represents brightness. Mangrove lizards had significantly lower PC2 scores, corresponding to higher ultraviolet reflectance and lower red reflectance. Coastal beach scrub lizards had lower scores on PC3, corresponding to lower ultraviolet reflectance and higher blue reflectance.

On Bimini, coastal beach scrub lizards had significantly brighter dewlaps than lizards from mangroves (PC1), but mangrove lizards had higher PC2 scores than beach scrub lizards, indicating higher violet and blue reflectance, and lower red reflectance. Lizards from primary coppice had higher PC3 scores overall, which correlated very positively with ultraviolet reflectance.

On Cayman Brac, coppice-lizard dewlaps were significantly less bright than lizards from the other habitats. Coastal beach scrub lizards had dewlaps that scored low on PC2, corresponding to lower violet-blue and more red, while the mangrove lizards exhibited the opposite: relatively higher levels of violet-blue and less red. In PC3 space we found that dewlaps from lizards in the coastal habitat had high ultraviolet reflectance, coppice lizards had intermediate levels, and mangrove lizards had relatively low levels.

On Little Cayman, the dewlaps of coppice lizards were significantly darker (PC1) than coastal-lizards. Mangrove lizards had less ultraviolet and redder dewlaps (PC2). The dewlaps of the coastal beach scrub lizards had higher levels of red and ultraviolet reflectance and less blue reflectance than the dewlaps of the other habitat-populations (PC3).

On Long Island, lizards from the coppice habitat had darker dewlaps than lizards from the other habitats (PC1). Coastal lizards had relatively more ultraviolet and less blue-green reflectance in their dewlaps (PC3). These coastal-habitat lizards also scored lower on PC4, corresponding to slightly more violet and green-yellow dewlaps, and less blue dewlaps, than the mangrove lizards on the island.

Sensitivity analyses on classifiers suggested an overall higher relative importance for PC2 and PC3 in determining between-group differences on Abaco, both in SVM and LDA classifiers (Fig. S15, S16), consistent with our ANOVA results (Fig. 3B). There was no strong signal of differences in relative importance among principal components on the other islands. Sensitivity analyses of SVMs trained on reflectance scores rather than principal components revealed, however, a consistently higher importance of ultraviolet reflectance in between-group differences on all islands (Fig. S17). This pattern was not recovered for LDAs trained on reflectance scores (Fig. S18).

We did not find significant spatial autocorrelation between the sampling sites on the islands where we detected a significant habitat effect. We did, however, detect a significant positive signal of autocorrelation on Eleuthera (Table S11), suggesting possible color differentiation through isolation-by-distance on this island.

In contrast, differences in dewlap coloration between habitats were often detected in close ge-

ographical proximity. For example, Bimini, Cayman Brac, and Little Cayman were among the
366 smallest islands in our study (Fig. S1). Indeed, most significant differences in dewlap coloration
367 involved sites that were 5-10km apart. Our most extreme case of significant differences occurred
368 for PC3 between a beach scrub site and a coppice site, separated from each other by a few hundreds
369 of meters at most on Bimini (multiple pairwise Wilcoxon-Mann-Whitney tests, Fig. S19).

370 Patterns of differentiation were inconsistent across the five most significant islands. Contrasts
371 in principal components between habitats, calculated on pooled data from the whole archipelago,
372 were not similar, for any component, among islands (Fig. S20; MANOVA, Pillai's trace = 0.354,
373 $F(6, 32) = 1$, P = 0.36). No pattern of variation was shared by all five significant islands, along
374 any dimension. Some patterns did seem more common however, such as darker dewlaps among
375 coppice lizards (Cayman Brac, Little Cayman, and Long Island, Fig. 3) or the intermediate posi-
376 tion of coppice lizards in chromatic color space (Cayman Brac and Long Island). In other cases,
377 patterns of differentiation were reversed from one island to another, with more ultraviolet reflecting
378 dewlaps in mangroves than in coastal habitat on Abaco and Cayman Brac, but the opposite on
379 Little Cayman and Long Island. Overall, it seemed that patterns of heterogeneity of variance were
380 often driven by higher variances in coloration within beach scrub lizards (Fig. 3, Table 1). Yet
381 other patterns were idiosyncratic, such as the combination of higher red and ultraviolet reflectance
382 in coastal lizards on Little Cayman, where the rule seemed to be a negative correlation between
383 ultraviolet and red reflectance across every other island.

385

386 Discussion

387 **Dewlap coloration differs between habitat-types** We found that male dewlap coloration in
388 *A. sagrei* significantly varied between fine-scale habitat-types (beach scrub bush, primary coppice
389 forest and mangrove forest) on five islands of the West Indies: Abaco, Bimini, Cayman Brac, Little
390 Cayman and Long Island. However, the habitat-specific variation in dewlaps was not consistent
391 between these islands. Although those results are consistent with adaptation at a very local scale,
392 other evolutionary drivers could be at work, such as phenotypic plasticity, random drift, or multiple
393 colonization events. We reject this last explanation because all of the island populations in this
394 study are strictly monophyletic, implying a single colonization event per island (van de Schoot,
395 unpublished thesis; Driessens et al. 2017; Reynolds et al. 2020).

396 **A role of neutral drift is unlikely** Differences in organismal traits between environments
397 are not necessarily proof of adaptation or selection, and genetic drift may result in patterns sim-
398 ilar to local adaptation (Miles et al., 2019). Nevertheless, two lines of evidence from our data
399 suggest that this scenario may be implausible. First, we found little evidence for a role of phe-
400 notypic isolation-by-distance (spatial autocorrelation) in explaining the differences we report. We
401 did detect a significant signal of isolation-by-distance on Eleuthera, but there were no differences
402 in dewlap coloration between habitats on this island. Second, we detected differences between
403 habitats at relatively small spatial scales, most of the time between sites 5-10km apart, sometimes
404 a few hundred meters away, on Bimini for example. Such small-scale differences would be unlikely
405 under strong gene flow (Richardson et al., 2014). Our study islands lack geographic barriers to the
406 movement of *A. sagrei*, which have been shown to be highly mobile (Kamath and Losos, 2018),
407 implying widespread gene flow across sites and habitats. Moreover, habitat-populations within
408 each island were found to be non-monophyletic and often share identical haplotypes, based on
409 phylogenetic analysis of mitochondrial DNA sequences (van de Schoot et al. unpublished thesis),
410 suggesting gene flow between habitats may be widespread.

411

412 Our results align with previous documented cases of persistent dewlap color divergence despite
413 gene flow in multiple species of anoles, sometimes in relation to environmental conditions. Ng et al.
414 (2012) and Ng et al. (2016) found divergent dewlap coloration in the face of gene flow between
415 subspecies of *A. distichus* across Hispaniola, and proposed this as a mechanism of reproductive
416 isolation in the early stages of speciation (Ng and Glor, 2011; Lambert et al., 2013; Ng et al.,
417 2017). Stapley et al. (2011) found that dewlap color polymorphism was maintained in the absence
418 of genetic structure between populations of *A. apletophallus* from Panama. Thorpe and Stenson
419 (2002) found that divergence in dewlap coloration matched habitat-type better than mitochondrial

420 lineage in *A. roquet* on Martinique, and a convergent pattern was found in *A. trinitatis* on the
featureless island of St Vincent (Thorpe, 2002). Finally, regionally-distinct body coloration, but
not dewlap coloration, is present in *A. conspersus* on another small island, Grand Cayman, where
no physical barriers to gene flow exist (Macedonia, 2001).

421
422
423
424

Dewlap coloration could be locally adapted Although phenotypic divergence at small spatial scales may be an indicator of local adaptation, the inconsistency of the between-habitat divergence patterns we observed across islands implies that neutral processes cannot be definitely ruled out (Losos, 2011). Conversely, the absence of parallel divergence across islands does not rule out local adaptation either (Losos, 2011). If selection has an effect, it may be dependent on aspects of the environment that are not encompassed by our broad habitat-type classification into coastal dry scrub, primary coppice forest and mangrove forest, and dewlap coloration may be influenced by a mosaic of interacting local environmental factors, which need not be the same across islands.

425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455

Previous studies have described convergent patterns of dewlap color evolution in similar environments across islands and species (Thorpe and Stenson, 2002; Thorpe, 2002). Others have suggested that dewlap coloration may have evolved to be maximally detectable under local light conditions imposed by the environment, primarily through UV contrast (i.e. UV-brighter dewlaps in UV-dark, mesic habitats and UV-darker dewlaps in UV-bright, xeric habitats), in *A. cristatellus* and *A. cooki* from Puerto Rico (Leal and Fleishman, 2002, 2004). Although UV reflectance was commonly involved in between-habitat divergence in *A. cooki* and *A. cristatellus*, we found no such patterns in *A. sagrei*, where instead, we found the darkest dewlaps in the dark, mesic habitat – primary coppice forest – on three islands, and dewlaps often differed the most between beach scrub and mangrove forest, two xeric habitats with similar, high irradiance levels (Howard, 1950; Schoener, 1968). The inconsistent and idiosyncratic patterns we observed suggest that dewlap color variation between habitats cannot be predicted by habitat identity alone. Studies of Jamaican and Hispaniolan anoles similarly found between-habitat differences in dewlap coloration but no evidence for higher dewlap detectability in different habitats (Fleishman et al., 2009; Ng et al., 2012). Habitats on different islands may also differ in aspects other than light conditions, such as densities of predators or congeners, which have been shown to affect among-island dewlap diversity (Vanhooydonck et al., 2009; Baeckens et al., 2018). In particular, Baeckens et al. (2018) recently showed that dewlaps with spotted patterns occurred more often in *A. sagrei* on islands with more coexisting species of anoles. Our data are consistent with those previous results in suggesting that adaptation to local light conditions, or at least broad habitat types, may not be a major driver of the within-island variation in dewlap coloration in *A. sagrei*.

Sexual selection could be at play Sexual selection is an important component of an adaptive, sensory drive hypothesis of dewlap color divergence across environmental conditions, because in this scenario habitat-dependent selection is executed by the signal recipients (Endler and McLellan, 1988), such as mating partners (Driessens et al., 2014). An alternative yet non-mutually exclusive explanation for the observed idiosyncratic patterns of within-island divergence is that of "Fisherian" sexual selection (Andersson, 1994). Under the Fisherian model of sexual selection, arbitrary female preferences (i.e. independent of the environment) for certain male ornaments may drive divergent evolution of male signals, such as dewlap coloration, if female preferences differ between localities for other reasons than environment-dependent perception abilities (a situation that could have arisen e.g. in an initial phase of genetic drift). Substantial levels of promiscuity in *A. sagrei* suggest ample opportunity for female mate choice (Kamath and Losos, 2018), and are in line with this explanation. Therefore, arbitrary, habitat-independent female preferences could further explain the inconsistent patterns of divergence across islands. In contrast with this, however, Baeckens et al. (2018) found no link between *A. sagrei* dewlap coloration and size dimorphism (a proxy for sexual selection) in an among-island study of the same archipelagos.

456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475

Besides, although this process is well-established in some systems such as African cichlids (e.g. Seehausen 1997), previous work suggests that dewlap coloration in *Anolis* may act as a mating cue according to a "good genes" model of sexual selection (Andersson, 1994), rather than a Fisherian process. In the good genes model, the cue under sexual selection is an indicator of individual qual-

ity (e.g. better immune response to disease) and of indirect benefit to the offspring. Several studies suggest that this possibility could be the case for anoles. For example, Cook et al. (2013) found lower orange reflectance in dewlaps with heavily parasitized *A. brevirostris*, suggesting a trade-off in carotenoid use between the immune response and pigment deposition. Steffen and Guyer (2014) found that lower UV and orange-red reflectance predict contest-winning success between *A. sagrei* males, while Driessens et al. (2015) further found that more yellow and red dewlaps (relative to UV) predict better body condition, and that higher yellow and UV reflectance at the margin of the dewlap predict higher hematocrit (the concentration of red blood cells), indicating a better health. Other aspects of the dewlap than color have also been found to be indicators of individual quality, such as dewlap size (Vanhoydonck et al., 2005, 2009), but not dewlap display frequency (Tokarz, 2002; Tokarz et al., 2005; Driessens et al., 2015).

487

488 A role of phenotypic plasticity is unlikely Differences in coloration between habitat populations may not be genetically determined and instead may be influenced by environmental factors such as parasite load, as mentioned above (Cook et al., 2013). The yellow, orange and red coloration in anoline dewlaps are produced by pterins and carotenoids (Ortiz, 1962; Ortiz et al., 1962; Ortiz and Williams-Ashman, 1963; Ortiz and Maldonado, 1966; Macedonia et al., 2000; Steffen and McGraw, 2007, 2009). Animals lack the ability to synthesize carotenoids, and those must therefore be found in the diet, while pterins are synthesized from nucleotides (Goodwin, 1984; Hill et al., 2002; Hill and McGraw, 2006). However, experimental manipulation of dietary carotenoid content showed no effect on dewlap coloration in *A. sagrei* (Steffen et al., 2010) nor in *A. distichus* (Ng et al., 2013), which also has an orange-based dewlap. This makes a plastic response to differences in diet across habitats unlikely. Furthermore, developmental plasticity during the ontogeny is also unlikely because dewlap coloration develops at sexual maturity in anoles (Ng et al., 2013). The differences we observed could therefore be genetically based. This hypothesis is further supported by Cox et al. (2017), who found a high degree of heritability of dewlap coloration in *A. sagrei*. Although most studies used one or two-generation common garden experiments and thus could not rule out transgenerational plastic effects (Tariel et al., 2020), dewlap coloration generally seems to not be a plastic trait. This further reinforces an adaptive explanation, where dewlap color could be under differential natural and/or sexual selection in these different habitats.

506

507 Implications in the context of speciation Local adaptation can be a precursor to ecological 508 speciation, a process that may have given rise to the adaptive radiation of *Anolis* lizards (Harmon et al., 2003; Gavrilets and Losos, 2009). Ecologically-mediated divergence of a sexual signal may 510 be a potent path to the evolution of reproductive isolation through divergent sexual selection 511 (Reynolds and Fitzpatrick, 2007; Servedio et al., 2011). Evidence suggests that dewlap coloration 512 could take this role in anoles (Ng and Glor, 2011; Lambert et al., 2013; Geneva et al., 2015; Ng et al., 2017), or at least that it is frequently involved in species recognition (Williams, 1969; Williams and Rand, 1977; Losos, 1985; Macedonia and Stamps, 1994; Fleishman, 2000; Macedonia et al., 2013; Ingram et al., 2016; Baekens et al., 2018). Although this signal is not detected at 516 the phylogenetic scale of the whole genus (Nicholson et al., 2007; Harrison and Poe, 2012; Ingram et al., 2016), sexual signals are often evolutionarily very labile (Kraaijeveld et al., 2011), and 518 the anole dewlap in particular is capable of rapid macroevolution; for example, *A. conspersus* on Grand Cayman evolved a UV-blue dewlap from an ancestral orange dewlap in 2 to 3 million years (Macedonia, 2001). We present evidence of multiple cases of potentially adaptive maintenance of 520 habitat-associated dewlap divergence over small geographical scales in *A. sagrei* across the West Indies. While these intra-island populations do not appear to be in the process of speciation, 522 our results suggest that the anoline dewlap has enough micro-scale, local adaptive potential to 523 contribute to reproductive isolation, should it be recruited for assortative mating.

525

Acknowledgements

526 Collection permission was granted by the Bahamas Environment, Science and Technology Commission, 527 the Bahamas National Trust, and the Cayman Islands Department of the Environment. 528 The authors thank Sofia Prado-Irwin, Pavitra Muralidhar, Nicholas Herrmann, Richard E. Glor, 529 Alberto R. Puente-Rolón, Kevin Aviles-Rodriguez, Kristin Winchell, Jason Fredette and Melissa

Kemp for assistance in the field and Max Lambert and James Stroud for helpful discussions. Funding for this work was provided by the Templeton Foundation (to JBL), NSF DEB #1927194 (to JBL and AJG), NSF DEB #1500761 (to AJG), NSF DBI #1609284 (to CMD), and a Harvard Museum of Comparative Zoology Putnam Expedition Grant (to RGR).

530
531
532
533

Figures

534




Figure 1: Map of the West Indies with sampled islands highlighted in black.




Figure 2: SVM classification accuracy across islands based on principal component data. Histograms show accuracy distributions over 100 replicates for each five cross-validation bins per island. The dashed line is the density of a corresponding null binomial distribution, which would be expected under random guessing (testing sets with 20% of the observations for each island and success probability of 1/3). Inset plots show the corresponding average confusion matrices and represent the proportion of lizards from each habitat (columns) reassigned in each other habitat (rows), with an interpretation guide in the right panel. Binomial test P-values indicate deviations of the mean classification accuracy to the null distribution. *, $P < 0.05$.




Figure 3: Dewlap color variation between habitat-types across the most significant islands. (A) Mapping of reflectance at various wavelengths onto the principal components (loadings from the PCA rotation matrix). (B) Distribution of PC scores between habitats along the first four PCs on each island where significant between-habitat differences were detected using SVMs. P-values are reported for univariate ANOVA (or Kruskal-Wallis tests when applicable, see Methods). Post hoc significant differences at a 0.05 error rate are indicated with horizontal bars. *, $P < 0.05$.

Tables

535

Table 1: Significance of habitat differences in dewlap coloration, using ANOVA for all islands where significant multivariate differences in dewlap coloration were detected by SVMs. Best fitting model: 1, OLS; 2, GLS. df, degrees of freedom. ΔAICc , difference in AICc between the best fitting model and the OLS-model. AICcw, AICc weight. LRT, likelihood ratio test. Log-lik., log-likelihood. χ^2 , likelihood ratio. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$.

Island	Variable	Best fit	df	AICc	ΔAICc	AICcw	ΔAICc	dfLRT	Log-lik.	χ^2	P
Abaco	PC1	1	4	710.4	0.0	0.746	2	-357.0	0.14	0.9308	***
Abaco	PC2	1	4	620.1	0.0	0.882	2	-310.2	31.74	0.0000	***
Abaco	PC3	1	4	517.8	0.0	0.732	2	-257.2	27.37	0.0000	***
Abaco	PC4	1	4	440.6	0.0	0.596	2	-217.2	1.36	0.5070	*
Bimini	PC1	1	4	561.3	0.0	0.595	2	-283.1	7.40	0.0248	*
Bimini	PC2	1	4	448.1	0.0	0.656	2	-223.8	8.09	0.0175	*
Bimini	PC3	2	6	405.3	-0.2	0.529	2	-199.2	10.39	0.0056	**
Bimini	PC4	1	4	274.2	0.0	0.854	2	-132.7	0.33	0.8499	***
Cayman Brac	PC1	2	6	402.8	-4.1	0.884	2	-200.9	13.81	0.0010	**
Cayman Brac	PC2	1	4	332.1	0.0	0.853	2	-165.9	8.41	0.0149	*
Cayman Brac	PC3	1	4	295.8	0.0	0.800	2	-146.6	27.16	0.0000	***
Cayman Brac	PC4	1	4	279.2	0.0	0.897	2	-137.8	5.63	0.0600	*
Little Cayman	PC1	1	4	367.2	0.0	0.777	2	-186.0	8.18	0.0167	*
Little Cayman	PC2	2	6	287.6	-3.6	0.859	2	-140.5	29.76	0.0000	***
Little Cayman	PC3	1	4	277.7	0.0	0.669	2	-138.1	21.34	0.0000	***
Little Cayman	PC4	1	4	226.7	0.0	0.780	2	-110.7	2.85	0.2410	
Long Island	PC1	2	6	442.3	-2.1	0.740	2	-221.2	2.91	0.2331	
Long Island	PC2	2	6	351.4	-3.1	0.823	2	-172.6	4.52	0.1043	
Long Island	PC3	1	4	322.1	0.0	0.862	2	-160.0	11.24	0.0036	**
Long Island	PC4	1	4	195.5	0.0	0.767	2	-92.9	6.46	0.0395	*

References

- Andersson, M. B. (1994). *Sexual Selection*. Monographs in Behavior and Ecology. Princeton University Press, Princeton, N.J. 536
- Arnold, S. J. (1983). Morphology, Performance and Fitness. *American Zoologist*, 23(2):347–361. 537
- Baeckens, S., Driessens, T., and Van Damme, R. (2018). The brown anole dewlap revisited: Do predation pressure, sexual selection, and species recognition shape among-population signal diversity? *PeerJ*, 6:e4722. 538
- Bartoń, K. (2019). MuMIn: Multi-Model Inference. 543
- Benjamini, Y. and Hochberg, Y. (1995). Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society Series B*, 57:289–300. 544
- Box, G. E. P. (1949). A General Distribution Theory for a Class of Likelihood Criteria. *Biometrika*, 36(3/4):317. 545
- Bradbury, J. W. and Vehrencamp, S. L. (2011). *Principles of Animal Communication*. Sinauer Associates, Sunderland, Mass, 2nd ed edition. 546
- Cook, E. G., Murphy, T. G., and Johnson, M. A. (2013). Colorful displays signal male quality in a tropical anole lizard. *Naturwissenschaften*, 100(10):993–996. 549
- Cortes, C. and Vapnik, V. (1995). Support-vector networks. *Machine Learning*, 20(3):273–297. 550
- Cortez, P. (2010). Data Mining with Neural Networks and Support Vector Machines Using the R/rminer Tool. In Perner, P., editor, *Advances in Data Mining - Applications and Theoretical Aspects 10th Industrial Conference on Data Mining (ICDM 2010), Lecture Notes in Artificial Intelligence 6171*, number 6171 in Lecture Notes in Computer Science Lecture Notes in Artificial Intelligence, pages 572–583. Springer, Berlin. 551
- Cortez, P. (2016). Rminer: Data Mining Classification and Regression Methods. 552
- Cortez, P. and Embrechts, M. J. (2013). Using sensitivity analysis and visualization techniques to open black box data mining models. *Information Sciences*, 225:1–17. 553
- Cox, R. M., Costello, R. A., Camber, B. E., and McGlothlin, J. W. (2017). Multivariate genetic architecture of the *Anolis* dewlap reveals both shared and sex-specific features of a sexually dimorphic ornament. *Journal of Evolutionary Biology*, 30(7):1262–1275. 554
- Cristianini, N. and Shawe-Taylor, J. (2000). *An Introduction to Support Vector Machines and Other Kernel-Based Learning Methods*. Cambridge University Press, first edition. 555
- Cuthill, I. C., Bennett, A. T. D., Partridge, J. C., and Maier, E. J. (1999). Plumage Reflectance and the Objective Assessment of Avian Sexual Dichromatism. *The American Naturalist*, 153(2):183–200. 556
- Dieckmann, U. and Doebeli, M. (1999). On the origin of species by sympatric speciation. *Nature*, 400(6742):354–357. 557
- Driessens, T., Baeckens, S., Balzarolo, M., Vanhooydonck, B., Huyghe, K., and Van Damme, R. (2017). Climate-related environmental variation in a visual signalling device: The male and female dewlap in *Anolis sagrei* lizards. *Journal of Evolutionary Biology*, 30(10):1846–1861. 558
- Driessens, T., Huyghe, K., Vanhooydonck, B., and Van Damme, R. (2015). Messages conveyed by assorted facets of the dewlap, in both sexes of *Anolis sagrei*. *Behavioral Ecology and Sociobiology*, 69(8):1251–1264. 559
- Driessens, T., Vanhooydonck, B., and Van Damme, R. (2014). Deterring predators, daunting opponents or drawing partners? Signaling rates across diverse contexts in the lizard *Anolis sagrei*. *Behavioral Ecology and Sociobiology*, 68(2):173–184. 560

- Dunnett, C. W. (1980). Pairwise Multiple Comparisons in the Unequal Variance Case. *Journal of the American Statistical Association*, 75(372):796–800. 581
582
- Endler, J. (1998). Sensory ecology, receiver biases and sexual selection. *Trends in Ecology & Evolution*, 13(10):415–420. 583
584
- Endler, J. A. (1984). Natural and sexual selection on color patterns in poeciliid fishes. In Balon, E. K. and Zaret, T. M., editors, *Evolutionary Ecology of Neotropical Freshwater Fishes*, volume 3, pages 95–111. Springer Netherlands, Dordrecht. 585
- Endler, J. A. (1990). On the measurement and classification of colour in studies of animal colour patterns. *Biological Journal of the Linnean Society*, 41(4):315–352. 588
589
- Endler, J. A. (1991). Variation in the appearance of guppy color patterns to guppies and their predators under different visual conditions. *Vision Research*, 31(3):587–608. 590
591
- Endler, J. A. (1992). Signals, Signal Conditions, and the Direction of Evolution. *The American Naturalist*, 139:S125–S153. 592
593
- Endler, J. A. (1993a). The Color of Light in Forests and Its Implications. *Ecological Monographs*, 63(1):1–27. 594
595
- Endler, J. A. (1993b). Some general comments on the evolution and design of animal communication systems. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 340(1292):215–225. 596
597
598
- Endler, J. A. and McLellan, T. (1988). The Processes of Evolution: Toward a Newer Synthesis. *Annual Review of Ecology and Systematics*, 19(1):395–421. 599
600
- Felsenstein, J. (1976). The Theoretical Population Genetics of Variable Selection and Migration. *Annual Review of Genetics*, 10(1):253–280. 601
602
- Fleishman, L. J. (2000). *Signal Function, Signal Efficiency and the Evolution of Anoline Lizard Dewlap Color*, pages 209–236. Tapir Academic, Trondheim. 603
604
- Fleishman, L. J., Leal, M., and Persons, M. H. (2009). Habitat light and dewlap color diversity in four species of Puerto Rican anoline lizards. *Journal of Comparative Physiology A*, 195(11):1043–1060. 605
606
607
- Fleishman, L. J. and Persons, M. (2001). The influence of stimulus and background colour on signal visibility in the lizard *Anolis cristatellus*. *The Journal of Experimental Biology*, 204(Pt 9):1559–1575. 608
609
610
- Fox, J., Friendly, M., and Monette, G. (2018). Heplots: Visualizing Tests in Multivariate Linear Models. 611
612
- Friedrich, S., Konietzschke, F., and Pauly, M. (2018). Analysis of Multivariate Data and Repeated Measures Designs with the R Package MANOVA.RM. *arXiv:1801.08002 [stat]*. 613
614
- García-Ramos, G. and Kirkpatrick, M. (1997). Genetic Models of Adaptation and Gene Flow in Peripheral Populations. *Evolution*, 51(1):21–28. 615
616
- Gavrilets, S. and Losos, J. B. (2009). Adaptive Radiation: Contrasting Theory with Data. *Science*, 323(5915):732–737. 617
618
- Geneva, A. J., Hilton, J., Noll, S., and Glor, R. E. (2015). Multilocus phylogenetic analyses of Hispaniolan and Bahamian trunk anoles (*distichus* species group). *Molecular Phylogenetics and Evolution*, 87:105–117. 619
620
621
- Gittleman, J. L. and Kot, M. (1990). Adaptation: Statistics and a Null Model for Estimating Phylogenetic Effects. *Systematic Zoology*, 39(3):227. 622
623
- Goodwin, T. W. (1984). *The Biochemistry of the Carotenoids*. Springer Netherlands, Dordrecht. 624
- Halfwerk, W., Jones, P. L., Taylor, R. C., Ryan, M. J., and Page, R. A. (2014). Risky Ripples Allow Bats and Frogs to Eavesdrop on a Multisensory Sexual Display. *Science*, 343(6169):413–416. 625
626

- 627 Harmon, L. J., Schulte, J. A., Larson, A., and Losos, J. B. (2003). Tempo and Mode of Evolutionary
628 Radiation in Iguanid Lizards. *Science*, 301(5635):961–964.
- 629 Harrison, A. and Poe, S. (2012). Evolution of an ornament, the dewlap, in females of the lizard
630 genus *Anolis*. *Biological Journal of the Linnean Society*, 106(1):191–201.
- 631 Hendry, A. P., Day, T., and Taylor, E. B. (2007a). Population Mixing and the Adaptive Divergence
of Quantitative Traits in Discrete Populations: A Theoretical Framework for Empirical Tests. *Evolution*,
632 55(3):459–466.
633
- Hendry, A. P., Taylor, E. B., and McPhail, J. D. (2007b). Adaptive Divergence and the Balance
Between Selection and Gene Flow: Lake and Stream Stickleback in the Misty System. *Evolution*,
634 56(6):1199–1216.
635
636
- Henze, N. and Zirkler, B. (1990). A class of invariant consistent tests for multivariate normality.
Communications in Statistics - Theory and Methods, 19(10):3595–3617.
637
638
- Hijmans, R. J. (2019). Geosphere: Spherical Trigonometry.
639
- Hill, G. E., Inouye, C. Y., and Montgomerie, R. (2002). Dietary carotenoids predict plumage
coloration in wild house finches. *Proceedings of the Royal Society of London. Series B: Biological
Sciences*, 269(1496):1119–1124.
640
641
642
- Hill, G. E. and McGraw, K. J., editors (2006). *Bird Coloration*. Harvard University Press, Cam-
bridge, Mass.
643
644
- Hollander, M., Wolfe, D. A., and Chicken, E. (2013). *Nonparametric Statistical Methods*. Wiley
Series in Probability and Statistics. John Wiley & Sons, Inc, Hoboken, New Jersey, third edition
edition.
645
646
647
- Howard, R. A. (1950). Vegetation of the Bimini Island Group: Bahamas, B. W. I. *Ecological
Monographs*, 20(4):317–349.
648
649
- Ingram, T., Harrison, A., Mahler, D. L., Castañeda, M. d. R., Glor, R. E., Herrel, A., Stuart,
Y. E., and Losos, J. B. (2016). Comparative tests of the role of dewlap size in *Anolis* lizard
speciation. *Proceedings of the Royal Society B: Biological Sciences*, 283(1845):20162199.
650
651
652
- Kamath, A. and Losos, J. B. (2018). Estimating encounter rates as the first step of sexual se-
lection in the lizard *Anolis sagrei*. *Proceedings of the Royal Society B: Biological Sciences*,
285(1873):20172244.
653
654
655
- Kim, B. and von Oertzen, T. (2018). Classifiers as a model-free group comparison test. *Behavior
Research Methods*, 50(1):416–426.
656
657
- Kolbe, J. J., Larson, A., Losos, J. B., and de Queiroz, K. (2008). Admixture determines genetic
diversity and population differentiation in the biological invasion of a lizard species. *Biology
Letters*, 4(4):434–437.
658
659
660
- Kolbe, J. J., Leal, M., Schoener, T. W., Spiller, D. A., and Losos, J. B. (2012). Founder Ef-
fects Persist Despite Adaptive Differentiation: A Field Experiment with Lizards. *Science*,
335(6072):1086–1089.
661
662
663
- Korkmaz, S., Goksuluk, D., and Zararsiz, G. (2014). MVN: An R Package for Assessing Multi-
variate Normality. *The R Journal*, 6(2):151–162.
664
665
- Kraaijeveld, K., Kraaijeveld-Smit, F. J. L., and Maan, M. E. (2011). Sexual selection and specia-
tion: The comparative evidence revisited. *Biological Reviews*, 86(2):367–377.
666
667
- Lambert, S. M., Geneva, A. J., Luke Mahler, D., and Glor, R. E. (2013). Using genomic data
to revisit an early example of reproductive character displacement in Haitian *Anolis* lizards.
Molecular Ecology, 22(15):3981–3995.
668
669
670
- Lazareva, O. F., Shimizu, T., and Wasserman, E. A. (2012). *How Animals See the WorldCompar-
ative Behavior, Biology, and Evolution of Vision*. Oxford University Press.
671
672

- Leal, M. and Fleishman, L. J. (2002). Evidence for habitat partitioning based on adaptation to environmental light in a pair of sympatric lizard species. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, 269(1489):351–359. 673
674
675
- Leal, M. and Fleishman, L. J. (2004). Differences in Visual Signal Design and Detectability between Allopatric Populations of *Anolis* Lizards. *The American Naturalist*, 163(1):26–39. 676
677
- ⁶⁷⁸ Leal, M. and Rodríguez-Robles, J. A. (1995). Antipredator Responses of *Anolis cristatellus* (Sauria: Polychrotidae). *Copeia*, 1995(1):155–161.
- ⁶⁸⁰ Leal, M. and Rodriguez-Robles, J. A. (1997). Antipredator Responses of the Puerto Rican Giant Anole, *Anolis cuvieri* (Squamata: Polychrotidae). *Biotropica*, 29(3):372–375. 675
- ⁶⁸² Leal, M. and Rodríguez-Robles, J. A. (1997). Signalling displays during predator–prey interactions in a Puerto Rican anole, *Anolis cristatellus*. *Animal Behaviour*, 54(5):1147–1154. 676
- ⁶⁸⁴ Lenormand, T. (2002). Gene flow and the limits to natural selection. *Trends in Ecology & Evolution*, 17(4):183–189. 677
- ⁶⁸⁶ Losos, J. B. (1985). An Experimental Demonstration of the Species-Recognition Role of *Anolis* Dewlap Color. *Copeia*, 1985(4):905–910. 678
- ⁶⁸⁸ Losos, J. B. (2009). *Lizards in an Evolutionary Tree: Ecology and Adaptive Radiation of Anoles*. University of California Press. 679
- ⁶⁹⁰ Losos, J. B. (2011). Convergence, Adaptation, and Constraint. *Evolution*, 65(7):1827–1840. 680
- ⁶⁹¹ Losos, J. B., Irschick, D. J., and Schoener, T. W. (1994). Adaptation and Constraint in the Evolution of Specialization of Bahamian *Anolis* Lizards. *Evolution*, 48(6):1786–1798. 681
- ⁶⁹³ Losos, J. B., Schoener, T. W., Warheit, K. I., and Creer, D. (2001). Experimental studies of adaptive differentiation in Bahamian *Anolis* lizards. *Genetica*, 112–113:399–415. 682
- ⁶⁹⁵ Losos, J. B., Warheit, K. I., and Schoener, T. W. (1997). Adaptive differentiation following experimental island colonization in *Anolis* lizards. *Nature*, 387(6628):70–73. 683
- ⁶⁹⁷ Macedonia, J. M. (2001). Habitat light, colour variation, and ultraviolet reflectance in the Grand Cayman anole, *Anolis conspersus*. *Biological Journal of the Linnean Society*, 73(3):299–320. 684
- ⁶⁹⁹ Macedonia, J. M., Clark, D. L., Riley, R. G., and Kemp, D. J. (2013). Species recognition of color and motion signals in *Anolis grahami*: Evidence from responses to lizard robots. *Behavioral Ecology*, 24(4):846–852. 685
- ⁷⁰² Macedonia, J. M., Clark, D. L., and Tamasi, A. L. (2014). Does Selection Favor Dewlap Colors that Maximize Detectability? A Test with Five Species of Jamaican *Anolis* Lizards. *Herpetologica*, 70(2):157–170. 686
- ⁷⁰⁵ Macedonia, J. M., James, S., Wittle, L. W., and Clark, D. L. (2000). Skin Pigments and Coloration in the Jamaican Radiation of *Anolis* Lizards. *Journal of Herpetology*, 34(1):99–109. 687
- ⁷⁰⁷ Macedonia, J. M. and Stamps, J. A. (1994). Species Recognition in *Anolis grahami* (Sauria, Iguanidae): Evidence from Responses to Video Playbacks of Conspecific and Heterospecific Displays. *Ethology*, 98(3–4):246–264. 688
- ⁷¹⁰ Maia, R., Eliason, C. M., Bitton, P.-P., Doucet, S. M., and Shawkey, M. D. (2013). Pavo: An R package for the analysis, visualization and organization of spectral data. *Methods in Ecology and Evolution*, pages n/a–n/a. 689
- ⁷¹³ Miles, L. S., Rivkin, L. R., Johnson, M. T. J., Munshi-South, J., and Verrelli, B. C. (2019). Gene flow and genetic drift in urban environments. *Molecular Ecology*, 28(18):4138–4151. 690
- ⁷¹⁵ Morrison, D. F. (1988). *Multivariate Statistical Methods*. McGraw-Hill Series in Probability and Statistics. McGraw-Hill, Hamburg Auckland. 691
- ⁷¹⁷ Nemenyi, P. (1963). *Distribution-Free Multiple Comparisons*. Ph. D. dissertation, Princeton University, Princeton, NJ. 692

- ⁷¹⁹ Ng, J., Geneva, A. J., Noll, S., and Glor, R. E. (2017). Signals and Speciation: *Anolis* Dewlap
⁷²⁰ Color as a Reproductive Barrier. *Journal of Herpetology*, 51(3):437–447.
- ⁷²¹ Ng, J. and Glor, R. E. (2011). Genetic differentiation among populations of a Hispaniolan trunk
⁷²² anole that exhibit geographical variation in dewlap colour. *Molecular Ecology*, 20(20):4302–4317.
- ⁷²³ Ng, J., Kelly, A. L., MacGuigan, D. J., and Glor, R. E. (2013). The Role of Heritable and Dietary
Factors in the Sexual Signal of a Hispaniolan *Anolis* Lizard, *Anolis distichus*. *Journal of Heredity*,
104(6):862–873. ⁷²⁴
⁷²⁵
- Ng, J., Landeen, E. L., Logsdon, R. M., and Glor, R. E. (2012). Correlation Between *Anolis*
Lizard Dewlap Phenotype and Environmental Variation Indicates Adaptive Divergence of a
Signal Important to Sexual Selection and Species Recognition. *Evolution*, 67(2):573–582. ⁷²⁶
⁷²⁷
⁷²⁸
- Ng, J., Ossip-Klein, A. G., and Glor, R. E. (2016). Adaptive signal coloration maintained in the
face of gene flow in a Hispaniolan *Anolis* Lizard. *BMC Evolutionary Biology*, 16(1):193. ⁷²⁹
⁷³⁰
- Nicholson, K. E., Harmon, L. J., and Losos, J. B. (2007). Evolution of *Anolis* Lizard Dewlap
Diversity. *PLoS ONE*, 2(3):e274. ⁷³¹
⁷³²
- Nosil, P. and Crespi, B. J. (2004). Does Gene Flow Constrain Adaptive Divergence or Vice Versa? A
Test Using Ecomorphology and Sexual Isolation in *Timema cristinae* Walking-Sticks. *Evolution*,
58(1):102–112. ⁷³³
⁷³⁴
⁷³⁵
- Ortiz, E. (1962). Drosopterins in the dewlap of some anoline lizards. *American Zoologist*, 2:545–546. ⁷³⁶
- Ortiz, E. and Maldonado, A. A. (1966). Pteridine accumulation in lizards of the genus *Anolis*.
Caribbean Journal of Science, 6:9–13. ⁷³⁷
⁷³⁸
- Ortiz, E., Throckmorton, L. H., and Williams-Ashman, H. G. (1962). Drosopterins in the Throat-
Fans of Some Puerto Rican Lizards. *Nature*, 196(4854):595–596. ⁷³⁹
⁷⁴⁰
- Ortiz, E. and Williams-Ashman, H. (1963). Identification of skin pteridines in the pasture lizard
Anolis pulchellus. *Comparative Biochemistry and Physiology*, 10(3):181–190. ⁷⁴¹
⁷⁴²
- Pigot, A. L., Sheard, C., Miller, E. T., Bregman, T. P., Freeman, B. G., Roll, U., Seddon, N.,
Trisos, C. H., Weeks, B. C., and Tobias, J. A. (2020). Macroevolutionary convergence connects
morphological form to ecological function in birds. *Nature Ecology & Evolution*, 4(2):230–239. ⁷⁴³
⁷⁴⁴
⁷⁴⁵
- Pinheiro, J. and Bates, D. (2000). *Mixed-Effects Models in S and S-PLUS*. Statistics and Com-
puting. Springer-Verlag, New York. ⁷⁴⁶
⁷⁴⁷
- Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., and Team, R. C. (2020). Nlme: Linear and
Nonlinear Mixed Effects Models. ⁷⁴⁸
⁷⁴⁹
- Pohlert, T. (2020). PMCMRplus: Calculate Pairwise Multiple Comparisons of Mean Rank Sums
Extended. R package version 1.4.4. ⁷⁵⁰
⁷⁵¹
- R Core Team (2019). R: A Language and Environment for Statistical Computing. R Foundation
for Statistical Computing. ⁷⁵²
⁷⁵³
- Rand, A. S. and Williams, E. E. (1970). An Estimation of Redundancy and Information Content
of Anole Dewlaps. *The American Naturalist*, 104(935):99–103. ⁷⁵⁴
⁷⁵⁵
- Reynolds, R. G. and Fitzpatrick, B. M. (2007). Assortative Mating in Poison-Dart Frogs Based
on an Ecologically Important Trait. *Evolution*, 61(9):2253–2259. ⁷⁵⁶
⁷⁵⁷
- Reynolds, R. G., Kolbe, J. J., Glor, R. E., López-Darias, M., Gómez Pourroy, C. V., Harrison,
A. S., Queiroz, K., Revell, L. J., and Losos, J. B. (2020). Phylogeographic and phenotypic
outcomes of brown anole colonization across the Caribbean provide insight into the beginning
stages of an adaptive radiation. *Journal of Evolutionary Biology*, 33(4):468–494. ⁷⁵⁸
⁷⁵⁹
⁷⁶⁰
⁷⁶¹
- Richardson, J. L. and Urban, M. C. (2013). Strong Selection Barriers Explain Microgeographic
Adaptation in Wild Salamander Populations. *Evolution*, 67(6):1729–1740. ⁷⁶²
⁷⁶³

- Richardson, J. L., Urban, M. C., Bolnick, D. I., and Skelly, D. K. (2014). Microgeographic adaptation and the spatial scale of evolution. *Trends in Ecology & Evolution*, 29(3):165–176. 764
765
- Ripley, B. D. (1996). *Pattern Recognition and Neural Networks*. Cambridge University Press, first edition. 766
767
- Schoener, T. W. (1968). The *Anolis*Lizards of Bimini: Resource Partitioning in a Complex Fauna. *Ecology*, 49(4):704–726. 768
- Seehausen, O. (1997). Cichlid Fish Diversity Threatened by Eutrophication That Curbs Sexual Selection. *Science*, 277(5333):1808–1811.
- Servedio, M. R., Doorn, G. S. V., Kopp, M., Frame, A. M., and Nosil, P. (2011). Magic traits in speciation: ‘magic’ but not rare? *Trends in Ecology & Evolution*, 26(8):389–397.
- Sigmund, W. R. (1983). Female Preference for *Anolis carolinensis* Males as a Function of Dewlap Color and Background Coloration. *Journal of Herpetology*, 17(2):137–143.
- Stapley, J., Wordley, C., and Slate, J. (2011). No Evidence of Genetic Differentiation Between Anoles With Different Dewlap Color Patterns. *Journal of Heredity*, 102(1):118–124.
- Steffen, J. E. and Guyer, C. C. (2014). Display behaviour and dewlap colour as predictors of contest success in brown anoles: Dewlap Colour and Behaviour in Contests. *Biological Journal of the Linnean Society*, 111(3):646–655.
- Steffen, J. E., Hill, G. E., and Guyer, C. (2010). Carotenoid Access, Nutritional Stress, and the Dewlap Color of Male Brown Anoles. *Copeia*, 2010(2):239–246.
- Steffen, J. E. and McGraw, K. J. (2007). Contributions of pterin and carotenoid pigments to dewlap coloration in two anole species. *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology*, 146(1):42–46.
- Steffen, J. E. and McGraw, K. J. (2009). How dewlap color reflects its carotenoid and pterin content in male and female brown anoles (*Norops sagrei*). *Comparative Biochemistry and Physiology Part B: Biochemistry and Molecular Biology*, 154(3):334–340.
- Tariel, J., Plénet, S., and Luquet, É. (2020). Transgenerational plasticity of inducible defences: Combined effects of grand-parental, parental and current environments. *Ecology and Evolution*, 10(5):2367–2376.
- Thorpe, R. S. (2002). Analysis of Color Spectra in Comparative Evolutionary Studies: Molecular Phylogeny and Habitat Adaptation in the St. Vincent Anole (*Anolis trinitatis*). *Systematic Biology*, 51(4):554–569.
- Thorpe, R. S. and Stenson, A. G. (2002). Phylogeny, Paraphyly and Ecological Adaptation of the Colour and Pattern in the Anolis Roquet Complex on Martinique: Interaction Between Phylogeny and Adaptation. *Molecular Ecology*, 12(1):117–132.
- Tokarz, R. R. (2002). An Experimental Test of the Importance of the Dewlap in Male Mating Success in the Lizard *Anolis sagrei*. *Herpetologica*, 58(1):87–94.
- Tokarz, R. R. (2006). Importance of Prior Physical Contact with Familiar Females in the Development of a Male Courtship and Mating Preference for Unfamiliar Females in the Lizard *Anolis Sagrei*. *Herpetologica*, 62(2):115–124.
- Tokarz, R. R., Paterson, A. V., and McMann, S. (2005). Importance of Dewlap Display in Male Mating Success in Free-Ranging Brown Anoles (*Anolis sagrei*). *Journal of Herpetology*, 39(1):174–177.
- Tukey, J. W. (1949). Comparing Individual Means in the Analysis of Variance. *Biometrics*, 5(2):99.
- Vanhooydonck, B., Herrel, A., Meyers, J. J., and Irschick, D. J. (2009). What determines dewlap diversity in *Anolis* lizards? An among-island comparison. *Journal of Evolutionary Biology*, 22(2):293–305. 806
807
808
809

- 810 Vanhooydonck, B., Herrel, A. Y., Van Damme, R., and Irschick, D. J. (2005). Does dewlap size
811 predict male bite performance in Jamaican *Anolis* lizards? *Functional Ecology*, 19(1):38–42.
- 812 Venables, W. N. and Ripley, B. D. (2002). *Modern Applied Statistics with S*. Statistics and
813 Computing. Springer, New York, 4th ed edition.
- 814 Willi, Y. and Hoffmann, A. A. (2012). Microgeographic adaptation linked to forest fragmentation
and habitat quality in the tropical fruit fly *Drosophila birchii*. *Oikos*, 121(10):1627–1637. 815
- Williams, E. E. (1969). The Ecology of Colonization as Seen in the Zoogeography of Anoline 816
Lizards on Small Islands. *The Quarterly Review of Biology*, 44(4):345–389. 817
- Williams, E. E. and Rand, A. S. (1977). Species Recognition, Dewlap Function and Faunal Size. 818
American Zoologist, 17(1):261–270. 819
- Zuur, A. F., editor (2009). *Mixed Effects Models and Extensions in Ecology with R*. Statistics for 820
Biology and Health. Springer, New York, NY. 821

Supplementary Figures

822




Figure S1: Map of the sampling sites and corresponding habitats across nine islands of the West Indies.




Figure S2: Correlation between dewlap brightness (as measured by the mean reflectance from 300 to 700nm in wavelength) and PC1 score for each island. Pearson's squared correlation coefficients are reported. *, $P < 0.05$.




Figure S3: Correlation between dewlap brightness (as measured by the mean reflectance from 300 to 700nm in wavelength) and PC1 score across the whole archipelago. Pearson's squared correlation coefficient is reported. *, $P < 0.05$.




Figure S4: (A) Principal component scores and 5% confidence ellipses across habitats for the whole archipelago. The principal component analysis was performed on reflectance data from all islands pooled together. (B) PCA rotation matrix showing the loadings of each wavelength from 300 to 700nm onto the principal components.




Figure S5: 5-95th percentile range of lizard dewlap reflectance values (in % of incoming light) across wavelengths for each island and each habitat.




Figure S6: Archipelago-wide SVM classification accuracy based on principal component data. Machines were trained on individual dewlaps regardless of island identity. The histogram shows the accuracy distribution over 100 replicates for each five cross-validation bins. The legend is the same as in Figure 2.




Figure S7: Archipelago-wide SVM classification accuracy based on reflectance data at 50nm-intervals in wavelength (see Methods). Machines were trained on individual dewlaps regardless of island identity. The histogram shows the accuracy distribution over 100 replicates for each five cross-validation bins. The legend is the same as in Figure 2.




Figure S8: Sensitivity analyses of the different input variables in the archipelago-wide SVM classification on principal component data (Figure S6), with relative importance computed for every machine.




Figure S9: Sensitivity analyses of the different input variables in the archipelago-wide SVM classification on reflectance data at 50nm-intervals in wavelength (Figure S7), with relative importance computed for every machine.




Figure S10: Archipelago-wide LDA classification accuracy based on principal component data. Machines were trained on individual dewlaps regardless of island identity. The histogram shows the accuracy distribution over 100 replicates for each five cross-validation bins. The legend is the same as in Figure 2.




Figure S11: Archipelago-wide LDA classification accuracy based on reflectance data at 50nm intervals in wavelength (see Methods). Machines were trained on individual dewlaps regardless of island identity. The histogram shows the accuracy distribution over 100 replicates for each five cross-validation bins. The legend is the same as in Figure 2.




Figure S12: LDA classification accuracy across islands based on principal component data. Histograms show accuracy distributions over 100 replicates for each five cross-validation bins per island. The legend is the same as in Figure 2.




Figure S13: SVM classification accuracy across islands based on reflectance data at 50nm-intervals in wavelength (see Methods). Histograms show accuracy distributions over 100 replicates for each five cross-validation bins per island. The legend is the same as in Figure 2.




Figure S14: LDA classification accuracy across islands based on reflectance data at 50nm-intervals in wavelength (see Methods). Histograms show accuracy distributions over 100 replicates for each five cross-validation bins per island. The legend is the same as in Figure 2.




Figure S15: Sensitivity analyses of the different input variables in the within-island SVM classification on principal component data (Figure ??), with relative importance computed for every machine.




Figure S16: Sensitivity analyses of the different input variables in the within-island LDA classification on principal component data (Figure S12), with relative importance computed for every machine.




Figure S17: Sensitivity analyses of the different input variables in the archipelago-wide SVM classification on reflectance at 50nm-intervals in wavelength (Figure S13), with relative importance computed for every machine.




Figure S18: Sensitivity analyses of the different input variables in the archipelago-wide LDA classification on reflectance at 50nm-intervals in wavelength (Figure S14), with relative importance computed for every machine.




Figure S19: Spatial scale of between-habitat differences in dewlap coloration. For each variable and each pair of habitats where significant differences were detected (Figure 3), we performed multiple post hoc pairwise comparisons between the sites involved (Figure S1, Table S2), using nonparametric Wilcoxon-Mann-Whitney tests. Here we report, for each pair of habitats, the distances between sites that significantly differed in dewlap coloration at an error rate of 0.05 (P-values corrected with the Benjamini-Hochberg procedure for multiple testing).




Figure S20: Test of parallel divergence between islands. Differences in habitat-means, or contrasts, are shown for two pairs of habitats for each principal component on each island, rescaled so the standard deviation of the means along each principal component is one. The contrasts represent the patterns of between-habitat variation on each island, for a given principal component. The absence of clustering of islands by variable indicates that islands differ in their between-habitat divergence patterns. This is confirmed by a non-significant MANOVA test of the between versus within-variable variance in contrasts.

Supplementary Tables

823

Table S1: Number of lizards sampled in each habitat on each island.

	coastal	coppice	mangrove
Abaco	41	24	21
Bimini	38	14	15
Cayman Brac	15	18	17
Eleuthera	22	25	9
Little Cayman	17	12	16
Long Island	26	14	13
North Andros	9	9	10
Ragged Island	18	15	17
South Andros	10	9	12

Table S2: Locations of the sampling sites across islands, with mean principal component scores per site.

Island	Longitude	Latitude	Habitat	PC1	PC2	PC3	PC4
Abaco	-77.7256	26.9083	mangrove	-5.4905	1.3541	-0.4741	0.0083
Abaco	-77.5800	26.9020	coastal	1.8633	0.0365	-0.4475	0.0033
Abaco	-77.5763	26.9128	coppice	-1.6738	-1.7793	-0.0499	0.0012
Abaco	-77.1784	26.1045	coastal	1.1863	2.0408	-0.3468	0.0022
Abaco	-77.0055	26.3254	mangrove	-9.0319	-2.7460	0.4687	0.0077
Abaco	-77.0039	26.3170	coppice	0.9967	0.5161	-0.0267	-0.0118
Abaco	-76.9968	26.3260	coastal	7.6077	0.3186	0.1771	-0.0008
Bimini	-79.3022	25.5859	coastal	5.7537	-0.1593	-0.2505	0.0001
Bimini	-79.3014	25.7052	coastal	-3.1822	1.6617	-0.0460	0.0024
Bimini	-79.3002	25.7042	coppice	-1.3514	-3.8786	0.1027	-0.0027
Bimini	-79.2709	25.7066	mangrove	3.3656	0.6244	0.1569	-0.0021
Cayman Brac	-79.8627	19.6878	coastal	6.6606	-2.5670	0.0166	-0.0007
Cayman Brac	-79.8441	19.6949	mangrove	-1.0914	4.3607	0.0855	0.0001
Cayman Brac	-79.7887	19.7209	coppice	-4.5197	-1.9793	-0.0946	0.0004
Eleuthera	-76.3347	24.8146	coppice	3.2669	-1.2404	0.1018	-0.0085
Eleuthera	-76.3058	24.8127	coastal	0.4216	-3.5133	-0.0567	0.0009
Eleuthera	-76.2901	24.7981	mangrove	2.1881	0.7517	0.3957	-0.0055
Eleuthera	-76.1616	24.9129	coppice	-1.9136	1.0868	-0.4978	-0.0092
Eleuthera	-76.1492	24.9335	coastal	-3.1863	2.4270	0.1881	0.0218
Little Cayman	-80.0660	19.6906	coppice	0.8021	-1.9569	-0.0760	-0.0068
Little Cayman	-80.0205	19.6865	coastal	-6.6917	-1.2615	0.0659	0.0057
Little Cayman	-79.9871	19.6986	mangrove	6.5083	2.8079	-0.0129	-0.0010
Long Island	-75.2299	23.4740	mangrove	-1.2873	1.9371	-0.1880	-0.0029
Long Island	-75.2063	23.4282	coastal	2.3686	-0.9033	0.0215	0.0096
Long Island	-75.1884	23.4292	coppice	-4.6266	0.5060	0.1049	-0.0070
Long Island	-75.1408	23.3883	coastal	3.6139	-1.4521	0.0475	0.0025
North Andros	-77.8908	24.8391	coastal	-2.1881	-1.1236	0.0397	-0.0060
North Andros	-77.8428	24.7516	coppice	-1.8115	0.0012	-0.1678	0.0024
North Andros	-77.7540	24.6644	mangrove	3.5997	1.0101	0.1153	0.0033
Ragged Island	-75.7364	22.1768	coppice	3.2851	-0.3274	0.1911	-0.0013
Ragged Island	-75.7314	22.2097	coastal	-0.6412	-0.8878	-0.1293	-0.0033
Ragged Island	-75.7276	22.2045	mangrove	-2.9188	1.5792	-0.0034	0.0099
Ragged Island	-75.7270	22.1973	mangrove	-1.2210	0.7285	-0.0721	-0.0028
South Andros	-77.6050	24.2027	mangrove	-3.9253	0.4734	0.0477	-0.0005
South Andros	-77.5936	24.1289	coppice	6.1152	-0.4925	0.0349	0.0012
South Andros	-77.5453	24.0764	coastal	-0.7933	-0.1248	-0.0887	-0.0004

Table S3: Proportion of variance explained by the first four principal components on each island, as well as across the whole archipelago.

Island	PC1	PC2	PC3	PC4	Total
Abaco	0.400	0.279	0.147	0.079	0.906
Bimini	0.502	0.208	0.160	0.051	0.921
Cayman Brac	0.438	0.190	0.155	0.105	0.888
Eleuthera	0.490	0.233	0.138	0.066	0.926
Little Cayman	0.441	0.212	0.176	0.078	0.907
Long Island	0.515	0.205	0.161	0.043	0.925
North Andros	0.560	0.170	0.152	0.054	0.937
Ragged Island	0.483	0.226	0.127	0.072	0.907
South Andros	0.488	0.247	0.146	0.067	0.948
Archipelago	0.473	0.197	0.164	0.079	0.913

Table S4: Pearson's correlation test between dewlap brightness, as measured by the average reflectance between 300 and 700nm in wavelength, and PC1 scores, for all islands and across the whole archipelago. ***, $P < 0.001$.

Island	r^2	P	
Abaco	0.908	< 0.0001	***
Bimini	0.999	< 0.0001	***
Cayman Brac	0.987	< 0.0001	***
Eleuthera	0.963	< 0.0001	***
Little Cayman	0.965	< 0.0001	***
Long Island	0.986	< 0.0001	***
North Andros	0.994	< 0.0001	***
Ragged Island	0.978	< 0.0001	***
South Andros	0.979	< 0.0001	***
Archipelago	0.976	< 0.0001	***

Table S5: Henze-Zirkler's test of multivariate normality, performed on principal components in each habitat and on each island. HZ, test statistic. *, P < 0.05; **, P < 0.01; ***, P < 0.001.

Island	Habitat	HZ	P	
Abaco	coastal	1.10	0.0027	**
Abaco	coppice	1.07	0.0022	**
Abaco	mangrove	1.06	0.0023	**
Bimini	coastal	1.28	0.0001	***
Bimini	coppice	0.85	0.0482	*
Bimini	mangrove	1.19	0.0001	***
Cayman Brac	coastal	0.65	0.5311	
Cayman Brac	coppice	0.70	0.3940	
Cayman Brac	mangrove	0.66	0.5357	
Eleuthera	coastal	1.61	0.0000	***
Eleuthera	coppice	1.48	0.0000	***
Eleuthera	mangrove	0.73	0.1423	
Little Cayman	coastal	0.62	0.6599	
Little Cayman	coppice	0.64	0.4867	
Little Cayman	mangrove	0.87	0.0413	*
Long Island	coastal	0.82	0.1468	
Long Island	coppice	0.92	0.0150	*
Long Island	mangrove	0.77	0.1289	
North Andros	coastal	0.66	0.3174	
North Andros	coppice	0.76	0.0900	
North Andros	mangrove	0.67	0.3185	
Ragged Island	coastal	0.76	0.2268	
Ragged Island	coppice	0.80	0.1115	
Ragged Island	mangrove	0.54	0.9022	
South Andros	coastal	0.66	0.3451	
South Andros	coppice	0.66	0.3154	
South Andros	mangrove	0.91	0.0144	*

Table S6: Box's M-test of homogeneity of covariance matrices across habitats on each island. χ^2 , test statistic. *, P < 0.05; **, P < 0.01; ***, P < 0.001.

Island	χ^2	df	P	
Abaco	47.1	20	0.0006	***
Bimini	36.0	20	0.0152	*
Cayman Brac	36.9	20	0.0120	*
Eleuthera	44.6	20	0.0013	**
Little Cayman	32.8	20	0.0356	*
Long Island	56.2	20	0.0000	***
North Andros	33.7	20	0.0283	*
Ragged Island	29.3	20	0.0824	
South Andros	46.5	20	0.0007	***

Table S7: Shapiro-Wilk's test of univariate normality performed on each island where significant differences were detected by SVM classification, in each habitat where deviations from multivariate normality were detected. W , test statistic. *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.001$.

Island	Habitat	Variable	W	P
Abaco	coastal	PC1	0.954	0.0941
Abaco	coastal	PC2	0.927	0.0112 *
Abaco	coastal	PC3	0.973	0.4228
Abaco	coastal	PC4	0.955	0.1027
Abaco	coppice	PC1	0.970	0.6776
Abaco	coppice	PC2	0.816	0.0005 ***
Abaco	coppice	PC3	0.930	0.0976
Abaco	coppice	PC4	0.941	0.1711
Abaco	mangrove	PC1	0.881	0.0155 *
Abaco	mangrove	PC2	0.869	0.0093 **
Abaco	mangrove	PC3	0.986	0.9873
Abaco	mangrove	PC4	0.939	0.2044
Bimini	coastal	PC1	0.821	0.0000 ***
Bimini	coastal	PC2	0.960	0.1854
Bimini	coastal	PC3	0.856	0.0002 ***
Bimini	coastal	PC4	0.945	0.0611
Bimini	coppice	PC1	0.911	0.1648
Bimini	coppice	PC2	0.958	0.6927
Bimini	coppice	PC3	0.953	0.6146
Bimini	coppice	PC4	0.971	0.8953
Bimini	mangrove	PC1	0.884	0.0536
Bimini	mangrove	PC2	0.976	0.9363
Bimini	mangrove	PC3	0.982	0.9805
Bimini	mangrove	PC4	0.975	0.9232
Eleuthera	coastal	PC1	0.909	0.0461 *
Eleuthera	coastal	PC2	0.886	0.0157 *
Eleuthera	coastal	PC3	0.906	0.0390 *
Eleuthera	coastal	PC4	0.962	0.5293
Eleuthera	coppice	PC1	0.922	0.0567
Eleuthera	coppice	PC2	0.954	0.3055
Eleuthera	coppice	PC3	0.781	0.0001 ***
Eleuthera	coppice	PC4	0.901	0.0188 *
Little Cayman	mangrove	PC1	0.907	0.1024
Little Cayman	mangrove	PC2	0.904	0.0924
Little Cayman	mangrove	PC3	0.739	0.0005 ***
Little Cayman	mangrove	PC4	0.973	0.8802
Long Island	coppice	PC1	0.686	0.0003 ***
Long Island	coppice	PC2	0.848	0.0210 *
Long Island	coppice	PC3	0.931	0.3188
Long Island	coppice	PC4	0.904	0.1280
South Andros	mangrove	PC1	0.787	0.0067 **
South Andros	mangrove	PC2	0.861	0.0500 *
South Andros	mangrove	PC3	0.697	0.0008 ***
South Andros	mangrove	PC4	0.950	0.6411

Table S8: Univariate ANOVAs performed on each principal component across the whole archipelago. Legend is the same as for Table 1, except that best fitting models 3 and 4 refer to the mixed effect equivalents to the OLS and GLS model, with island as a random effect (see Methods).

Variable	Best fit	df	AICc	ΔAICc	AICcw	df _{LRT}	Log-lik.	χ^2	P
PC1	3	5	3749.9	-228.3	0.613	2	-1874.7	8.69	0.0130 *
PC2	4	7	3002.2	-162.3	0.976	2	-1496.2	17.76	0.0001 ***
PC3	4	7	2826.3	-175.4	0.968	2	-1407.8	7.03	0.0298 *
PC4	4	7	2015.7	-305.8	0.519	2	-1000.1	0.47	0.7914

Table S9: Mean SVM classification accuracy per island, over all replicates and cross-validation bins. N , number of observations per island; p_{test} , proportion of the data sampled to form the training set; n_{test} , number of observations in the testing set. P-values indicate deviations from the expected null binomial distribution, with n_{test} events per island and random guess success probability 1/3. *, $P < 0.05$, **, $P < 0.01$, ***, $P < 0.001$.

Island	Accuracy	N	p_{test}	n_{test}	P
Abaco	0.612	86	0.2	17	0.0080 **
Bimini	0.547	67	0.2	13	0.0347 *
Cayman Brac	0.721	50	0.2	10	0.0034 **
Eleuthera	0.437	56	0.2	11	0.2890
Little Cayman	0.734	45	0.2	9	0.0083 **
Long Island	0.651	53	0.2	10	0.0197 *
North Andros	0.453	28	0.2	5	0.2099
Ragged Island	0.364	50	0.2	10	0.4407
South Andros	0.600	31	0.2	6	0.1001

Table S10: Results of nonparametric Kruskal-Wallis tests performed on each variable on each island where deviations from normality were detected.

Island	Variable	χ^2	df	P
Abaco	PC1	0.74	2	0.6924
Abaco	PC2	23.13	2	0.0000 ***
Bimini	PC1	7.38	2	0.0250 *
Bimini	PC3	15.17	2	0.0005 ***
Little Cayman	PC3	19.95	2	0.0000 ***
Long Island	PC1	10.98	2	0.0041 **
Long Island	PC2	4.02	2	0.1339

Table S11: Individual-based permutation tests of spatial autocorrelation within islands. P-values were computed from 1,000 permutations of individual site-labels. Pearson's coefficient r measures the correlation between distances in color space and geodesic distances among the sites. N , number of sites. *, $P < 0.05$.

Island	r	P	N
Abaco	-0.213	0.817	7
Bimini	0.044	0.510	4
Cayman Brac	-0.010	0.465	3
Eleuthera	0.816	0.015	5 *
Little Cayman	-0.688	0.684	3
Long Island	-0.189	0.579	4
North Andros	0.730	0.199	3
Ragged Island	0.706	0.114	4
South Andros	-0.852	0.776	3