

# High-Redshift Atomic Cooling Halos generally lead to Intermediate-Massive Black Hole Seeds

Ryan Hazlett<sup>1</sup>, Eli Visbal<sup>1</sup>, Greg L. Bryan<sup>2</sup>, Mihir Kulkarni<sup>3</sup>

<sup>1</sup>Ritter Astrophysical Research Center, Department of Physics and Astronomy, University of Toledo (ryan.hazlett@rockets.utoledo.edu)

<sup>2</sup>Department of Astronomy, Columbia University

<sup>3</sup>Institut für Astrophysik und Geophysik, Georg-August Universität Göttingen



## INTRODUCTION

- Recent observations by JWST have revealed a population of high- $z$  AGN hosting massive black holes with masses between  $10^6 - 10^8 M_\odot$  (Larson2023+; Greene2024+).
- These observations hint at the need for heavy black hole seeds with predicted masses  $\geq 10^5 M_\odot$  formed by the first stars.
- We explore two theoretical pathways that can form supermassive Pop III stars and heavy black hole seeds:
  - A high Lyman-Werner background,  $J_{21}$ , that suppresses cooling by dissociating  $H_2$ .
  - High halo assembly rate,  $\dot{M}$ , on to the halo that delays star formation through dynamical heating.

## OBJECTIVE

- Our goal is to determine how common it is expected to form heavy BH seeds by studying at high-resolution, the gas accretion rate on halos with varying  $J_{21}$  and accretion rate
- We explore a parameter space that spans a range of  $J_{21}$  and  $\dot{M}$  for a large sample of halos and determine the impact on the ultimate predicted Pop III mass.


## SIMULATION & RESULTS

- Hydrodynamical simulations were carried out using adaptive mesh refinement code *ENZO* (Bryan2014+) using a dark matter resolution of  $7000 M_\odot$  and a maximum level of refinement of  $10^{-4}$  pc. See top-center figure for an example.
- The average radial profiles (see bottom-middle figure) are generally similar for different  $J_{21}$ . A notable difference are the elevated average inflow rates for the  $J_{21} \geq 10$  simulations.
- We also find little correlation for different  $\dot{M}$ , with variations in  $J_{21}$  having a more dominant effect.


## REFERENCES

- Bryan et al. 2014, *ApJS*, 211, 2  
 Greene et al. 2024, *ApJ*, 964, 39  
 Hammerlé et al. 2018, *MNRAS*, 474, 2757  
 Larson et al. 2023, *ApJ*, 953, L29  
 Latif et al. 2021, *MNRAS*, 508, 1756  
 Regan et al. 2015, *MNRAS*, 449, 3766  
 Regan et al. 2020, *OJAp*, 3, 5  
 Regan et al. 2023, *OJAp*, 6, 12  
 Suazo et al. 2019, *ApJ*, 885, 127  
 Wise et al. 2019, *Nature*, 566, 7742

## GAS DENSITY PROJECTIONS OF SIMULATIONS AT COLLAPSE




## AVERAGE RADIAL PROFILES FROM ~50 SIMULATIONS WITH VARYING $J_{21}$



## PREDICTING POP III STELLAR MASS

- Instead of modeling the accretion on to the protostar, we stop the simulation at collapse & predict the accretion using the radial profile for infall rate to estimate a final Pop III mass.
- We combine spherically averaged inward velocities, estimate inflow rates, & assume everything is accreted until the inflow rates falls below a critical supermassive star (SMS) accretion rate.
- We find agreement within a factor of  $\sim 3$  with simulations using a sink particle approach (Suazo2019+) by using a critical SMS rate of  $6.7 \times 10^{-3} M_\odot \text{ yr}^{-1}$ .

## CONCLUSIONS



Note:  $J_{21} = 1$  is set to  $10^{-1}$  in this plot &  $J_{21}$  is  $10^{-21} \text{ erg cm}^{-2} \text{ s}^{-1} \text{ Hz}^{-1} \text{ sr}^{-1}$

- We find that the inflow rates are above our critical SMS rate for  $J_{21} \geq 10$ , if Jeans instability is not a factor, we would expect these halos to form massive BH seeds (see above figure).
- With Jeans instability, this generically results in fragments with BH seeds that are at least  $\sim 10^3 M_\odot$ . These fragments could potentially merge at later times.
- Whatever form the BH seeds take, they would be orders of magnitude more numerous than BH seeds produced by rare halos at  $J_{crit} \sim 10^4 J_{21}$ . **These simulations point to more abundant massive BH seed formation than previously suggested.**

## ACKNOWLEDGEMENTS



This work is supported in part by NSF grant AST-2009309 and NASA ATP grant 80NSSC22K0629. All numerical simulations were carried out using the NASA Pleiades Supercomputer.