

# Formale Semantik

## 04. Aussagenlogik

Roland Schäfer

Institut für Germanistische Sprachwissenschaft  
Friedrich-Schiller-Universität Jena

stets aktuelle Fassungen: <https://github.com/rsling/VL-Deutsche-Syntax>

- 1 What logic is about
  - On reasoning
  - Where we need logic
- 2 Statement calculus

- Formalization: Recursive Syntax
- Interpretation
- Laws of the PropC
- Rules of Inference
- Proof



What logic is about

- a collection of statements (propositions)

- a collection of statements (propositions)
- axioms (statements accepted to be true)

- a collection of statements (propositions)
- axioms (statements accepted to be true)
- maybe based on observations (induction)

- a collection of statements (propositions)
- axioms (statements accepted to be true)
- maybe based on observations (induction)
- statements that follow from the axioms (deduction)



- a collection of statements (propositions)
- axioms (statements accepted to be true)
- maybe based on observations (induction)
- statements that follow from the axioms (deduction)
- predictions beyond the axioms

- a collection of statements (propositions)
- axioms (statements accepted to be true)
- maybe based on observations (induction)
- statements that follow from the axioms (deduction)
- predictions beyond the axioms
- rechecking for usability: e.g., Russell's paradox

- **axioms**: atomic truths of your theory

- **axioms**: atomic truths of your theory
- **theorem**: a proposition you want to prove

- **axioms**: atomic truths of your theory
- **theorem**: a proposition you want to prove
- **lemma**: subsidiary propositions (used to prove the theorem)

- **axioms**: atomic truths of your theory
- **theorem**: a proposition you want to prove
- **lemma**: subsidiary propositions (used to prove the theorem)
- **corollary**: propositions proved while proving some axiom

# A method of reasoning

- logic does not generate truths

# A method of reasoning

- logic does not generate truths
- formalizing statements, predications etc.



# A method of reasoning

- logic does not generate truths
- formalizing statements, predications etc.
- rules of deduction from axioms to theorems

# A method of reasoning

- logic does not generate truths
- formalizing statements, predications etc.
- rules of deduction from axioms to theorems
- empirical (induction) and exact (deduction) science

# A method of reasoning

- logic does not generate truths
- formalizing statements, predications etc.
- rules of deduction from axioms to theorems
- empirical (induction) and exact (deduction) science
- aiming at an adequate model of the world (e.g., heliocentric universe)

# Why logic for semantics?

- truth-conditional

# Why logic for semantics?

- truth-conditional
- compositional behavior of propositions and connectives

# Why logic for semantics?

- truth-conditional
- compositional behavior of propositions and connectives
- a logic for entailments

# Why logic for semantics?

- truth-conditional
- compositional behavior of propositions and connectives
- a logic for entailments
- why, e.g.: *It is not the case that someone is happy.*  $\rightarrow$  *Nobody is happy.*

## Statement calculus



# Atomic formulas: statements

- statements/propositions = the atoms

# Atomic formulas: statements

- statements/propositions = the atoms
- a propositional symbol  $p$ : a well-formed formula (wff)

# Atomic formulas: statements

- statements/propositions = the **atoms**
- a propositional symbol  $p$ : a well-formed formula (**wff**)
- ex.: *Herr Keydana is a passionate cyclist.:  $k$*

# Atomic formulas: statements

- statements/propositions = the **atoms**
- a propositional symbol  $p$ : a well-formed formula (**wff**)
- ex.: *Herr Keydana is a passionate cyclist.:  $k$*
- $\llbracket k \rrbracket = 1$  or  $0$  (depending on corresponding **model**)

# Complex (molecular) formulas

- **syntax**: restricts the forms of wff's to make them interpretable

# Complex (molecular) formulas

- **syntax**: restricts the forms of wff's to make them interpretable
- define functors: functions in  $\{0, 1\}$

# Complex (molecular) formulas

- **syntax**: restricts the forms of wff's to make them interpretable
- define functors: functions in  $\{0, 1\}$
- If  $p$  and  $q$  are wff's, then

is also a wff (a **molecular term**).

# Complex (molecular) formulas

- **syntax**: restricts the forms of wff's to make them interpretable
- define functors: functions in  $\{0, 1\}$
- If  $p$  and  $q$  are wff's, then
  - ▶  $\neg p$

is also a wff (a **molecular term**).



# Complex (molecular) formulas

- **syntax**: restricts the forms of wff's to make them interpretable
- define functors: functions in  $\{0, 1\}$
- If  $p$  and  $q$  are wff's, then
  - ▶  $\neg p$
  - ▶  $p \vee q$

is also a wff (a **molecular term**).

# Complex (molecular) formulas

- **syntax**: restricts the forms of wff's to make them interpretable
- define functors: functions in  $\{0, 1\}$
- If  $p$  and  $q$  are wff's, then
  - ▶  $\neg p$
  - ▶  $p \vee q$
  - ▶  $p \wedge q$

is also a wff (a **molecular term**).

# Complex (molecular) formulas

- **syntax**: restricts the forms of wff's to make them interpretable
- define functors: functions in  $\{0, 1\}$
- If  $p$  and  $q$  are wff's, then
  - ▶  $\neg p$
  - ▶  $p \vee q$
  - ▶  $p \wedge q$
  - ▶  $p \rightarrow q$

is also a wff (a **molecular term**).

# Complex (molecular) formulas

- **syntax**: restricts the forms of wff's to make them interpretable
- define functors: functions in  $\{0, 1\}$
- If  $p$  and  $q$  are wff's, then
  - ▶  $\neg p$
  - ▶  $p \vee q$
  - ▶  $p \wedge q$
  - ▶  $p \rightarrow q$
  - ▶  $p \leftrightarrow q$

is also a wff (a **molecular term**).

# Complex (molecular) formulas

- **syntax**: restricts forms of wff's to make them interpretable
- define functors: functions in  $\{\langle 0, 1 \rangle, \langle 1, 0 \rangle, 0, 1\}$
- If  $p$  and  $q$  are wff's, then
  - ▶  $\neg p$  (negation)
  - ▶  $p \vee q$  (disjunction)
  - ▶  $p \wedge q$  (conjunction)
  - ▶  $p \rightarrow q$  (conditional)
  - ▶  $p \leftrightarrow q$  (biconditional)

is also a wff.

# Complex (molecular) formulas

- **syntax**: restricts forms of wff's to make them interpretable
- define functors: functions in  $\{\langle 0, 1 \rangle, \langle 1, 0 \rangle, 0, 1\}$
- If  $p$  and  $q$  are wff's, then
  - ▶  $\neg p$  (negation - 'not')
  - ▶  $p \vee q$  (disjunction - 'or')
  - ▶  $p \wedge q$  (conjunction - 'and')
  - ▶  $p \rightarrow q$  (conditional - 'if')
  - ▶  $p \leftrightarrow q$  (biconditional - 'iff')

is also a wff.

- standard definition:

$$\llbracket \neg \rrbracket = \begin{bmatrix} 1 \rightarrow 0 \\ 0 \rightarrow 1 \end{bmatrix}$$

# Functions and truth tables

- standard definition:

$$\llbracket \neg \rrbracket = \begin{bmatrix} 1 \rightarrow 0 \\ 0 \rightarrow 1 \end{bmatrix}$$

- but most widely used: **truth tables**

$\neg$	$p$
0	1
1	0



# Disjunction

$p$	$\vee$	$q$
1	1	1
1	1	0
0	1	1
0	0	0

# Disjunction

$p$	$\vee$	$q$
1	1	1
1	1	0
0	1	1
0	0	0

- *Herr Keydana is a passionate cyclist **or** we all love logic.*

# Disjunction

$p$	$\vee$	$q$
1	1	1
1	1	0
0	1	1
0	0	0

- *Herr Keydana is a passionate cyclist **or** we all love logic.*
- $K \vee L$

# Conjunction

$p$	$\wedge$	$q$
1	1	1
1	0	0
0	0	1
0	0	0

# Conjunction

$p$	$\wedge$	$q$
1	1	1
1	0	0
0	0	1
0	0	0

- Herr Keydana is a passionate cyclist **and** we all love logic.

# Conjunction

$p$	$\wedge$	$q$
1	1	1
1	0	0
0	0	1
0	0	0

- *Herr Keydana is a passionate cyclist **and** we all love logic.*
- $K \wedge L$

# Conditional

$p$	$\rightarrow$	$q$
1	1	1
1	0	0
0	1	1
0	1	0

# Conditional

$p$	$\rightarrow$	$q$
1	1	1
1	0	0
0	1	1
0	1	0

- *If it rains, **then** the streets get wet.*



# Conditional

$p$	$\rightarrow$	$q$
1	1	1
1	0	0
0	1	1
0	1	0

- *If it rains, **then** the streets get wet.*
- $R \rightarrow S$

# Any problems with that?

***If it rains, the streets get wet.***

- it is raining (1) , the streets are wet 1 : 1

# Any problems with that?

***If it rains, the streets get wet.***

- it is raining (1) , the streets are wet 1 : 1
- it is raining (1) , the streets are dry 0 : 0

# Any problems with that?

***If it rains, the streets get wet.***

- it is raining (1) , the streets are wet 1 : 1
- it is raining (1) , the streets are dry 0 : 0
- it is not raining (0) , the streets are wet 1 : 1

# Any problems with that?

***If it rains, the streets get wet.***

- it is raining (1) , the streets are wet 1 : 1
- it is raining (1) , the streets are dry 0 : 0
- it is not raining (0) , the streets are wet 1 : 1
- it is not raining (0) , the streets are dry 0 : 1

# Any problems with that?

***If it rains, the streets get wet.***

- it is raining (1) , the streets are wet 1 : 1
- it is raining (1) , the streets are dry 0 : 0
- it is not raining (0) , the streets are wet 1 : 1
- it is not raining (0) , the streets are dry 0 : 1
- ex vero non sequitur falsum

# Biconditional

$p$	$\leftrightarrow$	$q$
1	1	1
1	0	0
0	0	1
0	1	0

$p$	$\leftrightarrow$	$q$
1	1	1
1	0	0
0	0	1
0	1	0

- ***If and only if*** your score is above 50, ***then*** you pass the semantics exam.



# Biconditional

$p$	$\leftrightarrow$	$q$
1	1	1
1	0	0
0	0	1
0	1	0

- ***If and only if*** your score is above 50, ***then*** you pass the semantics exam.
- $S \leftrightarrow P$

- brackets are facultative

# Scope of functors

- brackets are facultative
- or set non-default functor scope

# Scope of functors

- brackets are facultative
- or set non-default functor scope
- default scope



# An example

- $p \wedge \neg q \vee r \rightarrow \neg s$

# An example

- $p \wedge \neg q \vee r \rightarrow \neg s$
- $p \wedge (\neg q) \vee r \rightarrow (\neg s)$

# An example

- $p \wedge \neg q \vee r \rightarrow \neg s$
- $p \wedge (\neg q) \vee r \rightarrow (\neg s)$
- $(p \wedge (\neg q)) \vee r \rightarrow (\neg s)$

# An example

- $p \wedge \neg q \vee r \rightarrow \neg s$
- $p \wedge (\neg q) \vee r \rightarrow (\neg s)$
- $(p \wedge (\neg q)) \vee r \rightarrow (\neg s)$
- $((p \wedge (\neg q)) \vee r) \rightarrow (\neg s)$



# An example

- $p \wedge \neg q \vee r \rightarrow \neg s$
- $p \wedge (\neg q) \vee r \rightarrow (\neg s)$
- $(p \wedge (\neg q)) \vee r \rightarrow (\neg s)$
- $((p \wedge (\neg q)) \vee r) \rightarrow (\neg s)$
- $((p \wedge (\neg q)) \vee r) \rightarrow (\neg s)$

# Large truth tables

- for  $n$  atoms in the term:  $2^n$  lines

# Large truth tables

- for  $n$  atoms in the term:  $2^n$  lines
- alternating blocks of 1's and 0's under every atom

# Large truth tables

- for  $n$  atoms in the term:  $2^n$  lines
- alternating blocks of 1's and 0's under every atom
- $2^{(m-1)}$  times '1' followed by  $2^{(m-1)}$  times '0' for the  $m$ -th atom from the right

# Large truth tables

- for  $n$  atoms in the term:  $2^n$  lines
- alternating blocks of 1's and 0's under every atom
- $2^{(m-1)}$  times '1' followed by  $2^{(m-1)}$  times '0' for the  $m$ -th atom from the right
- until  $2^n$  lines are reached

# An example

$p$	$\wedge$	$\neg$	$q$	$\vee$	$r$	$\rightarrow$	$\neg$	$s$
1			1		1			1
1			1		1			0
1			1		0			1
1			1		0			0
1			0		1			1
1			0		1			0
1			0		0			1
1			0		0			0
0			1		1			1
0			1		1			0
0			1		0			1
0			1		0			0
0			0		1			1
0			0		1			0
0			0		0			1
0			0		0			0

# An example

$p$	$\wedge$	$\neg$	$q$	$\vee$	$r$	$\rightarrow$	$\neg$	$s$
1		0	1		1		0	1
1		0	1		1		1	0
1		0	1		0		0	1
1		0	1		0		1	0
1		1	0		1		0	1
1		1	0		1		1	0
1		1	0		0		0	1
1		1	0		0		1	0
1		1	0		0		1	0
0		0	1		1		0	1
0		0	1		1		1	0
0		0	1		0		0	1
0		0	1		0		1	0
0		1	0		1		0	1
0		1	0		1		1	0
0		1	0		0		0	1
0		1	0		0		1	0

# An example

$p$	$\wedge$	$\neg$	$q$	$\vee$	$r$	$\rightarrow$	$\neg$	$s$
1	0	0	1		1		0	1
1	0	0	1		1		1	0
1	0	0	1		0		0	1
1	0	0	1		0		1	0
1	1	1	0		1		0	1
1	1	1	0		1		1	0
1	1	1	0		0		0	1
1	1	1	0		0		1	0
0	0	0	1		1		0	1
0	0	0	1		1		1	0
0	0	0	1		0		0	1
0	0	0	1		0		1	0
0	0	1	0		1		0	1
0	0	1	0		1		1	0
0	0	1	0		0		0	1
0	0	1	0		0		1	0



# An example

$p$	$\wedge$	$\neg$	$q$	$\vee$	$r$	$\rightarrow$	$\neg$	$s$
1	0	0	1	1	1		0	1
1	0	0	1	1	1		1	0
1	0	0	1	0	0		0	1
1	0	0	1	0	0		1	0
1	1	1	0	1	1		0	1
1	1	1	0	1	1		1	0
1	1	1	0	1	0		0	1
1	1	1	0	1	0		1	0
0	0	0	1	1	1		0	1
0	0	0	1	1	1		1	0
0	0	0	1	0	0		0	1
0	0	0	1	0	0		1	0
0	0	1	0	1	1		0	1
0	0	1	0	1	1		1	0
0	0	1	0	0	0		0	1
0	0	1	0	0	0		1	0

# An example

$p$	$\wedge$	$\neg$	$q$	$\vee$	$r$	$\rightarrow$	$\neg$	$s$
1	0	0	1	1	1	0	0	1
1	0	0	1	1	1	1	1	0
1	0	0	1	0	0	1	0	1
1	0	0	1	0	0	1	1	0
1	1	1	0	1	1	0	0	1
1	1	1	0	1	1	1	1	0
1	1	1	0	1	0	0	0	1
1	1	1	0	1	0	1	1	0
0	0	0	1	1	1	0	0	1
0	0	0	1	1	1	1	1	0
0	0	0	1	0	0	1	0	1
0	0	0	1	0	0	1	1	0
0	0	1	0	1	1	0	0	1
0	0	1	0	1	1	1	1	0
0	0	1	0	0	0	1	0	1
0	0	1	0	0	0	1	1	0

# An example

$p$	$\wedge$	$\neg$	$q$	$\vee$	$r$	$\rightarrow$	$\neg$	$s$
1	0	0	1	1	1	0	0	1
1	0	0	1	1	1	1	1	0
1	0	0	1	0	0	1	0	1
1	0	0	1	0	0	1	1	0
1	1	1	0	1	1	0	0	1
1	1	1	0	1	1	1	1	0
1	1	1	0	1	0	0	0	1
1	1	1	0	1	0	1	1	0
0	0	0	1	1	1	0	0	1
0	0	0	1	1	1	1	1	0
0	0	0	1	0	0	1	0	1
0	0	0	1	0	0	1	1	0
0	0	1	0	1	1	0	0	1
0	0	1	0	1	1	1	1	0
0	0	1	0	0	0	1	0	1
0	0	1	0	0	0	1	1	0

# Assignments: a contingent example

$p$	$\wedge$	$\neg$	$q$	$\vee$	$r$	$\rightarrow$	$\neg$	$s$
1	0	0	1	1	1	0	0	1
1	0	0	1	1	1	1	1	0
1	0	0	1	0	0	1	0	1
1	0	0	1	0	0	1	1	0
1	1	1	0	1	1	0	0	1
1	1	1	0	1	1	1	1	0
1	1	1	0	1	0	0	0	1
1	1	1	0	1	0	1	1	0
0	0	0	1	1	1	0	0	1
0	0	0	1	1	1	1	1	0
0	0	0	1	0	0	1	0	1
0	0	0	1	0	0	1	1	0
0	0	1	0	1	1	0	0	1
0	0	1	0	1	1	1	1	0
0	0	1	0	0	0	1	0	1
0	0	1	0	0	0	1	1	0

- take  $p \vee \neg p$

# Tautology

- take  $p \vee \neg p$

- truth-table: 

$p$	$\vee$	$\neg$	$p$
1	1	0	1
0	1	1	0

# Tautology

- take  $p \vee \neg p$

- truth-table: 

$p$	$\vee$	$\neg$	$p$
1	1	0	1
0	1	1	0

- true under every assignment, it is **valid**

- take  $p \vee \neg p$

- truth-table: 

$p$	$\vee$	$\neg$	$p$
1	1	0	1
0	1	1	0

- true under every assignment, it is **valid**
- by *law of excluded middle*: for every  $P$ ,  $P \vee \neg P$  is true



# Contradiction

- take  $p \wedge \neg p$

# Contradiction

- take  $p \wedge \neg p$

- truth-table:

$p$	$\wedge$	$\neg$	$p$
1	0	0	1
0	0	1	0

# Contradiction

- take  $p \wedge \neg p$

- truth-table: 

$p$	$\wedge$	$\neg$	$p$
1	0	0	1
0	0	1	0

- false under every assignment, called **contradictory**

- take  $p \wedge p$

- take  $p \wedge p$

- truth-table: 

$p$	$\wedge$	$p$
1	1	1
0	0	0

- take  $p \wedge p$

- truth-table: 

$p$	$\wedge$	$p$
1	1	1
0	0	0

- the truth value depends on the assignment

# What are laws?

- notice: similarities of set theory and logic

# What are laws?

- notice: similarities of set theory and logic
- non-trivial exact nature of their equivalence



# What are laws?

- notice: similarities of set theory and logic
- non-trivial exact nature of their equivalence
- laws state equivalences of (types of) wff

# What are laws?

- notice: similarities of set theory and logic
- non-trivial exact nature of their equivalence
- laws state equivalences of (types of) wff
- truth-conservative rewriting of wff's

# What are laws?

- notice: similarities of set theory and logic
- non-trivial exact nature of their equivalence
- laws state equivalences of (types of) wff
- truth-conservative rewriting of wff's
- any subformula which is a tautology (T) or contradiction (F):  
ignore by Identity Laws (Id.):

# What are laws?

- notice: similarities of set theory and logic
- non-trivial exact nature of their equivalence
- laws state equivalences of (types of) wff
- truth-conservative rewriting of wff's
- any subformula which is a tautology (T) or contradiction (F):

ignore by Identity Laws (Id.):

▶  $(P \vee F) \Leftrightarrow P, (P \vee T) \Leftrightarrow P$

# What are laws?

- notice: similarities of set theory and logic
- non-trivial exact nature of their equivalence
- laws state equivalences of (types of) wff
- truth-conservative rewriting of wff's
- any subformula which is a tautology (T) or contradiction (F):

ignore by Identity Laws (Id.):

- ▶  $(P \vee F) \Leftrightarrow P, (P \vee T) \Leftrightarrow T$
- ▶  $(P \wedge F) \Leftrightarrow F, (P \wedge T) \Leftrightarrow P$

# Equivalences: $\Leftrightarrow$

- $X \Leftrightarrow Y$ : X has the same truth-conditions as Y

# Equivalences: $\Leftrightarrow$

- $X \Leftrightarrow Y$ : X has the same truth-conditions as Y
- derivability of laws and rules (convenient redundancies)

# Equivalences: $\Leftrightarrow$

- $X \Leftrightarrow Y$ : X has the same truth-conditions as Y
- derivability of laws and rules (convenient redundancies)
- Idempotency (Idemp.):



# Equivalences: $\Leftrightarrow$

- $X \Leftrightarrow Y$ : X has the same truth-conditions as Y
- derivability of laws and rules (convenient redundancies)
- Idempotency (Idemp.):
  - ▶  $(P \vee P) \Leftrightarrow P$

# Equivalences: $\Leftrightarrow$

- $X \Leftrightarrow Y$ : X has the same truth-conditions as Y
- derivability of laws and rules (convenient redundancies)
- Idempotency (Idemp.):
  - ▶  $(P \vee P) \Leftrightarrow P$
  - ▶  $(P \wedge P) \Leftrightarrow P$

# Equivalences: $\Leftrightarrow$

- $X \Leftrightarrow Y$ : X has the same truth-conditions as Y
- derivability of laws and rules (convenient redundancies)
- Idempotency (Idemp.):
  - ▶  $(P \vee P) \Leftrightarrow P$
  - ▶  $(P \wedge P) \Leftrightarrow P$
  - ▶ *Peter walks and Peter walks.  $\Leftrightarrow$  Peter walks.*

- Associative Laws for  $\vee$  and  $\wedge$  (Assoc.):

- Associative Laws for  $\vee$  and  $\wedge$  (Assoc.):
  - ▶  $((P \vee Q) \vee R) \Leftrightarrow (P \vee (Q \vee R))$

- Associative Laws for  $\vee$  and  $\wedge$  (Assoc.):
  - ▶  $((P \vee Q) \vee R) \Leftrightarrow (P \vee (Q \vee R))$
  - ▶  $((\text{He walks or she talks}) \text{ or we walk.}) \Leftrightarrow (\text{He walks or } (\text{she talks or we walk.}))$

- Associative Laws for  $\vee$  and  $\wedge$  (Assoc.):
  - ▶  $((P \vee Q) \vee R) \Leftrightarrow (P \vee (Q \vee R))$
  - ▶  $((\text{He walks or she talks}) \text{ or we walk.}) \Leftrightarrow (\text{He walks or (she talks or we walk.)})$
- Commutative Laws for  $\vee$  and  $\wedge$  (Comm.):

- Associative Laws for  $\vee$  and  $\wedge$  (Assoc.):
  - ▶  $((P \vee Q) \vee R) \Leftrightarrow (P \vee (Q \vee R))$
  - ▶  $((\text{He walks or she talks}) \text{ or we walk.}) \Leftrightarrow (\text{He walks or (she talks or we walk.)})$
- Commutative Laws for  $\vee$  and  $\wedge$  (Comm.):
  - ▶  $(P \vee Q) \Leftrightarrow (Q \vee P)$



- Associative Laws for  $\vee$  and  $\wedge$  (Assoc.):
  - ▶  $((P \vee Q) \vee R) \Leftrightarrow (P \vee (Q \vee R))$
  - ▶  $((\text{He walks or she talks}) \text{ or we walk.}) \Leftrightarrow (\text{He walks or (she talks or we walk.)})$
- Commutative Laws for  $\vee$  and  $\wedge$  (Comm.):
  - ▶  $(P \vee Q) \Leftrightarrow (Q \vee P)$
  - ▶  $\text{Peter walks or Sue snores.} \Leftrightarrow \text{Sue snores or Peter walks.}$

- Associative Laws for  $\vee$  and  $\wedge$  (Assoc.):
  - ▶  $((P \vee Q) \vee R) \Leftrightarrow (P \vee (Q \vee R))$
  - ▶  $((\text{He walks or she talks}) \text{ or we walk.}) \Leftrightarrow (\text{He walks or (she talks or we walk.)})$
- Commutative Laws for  $\vee$  and  $\wedge$  (Comm.):
  - ▶  $(P \vee Q) \Leftrightarrow (Q \vee P)$
  - ▶  $\text{Peter walks or Sue snores.} \Leftrightarrow \text{Sue snores or Peter walks.}$
- Distributive Laws for  $\vee \wedge$  and  $\wedge \vee$  (Distr.):

- Associative Laws for  $\vee$  and  $\wedge$  (Assoc.):
  - ▶  $((P \vee Q) \vee R) \Leftrightarrow (P \vee (Q \vee R))$
  - ▶  $((\text{He walks or she talks}) \text{ or we walk.}) \Leftrightarrow (\text{He walks or (she talks or we walk.)})$
- Commutative Laws for  $\vee$  and  $\wedge$  (Comm.):
  - ▶  $(P \vee Q) \Leftrightarrow (Q \vee P)$
  - ▶  $\text{Peter walks or Sue snores.} \Leftrightarrow \text{Sue snores or Peter walks.}$
- Distributive Laws for  $\vee \wedge$  and  $\wedge \vee$  (Distr.):
  - ▶  $(P \vee (Q \wedge R)) \Leftrightarrow ((P \vee Q) \wedge (P \vee R))$

- **Associative Laws for  $\vee$  and  $\wedge$  (Assoc.):**
  - ▶  $((P \vee Q) \vee R) \Leftrightarrow (P \vee (Q \vee R))$
  - ▶  $((\text{He walks or she talks}) \text{ or we walk.}) \Leftrightarrow (\text{He walks or (she talks or we walk.)})$
- **Commutative Laws for  $\vee$  and  $\wedge$  (Comm.):**
  - ▶  $(P \vee Q) \Leftrightarrow (Q \vee P)$
  - ▶  $\text{Peter walks or Sue snores.} \Leftrightarrow \text{Sue snores or Peter walks.}$
- **Distributive Laws for  $\vee \wedge$  and  $\wedge \vee$  (Distr.):**
  - ▶  $(P \vee (Q \wedge R)) \Leftrightarrow ((P \vee Q) \wedge (P \vee R))$
  - ▶  $(\text{Sue snores}) \text{ and } (\text{Peter walks or we talk.}) \Leftrightarrow (\text{Sue snores and Peter walks}) \text{ or } (\text{Sue snores and we talk.})$

# Laws dealing with tautology and contradiction

- Complement Laws:

# Laws dealing with tautology and contradiction

- Complement Laws:
  - ▶ Tautology (T):  $(P \vee \neg P) \Leftrightarrow \mathbf{T}$

# Laws dealing with tautology and contradiction

- Complement Laws:

- ▶ Tautology (T):  $(P \vee \neg P) \Leftrightarrow \mathbf{T}$
- ▶ Contradiction (F):  $(P \wedge \neg P) \Leftrightarrow \mathbf{F}$

# Laws dealing with tautology and contradiction

- Complement Laws:

- ▶ Tautology (T):  $(P \vee \neg P) \Leftrightarrow \mathbf{T}$
- ▶ Contradiction (F):  $(P \wedge \neg P) \Leftrightarrow \mathbf{F}$
- ▶ Double Negation (DN):  $(\neg\neg P) \Leftrightarrow P$



# Laws dealing with tautology and contradiction

- Complement Laws:

- ▶ Tautology (T):  $(P \vee \neg P) \Leftrightarrow \mathbf{T}$
- ▶ Contradiction (F):  $(P \wedge \neg P) \Leftrightarrow \mathbf{F}$
- ▶ Double Negation (DN):  $(\neg \neg P) \Leftrightarrow P$
- ▶ *It is not the case that Sandy is not walking.*  
 $\Leftrightarrow$  *Sandy is walking.*

- **Implication** (Impl.):

$P$	$\rightarrow$	$Q$	$\Leftrightarrow$	$\neg$	$P$	$\vee$	$Q$
1	1	1		0	1	1	1
1	0	0		0	1	0	0
0	1	1		1	0	1	1
0	1	0		1	0	1	0

- **Implication (Impl.):**

$P$	$\rightarrow$	$Q$	$\Leftrightarrow$	$\neg$	$P$	$\vee$	$Q$
1	1	1		0	1	1	1
1	0	0		0	1	0	0
0	1	1		1	0	1	1
0	1	0		1	0	1	0

- **Contraposition (Contr.):**

$P$	$\rightarrow$	$Q$	$\Leftrightarrow$	$\neg$	$Q$	$\rightarrow$	$\neg$	$P$
1	1	1		0	1	1	0	1
1	0	0		1	0	0	0	1
0	1	1		0	1	1	1	0
0	1	0		1	0	1	1	0

- DeMorgan's Laws:

- DeMorgan's Laws:

- ▶  $\neg(P \vee Q) \Leftrightarrow (\neg P \wedge \neg Q)$

- DeMorgan's Laws:

- ▶  $\neg(P \vee Q) \Leftrightarrow (\neg P \wedge \neg Q)$
- ▶ alternatively:  $\overline{P \vee Q} \Leftrightarrow \overline{P} \wedge \overline{Q}$

# DeMorgan (DeM)

- DeMorgan's Laws:

- ▶  $\neg(P \vee Q) \Leftrightarrow (\neg P \wedge \neg Q)$
- ▶ alternatively:  $\overline{P \vee Q} \Leftrightarrow \overline{P} \wedge \overline{Q}$
- ▶  $\neg(P \wedge Q) \Leftrightarrow (\neg P \vee \neg Q)$

- DeMorgan's Laws:

- ▶  $\neg(P \vee Q) \Leftrightarrow (\neg P \wedge \neg Q)$
- ▶ alternatively:  $\overline{P \vee Q} \Leftrightarrow \overline{P} \wedge \overline{Q}$
- ▶  $\neg(P \wedge Q) \Leftrightarrow (\neg P \vee \neg Q)$
- ▶ consequently:  $\overline{\overline{P} \vee \overline{Q}} \Leftrightarrow \overline{\overline{P}} \wedge \overline{\overline{Q}} \Leftrightarrow P \wedge Q$



# The Modus Ponens (MP)

- Definition:

$P \rightarrow Q$	premise 1
$P$	premise 2
$Q$	conclusion

# The Modus Ponens (MP)

- Definition:

$P \rightarrow Q$	premise 1
$P$	premise 2
$Q$	conclusion

- or:  $(P \rightarrow Q) \wedge (P) \rightarrow (Q)$

# The Modus Ponens (MP)

- Definition:

$P \rightarrow Q$	premise 1
$P$	premise 2
$Q$	conclusion

- or:  $(P \rightarrow Q) \wedge (P) \rightarrow (Q)$
- (1) *If It rains, the streets get wet.* (2) *It is raining.*  
 $\rightarrow$  *The streets are getting wet.*

# MP: a truth table illustration

- Premises are always set to be true!

# MP: a truth table illustration

- Premises are always set to be true!
- the table:

$P$	$\rightarrow$	$Q$
1	1	1
1	0	0
0	1	1
0	1	0

# MP: a truth table illustration

- The conditional must be true.
- cancel the 'false' row

$P$	$\rightarrow$	$Q$
1	1	1
1	0	0
0	1	1
0	1	0

# MP: a truth table illustration

- $P$  must be true.
- cancel the 'false' rows,  $Q$  can only be true:

$P$	$\rightarrow$	$Q$
1	1	1
1	0	0
0	1	1
0	1	0

# The Modus Tollens (MT)

- Definition:

$P$	$\rightarrow$	$Q$
		$\neg Q$
$\neg P$		



# The Modus Tollens (MT)

- Definition:

$P$	$\rightarrow$	$Q$
		$\neg Q$
$\neg P$		

- the table illustration:

$P$	$\rightarrow$	$Q$	
1	1	1	(by premise 2)
1	0	0	(by premise 1)
0	1	1	(by premise 2)
0	1	0	

- Hypothetical Syllogism (HS):
  - ▶  $((P \rightarrow Q) \wedge (Q \rightarrow R)) \rightarrow (P \rightarrow R)$
  - ▶ (1) *If it rains, the streets get wet.* (2) *If the streets get wet, it smells nice.*  $\rightarrow$  *If it rains, it smells nice.*

- Hypothetical Syllogism (HS):

- ▶  $((P \rightarrow Q) \wedge (Q \rightarrow R)) \rightarrow (P \rightarrow R)$
- ▶ (1) *If it rains, the streets get wet.* (2) *If the streets get wet, it smells nice.*  $\rightarrow$  *If it rains, it smells nice.*

- Disjunctive Syllogism (DS):

- ▶  $((P \vee Q) \wedge (\neg P)) \rightarrow (Q)$
- ▶ (1) *Either Peter sleeps or Peter is awake.* (2) *Peter isn't awake.*  $\rightarrow$  *Peter sleeps.*

- Simplification (Simp.):
  - ▶  $(P \wedge Q) \rightarrow P$
  - ▶ (1) *It is raining and the sun is shining.*  $\rightarrow$  *It is raining.*

- Simplification (Simp.):
  - ▶  $(P \wedge Q) \rightarrow P$
  - ▶ (1) *It is raining and the sun is shining.*  $\rightarrow$  *It is raining.*
- Conjunction (Conj.):
  - ▶  $(P) \wedge (Q) \rightarrow (P \wedge Q)$
  - ▶ (1) *It is raining.* (2) *The sun is shining.*  $\rightarrow$  *It is raining and the sun is shining.*

- **Simplification (Simp.):**
  - ▶  $(P \wedge Q) \rightarrow P$
  - ▶ (1) *It is raining and the sun is shining.*  $\rightarrow$  *It is raining.*
- **Conjunction (Conj.):**
  - ▶  $(P) \wedge (Q) \rightarrow (P \wedge Q)$
  - ▶ (1) *It is raining.* (2) *The sun is shining.*  $\rightarrow$  *It is raining and the sun is shining.*
- **Addition (Add.):**
  - ▶  $(P) \rightarrow (P \vee Q)$
  - ▶ (1) *It is raining.*  $\rightarrow$  *It is raining or the sun is shining.*
  - ▶ What if Q is instantiated as true or false by another premise?

# A sample proof

- Prove  $p \vee q$  from  $(p \vee q) \rightarrow \neg(r \wedge \neg s)$  and  $r \wedge \neg s$

# A sample proof

- Prove  $p \vee q$  from  $(p \vee q) \rightarrow \neg(r \wedge \neg s)$  and  $r \wedge \neg s$
- The proof:

		$p \vee q$
1	$(p \vee q) \rightarrow \neg(r \wedge \neg s)$	
2	$r \wedge \neg s$	
<hr/>		
	$p \vee q$	1,2,MT





## Kontakt

Prof. Dr. Roland Schäfer  
Institut für Germanistische Sprachwissenschaft  
Friedrich-Schiller-Universität Jena  
Fürstengraben 30  
07743 Jena

<https://rolandschaefer.net>  
[roland.schaefer@uni-jena.de](mailto:roland.schaefer@uni-jena.de)

## Creative Commons BY-SA-3.0-DE

Dieses Werk ist unter einer Creative Commons Lizenz vom Typ *Namensnennung - Weitergabe unter gleichen Bedingungen 3.0 Deutschland* zugänglich. Um eine Kopie dieser Lizenz einzusehen, konsultieren Sie

<http://creativecommons.org/licenses/by-sa/3.0/de/> oder wenden Sie sich brieflich an Creative Commons, Postfach 1866, Mountain View, California, 94042, USA.