1	A search for sparticles in zero lepton final states
2	Russell W. Smith

3	Submitted in partial fulfillment of the
1	requirements for the degree of
5	Doctor of Philosophy
ñ	in the Graduate School of Arts and Sciences

7 COLUMBIA UNIVERSITY 8 2016

9

10

11

© 2016

Russell W. Smith

All rights reserved

12	ABSTRACT
13	A search for sparticles in zero lepton final states
14	Russell W. Smith
15	TODO: Here's where your abstract will eventually go. The above text is all in the
16	center, but the abstract itself should be written as a regular paragraph on the page
17	and it should not have indentation. Just replace this text.

# 18 Contents

19	Contents			i
20	1	1		
21	2	The	e Standard Model	5
22		2.1	Quantum Field Theory	5
23		2.2	Symmetries	5
24		2.3	The Standard Model	5
25		2.4	Electroweak Symmetry breaking	6
26		2.5	Deficiencies of the Standard Model	6
27	3	Sup	persymmetry	7
28		3.1	Motivation	7
29		3.2	Supersymmetry	7
30		3.3	Additional particle content	7
31		3.4	Phenomenology	7
32	4	The	e Large Hadron Collider	9
33		4.1	Magnets	9
34	5	The	e ATLAS detector	11
35		5.1	Inner Detector	11
36		5.2	Calorimeter	13

37		5.3	Muon Spectrometer	13
38	6	The	Recursive Jigsaw Technique	15
39		6.1	Razor variables	15
40		6.2	SuperRazor variables	15
41		6.3	The Recursive Jigsaw Technique	15
42		6.4	Variables used in the search for zero lepton SUSY	15
43	7	Tab	le of Contents Title	17
44	8	A so	earch for supersymmetric particles in zero lepton final states	
45		with	n the Recursive Jigsaw Technique	19
46		8.1	Object reconstruction	19
47		8.2	Signal regions	20
48		8.3	Background estimation	20
49	9	Res	ults	21
50		9.1	Statistical Analysis	21
51		9.2	Signal Region distributions	21
52		9.3	Pull Plots	21
53		9.4	Systematic Uncertainties	21
54		9.5	Exclusion plots	21
55	Co	onclu	sion	23
56		9.6	New Section	23
57	Bi	ibliog	graphy	25

# Acknowledgements

Dedication

61

60

#### Introduction

Particle physics is a remarkably successful field of scientific inquiry. The ability to precisely predict the properties of a exceedingly wide range of physical phenomena, such as the description of the cosmic microwave background (cite planck) anomalous magnetic moment of the muon (cite paper on this), and the measurement of the number of weakly-interacting neutrino flavors is truly amazing.

The theory that has allowed this range of predictions is the Standard Model of particle physics (SM). The Standard Model combines the electroweak theory of Glashow, Weinberg, and Salam [Weinberg:1967tq, 1, 2] with the theory of the strong interactions, as first envisioned by Gell-Mann and Zweig [GellMann:1964nj,

Zweig:1964jf]. This quantum field theory (QFT) contains a tiny number of particles,
 whose interactions describe phenomena up to at least the TeV scale. These particles

are manifestations of the fields of the Standard Model, after application of the Higgs

Mechanism. The particle content of the SM consists only of the six quarks, six leptons,

75 the four gauge bosons, and the scalar Higgs boson.

Despite its impressive range of described phenomena, the Standard Model has some theoretical and experimental deficiencies. The SM contains 26 free parameters <sup>1</sup>. It would be more theoretically pleasing to understand these free parameters in terms of a more fundamental theory. The major theoretical concern of the Standard Model, as it pertains to this thesis, is the "hierarchy problem" [Weinberg:1979bn, 3–6]. The light mass of the Higgs boson (125 GeV) should be quadratically dependent on the scale of UV physics, due to the quantum corrections from high-energy physics

processes. The most perplexing experimental issue is the existence of "dark matter", as demonstrated by galactic rotation curves [darkMatterPrimer, 7–12]. From cosmological data, it has been shown that there exists additional matter which has not yet been seen interacting with the particles of the Standard Model. There is no particle in the SM which can act as a candidate for dark matter.

Both of these major issues, as well as numerous others, can be solved 88 by the introduction of "supersymmetry" [Gervais:1971xj, Golfand:1971iw, 89 Volkov:1973ix, Ferrara:1974ac, 6, 13–19]. In supersymmetric theories, each 90 SM particles has a so-called "superpartner", or sparticle partner, differing from given 91 SM particle by 1/2 in spin. These theories solve the hierarchy problem, since the 92 quantum corrections induced from the superpartners exactly cancel those induced by the SM particles. In addition, these theories are usually constructed assuming R-parity, which can be thought of as the "charge" of supersymmetry, with SM particles having R = 1 and sparticles having R = -1. In collider experiments, since 96 the incoming SM particles have total R = 1, the resulting sparticles are produced 97 in pairs. This produces a rich phenomenology, which is often characterized by large 98 missing transverse energy  $(E_T^{\text{miss}})$ , which provides significant discrimination against 99 SM backgrounds [20]. 100

Despite the power of searches for supersymmetry where  $E_{\rm T}^{\rm miss}$  is a primary discriminating variable, there has been significant interest in the use of other variables to discriminate against SM backgrounds. These include searches employing variables such as  $\alpha T$ ,  $M_{T,2}$ , and the razor variables  $(M_R, R^2)$  [SUSY-2014-06, ATLAS-CONF-2016-009, 21–31]. In this thesis, we will present the first search for supersymmetry using the novel Recursive Jigsaw Reconstruction (RJR) technique.

 $<sup>^1{\</sup>rm This}$  is the Standard Model corrected to include neutrino masses. These parameters are the fermion masses (6 leptons, 6 quarks), CKM and PMNS mixing angles (8 angles, 2 CP-violating phases), W/Z/Higgs masses (3) , the Higgs field expectation value, and the couplings of the strong, weak, and electromagnetic forces (3  $\alpha_{force}$ ) .

RJR can be considered the conceptual successor of the razor variables. We impose a particular final state "decay tree" on an event, which roughly corresponds to a simplified Feynmann diagram. This allows an understanding of internal decay structure of an event, as well as additional rejection of SM backgrounds.

This thesis details a search for the superpartners of the gluons and quarks, the 111 gluinos and squarks, in final states with zero leptons, with of data using the AT-r 112 LAS detector. This thesis is organized as follows. The theoretical motivation of the 113 Standard Model and supersymmetry are described in Chapters 2 and 3. The Large 114 Hadron Collider and the ATLAS detector are presented in Chapters 4 and 5. Chap-115 ter 5 provides a detailed description of Recursive Jigsaw Reconstruction, as well as 116 a description of the variables used for the particular search presented in this thesis. 117 Chapter 6 presents the details of the analysis, including the dataset, object recon-118 struction, and selections used by the analysis. In Chapter 7, the final results are 119 presented; since there is no evidence of a supersymmetric signal in the analysis, we 120 present the final exclusion curves in simplified supersymmetric models. 121

122

123

#### The Standard Model

- Here you can write some introductory remarks about your chapter. I like to give each sentence its own line.
- When you need a new paragraph, just skip an extra line.

### 2.1 Quantum Field Theory

### 128 2.2 Symmetries

### 129 2.3 The Standard Model

#### 130 Overview

- 131 By using the asterisk to start a new section, I keep the section from appearing in the
- table of contents. If you want your sections to be numbered and to appear in the
- table of contents, remove the asterisk.

#### 134 Fermions

- By using the asterisk to start a new section, I keep the section from appearing in the
- table of contents. If you want your sections to be numbered and to appear in the
- table of contents, remove the asterisk.

#### 138 Bosons

By using the asterisk to start a new section, I keep the section from appearing in the

table of contents. If you want your sections to be numbered and to appear in the

table of contents, remove the asterisk.

### 142 2.4 Electroweak Symmetry breaking

143 By using the asterisk to start a new section, I keep the section from appearing in the

table of contents. If you want your sections to be numbered and to appear in the

table of contents, remove the asterisk.

### 146 2.5 Deficiencies of the Standard Model

By using the asterisk to start a new section, I keep the section from appearing in the

table of contents. If you want your sections to be numbered and to appear in the

table of contents, remove the asterisk.

Char	oter	3

150

151

## Supersymmetry

- 152 Here you can write some introductory remarks about your chapter. I like to give each
- 153 sentence its own line.
- When you need a new paragraph, just skip an extra line.

### 3.1 Motivation

- 156 Only Additional allowed Lorentz invariant symmetry
- 157 Dark Matter
- 158 Cancellation of quadratic divergences in corrections to the
- 159 Higgs Mass
- 160 3.2 Supersymmetry
- 161 3.3 Additional particle content
- 162 3.4 Phenomenology
- 163 R parity Consequences for sq/gl decays

164

165

167

## The Large Hadron Collider

- Here you can write some introductory remarks about your chapter. I like to give each sentence its own line.
- When you need a new paragraph, just skip an extra line. 168

#### Magnets 4.1


- By using the asterisk to start a new section, I keep the section from appearing in the 170
- table of contents. If you want your sections to be numbered and to appear in the 171
- table of contents, remove the asterisk. 172

174

### The ATLAS detector

Here you can write some introductory remarks about your chapter. I like to give each sentence its own line.

177 When you need a new paragraph, just skip an extra line.




178

179

### 5.1 Inner Detector

By using the asterisk to start a new section, I keep the section from appearing in the table of contents. If you want your sections to be numbered and to appear in the table of contents, remove the asterisk.


### 184 Pixel Detector



186

#### 187 Insertable B-Layer


188 Qualification task, so add a bit more.



190


189

## Semiconductor Tracker



192

### 194 Transition Radiation Tracker



196

## <sup>197</sup> 5.2 Calorimeter




198 199

200 Electromagnetic Calorimeter

201 Hadronic Calorimeter

## 202 5.3 Muon Spectrometer



203

205

206

### The Recursive Jigsaw Technique

- Here you can write some introductory remarks about your chapter. I like to give each sentence its own line.
- When you need a new paragraph, just skip an extra line.

### 210 6.1 Razor variables

- 211 By using the asterisk to start a new section, I keep the section from appearing in the
- 212 table of contents. If you want your sections to be numbered and to appear in the
- 213 table of contents, remove the asterisk.

### 214 6.2 SuperRazor variables

- 215 6.3 The Recursive Jigsaw Technique
- <sup>216</sup> 6.4 Variables used in the search for zero lepton
- SUSY

218	Chapter 7			
219	Title of Chapter 1			

220

221

Title of Chapter 1

- 222 Here you can write some introductory remarks about your chapter. I like to give each
- 223 sentence its own line.
- When you need a new paragraph, just skip an extra line.

## 225 8.1 Object reconstruction

226 Photons, Muons, and Electrons

227 Jets

### 228 Missing transverse momentum

229 Probably longer, show some plots from the PUB note that we worked on

# 230 8.2 Signal regions

- Gluino signal regions
- 232 Squark signal regions
- 233 Compressed signal regions

## 8.3 Background estimation

- 235 **Z vv**
- 236 **W** ev
- 237 ttbar

238

239

## Title of Chapter 1

- 240 Here you can write some introductory remarks about your chapter. I like to give each
- 241 sentence its own line.
- 242 When you need a new paragraph, just skip an extra line.

## 243 9.1 Statistical Analysis

244 maybe to be moved to an appendix

## 9.2 Signal Region distributions

- 9.3 Pull Plots
- 9.4 Systematic Uncertainties
- 248 9.5 Exclusion plots

### Conclusion

- 250 Here you can write some introductory remarks about your chapter. I like to give each
- 251 sentence its own line.

249

252 When you need a new paragraph, just skip an extra line.

## 9.6 New Section

- 254 By using the asterisk to start a new section, I keep the section from appearing in the
- 255 table of contents. If you want your sections to be numbered and to appear in the
- table of contents, remove the asterisk.

### **Bibliography**

- 258 [1] S. L. Glashow, Partial Symmetries of Weak Interactions, Nucl. Phys. **22** (1961) p. 579.
- 260 [2] A. Salam, Weak and Electromagnetic Interactions, 261 Conf. Proc. **C680519** (1968) p. 367.

- 262 [3] S. Weinberg, Implications of Dynamical Symmetry Breaking, 263 Phys. Rev. **D13** (1976) p. 974.
- 264 [4] E. Gildener, Gauge Symmetry Hierarchies, Phys. Rev. **D14** (1976) p. 1667.
- [5] L. Susskind,
   Dynamics of Spontaneous Symmetry Breaking in the Weinberg-Salam Theory,
   Phys. Rev. **D20** (1979) p. 2619.
- 268 [6] S. P. Martin, "A Supersymmetry Primer," 1997, eprint: arXiv:hep-ph/9709356.
- 270 [7] V. C. Rubin and W. K. Ford Jr., Rotation of the Andromeda Nebula from a 271 Spectroscopic Survey of Emission Regions, Astrophys. J. **159** (1970) p. 379.
- 272 [8] M. S. Roberts and R. N. Whitehurst, 273 "The rotation curve and geometry of M31 at large galactocentric distances, 274 Astrophys. J. **201** (1970) p. 327.
- [9] V. C. Rubin, N. Thonnard, and W. K. Ford Jr.,
   Rotational properties of 21 SC galaxies with a large range of luminosities and
   radii, from NGC 4605 /R = 4kpc/ to UGC 2885 /R = 122 kpc/,
   Astrophys. J. 238 (1980) p. 471.
- V. C. Rubin et al., Rotation velocities of 16 SA galaxies and a comparison of Sa, Sb, and SC rotation properties, Astrophys. J. **289** (1985) p. 81.
- 281 [11] A. Bosma, 282 21-cm line studies of spiral galaxies. 2. The distribution and kinematics of

- neutral hydrogen in spiral galaxies of various morphological types., 283
- Astron. J. **86** (1981) p. 1825. 284

- M. Persic, P. Salucci, and F. Stel, The Universal rotation curve of spiral 285 galaxies: 1. The Dark matter connection,
- Mon. Not. Roy. Astron. Soc. **281** (1996) p. 27, 287
- arXiv: astro-ph/9506004 [astro-ph]. 288
- H. Miyazawa, Baryon Number Changing Currents, |13|289 Prog. Theor. Phys. **36** (1966) p. 1266. 290
- J.-L. Gervais and B. Sakita, 291
- Field Theory Interpretation of Supergauges in Dual Models, 292
- Nucl. Phys. **B34** (1971) p. 632. 293
- [15]A. Neveu and J. H. Schwarz, Factorizable dual model of pions, 294 Nucl. Phys. **B31** (1971) p. 86. 295
- A. Neveu and J. H. Schwarz, Quark Model of Dual Pions, 296 Phys. Rev. **D4** (1971) p. 1109. 297
- J. Wess and B. Zumino, 298
- A Lagrangian Model Invariant Under Supergauge Transformations, 299
- Phys. Lett. **B49** (1974) p. 52. 300
- A. Salam and J. A. Strathdee, Supersymmetry and Nonabelian Gauges, [18]301 Phys. Lett. **B51** (1974) p. 353. 302
- J. Wess and B. Zumino, Supergauge Transformations in Four-Dimensions, [19]303 Nucl. Phys. **B70** (1974) p. 39. 304
- G. R. Farrar and P. Fayet, Phenomenology of the Production, Decay, and [20]305 Detection of New Hadronic States Associated with Supersymmetry, 306 Phys. Lett. **B76** (1978) p. 575. 307
- CMS Collaboration, 308
- Search for supersymmetry with razor variables in pp collisions at  $\sqrt{s} = 7$  TeV, 309
- Phys. Rev. D **90** (2014) p. 112001, arXiv: 1405.3961 [hep-ex]. 310
- CMS Collaboration, 311
- Search for supersymmetry with photons in pp collisions at  $\sqrt{s} = 8$  TeV, 312
- Phys. Rev. D **92** (2015) p. 072006, arXiv: 1507.02898 [hep-ex]. 313

- CMS Collaboration, Inclusive search for supersymmetry using razor variables in pp collisions at  $\sqrt{s}=7$  TeV, Phys. Rev. Lett. **111** (2013) p. 081802, arXiv: 1212.6961 [hep-ex].
- CMS Collaboration, Search for Supersymmetry in pp Collisions at 7 TeV in Events with Jets and Missing Transverse Energy, Phys. Lett. B **698** (2011) p. 196, arXiv: 1101.1628 [hep-ex].
- 320 [25] CMS Collaboration, Search for Supersymmetry at the LHC in Events with
  321 Jets and Missing Transverse Energy, Phys. Rev. Lett. **107** (2011) p. 221804,
  322 arXiv: 1109.2352 [hep-ex].
- 232 [26] CMS Collaboration, Search for supersymmetry in hadronic final states using  $M_{T2}$  in pp collisions at  $\sqrt{s} = 7$  TeV, JHEP 10 (2012) p. 018, arXiv: 1207.1798 [hep-ex].
- CMS Collaboration, Searches for supersymmetry using the  $M_{T2}$  variable in hadronic events produced in pp collisions at 8 TeV, JHEP **05** (2015) p. 078, arXiv: 1502.04358 [hep-ex].
- CMS Collaboration, Search for new physics with the  $M_{T2}$  variable in all-jets final states produced in pp collisions at  $\sqrt{s} = 13$  TeV (2016), arXiv: 1603.04053 [hep-ex].
- 332 [29] ATLAS Collaboration, Multi-channel search for squarks and gluinos in  $\sqrt{s} = 7$  TeV pp collisions with the ATLAS detector at the LHC, 334 Eur. Phys. J. C **73** (2013) p. 2362, arXiv: 1212.6149 [hep-ex].
- 335 [30] ATLAS Collaboration, 336 Search for the electroweak production of supersymmetric particles in 337  $\sqrt{s} = 8 \ TeV \ pp \ collisions \ with \ the \ ATLAS \ detector$ , 338 Phys. Rev. D **93** (2016) p. 052002, arXiv: 1509.07152 [hep-ex].
- 339 [31] ATLAS Collaboration, ATLAS Run 1 searches for direct pair production of 340 third-generation squarks at the Large Hadron Collider, 341 Eur. Phys. J. C **75** (2015) p. 510, arXiv: 1506.08616 [hep-ex].