1	A search for sparticles in zero lepton final states
2	Russell W. Smith

3	Submitted in partial fulfillment of the
4	requirements for the degree of
5	Doctor of Philosophy
6	in the Graduate School of Arts and Sciences

7 COLUMBIA UNIVERSITY

8 2016

9

10

11

© 2016

Russell W. Smith

All rights reserved

12	ABSTRACT
13	A search for sparticles in zero lepton final states
14	Russell W. Smith
15	TODO : Here's where your abstract will eventually go. The above text is all in the
16	center, but the abstract itself should be written as a regular paragraph on the page
17	and it should not have indentation. Just replace this text.

# 18 Contents

19	Contents			
20	1	Intr	1	
21	2	The	e Standard Model	5
22		2.1	Quantum Field Theory	5
23		2.2	Symmetries	5
24		2.3	The Standard Model	5
25		2.4	Electroweak Symmetry breaking	6
26		2.5	Deficiencies of the Standard Model	6
27	3	Sup	persymmetry	7
28		3.1	Motivation	7
29		3.2	Supersymmetry	7
30		3.3	Additional particle content	7
31		3.4	Phenomenology	7
32	4	The	e Large Hadron Collider	9
33		4.1	Magnets	9
34	5	The	e ATLAS detector	11
35		5.1	Inner Detector	11
36		5.2	Calorimeter	13

37		5.3	Muon Spectrometer	13
38	6	The	e Recursive Jigsaw Technique	15
39		6.1	Razor variables	15
40		6.2	SuperRazor variables	15
41		6.3	The Recursive Jigsaw Technique	15
42		6.4	Variables used in the search for zero lepton SUSY	15
43	7	Tab	le of Contents Title	17
44	8	A s	earch for supersymmetric particles in zero lepton final states	
45		witl	h the Recursive Jigsaw Technique	19
46		8.1	Object reconstruction	19
47		8.2	Signal regions	20
48		8.3	Background estimation	20
49	9	Res	ults	21
50		9.1	Statistical Analysis	21
51		9.2	Signal Region distributions	21
52		9.3	Pull Plots	21
53		9.4	Systematic Uncertainties	21
54		9.5	Exclusion plots	21
55	Co	onclu	ısion	23
56		9.6	New Section	23
57	Bi	iblios	vraphy	25

# Acknowledgements

58

Dedication

61

60

#### Introduction

Particle physics is a remarkably successful field of scientific inquiry. The ability to precisely predict the properties of a exceedingly wide range of physical phenomena, such as the description of the cosmic microwave background [1, 2], the understanding of the anomalous magnetic dipole moment of the electron [3, 4], and the measurement of the number of weakly-interacting neutrino flavors [5] is truly amazing.

The theory that has allowed this range of predictions is the Standard Model 67 of particle physics (SM). The Standard Model combines the electroweak theory of Glashow, Weinberg, and Salam [6–8] with the theory of the strong interactions, as 69 first envisioned by Gell-Mann and Zweig [9, 10]. This quantum field theory (QFT) 70 contains a tiny number of particles, whose interactions describe phenomena up to at 71 least the TeV scale. These particles are manifestations of the fields of the Standard 72 Model, after application of the Higgs Mechanism. The particle content of the SM 73 consists only of the six quarks, six leptons, the four gauge bosons, and the scalar 74 Higgs boson. 75

Despite its impressive range of described phenomena, the Standard Model has some theoretical and experimental deficiencies. The SM contains 26 free parameters <sup>1</sup>. It would be more theoretically pleasing to understand these free parameters in terms of a more fundamental theory. The major theoretical concern of the Standard Model, as it pertains to this thesis, is the "hierarchy problem" [11–15]. The light mass of the Higgs boson (125 GeV) should be quadratically dependent on the scale of UV physics, due to the quantum corrections from high-energy physics processes. The most

perplexing experimental issue is the existence of "dark matter", as demonstrated by galactic rotation curves [16–22]. From cosmological data, it has been shown that there exists additional matter which has not yet been seen interacting with the particles of the Standard Model. There is no particle in the SM which can act as a candidate for dark matter.

Both of these major issues, as well as numerous others, can be solved by the 88 introduction of "supersymmetry" [15, 23-33]. In supersymmetric theories, each SM 89 particles has a so-called "superpartner", or sparticle partner, differing from given 90 SM particle by 1/2 in spin. These theories solve the hierarchy problem, since the 91 quantum corrections induced from the superpartners exactly cancel those induced 92 by the SM particles. In addition, these theories are usually constructed assuming 93 R-parity, which can be thought of as the "charge" of supersymmetry, with SM particles having R=1 and sparticles having R=-1. In collider experiments, since the incoming SM particles have total R=1, the resulting sparticles are produced 96 in pairs. This produces a rich phenomenology, which is often characterized by large 97 missing transverse energy  $(E_{\rm T}^{\rm miss})$ , which provides significant discrimination against 98 SM backgrounds [34]. 99

Despite the power of searches for supersymmetry where  $E_{\rm T}^{\rm miss}$  is a primary discriminating variable, there has been significant interest in the use of other variables to discriminate against SM backgrounds. These include searches employing variables such as  $\alpha T$ ,  $M_{T,2}$ , and the razor variables  $(M_R, R^2)$  [35–45].In this thesis, we will present the first search for supersymmetry using the novel Recursive Jigsaw Reconstruction (RJR) technique. RJR can be considered the conceptual successor of the razor variables. We impose a particular final state "decay tree" on an event, which

 $<sup>^1\</sup>mathrm{This}$  is the Standard Model corrected to include neutrino masses. These parameters are the fermion masses (6 leptons, 6 quarks), CKM and PMNS mixing angles (8 angles, 2 CP-violating phases), W/Z/Higgs masses (3), the Higgs field expectation value, and the couplings of the strong, weak, and electromagnetic forces (3  $\alpha_{force}$ ) .

107 roughly corresponds to a simplified Feynmann diagram. This allows an understand-108 ing of internal decay structure of an event, as well as additional rejection of SM 109 backgrounds.

This thesis details a search for the superpartners of the gluons and quarks, the 110 gluinos and squarks, in final states with zero leptons, with of data using the AT-111 LAS detector. This thesis is organized as follows. The theoretical motivation of the 112 Standard Model and supersymmetry are described in Chapters 2 and 3. The Large 113 Hadron Collider and the ATLAS detector are presented in Chapters 4 and 5. Chap-114 ter 5 provides a detailed description of Recursive Jigsaw Reconstruction, as well as 115 a description of the variables used for the particular search presented in this thesis. 116 Chapter 6 presents the details of the analysis, including the dataset, object recon-117 struction, and selections used by the analysis. In Chapter 7, the final results are 118 presented; since there is no evidence of a supersymmetric signal in the analysis, we 119 present the final exclusion curves in simplified supersymmetric models. 120

121

122

#### The Standard Model

- Here you can write some introductory remarks about your chapter. I like to give each sentence its own line.
- 125 When you need a new paragraph, just skip an extra line.

### 2.1 Quantum Field Theory

### 127 2.2 Symmetries

#### 128 2.3 The Standard Model

#### 129 Overview

- 130 By using the asterisk to start a new section, I keep the section from appearing in the
- table of contents. If you want your sections to be numbered and to appear in the
- table of contents, remove the asterisk.

#### 33 Fermions

- By using the asterisk to start a new section, I keep the section from appearing in the
- table of contents. If you want your sections to be numbered and to appear in the
- table of contents, remove the asterisk.

#### 137 Bosons

- By using the asterisk to start a new section, I keep the section from appearing in the
- table of contents. If you want your sections to be numbered and to appear in the
- table of contents, remove the asterisk.

### 141 2.4 Electroweak Symmetry breaking

- 142 By using the asterisk to start a new section, I keep the section from appearing in the
- table of contents. If you want your sections to be numbered and to appear in the
- table of contents, remove the asterisk.

### 2.5 Deficiencies of the Standard Model

- By using the asterisk to start a new section, I keep the section from appearing in the
- table of contents. If you want your sections to be numbered and to appear in the
- table of contents, remove the asterisk.

Chapter	3
---------	---

150

149

### Supersymmetry

- 151 Here you can write some introductory remarks about your chapter. I like to give each
- 152 sentence its own line.
- 153 When you need a new paragraph, just skip an extra line.

### 3.1 Motivation

- Only Additional allowed Lorentz invariant symmetry
- Dark Matter
- 157 Cancellation of quadratic divergences in corrections to the
- 158 Higgs Mass
- 3.2 Supersymmetry
- 160 3.3 Additional particle content
- $_{\scriptscriptstyle 161}$  3.4 Phenomenology
- 162 R parity Consequences for sq/gl decays

164

163

## The Large Hadron Collider

- 165 Here you can write some introductory remarks about your chapter. I like to give each
- sentence its own line.
- When you need a new paragraph, just skip an extra line.

### 4.1 Magnets


- 169 By using the asterisk to start a new section, I keep the section from appearing in the
- 170 table of contents. If you want your sections to be numbered and to appear in the
- table of contents, remove the asterisk.

173

### The ATLAS detector

Here you can write some introductory remarks about your chapter. I like to give each sentence its own line.

176 When you need a new paragraph, just skip an extra line.




177

178

### 5.1 Inner Detector

By using the asterisk to start a new section, I keep the section from appearing in the table of contents. If you want your sections to be numbered and to appear in the table of contents, remove the asterisk.


### 183 Pixel Detector



185

#### 86 Insertable B-Layer

187 Qualification task, so add a bit more.



188 189


## Semiconductor Tracker



191

192

### 193 Transition Radiation Tracker



195

### <sup>196</sup> 5.2 Calorimeter




197 198

199 Electromagnetic Calorimeter

200 Hadronic Calorimeter

## 201 5.3 Muon Spectrometer



202

203

204

205

### The Recursive Jigsaw Technique

- Here you can write some introductory remarks about your chapter. I like to give each
- 207 sentence its own line.
- 208 When you need a new paragraph, just skip an extra line.

### 209 6.1 Razor variables

- 210 By using the asterisk to start a new section, I keep the section from appearing in the
- 211 table of contents. If you want your sections to be numbered and to appear in the
- 212 table of contents, remove the asterisk.

### 213 6.2 SuperRazor variables

- 214 6.3 The Recursive Jigsaw Technique
- <sup>215</sup> 6.4 Variables used in the search for zero lepton
- SUSY

217	Chapter 7
218	Title of Chapter 1

220

## Title of Chapter 1

- 221 Here you can write some introductory remarks about your chapter. I like to give each
- 222 sentence its own line.
- 223 When you need a new paragraph, just skip an extra line.

### 224 8.1 Object reconstruction

225 Photons, Muons, and Electrons

226 **Jets** 

#### Missing transverse momentum

Probably longer, show some plots from the PUB note that we worked on

## 229 8.2 Signal regions

- 230 Gluino signal regions
- 231 Squark signal regions
- 232 Compressed signal regions

## 233 8.3 Background estimation

- 234 **Z** vv
- 235 **W** ev
- 236 ttbar

238

237

## Title of Chapter 1

- 239 Here you can write some introductory remarks about your chapter. I like to give each
- 240 sentence its own line.
- When you need a new paragraph, just skip an extra line.

### 242 9.1 Statistical Analysis

- 243 maybe to be moved to an appendix
- 9.2 Signal Region distributions
- 9.3 Pull Plots
- 9.4 Systematic Uncertainties
- 9.5 Exclusion plots

### Conclusion

- $\,$  Here you can write some introductory remarks about your chapter. I like to give each
- 250 sentence its own line.

248

When you need a new paragraph, just skip an extra line.

## 9.6 New Section

- 253 By using the asterisk to start a new section, I keep the section from appearing in the
- 254 table of contents. If you want your sections to be numbered and to appear in the
- table of contents, remove the asterisk.

### **Bibliography**

256

279

280

281

O. Perdereau, Planck 2015 cosmological results, 257 AIP Conf. Proc. 1743 (2016) p. 050014. 258 N. Aghanim et al., 259 Planck 2016 intermediate results. LI. Features in the cosmic microwave 260 background temperature power spectrum and shifts in cosmological parameters 261 (2016), arXiv: 1608.02487 [astro-ph.CO]. 262 J. S. Schwinger, 263 On Quantum electrodynamics and the magnetic moment of the electron, 264 Phys. Rev. **73** (1948) p. 416. 265 S. Laporta and E. Remiddi, 266 The Analytical value of the electron (q-2) at order alpha\*\*3 in QED, 267 Phys. Lett. **B379** (1996) p. 283, arXiv: hep-ph/9602417 [hep-ph]. 268 S. Schael et al., Precision electroweak measurements on the Z resonance, 269 Phys. Rept. **427** (2006) p. 257, arXiv: hep-ex/0509008 [hep-ex]. 270 S. L. Glashow, Partial Symmetries of Weak Interactions, 271 Nucl. Phys. **22** (1961) p. 579. 272 S. Weinberg, A Model of Leptons, Phys. Rev. Lett. 19 (1967) p. 1264. [7]273 A. Salam, Weak and Electromagnetic Interactions, 274 Conf. Proc. **C680519** (1968) p. 367. 275 M. Gell-Mann, A Schematic Model of Baryons and Mesons, 276 Phys. Lett. 8 (1964) p. 214. 277 G. Zweig, "An SU(3) model for strong interaction symmetry and its breaking. |10|278

Version 2," DEVELOPMENTS IN THE QUARK THEORY OF HADRONS.

VOL. 1. 1964 - 1978, ed. by D. Lichtenberg and S. P. Rosen, 1964 p. 22,

URL: http://inspirehep.net/record/4674/files/cern-th-412.pdf.

- 282 [11] S. Weinberg, Implications of Dynamical Symmetry Breaking, 283 Phys. Rev. **D13** (1976) p. 974.
- 284 [12] S. Weinberg, Implications of Dynamical Symmetry Breaking: An Addendum, 285 Phys. Rev. **D19** (1979) p. 1277.
- 286 [13] E. Gildener, Gauge Symmetry Hierarchies, Phys. Rev. **D14** (1976) p. 1667.
- 287 [14] L. Susskind,
- Dynamics of Spontaneous Symmetry Breaking in the Weinberg-Salam Theory, Phys. Rev. **D20** (1979) p. 2619.
- 290 [15] S. P. Martin, "A Supersymmetry Primer," 1997,
   291 eprint: arXiv:hep-ph/9709356.
- V. C. Rubin and W. K. Ford Jr., Rotation of the Andromeda Nebula from a Spectroscopic Survey of Emission Regions, Astrophys. J. **159** (1970) p. 379.
- 294 [17] M. S. Roberts and R. N. Whitehurst, 295 "The rotation curve and geometry of M31 at large galactocentric distances, 296 Astrophys. J. **201** (1970) p. 327.
- V. C. Rubin, N. Thonnard, and W. K. Ford Jr.,

  Rotational properties of 21 SC galaxies with a large range of luminosities and
  radii, from NGC 4605 /R = 4kpc/ to UGC 2885 /R = 122 kpc/,

  Astrophys. J. 238 (1980) p. 471.
- V. C. Rubin et al., Rotation velocities of 16 SA galaxies and a comparison of Sa, Sb, and SC rotation properties, Astrophys. J. **289** (1985) p. 81.
- A. Bosma,

  21-cm line studies of spiral galaxies. 2. The distribution and kinematics of
  neutral hydrogen in spiral galaxies of various morphological types.,

  Astron. J. 86 (1981) p. 1825.
- M. Persic, P. Salucci, and F. Stel, The Universal rotation curve of spiral galaxies: 1. The Dark matter connection,
   Mon. Not. Roy. Astron. Soc. 281 (1996) p. 27,
   arXiv: astro-ph/9506004 [astro-ph].
- 311 [22] M. Lisanti, "Lectures on Dark Matter Physics," 2016, aprint: arXiv:1603.03797.
- H. Miyazawa, Baryon Number Changing Currents, Prog. Theor. Phys. **36** (1966) p. 1266.

- 315 [24] J.-L. Gervais and B. Sakita, Generalizations of dual models, 316 Nucl. Phys. **B34** (1971) p. 477.
- 317 [25] J.-L. Gervais and B. Sakita, 318 Field Theory Interpretation of Supergauges in Dual Models,
- Nucl. Phys. **B34** (1971) p. 632.
- Yu. A. Golfand and E. P. Likhtman, Extension of the Algebra of Poincare
   Group Generators and Violation of p Invariance,
   JETP Lett. 13 (1971) p. 323, [Pisma Zh. Eksp. Teor. Fiz.13,452(1971)].
- 323 [27] A. Neveu and J. H. Schwarz, Factorizable dual model of pions, Nucl. Phys. **B31** (1971) p. 86.
- [28] A. Neveu and J. H. Schwarz, Quark Model of Dual Pions,
   Phys. Rev. **D4** (1971) p. 1109.
- D. V. Volkov and V. P. Akulov, *Is the Neutrino a Goldstone Particle?*Phys. Lett. **B46** (1973) p. 109.
- [30] J. Wess and B. Zumino,
   A Lagrangian Model Invariant Under Supergauge Transformations,
   Phys. Lett. B49 (1974) p. 52.
- 332 [31] A. Salam and J. A. Strathdee, Supersymmetry and Nonabelian Gauges, 333 Phys. Lett. **B51** (1974) p. 353.
- 334 [32] S. Ferrara, J. Wess, and B. Zumino, Supergauge Multiplets and Superfields, 335 Phys. Lett. **B51** (1974) p. 239.
- J. Wess and B. Zumino, Supergauge Transformations in Four-Dimensions, Nucl. Phys. **B70** (1974) p. 39.
- 338 [34] G. R. Farrar and P. Fayet, Phenomenology of the Production, Decay, and
  339 Detection of New Hadronic States Associated with Supersymmetry,
  340 Phys. Lett. **B76** (1978) p. 575.
- 341 [35] ATLAS Collaboration, 342 Search for the electroweak production of supersymmetric particles in 343  $\sqrt{s} = 8 \ TeV \ pp \ collisions \ with \ the \ ATLAS \ detector$ , 344 Phys. Rev. D **93** (2016) p. 052002, arXiv: 1509.07152 [hep-ex].
- 345 [36] ATLAS Collaboration, Summary of the searches for squarks and gluinos using  $\sqrt{s} = 8 \, TeV \, pp \, collisions \, with \, the \, ATLAS \, experiment \, at \, the \, LHC,$  347 JHEP 10 (2015) p. 054, arXiv: 1507.05525 [hep-ex].

- 348 [37] ATLAS Collaboration, ATLAS Run 1 searches for direct pair production of 449 third-generation squarks at the Large Hadron Collider, 350 Eur. Phys. J. C **75** (2015) p. 510, arXiv: 1506.08616 [hep-ex].
- CMS Collaboration, Search for supersymmetry with razor variables in pp collisions at  $\sqrt{s} = 7$  TeV, Phys. Rev. D **90** (2014) p. 112001, arXiv: 1405.3961 [hep-ex].
- CMS Collaboration, Inclusive search for supersymmetry using razor variables in pp collisions at  $\sqrt{s}=7$  TeV, Phys. Rev. Lett. **111** (2013) p. 081802, arXiv: 1212.6961 [hep-ex].
- CMS Collaboration, Search for Supersymmetry in pp Collisions at 7 TeV in Events with Jets and Missing Transverse Energy,

  Phys. Lett. B **698** (2011) p. 196, arXiv: 1101.1628 [hep-ex].
- 360 [41] CMS Collaboration, Search for Supersymmetry at the LHC in Events with
  361 Jets and Missing Transverse Energy, Phys. Rev. Lett. 107 (2011) p. 221804,
  362 arXiv: 1109.2352 [hep-ex].
- CMS Collaboration, Search for supersymmetry in hadronic final states using  $M_{T2}$  in pp collisions at  $\sqrt{s} = 7$  TeV, JHEP **10** (2012) p. 018, arXiv: 1207.1798 [hep-ex].
- CMS Collaboration, Searches for supersymmetry using the  $M_{T2}$  variable in hadronic events produced in pp collisions at 8 TeV, JHEP **05** (2015) p. 078, arXiv: 1502.04358 [hep-ex].
- CMS Collaboration, Search for new physics with the  $M_{T2}$  variable in all-jets final states produced in pp collisions at  $\sqrt{s} = 13 \, TeV \, (2016)$ , arXiv: 1603.04053 [hep-ex].
- 372 [45] ATLAS Collaboration, Multi-channel search for squarks and gluinos in  $\sqrt{s} = 7$  TeV pp collisions with the ATLAS detector at the LHC, 374 Eur. Phys. J. C **73** (2013) p. 2362, arXiv: 1212.6149 [hep-ex].