1	A search for sparticles in zero lepton final states
2	Russell W. Smith

3	Submitted in partial fulfillment of the
4	requirements for the degree of
5	Doctor of Philosophy
6	in the Graduate School of Arts and Sciences

7 COLUMBIA UNIVERSITY

8 2016

10

11

© 2016

Russell W. Smith

All rights reserved

12	ABSTRACT
13	A search for sparticles in zero lepton final states
14	Russell W. Smith
15	TODO : Here's where your abstract will eventually go. The above text is all in the
16	center, but the abstract itself should be written as a regular paragraph on the page
17	and it should not have indentation. Just replace this text.

18 Contents

19	Contents				
20	1	1			
21	2	The	e Standard Model	5	
22		2.1	Quantum Field Theory	5	
23		2.2	Symmetries	5	
24		2.3	The Standard Model	5	
25		2.4	Electroweak Symmetry breaking	6	
26		2.5	Deficiencies of the Standard Model	6	
27	3	Sup	persymmetry	7	
28		3.1	Motivation	7	
29		3.2	Supersymmetry	7	
30		3.3	Additional particle content	7	
31		3.4	Phenomenology	7	
32	4	The	e Large Hadron Collider	9	
33		4.1	Magnets	9	
34	5	The	e ATLAS detector	11	
35		5.1	Inner Detector	11	
36		5.2	Calorimeter	13	

37		5.3	Muon Spectrometer	13
38	6	The	e Recursive Jigsaw Technique	15
39		6.1	Razor variables	15
40		6.2	SuperRazor variables	15
41		6.3	The Recursive Jigsaw Technique	15
42		6.4	Variables used in the search for zero lepton SUSY	15
43	7	Tab	le of Contents Title	17
44	8	A s	earch for supersymmetric particles in zero lepton final states	
45		witl	h the Recursive Jigsaw Technique	19
46		8.1	Object reconstruction	19
47		8.2	Signal regions	20
48		8.3	Background estimation	20
49	9	Res	ults	21
50		9.1	Statistical Analysis	21
51		9.2	Signal Region distributions	21
52		9.3	Pull Plots	21
53		9.4	Systematic Uncertainties	21
54		9.5	Exclusion plots	21
55	Co	onclu	ısion	23
56		9.6	New Section	23
57	Bi	iblios	vraphy	25

Acknowledgements

58

Dedication

Introduction

Particle physics is a remarkably successful field of scientific inquiry. The ability to precisely predict the properties of a exceedingly wide range of physical phenomena, 63 such as the description of the cosmic microwave background [1, 2], the understanding 64 of the anomalous magnetic dipole moment of the electron [3, 4], and the measurement 65 of the number of weakly-interacting neutrino flavors [5] is truly amazing. 66 The theory that has allowed this range of predictions is the Standard Model of par-67 ticle physics (SM). The Standard Model combines the electroweak theory of Glashow, 68 Weinberg, and Salam [6–8] with the theory of the strong interactions, as first envi-69 sioned by Gell-Mann and Zweig [9, 10]. This quantum field theory (QFT) contains 70 a tiny number of particles, whose interactions describe phenomena up to at least the 71 TeV scale. These particles are manifestations of the fields of the Standard Model, after application of the Higgs Mechanism. The particle content of the SM consists 73 only of the six quarks, the six leptons, the four gauge bosons, and the scalar Higgs boson. 75 Despite its impressive range of described phenomena, the Standard Model has 76 some theoretical and experimental deficiencies. The SM contains 26 free parameters 77 It would be more theoretically pleasing to understand these free parameters in 78 terms of a more fundamental theory. The major theoretical concern of the Standard Model, as it pertains to this thesis, is the $hierarchy\ problem[11-15]$. The light mass

 $^{^1\}mathrm{This}$ is the Standard Model corrected to include neutrino masses. These parameters are the fermion masses (6 leptons, 6 quarks), CKM and PMNS mixing angles (8 angles, 2 CP-violating phases), W/Z/Higgs masses (3), the Higgs field expectation value, and the couplings of the strong, weak, and electromagnetic forces (3 α_{force}) .

of the Higgs boson (125 GeV) should be quadratically dependent on the scale of UV physics, due to the quantum corrections from high-energy physics processes. The most perplexing experimental issue is the existence of dark matter, as demonstrated by galactic rotation curves [16-22]. This data has shown that there exists additional 84 matter which has not yet been seen interacting with the particles of the Standard 85 Model. There is no particle in the SM which can act as a candidate for dark matter. 86 Both of these major issues, as well as numerous others, can be solved by the 87 introduction of supersymmetry (SUSY) [15, 23–33]. In supersymmetric theories, each 88 SM particles has a so-called *superpartner*, or sparticle partner, differing from given SM 89 particle by 1/2 in spin. These theories solve the hierarchy problem, since the quantum corrections induced from the superpartners exactly cancel those induced by the SM particles. In addition, these theories are usually constructed assuming R-parity, 92 which can be thought of as the "charge" of supersymmetry, with SM particles having 93

R=1 and sparticles having R=-1. In collider experiments, since the incoming

SM particles have total R=1, the resulting sparticles are produced in pairs. This

produces a rich phenomenology, which is characterized by significant hadronic activity

and large missing transverse energy $(E_{\rm T}^{\rm miss})$, which provide significant discrimination

94

95

96

97

98

against SM backgrounds [34].

Despite the power of searches for supersymmetry where $E_{\mathrm{T}}^{\mathrm{miss}}$ is a primary dis-99 criminating variable, there has been significant interest in the use of other variables 100 to discriminate against SM backgrounds. These include searches employing variables 101 such as αT , $M_{T,2}$, and the razor variables (M_R, R^2) [35–45]. In this thesis, we will 102 present the first search for supersymmetry using the novel Recursive Jigsaw Recon-103 struction (RJR) technique. RJR can be considered the conceptual successor of the 104 razor variables. We impose a particular final state "decay tree" on an events, which 105 roughly corresponds to a simplified Feynmann diagram in decays containing weakly-106 interacting particles. We account for the missing degrees of freedom associated to 107

the weakly-interacting particles by a series of simplifying assumptions, which allow us to calculate our variables of interest at each step in the decay tree. This allows an unprecedented understanding of the internal structure of the decay and the ability to construct additional variables to reject Standard Model backgrounds.

This thesis details a search for the superpartners of the gluon and quarks, the 112 gluino and squarks, in final states with zero leptons, with $13.3~{\rm fb^{-1}of}$ data using the 113 ATLAS detector. We organzie the thesis as follows. The theoretical foundations of 114 the Standard Model and supersymmetry are described in Chapters 2 and 3. The 115 Large Hadron Collider and the ATLAS detector are presented in Chapters 4 and 5. 116 Chapter 5 provides a detailed description of Recursive Jigsaw Reconstruction and a 117 description of the variables used for the particular search presented in this thesis. 118 Chapter 6 presents the details of the analysis, including details of the dataset, object 119 reconstruction, and selections used. In Chapter 7, the final results are presented; 120 since there is no evidence of a supersymmetric signal in the analysis, we present the 121 final exclusion curves in simplified supersymmetric models. 122

Chapter 2

123

124

The Standard Model

- Here you can write some introductory remarks about your chapter. I like to give each sentence its own line.
- 127 When you need a new paragraph, just skip an extra line.

2.1 Quantum Field Theory

129 2.2 Symmetries

130 2.3 The Standard Model

131 Overview

- By using the asterisk to start a new section, I keep the section from appearing in the
- table of contents. If you want your sections to be numbered and to appear in the
- table of contents, remove the asterisk.

55 Fermions

- By using the asterisk to start a new section, I keep the section from appearing in the
- table of contents. If you want your sections to be numbered and to appear in the
- table of contents, remove the asterisk.

139 Bosons

140 By using the asterisk to start a new section, I keep the section from appearing in the

table of contents. If you want your sections to be numbered and to appear in the

table of contents, remove the asterisk.

143 2.4 Electroweak Symmetry breaking

144 By using the asterisk to start a new section, I keep the section from appearing in the

table of contents. If you want your sections to be numbered and to appear in the

table of contents, remove the asterisk.

147 2.5 Deficiencies of the Standard Model

By using the asterisk to start a new section, I keep the section from appearing in the

table of contents. If you want your sections to be numbered and to appear in the

table of contents, remove the asterisk.

Cha	oter	3

151

Supersymmetry

- 153 Here you can write some introductory remarks about your chapter. I like to give each
- 154 sentence its own line.
- When you need a new paragraph, just skip an extra line.

3.1 Motivation

- 157 Only Additional allowed Lorentz invariant symmetry
- 158 Dark Matter
- 159 Cancellation of quadratic divergences in corrections to the
- 160 Higgs Mass
- 161 3.2 Supersymmetry
- 162 3.3 Additional particle content
- 163 3.4 Phenomenology
- 164 R parity Consequences for sq/gl decays

Chapter 4

166

165

The Large Hadron Collider

- Here you can write some introductory remarks about your chapter. I like to give each sentence its own line.
- When you need a new paragraph, just skip an extra line.


170 4.1 Magnets

- 171 By using the asterisk to start a new section, I keep the section from appearing in the
- table of contents. If you want your sections to be numbered and to appear in the
- table of contents, remove the asterisk.

The ATLAS detector

Here you can write some introductory remarks about your chapter. I like to give each sentence its own line.

178 When you need a new paragraph, just skip an extra line.




179

180

5.1 Inner Detector

By using the asterisk to start a new section, I keep the section from appearing in the table of contents. If you want your sections to be numbered and to appear in the table of contents, remove the asterisk.


Pixel Detector



187

88 Insertable B-Layer

189 Qualification task, so add a bit more.



191


190

92 Semiconductor Tracker



194

195 Transition Radiation Tracker



197


¹⁹⁸ 5.2 Calorimeter



199 200

- 201 Electromagnetic Calorimeter
- 202 Hadronic Calorimeter

203 5.3 Muon Spectrometer



204

205

Chapter 6

206

207

The Recursive Jigsaw Technique

- 208 Here you can write some introductory remarks about your chapter. I like to give each
- 209 sentence its own line.
- 210 When you need a new paragraph, just skip an extra line.

211 6.1 Razor variables

- 212 By using the asterisk to start a new section, I keep the section from appearing in the
- table of contents. If you want your sections to be numbered and to appear in the
- table of contents, remove the asterisk.

215 6.2 SuperRazor variables

- 216 6.3 The Recursive Jigsaw Technique
- ²¹⁷ 6.4 Variables used in the search for zero lepton
- SUSY

219	Chapter 7			
220	Title of Chapter 1			

Title of Chapter 1

- 223 Here you can write some introductory remarks about your chapter. I like to give each
- 224 sentence its own line.
- 225 When you need a new paragraph, just skip an extra line.

226 8.1 Object reconstruction

- 227 Photons, Muons, and Electrons
- 228 **Jets**
- 229 Missing transverse momentum
- 230 Probably longer, show some plots from the PUB note that we worked on

²³¹ 8.2 Signal regions

- 232 Gluino signal regions
- 233 Squark signal regions
- 234 Compressed signal regions

235 8.3 Background estimation

- 236 **Z** vv
- 237 **W** ev
- 238 ttbar

Chapter 9

240

239

Title of Chapter 1

- 241 Here you can write some introductory remarks about your chapter. I like to give each
- 242 sentence its own line.
- 243 When you need a new paragraph, just skip an extra line.

244 9.1 Statistical Analysis

245 maybe to be moved to an appendix

²⁴⁶ 9.2 Signal Region distributions

- 9.3 Pull Plots
- 9.4 Systematic Uncertainties
- 9.5 Exclusion plots

Conclusion

- 251 Here you can write some introductory remarks about your chapter. I like to give each
- 252 sentence its own line.

250

253 When you need a new paragraph, just skip an extra line.

9.6 New Section

- 255 By using the asterisk to start a new section, I keep the section from appearing in the
- 256 table of contents. If you want your sections to be numbered and to appear in the
- table of contents, remove the asterisk.

Bibliography

258

282

283

O. Perdereau, Planck 2015 cosmological results, 259 AIP Conf. Proc. 1743 (2016) p. 050014. 260 N. Aghanim et al., 261 Planck 2016 intermediate results. LI. Features in the cosmic microwave 262 background temperature power spectrum and shifts in cosmological parameters 263 (2016), arXiv: 1608.02487 [astro-ph.CO]. 264 J. S. Schwinger, 265 On Quantum electrodynamics and the magnetic moment of the electron, 266 Phys. Rev. **73** (1948) p. 416. 267 S. Laporta and E. Remiddi, 268 The Analytical value of the electron (q-2) at order alpha**3 in QED, 269 Phys. Lett. **B379** (1996) p. 283, arXiv: hep-ph/9602417 [hep-ph]. 270 S. Schael et al., Precision electroweak measurements on the Z resonance, 271 Phys. Rept. **427** (2006) p. 257, arXiv: hep-ex/0509008 [hep-ex]. 272 S. L. Glashow, Partial Symmetries of Weak Interactions, 273 Nucl. Phys. **22** (1961) p. 579. 274 S. Weinberg, A Model of Leptons, Phys. Rev. Lett. 19 (1967) p. 1264. 275 A. Salam, Weak and Electromagnetic Interactions, 276 Conf. Proc. **C680519** (1968) p. 367. 277 M. Gell-Mann, A Schematic Model of Baryons and Mesons, 278 Phys. Lett. 8 (1964) p. 214. 279 G. Zweig, "An SU(3) model for strong interaction symmetry and its breaking. |10|280 Version 2," DEVELOPMENTS IN THE QUARK THEORY OF HADRONS. 281 VOL. 1. 1964 - 1978, ed. by D. Lichtenberg and S. P. Rosen, 1964 p. 22,

URL: http://inspirehep.net/record/4674/files/cern-th-412.pdf.

- 284 [11] S. Weinberg, Implications of Dynamical Symmetry Breaking, 285 Phys. Rev. **D13** (1976) p. 974.
- 286 [12] S. Weinberg, Implications of Dynamical Symmetry Breaking: An Addendum, 287 Phys. Rev. **D19** (1979) p. 1277.
- 288 [13] E. Gildener, Gauge Symmetry Hierarchies, Phys. Rev. **D14** (1976) p. 1667.
- 289 [14] L. Susskind,
- Dynamics of Spontaneous Symmetry Breaking in the Weinberg-Salam Theory, Phys. Rev. **D20** (1979) p. 2619.
- [15] S. P. Martin, "A Supersymmetry Primer," 1997,
 eprint: arXiv:hep-ph/9709356.
- V. C. Rubin and W. K. Ford Jr., Rotation of the Andromeda Nebula from a Spectroscopic Survey of Emission Regions, Astrophys. J. **159** (1970) p. 379.
- 296 [17] M. S. Roberts and R. N. Whitehurst, 297 "The rotation curve and geometry of M31 at large galactocentric distances, 298 Astrophys. J. **201** (1970) p. 327.
- V. C. Rubin, N. Thonnard, and W. K. Ford Jr.,

 Rotational properties of 21 SC galaxies with a large range of luminosities and
 radii, from NGC 4605 /R = 4kpc/ to UGC 2885 /R = 122 kpc/,

 Astrophys. J. 238 (1980) p. 471.
- V. C. Rubin et al., Rotation velocities of 16 SA galaxies and a comparison of Sa, Sb, and SC rotation properties, Astrophys. J. **289** (1985) p. 81.
- A. Bosma,
 21-cm line studies of spiral galaxies. 2. The distribution and kinematics of
 neutral hydrogen in spiral galaxies of various morphological types.,
 Astron. J. 86 (1981) p. 1825.
- 309 [21] M. Persic, P. Salucci, and F. Stel, The Universal rotation curve of spiral 310 galaxies: 1. The Dark matter connection, 311 Mon. Not. Roy. Astron. Soc. 281 (1996) p. 27, 312 arXiv: astro-ph/9506004 [astro-ph].
- 313 [22] M. Lisanti, "Lectures on Dark Matter Physics," 2016, arXiv:1603.03797.
- H. Miyazawa, Baryon Number Changing Currents, Prog. Theor. Phys. **36** (1966) p. 1266.

- 317 [24] J.-L. Gervais and B. Sakita, Generalizations of dual models, 318 Nucl. Phys. **B34** (1971) p. 477.
- 319 [25] J.-L. Gervais and B. Sakita, 320 Field Theory Interpretation of Supergauges in Dual Models,
- Nucl. Phys. **B34** (1971) p. 632.
- Yu. A. Golfand and E. P. Likhtman, Extension of the Algebra of Poincare Group Generators and Violation of p Invariance, JETP Lett. **13** (1971) p. 323, [Pisma Zh. Eksp. Teor. Fiz.13,452(1971)].
- 325 [27] A. Neveu and J. H. Schwarz, Factorizable dual model of pions, Nucl. Phys. **B31** (1971) p. 86.
- 327 [28] A. Neveu and J. H. Schwarz, *Quark Model of Dual Pions*, 328 Phys. Rev. **D4** (1971) p. 1109.
- 329 [29] D. V. Volkov and V. P. Akulov, *Is the Neutrino a Goldstone Particle?*330 Phys. Lett. **B46** (1973) p. 109.
- 331 [30] J. Wess and B. Zumino,

 A Lagrangian Model Invariant I
- A Lagrangian Model Invariant Under Supergauge Transformations, Phys. Lett. **B49** (1974) p. 52.
- 334 [31] A. Salam and J. A. Strathdee, Supersymmetry and Nonabelian Gauges, 335 Phys. Lett. **B51** (1974) p. 353.
- 336 [32] S. Ferrara, J. Wess, and B. Zumino, Supergauge Multiplets and Superfields, 337 Phys. Lett. **B51** (1974) p. 239.
- J. Wess and B. Zumino, Supergauge Transformations in Four-Dimensions, Nucl. Phys. **B70** (1974) p. 39.
- 340 [34] G. R. Farrar and P. Fayet, Phenomenology of the Production, Decay, and
 341 Detection of New Hadronic States Associated with Supersymmetry,
 342 Phys. Lett. **B76** (1978) p. 575.
- 343 [35] ATLAS Collaboration, 344 Search for the electroweak production of supersymmetric particles in 345 $\sqrt{s} = 8$ TeV pp collisions with the ATLAS detector, 346 Phys. Rev. D **93** (2016) p. 052002, arXiv: 1509.07152 [hep-ex].
- 347 [36] ATLAS Collaboration, Summary of the searches for squarks and gluinos using
 348 $\sqrt{s} = 8 \text{ TeV pp collisions with the ATLAS experiment at the LHC},$ 349 JHEP **10** (2015) p. 054, arXiv: 1507.05525 [hep-ex].

- 350 [37] ATLAS Collaboration, ATLAS Run 1 searches for direct pair production of 351 third-generation squarks at the Large Hadron Collider, 352 Eur. Phys. J. C **75** (2015) p. 510, arXiv: 1506.08616 [hep-ex].
- 353 [38] CMS Collaboration, 354 Search for supersymmetry with razor variables in pp collisions at $\sqrt{s} = 7$ TeV, 355 Phys. Rev. D **90** (2014) p. 112001, arXiv: 1405.3961 [hep-ex].
- CMS Collaboration, Inclusive search for supersymmetry using razor variables in pp collisions at $\sqrt{s}=7$ TeV, Phys. Rev. Lett. **111** (2013) p. 081802, arXiv: 1212.6961 [hep-ex].
- CMS Collaboration, Search for Supersymmetry in pp Collisions at 7 TeV in Events with Jets and Missing Transverse Energy,
 Phys. Lett. B **698** (2011) p. 196, arXiv: 1101.1628 [hep-ex].
- CMS Collaboration, Search for Supersymmetry at the LHC in Events with

 Jets and Missing Transverse Energy, Phys. Rev. Lett. 107 (2011) p. 221804,

 arXiv: 1109.2352 [hep-ex].
- CMS Collaboration, Search for supersymmetry in hadronic final states using M_{T2} in pp collisions at $\sqrt{s} = 7$ TeV, JHEP **10** (2012) p. 018, arXiv: 1207.1798 [hep-ex].
- 368 [43] CMS Collaboration, Searches for supersymmetry using the M_{T2} variable in 369 hadronic events produced in pp collisions at 8 TeV, JHEP **05** (2015) p. 078, 370 arXiv: 1502.04358 [hep-ex].
- CMS Collaboration, Search for new physics with the M_{T2} variable in all-jets final states produced in pp collisions at $\sqrt{s} = 13$ TeV (2016), arXiv: 1603.04053 [hep-ex].
- 374 [45] ATLAS Collaboration, Multi-channel search for squarks and gluinos in $\sqrt{s} = 7$ TeV pp collisions with the ATLAS detector at the LHC, 376 Eur. Phys. J. C **73** (2013) p. 2362, arXiv: 1212.6149 [hep-ex].