

Probabilistic Programming

Rajbir-Singh Nirwan
October 22, 2019

Outline

- Probabilistic Modelling
- Sampling
- Variational Inference
- Probabilistic Programming

Why do we need it?

- Uncertainty estimation
- Intrinsic Regularization
- Explicit assumptions
- More interpretable models

Recap

Observed Data

$$\mathcal{D} = \{(\mathbf{x}_n, y_n)\}_{n=1}^N$$

$$\mathbf{x}_n \in \mathbb{R}^D, y_n \in \mathbb{R}$$

Model

$$y_n = f_{\mathbf{w}}(\mathbf{x}_n) + \epsilon_n = \mathbf{w}^T \mathbf{x}_n + \epsilon_n$$

$$E_{\mathcal{D}}(\mathbf{w}) = \sum_{n=1}^N (y_n - \mathbf{w}^T \mathbf{x}_n)^2 + \lambda \|\mathbf{w}\|_2^2$$

Fit

$$\mathbf{w}^* = \min_{\mathbf{w}} E_{\mathcal{D}}(\mathbf{w})$$

Prediction

$$y_{new} = f_{\mathbf{w}^*}(\mathbf{x}_{new})$$

Recap

Observed Data

$$\mathcal{D} = \{(\mathbf{x}_n, y_n)\}_{n=1}^N$$

$$\mathbf{x}_n \in \mathbb{R}^D, y_n \in \mathbb{R}$$

Model

$$y_n = f_{\mathbf{w}}(\mathbf{x}_n) + \epsilon_n = \mathbf{w}^T \mathbf{x}_n + \epsilon_n$$

$$E_{\mathcal{D}}(\mathbf{w}) = \sum_{n=1}^N (y_n - \mathbf{w}^T \mathbf{x}_n)^2 + \lambda \|\mathbf{w}\|_2^2$$

Fit

$$\mathbf{w}^* = \min_{\mathbf{w}} E_{\mathcal{D}}(\mathbf{w})$$

Prediction

$$y_{new} = f_{\mathbf{w}^*}(\mathbf{x}_{new})$$

Observed Data

$$\mathcal{D} = \{(\mathbf{x}_n, y_n)\}_{n=1}^N$$

$$\mathbf{x}_n \in \mathbb{R}^D, y_n \in \mathbb{R}$$

Model

$$y_n = f_{\mathbf{w}}(\mathbf{x}_n) + \epsilon_n = \mathbf{w}^T \mathbf{x}_n + \epsilon_n$$

$$p(\mathbf{y} | \mathbf{w}, \sigma^2) = \prod_{n=1}^N \mathcal{N}(y_n | \mathbf{w}^T \mathbf{x}_n, \sigma^2)$$

$$p(\mathbf{w}) = \mathcal{N}(\mathbf{w} | \mathbf{0}, \mathbf{I})$$

Inference

$$p(\mathbf{w} | \mathbf{y}) = \frac{p(\mathbf{y} | \mathbf{w}) p(\mathbf{w})}{p(\mathbf{y})}$$

Prediction

$$p(y_{new} | \mathbf{y}) = \int p(y_{new} | \mathbf{w}) p(\mathbf{w} | \mathbf{y}) d\mathbf{w}$$

Probabilistic Modelling

$$p(x) = \int p(x, y) \, dy \quad \& \quad p(x, y) = p(y | x) p(x)$$

- Inference

$$p(\theta | \mathcal{D}) = \frac{p(\mathcal{D} | \theta) p(\theta)}{p(\mathcal{D})}$$

- Prediction

$$p(y | \mathcal{D}) = \int p(y | \theta) p(\theta | \mathcal{D}) \, d\theta$$

Probabilistic Modelling

$$p(x) = \int p(x, y) dy \quad \& \quad p(x, y) = p(y | x) p(x)$$

- Inference
$$p(\theta | \mathcal{D}) = \frac{p(\mathcal{D} | \theta) p(\theta)}{p(\mathcal{D})}$$
- Prediction
$$p(y | \mathcal{D}) = \int p(y | \theta) p(\theta | \mathcal{D}) d\theta$$
- Not tractable most of the time
 - ➔ Approximation Methods
 - Sampling
 - Variational Inference

Sampling (HMC)

- Approximate $p(\boldsymbol{\theta} | \mathcal{D})$ by N samples



$$\mathbb{E}_p[f] = \int p(\boldsymbol{\theta} | \mathcal{D}) f(\boldsymbol{\theta}) d\boldsymbol{\theta} \approx \frac{1}{N} \sum_{n=1}^N f(\boldsymbol{\theta}_n)$$

Sampling (HMC)

- Approximate $p(\boldsymbol{\theta} | \mathcal{D})$ by N samples

$$\mathbb{E}_p[f] = \int p(\boldsymbol{\theta} | \mathcal{D}) f(\boldsymbol{\theta}) d\boldsymbol{\theta} \approx \frac{1}{N} \sum_{n=1}^N f(\boldsymbol{\theta}_n)$$

- HMC: Particle moving in probability landscape according to Hamilton's equations with random Gaussian “kicks”



Sampling (HMC)

$$\mathbf{H}(\boldsymbol{\theta}, p) = \mathbf{U}(\boldsymbol{\theta}) + \mathbf{K}(p) = \text{const}$$

$$\mathbf{U}(\boldsymbol{\theta}) = -\log p(\boldsymbol{\theta} \mid \mathcal{D}) \quad \mathbf{K}(p) = \frac{p^2}{2m}$$

Sampling (HMC)

$$\mathbf{H}(\boldsymbol{\theta}, \mathbf{p}) = \mathbf{U}(\boldsymbol{\theta}) + \mathbf{K}(\mathbf{p}) = \text{const}$$

$$\mathbf{U}(\boldsymbol{\theta}) = -\log p(\boldsymbol{\theta} | \mathcal{D}) \quad \mathbf{K}(\mathbf{p}) = \frac{\mathbf{p}^2}{2m}$$

$$p(\boldsymbol{\theta}, \mathbf{p}) \propto \exp(-\mathbf{H}) = p(\boldsymbol{\theta} | \mathcal{D}) e^{-\mathbf{p}^2/2m}$$

→ $\boldsymbol{\theta}$ and \mathbf{p} are uncorrelated

→ \mathbf{p} is Gaussian

Sampling (HMC)

$$\mathbf{H}(\boldsymbol{\theta}, \mathbf{p}) = \mathbf{U}(\boldsymbol{\theta}) + \mathbf{K}(\mathbf{p}) = \text{const}$$

$$\mathbf{U}(\boldsymbol{\theta}) = -\log p(\boldsymbol{\theta} | \mathcal{D}) \quad \mathbf{K}(\mathbf{p}) = \frac{\mathbf{p}^2}{2m}$$

$$p(\boldsymbol{\theta}, \mathbf{p}) \propto \exp(-\mathbf{H}) = p(\boldsymbol{\theta} | \mathcal{D}) e^{-\mathbf{p}^2/2m}$$

→ $\boldsymbol{\theta}$ and \mathbf{p} are uncorrelated

→ \mathbf{p} is Gaussian

Hamilton's Equations:

$$\dot{\boldsymbol{\theta}} = \frac{\partial \mathbf{H}}{\partial \mathbf{p}} \quad \dot{\mathbf{p}} = -\frac{\partial \mathbf{H}}{\partial \boldsymbol{\theta}}$$

Sampling (HMC)

$$\mathbf{H}(\boldsymbol{\theta}, \mathbf{p}) = \mathbf{U}(\boldsymbol{\theta}) + \mathbf{K}(\mathbf{p}) = \text{const}$$

$$\mathbf{U}(\boldsymbol{\theta}) = -\log p(\boldsymbol{\theta} | \mathcal{D}) \quad \mathbf{K}(\mathbf{p}) = \frac{\mathbf{p}^2}{2m}$$

$$p(\boldsymbol{\theta}, \mathbf{p}) \propto \exp(-\mathbf{H}) = p(\boldsymbol{\theta} | \mathcal{D}) e^{-\mathbf{p}^2/2m}$$

→ $\boldsymbol{\theta}$ and \mathbf{p} are uncorrelated

→ \mathbf{p} is Gaussian


Hamilton's Equations:

$$\dot{\boldsymbol{\theta}} = \frac{\partial \mathbf{H}}{\partial \mathbf{p}} \quad \dot{\mathbf{p}} = -\frac{\partial \mathbf{H}}{\partial \boldsymbol{\theta}}$$

<https://github.com/PyMLVizard/PyMLViz>

Sampling (in Theory):

1. Choose some $\boldsymbol{\theta}$ at random
 2. N times:
 1. $\mathbf{p} \sim \mathcal{N}(0,1)$
 2. Solve for path with $\mathbf{H}(\boldsymbol{\theta}, \mathbf{p})$ for fixed amount of time
 3. Save $\boldsymbol{\theta}$ and \mathbf{p} as a sample
- Chain of $\boldsymbol{\theta}$ s converge to $p(\boldsymbol{\theta} | \mathcal{D})$



Variational Bayes

- Approximate $p(\boldsymbol{\theta} | \mathcal{D})$ by $q_{\nu}(\boldsymbol{\theta})$

Variational Bayes

- Approximate $p(\boldsymbol{\theta} | \mathcal{D})$ by $q_{\nu}(\boldsymbol{\theta})$
- Goodness is measured in e.g. Kullback-Leibler divergence


$$\mathbf{KL}[q_{\nu}(\boldsymbol{\theta}) || p(\boldsymbol{\theta} | \mathcal{D})] = \int q_{\nu}(\boldsymbol{\theta}) \ln \frac{q_{\nu}(\boldsymbol{\theta})}{p(\boldsymbol{\theta} | \mathcal{D})} d\boldsymbol{\theta}$$

Variational Bayes

- Approximate $p(\boldsymbol{\theta} | \mathcal{D})$ by $q_{\nu}(\boldsymbol{\theta})$
- Goodness is measured in e.g. Kullback-Leibler divergence

$$\mathbf{KL}[q_{\nu}(\boldsymbol{\theta}) || p(\boldsymbol{\theta} | \mathcal{D})] = \int q_{\nu}(\boldsymbol{\theta}) \ln \frac{q_{\nu}(\boldsymbol{\theta})}{p(\boldsymbol{\theta} | \mathcal{D})} d\boldsymbol{\theta}$$

**Turns Inference
Into Optimization**



Variational Bayes

$$\ln p(\mathcal{D}) = \mathbf{ELBO} + \mathbf{KL}[q_{\nu}(\boldsymbol{\theta}) \| p(\boldsymbol{\theta} | \mathcal{D})]$$

Variational Bayes

$$\ln p(\mathcal{D}) = \mathbf{ELBO} + \mathbf{KL}[q_{\nu}(\boldsymbol{\theta}) \| p(\boldsymbol{\theta} | \mathcal{D})]$$

$$\mathbf{ELBO} = \int q_{\nu}(\boldsymbol{\theta}) \ln \frac{p(\mathcal{D} | \boldsymbol{\theta}) p(\boldsymbol{\theta})}{q_{\nu}(\boldsymbol{\theta})} d\boldsymbol{\theta} = - \int q_{\nu}(\boldsymbol{\theta}) \ln \frac{q_{\nu}(\boldsymbol{\theta})}{\tilde{p}(\boldsymbol{\theta} | \mathcal{D})} d\boldsymbol{\theta}$$


$$\mathbf{KL}[q_{\nu}(\boldsymbol{\theta}) \| p(\boldsymbol{\theta} | \mathcal{D})] = \int q_{\nu}(\boldsymbol{\theta}) \ln \frac{q_{\nu}(\boldsymbol{\theta})}{p(\boldsymbol{\theta} | \mathcal{D})} d\boldsymbol{\theta}$$

Variational Bayes

$$\ln p(\mathcal{D}) = \mathbf{ELBO} + \mathbf{KL}[q_{\nu}(\boldsymbol{\theta}) \| p(\boldsymbol{\theta} | \mathcal{D})]$$

$$\mathbf{ELBO} = \int q_{\nu}(\boldsymbol{\theta}) \ln \frac{p(\mathcal{D} | \boldsymbol{\theta}) p(\boldsymbol{\theta})}{q_{\nu}(\boldsymbol{\theta})} d\boldsymbol{\theta} = - \int q_{\nu}(\boldsymbol{\theta}) \ln \frac{q_{\nu}(\boldsymbol{\theta})}{\tilde{p}(\boldsymbol{\theta} | \mathcal{D})} d\boldsymbol{\theta}$$

$$\mathbf{KL}[q_{\nu}(\boldsymbol{\theta}) \| p(\boldsymbol{\theta} | \mathcal{D})] = \int q_{\nu}(\boldsymbol{\theta}) \ln \frac{q_{\nu}(\boldsymbol{\theta})}{p(\boldsymbol{\theta} | \mathcal{D})} d\boldsymbol{\theta}$$



**Thanks for your
attention!**