

The influence of the Southland Current on the circulation within Pegasus Bay

Rafael Soutelino and Brett Beamsley

MetOcean Solutions

Objectives

Investigate the effect of the Southland Current (SC) on the coastal circulation at Pegasus Bay (PB) using a combination of long term realistic modelling and *in situ* measurements.

- 1 What is the contribution of the SC to the total currents on the inner shelf of PB?
- 2 What is the impact of remote forcing in modelling efforts at the inner shelf of PB?
- 3 What are the implications of underestimating the SC flow in the prediction of arbitrary passive tracers transport and dispersion in PB?

Introduction

The East coast of New Zealand is a notably complex area in terms of interaction between oceanic and coastal circulation, due to the existence of important western boundary current systems flowing along the continental slopes and outer shelves (Jannelle & Flemming, 2005). The SC is a northward/northeastward oceanic current that extends from Stewart Island in the south to the Chatham Rise in the north, formed at the subtropical convergence west of New Zealand (Chiswell, 1996). Most of it bends offshore towards the Chatham Rise (Figure 1), but a small portion bends towards PB, in the form of an anticyclonic eddy around BP (Carter & Herzer, 1979). A net southerly alongshore depth-averaged transport is reported in previous work, potentially associated with an eddy in PB.




Figure 1: Circulation features off the east coast of New Zealand South Island. The green mark is the location of the observations. The blue mark is an arbitrary location within the SC path.

Methods

A calibrated 10-year ROMS hindcast was carried out. Two different runs were undertaken. A CONTROL run consisted of full realistic hydrodynamical forcing (Figure 2 and Table 2) and a TEST run was deprived of most of the SC incoming flow (Table 1).




Figure 2: Hindcast nesting approach with ROMS. See Tables 1 and 2 for more details.

| Experiment | Nests |
|------------|----------------------------|
| CONTROL | CFSR → NZ → CESI → PEGASUS |
| TEST | CFSR → PEGASUS |

Table 1: Description of the numerical experiments.

| | NZ | CESI | PEGASUS |
|------------|---------|---------|---------|
| Resolution | 0.08° | 0.03° | 0.004° |
| Boundary | CFSR | NZ | CESI |
| Tides | No | No | Yes |
| Meteo | WRF12km | WRF12km | WRF12km |

Table 2: ROMS domains configuration summary.

Results

As seen in Figure 3, the northeasterly flowing SC clearly bifurcates after crossing the 44.42° S parallel. Some of the water recirculates into PB in the form of an anticyclonic eddy. There is a clear difference in the magnitude of the mean currents predicted by the different runs. The average flow from CONTROL explained 80% of the observations, while TEST, only 33%.




Figure 5: Comparison of the CONTROL and TEST experiments with *in situ* depth-averaged current measurements.

Plume propagation

A passive tracer point source was introduced in both experiments at a constant rate at the surface during 15 days, and then halted and allowed to evolve for 15 additional days. The plume extends over a greater spatial extent on CONTROL, especially in the SE direction. There is also more northward dispersion in CONTROL. The TEST run overestimates concentrations near the source and underestimates at larger distances.




Figure 6: (a,b): First 15 day average tracer concentration. (d,e): Daily-averaged tracer concentration 15 days after source halt. (c,f): [TEST - CONTROL] for each case.

Conclusions

- 1 The SC has a substantial influence in the circulation at PB, accounting for more than half of the total currents at times and controlling most of its variability.
- 2 The absence of proper open ocean forcing greatly affects model performance for this particular area.
- 3 Underestimating the SC implies in less efficient dispersion of passive tracer plumes at the PB inner shelf.

Acknowledgements

We thank the Lyttelton Port of Christchurch for making the measurements available and the ROMS scientific community.

Contact Information

- http://www.metoceanview.com
- http://www.rsoutelino.com
- Email: r.soutelino@metocean.co.nz