

Cell Forcing Chains

From sudokuwiki.org, the puzzle solver's site


2		
	3	6
5		7

Forcing Chains are a powerful extension of chain strategies. Given that a cell or unit contains certain numbers, and that these are the only alternatives, we can show that certain candidates must be eliminated. Specifically we show that **all** of the alternatives must eliminate the target. The chains *force* the removal as no alternative exists.

Dual Cell Forcing Chains

These are the different types of eliminations available from Cell Forcing Chains. The diagram shows starting cells with two candidates but the same types and principles apply to cells with three (Triple) or four (Quad) candidates - just add more chains

Cell Forcing Chains start with all cells ON in a cell



Cell Forcing Chain Family

In the first example, we can eliminate 3 in **E2** because of the numbers left in **G9**, the 8 and 9 there. The small purple chain shows that with two 9s in row G and column 2 a 9 in **G9** places a 9 in in **E2** and therefore removes the 3. The chain is +9[G9]-9[G2]+9[E2]-3[E2].

The 8 in **G9** takes a little longer but again it is just a matter of following pairs in units. We remove the 8 in **A9** giving 2 which forces a 3 in **A2** and that removes the 3 in **E2**.

Since we are arguing from a single cell and the cell has two candidates, this is the simplest Forcing Chain, a Dual Cell Forcing Chain.

	1	2	3	4	5	6	7	8	9
A	2 3 4	2 3 4 7	6	2 3 4 7	1	9	5	7 8	2 8
B	9	1 2 5	7	2 5	6	8	1 2	4	3
C	1 2 3 4	8	1 5	2 4 5	2 3 5	3 7	1 2 7	6 9	6 9
D	8	2 3 5 6	4	1	9	3 7	2 7	3 5 6	2 5 6
E	1 2 3 7	1 2 3 7	1 9	6	4	5	8	1 3 7	1 2
F	1 3 7	1 3 6	1 3 5	8	3 7	2	9	1 3 5 6	4
G	1 3	1 3	1 3	5 7	5 7	6	4	2	8 9
H	5	4	8 9	2 3	2 3	1	6	8 9	7
J	7 6	7 6	2	9	8	4	3	1 5	1 5

Dual Cell Forcing Chain : [Load Example](#) or : [From the Start](#)

Note: Dual Cell Forcing Chains are not found by the solver because the solver will find Digit Forcing Chains first. The reason is that one of the two candidates that would make a Dual Cell Forcing Chain would be part of the chain on the Digit Forcing Chain. In this example perhaps the 2 in **A2** would be considered first. Then $-2[A2]$ would imply $+3[A2]$ and the chain continues with the same consequences. Dual Cell Forcing Chains are documented here as an alternative approach as part of a family sequence. The example above can be artificially induced by unticking AIC and DFC in the solver.

Triple Cell Forcing Chains

So if bi-value cells can work this magic, let's go up an order of complexity and test a Triple Cell Forcing Chain. This one starts on **D1**. The first chain, in blue, is another short one-shot. 3 removes 3 from **D7**. Forcing chains don't always need a nice first deduction, all three chains could be very long.

The second chain $+5[D1]-5[D8]+3[D8]-3[D7]$ is a little more involved. The 5 in **D1** must remove the 3 in **D7**. 5 will force a 3 in **D8** which does the trick.

Now to force 9 in **D1** to remove the 3 in **D7** we have to use quite a complex chain. It reads $+9[D1]-9[H1]+9[H9]-9[E9]+3[E9|D8]-3[D7]$. The curly brackets is my symbol for an ALS (Almost Locked Set). *Documentation forthcoming.*

	1	2	3	4	5	6	7	8	9
A	5 ³	9	2	4	5 ₇	5 6 ₇	3 ₆	1	8
B	8	4	6	3	1	2	5	9	7
C	1	5 ³ ₇	5 ₇	9	5 ₈	5 6 ₈	4 ₃ 6 ₄	4 ₃ 6 ₆	2
D	5 ³ ₉	2 3	3	6	5 ₇	1	2 ₃ 3 ₇	5 ₉	4
E	6	1	5 ³ ₉	2 ₅	4	5 ₇	2 3	8	5 ₉
F	7	2 3 ₅	4	8	2 3 ₅	9	1 2 3 ₆	3 ₅ 6	1 ₅ 3
G	2	5 ₇ 8	5 ₇ 8 9	1 ₇	5 ₇ 8 9	3	1 ₄	4 5 ₉	6
H	5 ³ ₉	5 ₇	1	5 ₇	6	4	8	2	5 ₉
J	4	6	3 ₅ 8 9	1 2 ₅	2 ₅ 8 9	5 ₈	1 3 ₉	7	1 ₅ 3 ₉

Triple Cell Forcing Chain : [Load Example](#) or : [From the Start](#)

In red, the chain starts with a 9 in **D1** (+9). Removing the 9 in **H1** means the 9 must be in **H9** as it's the last in the row. If that is the case we must remove the 9 from **E9**. **E9** is interesting as it forms part of an Almost Locked Set with **D8**. An ALS is a set of N cells with N+1 candidates. In this case {3,5,9}. But by removing the 9 from **E9** we have reduced an Almost Locked Set to a **Locked Set**. A 2-cell locked set is a Naked Pair. 3 and 5 must exist in **D8** and **E9** - we just don't know which way round yet. If this is the case then 3 can't exist in **D7**.

Viola, we have forced 3 from **D7** using the contents of **D1**.

The second triple example is nicely complex one. It's improbable that you might detect it using pen and paper, but I include it to show the reach of the Forcing Chains. We show that the content of **J7** can remove the 7 in **B3**. The blue chain is easy. A 1 one **J7** leads to a 7 in **B7** and therefore no 7 in **B3**. Hence $+1[J7]-1[B7]+7[B7]-7[B3]$.

The purple chain, $+7[J7]-7[J1]+7[G3|H3]-7[B3]$ contains an interesting group, symbolised by $+7[G3|H3]$. All this means is that if **J1** is not a 7 then **G3** OR **H3** must be. If one of those must be 7 we can look along their alignment - the columns, and remove 7s, eg 7 in **B3**.

The last chain, $+9[J7]-9[H7]+9[H3]$

$-9[C3]+7\{C3|A1\}-7[B3]$ contains an ALS, which I described above. We have a {1,7,9} in two cells **A1** and **C3**. Removing 9 from **B3** leaves a locked set of 1/7 in that box. All other 1s and 7s can be removed, namely the 7 in **B3**.

Quad Cell Forcing Chains



Complex Triple Cell Forcing Chain : [Load Example](#) or : [From the Start](#)

Logic relentlessly obliges us to consider the Quad formation, although I have not and will not look for higher order forcing chains, although they are possible. Quads are pretty rare. I found 6 in a random stock of forty thousand puzzles.

In this example, the cell **D3** contains {1,2,4,5}. By a miracle were any one of these the solution (and one must be) we can show that there is no 1 in **F2**, no really giving us a great deal of help with the rest of the puzzle. Although somewhat crowded when overlain, all four chains are relatively simple - just one ALS in the green chain.

If there is a 1 in **D3** it follows that

$$+1[D3]-1[F2]$$

If there is a 2 in **D3** it follows that


$$+2[D3]-2[D9]+2[A9]-2[A2]+1[A2]-1[F2]$$

If there is a 4 in **D3** it follows that

$$+4[D3]-4[E1]+1[E1]-1[F2]$$

If there is a 5 in **D3** it follows that

$$+5[D3]-5[E2]+1[E2]-1[F2].$$



A Quad Cell Forcing Chain : [Load Example](#) or : [From the Start](#)

Back to [Digit Forcing Chains](#)

Continue to [Unit Forcing Chains...](#)

Go back to [Nishio Forcing Chains](#) Continue to [Unit Forcing Chains](#)

2		
	3	6
5		7