

- 2.21** Derive the normalization constant c_n in (2.5.28) by deriving the orthogonality relationship (2.5.29) using generating functions. Start by working out the integral

$$I = \int_{-\infty}^{\infty} g(x,t)g(x,s)e^{-x^2} dx,$$

and then consider the integral again with the generating functions in terms of series with Hermite polynomials.

- 2.24** Consider a particle in one dimension bound to a fixed center by a δ -function potential of the form

$$V(x) = -v_0\delta(x), \quad (v_0 \text{ real and positive}).$$

Find the wave function and the binding energy of the ground state. Are there excited bound states?

- 2.27** Derive an expression for the density of free-particle states in *two* dimensions, normalized with periodic boundary conditions inside a box of side length L . Your answer should be written as a function of k (or E) times $dE d\phi$, where ϕ is the polar angle that characterizes the momentum direction in two dimensions.

- 2.30** Using spherical coordinates, obtain an expression for \mathbf{j} for the ground and excited states of the hydrogen atom. Show, in particular, that for $m_l \neq 0$ states, there is a circulating flux in the sense that \mathbf{j} is in the direction of increasing or decreasing ϕ , depending on whether m_l is positive or negative.

- 2.33** The propagator in momentum space analogous to (2.6.26) is given by $\langle \mathbf{p}'', t | \mathbf{p}', t_0 \rangle$. Derive an explicit expression for $\langle \mathbf{p}'', t | \mathbf{p}', t_0 \rangle$ for the free-particle case.

- 2.36** Show that the wave-mechanical approach to the gravity-induced problem discussed in Section 2.7 also leads to phase-difference expression (2.7.17).