

BOUNDEDNESS QUESTIONS FOR CALABI-YAU'S

- ① GENTLE INTRO TO THE MMP
- ② BOUNDEDNESS
- ③ - ④ BOUNDEDNESS for CY's (elliptic)
+ log CY pairs

MINIMAL MODEL PROGRAM : A GENTLE INTRODUCTION

ONE OF THE MAIN GOALS OF ALGEBRAIC GEOMETRY
IS TO PRODUCE A COMPLETE CLASSIFICATION OF
PROJECTIVE VARIETIES.

MINIMAL MODEL PROGRAM : A GENTLE INTRODUCTION

ONE OF THE MAIN GOALS OF ALGEBRAIC GEOMETRY
IS TO PRODUCE A COMPLETE CLASSIFICATION OF
PROJECTIVE VARIETIES. TO THIS END, WE CAN
PARTITION VARIETIES INTO EQUIVALENCE CLASSES USING
2 POSSIBLE EQUIVALENCE RELATIONS :

① ISOMORPHISM : $X \xrightleftharpoons[f]{f^{-1}} Y$

② BIRATIONAL : $X \xrightleftharpoons[g]{g^{-1}} Y$
 ISOMORPHISM g, g^{-1} are
 rational
 maps -

$\mathbb{C}(X) =$ field of rat'l
 functions on X

MINIMAL MODEL PROGRAM : A GENTLE INTRODUCTION

X SMOOTH PROJECTIVE VARIETY / \mathbb{C}

CANONICAL BUNDLE: $\det(\Omega_X^1)$


CANONICAL DIVISOR: Any Weil divisor K_X s.t.
 $\mathcal{O}_X(K_X) \cong$ canonical bundle.

MINIMAL MODEL PROGRAM : A GENTLE INTRODUCTION

X SMOOTH PROJECTIVE VARIETY / \mathbb{C}

CONJECTURE

STARTING FROM X , THERE EXISTS AN
ALGORITHMIC WAY OF CONSTRUCTING A
DIAGRAM OF THE FOLLOWING FORM



SUCH THAT f SATISFIES
1 OF THE FOLLOWING:

- ① f is BIRATIONAL & K_2 is ample Z is a can model.
- ② f is a FIBRATION & $K_{X'} = f^* H$ ample on Z
- ③ f is a FIBR. & $-K_{X'}|_{f^{-1}(F)}$ is ample on $f^{-1}(F)$

MINIMAL MODEL PROGRAM : A GENTLE INTRODUCTION

HENCE IN THE BIRATIONAL CLASSIFICATION
OF ALGEBRAIC VARIETIES WE HAVE 3
IMPORTANT CLASSES OF VARIETIES THAT
ACT AS BUILDING BLOCKS

- ① CANONICAL MODELS (K is ample)
- ② K -TRIVIAL VAR'S ($K \equiv 0$)
- ③ FANO VAR'S ($-K$ is ample)

MINIMAL MODEL PROGRAM : A GENTLE INTRODUCTION

THE 2nd PART OF THE CLASSIFICATION AIMS
THEN TO CLASSIFY THOSE VARIETIES THAT BELONG
TO ONE OF THE 3 BUILDING BLOCKS


① CANONICAL MODELS (K IS AMPLE)

② K -TRIVIAL VAR'S ($K \equiv 0$)


③ FANO VAR'S (- K IS AMPLE)

MODULI SPACES :

MINIMAL MODEL PROGRAM : A GENTLE INTRODUCTION



MINIMAL MODEL PROGRAM : A GENTLE INTRODUCTION



IN ORDER TO COMPLETE SUCH CLASSIFICATION SCHEME , WE NEED ALSO TO WORK WITH SINGULAR VARIETIES .

MINIMAL MODEL PROGRAM : A GENTLE INTRODUCTION

X NORMAL PROJECTIVE VARIETY / \mathbb{C}

CANONICAL BUNDLE: $i_* \det(\Omega^1_{X^{sm}})$ rank 1 - sheaf

$$i: X^{sm} \hookrightarrow X$$

CANONICAL DIVISOR: Zer. closure of any

$$K_{X^{sm}}$$

K_X will be a Weil divisor
(not. nec. Cartier)

MINIMAL MODEL PROGRAM : A GENTLE INTRODUCTION

DEFINITION

- ① A PAIR IS THE DATUM OF :
- (i) X a normal variety
 - (ii) B an effective Weil divisor with coefficients $\in (0, 1]$
- ② A LOG PAIR IS A PAIR (X, B) SUCH THAT
 $K_X + B$ IS \mathbb{R} -CARTIER
(can be written as a \mathbb{R} -sum)
of Cartier divisors


MINIMAL MODEL PROGRAM : A GENTLE INTRODUCTION

WHY IS IT USEFUL TO CONSIDER LOG PAIRS?

MINIMAL MODEL PROGRAM : A GENTLE INTRODUCTION

WHY IS IT USEFUL TO CONSIDER LOG PAIRS?

CONSIDER FOR EXAMPLE



$$C = V(x_0x_1 - x_2^2) \subseteq \mathbb{P}^3, \quad H \subseteq \mathbb{P}^3 \text{ hyperpl.}$$

C is SING. at v

$$2L \in |H|_C \quad [2L = V(x_0, x_0x_1 - x_2^2)]$$

L is NOT CARTIER AT v

ADJ. FORM:


$$\begin{array}{ll} X & \text{SMOOTH} \\ D & \text{SMOOTH} \\ \text{codim } 1 & \end{array}$$

$$(K_X + D)|_D = K_D$$

MINIMAL MODEL PROGRAM : A GENTLE INTRODUCTION

WHY IS IT USEFUL TO CONSIDER LOG PAIRS?

CONSIDER FOR EXAMPLE



$$C = V(x_0x_1 - x_2^2) \subseteq \mathbb{P}^3$$

$H \subseteq \mathbb{P}^3$ hyperpl.
 $2L \sim H|_c$

ADJ. FORM:

X	SMOOTH
D	SMOOTH
with 1	

$$(K_X + D)|_D = K_D$$

$$(K_{C \setminus \{v\}} + L|_{C \setminus \{v\}})|_L = K_{L|_{C \setminus \{v\}}}$$

$$\begin{aligned} (K_C + L) \cdot L &= -\frac{3}{2} \\ (-2H|_c + \frac{1}{2}H|_c) \cdot L &= \end{aligned} \implies (K_C + L)|_L = K_L + \frac{1}{2}v$$

MINIMAL MODEL PROGRAM : A GENTLE INTRODUCTION

WHY IS IT USEFUL TO CONSIDER LOG PAIRS?

CONSIDER A FIBRATION IN K-TRIVIAL VARIETIES

$$f : Y \longrightarrow X$$

MINIMAL MODEL PROGRAM : A GENTLE INTRODUCTION

WE SAW THAT WE WILL NEED TO CONSIDER SINGULAR OBJECTS.
TO DO THAT, WE NEED TO MEASURE THEIR SINGULARITIES.
WE WILL DO SO USING (BIRATIONAL) RESOLUTIONS.

MINIMAL MODEL PROGRAM : A GENTLE INTRODUCTION

WE SAW THAT WE WILL NEED TO CONSIDER SINGULAR OBJECTS.
TO DO THAT, WE NEED TO MEASURE THEIR SINGULARITIES.
WE WILL DO SO USING (BIRATIONAL) RESOLUTIONS.

DEFINITION LET (X, B) BE A PAIR. A BIRAT'L MORPHISM

$$\pi: X' \longrightarrow X$$

IS A LOG RESOLUTION WHEN :

① π IS A PROPER RESOLUTION OF X

② $\tilde{B} + \text{Exc}(\pi)$ is

strict
transform
of B on X'

SIMPLE
NORMAL
CROSSING

S
N
C

$\forall x' \in X$
locally analytically
 $(X', (\tilde{B} + \text{Exc}(\pi)))$
 $(\mathbb{C}^n, \sum_{i=1}^k \{z_i = 0\})$

MINIMAL MODEL PROGRAM : A GENTLE INTRODUCTION

WE SAW THAT WE WILL NEED TO CONSIDER SINGULAR OBJECTS.
TO DO THAT, WE NEED TO MEASURE THEIR SINGULARITIES.
WE WILL DO SO USING (BIRATIONAL) RESOLUTIONS.

DEFINITION LET (X, B) BE A PAIR. A BIRAT'L MORPHISM

$$\pi : X' \longrightarrow X$$

IS A LOG RESOLUTION WHEN :

- ① X' IS SMOOTH;
- ② $(X', \tilde{B} + \text{Exc}(\pi))$ IS SNC.

THEOREM [HIRONAKA] LOG RESOLUTION EXIST IN CHAR 0.

MINIMAL MODEL PROGRAM : A GENTLE INTRODUCTION

DEFINITION

LET (X, B) BE A PAIR. A BIRAT'L MORPHISM


$$\pi : X' \longrightarrow X$$

IS A LOG RESOLUTION WHEN :

- ① X' IS SMOOTH ;
- ② $(X', \tilde{B} + \text{Exc}(\pi))$ IS SNC.

THEOREM [HIRONAKA] LOG RESOLUTION EXIST IN CHAR 0.

EXAMPLE



MINIMAL MODEL PROGRAM : A GENTLE INTRODUCTION

DEFINITION

LET (X, B) BE A PAIR. A BIRAT'L MORPHISM


$$\pi : X' \longrightarrow X$$

IS A LOG RESOLUTION WHEN :

- ① X' IS SMOOTH ;
- ② $(X', \tilde{B} + \text{Exc}(\pi))$ IS SNC.

THEOREM [HIRONAKA] LOG RESOLUTION EXIST IN CHAR 0.

EXAMPLE



MINIMAL MODEL PROGRAM : A GENTLE INTRODUCTION

DEFINITION

LET (X, B) BE A PAIR. A BIRAT'L MORPHISM


$$\pi : X' \longrightarrow X$$

IS A LOG RESOLUTION WHEN :

- ① X' IS SMOOTH ;
- ② $(X', \tilde{B} + \text{Exc}(\pi))$ IS SNC.

THEOREM [HIRONAKA] LOG RESOLUTION EXIST IN CHAR 0.

EXAMPLE



MINIMAL MODEL PROGRAM : A GENTLE INTRODUCTION

DEFINITION

LET (X, B) BE A PAIR. A BIRAT'L MORPHISM


$$\pi : X' \longrightarrow X$$

IS A LOG RESOLUTION WHEN :

- ① X' IS SMOOTH ;
- ② $(X', \tilde{B} + \text{Exc}(\pi))$ IS SNC.

THEOREM [HIRONAKA] LOG RESOLUTION EXIST IN CHAR 0.

EXAMPLE



MINIMAL MODEL PROGRAM : A GENTLE INTRODUCTION

DEFINITION

LET (X, B) BE A PAIR. A BIRAT'L MORPHISM


$$\pi: X' \longrightarrow X$$

IS A LOG RESOLUTION WHEN :

- ① X' IS SMOOTH;
- ② $(X', \tilde{B} + \text{Exc}(\pi))$ IS SNC.

THEOREM [HIRONAKA] LOG RESOLUTION EXIST IN CHAR 0.

EXAMPLE



MINIMAL MODEL PROGRAM : A GENTLE INTRODUCTION

DEFINITION

LET (X, B) BE A PAIR. A BIRAT'L MORPHISM


$$\pi : X' \longrightarrow X$$

IS A LOG RESOLUTION WHEN :

- ① X' IS SMOOTH ;
- ② $(X', \tilde{B} + \text{Exc}(\pi))$ IS SNC.

THEOREM [HIRONAKA] LOG RESOLUTION EXIST IN CHAR 0.

EXAMPLE



MINIMAL MODEL PROGRAM : A GENTLE INTRODUCTION

WE SAW THAT WE WILL NEED TO CONSIDER SINGULAR OBJECTS.
TO DO THAT, WE NEED TO MEASURE THEIR SINGULARITIES.
WE WILL DO SO USING (BIRATIONAL) RESOLUTIONS.

GIVEN (X, B) A LOG PAIR AND A LOG RESOL.
 $\pi: X' \longrightarrow X$

WE CAN ALWAYS WRITE IN A UNIQUE WAY

$$K_{X'} + \tilde{B} = \pi^*(K_X + B) + \sum_{\text{exc div.}} a_i E_i$$

MINIMAL MODEL PROGRAM : A GENTLE INTRODUCTION

WE SAW THAT WE WILL NEED TO CONSIDER SINGULAR OBJECTS.
TO DO THAT, WE NEED TO MEASURE THEIR SINGULARITIES.
WE WILL DO SO USING (BIRATIONAL) RESOLUTIONS.

GIVEN (X, B) A LOG PAIR AND A LOG RESOL.

$$\pi: X' \longrightarrow X$$

WE CAN ALWAYS WRITE IN A UNIQUE WAY

$$K_{X'} + B' = \pi^*(K_X + B)$$

$$\text{WHERE } B' = \pi_*^{-1} B + \sum_{\substack{\text{exc.} \\ \text{div.}}} a_i E_i$$

DEFINITION LET $D \subseteq X'$ BE A PRIME DIVISOR, THEN

$$\alpha(D; X, B) := 1 - \text{coeff. of } D \text{ in } B'$$

↑
log discrepancy

MINIMAL MODEL PROGRAM : A GENTLE INTRODUCTION

GIVEN (X, B) A LOG PAIR AND A LOG RESOL.

$$\pi: X' \longrightarrow X$$

WE CAN ALWAYS WRITE IN A UNIQUE WAY

$$K_{X'} + B' = \pi^*(K_X + B)$$

$$\text{WHERE } B' = \pi_*^{-1} B + \sum_{\substack{\text{exc.} \\ \text{div.}}} a_i E_i$$

DEFINITION ① LET $D \subseteq X'$ BE A PRIME DIVISOR, THEN

$$a(D; X, B) := 1 - \text{coeff. of } D \text{ in } B'.$$

[THIS IS CALLED THE LOG DISCREPANCY OF D wrt (X, B)]

② THE TOTAL LOG DISCREPANCY OF (X, B) is

$$a(X, B) := \inf a(D; X, B)$$

MINIMAL MODEL PROGRAM : A GENTLE INTRODUCTION

DEFINITION ① LET $D \subseteq X'$ BE A PRIME DIVISOR, THEN

$$a(D; X, B) := 1 - \text{coeff. of } D \text{ in } B'.$$

[THIS IS CALLED THE LOG DISCREPANCY OF D wrt (X, B)]

② THE TOTAL LOG DISCREPANCY OF (X, B) is

$$a(X, B) := \inf a(D; X, B)$$

SPECIAL DICHOTOMY: $a(X, B) = -\infty$

OR

$$a(X, B) \geq 0 \quad \text{MMP} \quad \smiley$$

MINIMAL MODEL PROGRAM : A GENTLE INTRODUCTION

DEFINITION LET (X, B) BE A LOG PAIR.

THEN (X, B) IS

LOG CANONICAL


KLT
LOG TERM.

CANONICAL

TERMINAL


$a(X, B) \geq 0$
 $a(X, B) > 0$
 $a(X, B) \geq -1$
 $a(X, B) > -1$

EXAMPLE



C_n = CONE OVER RAT'L NORMAL CURVE OF $\deg n$

EXAMPLE




MINIMAL MODEL PROGRAM : A GENTLE INTRODUCTION

~~X SMOOTH PROJECTIVE VARIETY / C~~ (X, B) LOG CANONICAL PAIR

CONJECTURE

STARTING FROM (X, B) , THERE EXISTS AN ALGORITHMIC WAY OF CONSTRUCTING A DIAGRAM OF THE FOLLOWING FORM

$X \xrightarrow{\text{composition of special birat'l maps}} (X', B')$



SUCH THAT f SATISFIES
1 OF THE FOLLOWING :

① f BIRAT'L & $K_{X'} + B'$ is AMPLE

② f FIBRATION &
 $K_X + B' = f^* H$ ample

③ f FIBRATION &
 $- (K_X + B')$ AMPLE on fibers

MINIMAL MODEL PROGRAM : A GENTLE INTRODUCTION

IN ORDER TO ACHIEVE SUCH RESULT , WE WILL TRY
TO CONTROL & TUNE THE POSITIVITY/NEGATIVITY OF
LOG DIVISORS .

(X, B) LOG PAIR

$K_X + B$ is \mathbb{R} -Cartier

$\forall C \subseteq X$ proper curve

$$(K_X + B) \cdot C$$

GOAL : either make $K_X + B$ nef after many
birational transformations

OR

die trying (but in finite time)

$$K_X + B \cdot C \geq 0$$

$\forall C \subseteq X$ proper curve

THE CONE THEOREM

THEOREM [Mori, Kollar, Shokurov, ...]

LET (X, B) BE A LOG CANONICAL PAIR.

WE HAVE THE FOLLOWING DECOMPOSITION

$$\overline{\text{NE}}(X) = \overline{\text{NE}}(X)_{K_X + B \geq 0} + \overline{\text{NE}}(X)_{K_X + B < 0}$$

the closure the one of
effective
cones

THE CONE THEOREM

THEOREM

LET (X, B) BE A LOG CANONICAL PAIR.

WE HAVE THE FOLLOWING DECOMPOSITION

$$\overline{\text{NE}}(X) = \overline{\text{NE}}(X)_{K_X+B \geq 0} + \overline{\text{NE}}(X)_{K_X+B < 0} =$$

$$= \overline{\text{NE}}(X)_{K_X+B \geq 0} + \sum_{i \in I} R_i + [C_i]$$

extremal rays
rat'l curves
 $-2\dim X \leq (K_X+B) \cdot C_i < 0$


locally discrete away from $(K_X+B)^\perp$

countable set

HORIZONTAL SLICE OF $\overline{\text{NE}}(x)$

extremal rays $\stackrel{\text{def}}{=} v_1 + v_2 \in R_i$

$$\downarrow \\ v_1, v_2 \in R_i$$



THE CONE THEOREM

THEOREM

LET (X, B) BE A LOG CANONICAL PAIR.

WE HAVE THE FOLLOWING DECOMPOSITION

$$\overline{\text{NE}}(X) = \overline{\text{NE}}(X)_{K_X+B \geq 0} + \overline{\text{NE}}(X)_{K_X+B < 0} =$$
$$= \overline{\text{NE}}(X)_{K_X+B \geq 0} + \sum_{i \in I} [R_i + [C_i]]$$

*R_i: extremal ray
rat'l curves
at most countable set
 $-2\dim X \leq (K_X+B) \cdot C_i < 0$*

MOREOVER, $\forall i \in I, \exists \text{cont}_{R_i}: X \rightarrow \mathbb{Z}_i$ SUCH THAT

cont_{R_i} HAS CONNECTED FIBERS & IF $\text{cont}_{R_i}(C) = \text{pt} \Rightarrow [C] \in R_i$.