

The structure of algebraic varieties: birational classification and boundedness questions

Roberto Svaldi

University of Cambridge

École Polytechnique Fédérale de Lausanne
September 26, 2018

Geometries

Klein's & Grothendieck's idea

A geometry is the datum of:

- A class \mathcal{C} of **objects** that we would like to classify, and
- **Functions** on these objects + **maps** between different objects in \mathcal{C} .

Geometries

Klein's & Grothendieck's idea

A geometry is the datum of:

- A class \mathcal{C} of **objects** that we would like to classify, and
- **Functions** on these objects + **maps** between different objects in \mathcal{C} .

Topology \longleftrightarrow sets & topologies + C^0 functions & maps

Differential Geometry \longleftrightarrow manifolds + C^∞ functions & maps

Analytic Geometry \longleftrightarrow \mathbb{C} -manifolds + C^ω functions & maps

Algebraic Geometry \longleftrightarrow alg. varieties + polynomial functions & maps

Algebraic geometry: rigid as there are fewer functions;
powerful as we can use algebra & geometry together.

Algebraic Varieties

Let k be a field, e.g., $k = \mathbb{Q}, \mathbb{R}, \mathbb{C}$.

$k[x_1, \dots, x_n]$ = ring of polynomials on the variables t_i with coeffs in k .

Algebraic Varieties

Let k be a field, e.g., $k = \mathbb{Q}, \mathbb{R}, \mathbb{C}$.

$k[x_1, \dots, x_n]$ = ring of polynomials on the variables t_i with coeffs in k .

(Affine) algebraic varieties are sets that are defined as simultaneous zeros of polynomial equations:

$$X = \{(a_1, \dots, a_n) \in k^n \mid f_1(a_1, \dots, a_n) = \dots = f_k(a_1, \dots, a_n) = 0\}$$

for a finite collection $f_1, \dots, f_k \in k[x_1, \dots, x_n]$.

Algebraic Varieties

Let k be a field, e.g., $k = \mathbb{Q}, \mathbb{R}, \mathbb{C}$.

$k[x_1, \dots, x_n]$ = ring of polynomials on the variables t_i with coeffs in k .

(Affine) algebraic varieties are sets that are defined as simultaneous zeros of polynomial equations:

$$X = \{(a_1, \dots, a_n) \in k^n \mid f_1(a_1, \dots, a_n) = \dots = f_k(a_1, \dots, a_n) = 0\}$$

for a finite collection $f_1, \dots, f_k \in k[x_1, \dots, x_n]$.

We will consider **irreducible** varieties, i.e. those that cannot further decomposed into unions of affine varieties.


Algebraic Varieties, II


Question

What does X look like? For example is $X \neq \emptyset$?


Consider $f(x_1, x_2) = x_1^2 + x_2^2 + 1 \in k[x_1, x_2]$. $X := \{f = 0\}$

If $k = \mathbb{R}$ then X is empty, while if $k = \mathbb{C}$ then X contains infinitely many points – it is a sphere minus 2 points.






$$f(x_1, x_2) = x_1^2 + x_2^2 + 1$$



$$f(x_1, x_2) = x_1^2 + x_2^2 - 1$$



$$f(x_1, x_2) = x_1^2 + x_2^2$$

Algebraic Varieties, II

Question

What does X look like? For example is $X \neq \emptyset$?

Consider $f(x_1, x_2) = x_1^2 + x_2^2 + 1 \in k[x_1, x_2]$. $X := \{f = 0\}$

If $k = \mathbb{R}$ then X is empty, while if $k = \mathbb{C}$ then X contains infinitely many points.

In general, it is a hard problem to decide whether a variety X is non-empty.

Algebraic Varieties, II

Question

What does X look like? For example is $X \neq \emptyset$?

Consider $f(x_1, x_2) = x_1^2 + x_2^2 + 1 \in k[x_1, x_2]$. $X := \{f = 0\}$

If $k = \mathbb{R}$ then X is empty, while if $k = \mathbb{C}$ then X contains infinitely many points.

In general, it is a hard problem to decide whether a variety X is non-empty. For example, consider $X = \{t_1^m + t_2^m - 1 = 0\}$, $m \in \mathbb{N}$. When $k = \mathbb{Q}$, deciding this is equivalent to Fermat Last Theorem.

This problem becomes much simpler if k is algebraically closed, e.g. $k = \mathbb{C}$. So, from now on we will assume that $k = \mathbb{C}$.

Rational functions and maps

Given an affine variety $X \subset \mathbb{C}^n$, each $p \in \mathbb{C}[x_1, \dots, x_n]$ defines a **function** on X .

Rational functions and maps

Given an affine variety $X \subset \mathbb{C}^n$, each $p \in \mathbb{C}[x_1, \dots, x_n]$ defines a **function** on X .

Moreover, if we take $s, t \in \mathbb{C}[x_1, \dots, x_n]$ and $t \not\equiv 0$ on X , then $\frac{s}{t}$ defines a **rational function** which is well defined at all points $x \in X$ where $h(x) \neq 0$.

Rational functions and maps

Given an affine variety $X \subset \mathbb{C}^n$, each $p \in \mathbb{C}[x_1, \dots, x_n]$ defines a **function** on X .

Moreover, if we take $s, t \in \mathbb{C}[x_1, \dots, x_n]$ and $t \not\equiv 0$ on X , then $\frac{s}{t}$ defines a **rational function** which is well defined at all points $x \in X$ where $h(x) \neq 0$.

The set of rational functions on an affine variety X forms a field $\mathbb{C}(X)$, the **field of rational functions** of X .

Rational functions and maps

Given an affine variety $X \subset \mathbb{C}^n$, each $p \in \mathbb{C}[x_1, \dots, x_n]$ defines a **function** on X .

Moreover, if we take $s, t \in \mathbb{C}[x_1, \dots, x_n]$ and $t \not\equiv 0$ on X , then $\frac{s}{t}$ defines a **rational function** which is well defined at all points $x \in X$ where $h(x) \neq 0$.

The set of rational functions on an affine variety X forms a field $\mathbb{C}(X)$, the **field of rational functions** of X .

Given $X \subset \mathbb{C}^n$, $Y \subset \mathbb{C}^m$ affine varieties a **rational map** $\pi: X \dashrightarrow Y$ is

$$x \mapsto y = (h_1(x), \dots, h_m(x)), \quad h_i \in \mathbb{C}(X), \text{ and } y \in Y.$$

Rational functions and maps

Given an affine variety $X \subset \mathbb{C}^n$, each $p \in \mathbb{C}[x_1, \dots, x_n]$ defines a **function** on X .

Moreover, if we take $s, t \in \mathbb{C}[x_1, \dots, x_n]$ and $t \not\equiv 0$ on X , then $\frac{s}{t}$ defines a **rational function** which is well defined at all points $x \in X$ where $h(x) \neq 0$.

The set of rational functions on an affine variety X forms a field $\mathbb{C}(X)$, the **field of rational functions** of X .

Given $X \subset \mathbb{C}^n$, $Y \subset \mathbb{C}^m$ affine varieties a **rational map** $\pi: X \dashrightarrow Y$ is

$$x \mapsto y = (h_1(x), \dots, h_m(x)), \quad h_i \in \mathbb{C}(X), \text{ and } y \in Y.$$

Rational maps that are defined at every point of X are called **morphisms**.

Examples

$$i: \mathbb{C} \rightarrow \mathbb{C}$$

$$t \mapsto \frac{t^2+1}{t}$$

i is rational but not a morphism!

$$\phi: \mathbb{C} \rightarrow \mathbb{C}$$

$$t \mapsto (t^3, t^2)$$

$$C := \{x_1^2 - x_2^3 = 0\} \subset \mathbb{C}^2$$

ϕ is a morphism!

$$c: \mathbb{C}^2 \rightarrow \mathbb{C}^2$$

$$(x_1, x_2) \mapsto \left(\frac{1}{x_1}, \frac{1}{x_2}\right)$$

c is rational but not a morphism!

$$c \circ c = \text{id}$$

Projective varieties

It is often easier to work with compact sets. Affine varieties are not compact.

Projective varieties

It is often easier to work with compact sets. Affine varieties are not compact. To compactify them, we need to introduce **projective space**

$$\mathbb{P}^n = \{(a_0 : \cdots : a_n) \mid a_i \in k, \text{ and } a_j \neq 0 \text{ for some } j \in \{0, \dots, n\}\}$$

Projective varieties

It is often easier to work with compact sets. Affine varieties are not compact. To compactify them, we need to introduce **projective space**

$$\mathbb{P}^n = \{(a_0 : \cdots : a_n) \mid a_i \in k, \text{ and } a_j \neq 0 \text{ for some } j \in \{0, \dots, n\}\}$$

with the condition that $(a_0 : \cdots : a_n) = (\lambda a_0 : \cdots : \lambda a_n)$ for any $\lambda \neq 0$.

Projective varieties

It is often easier to work with compact sets. Affine varieties are not compact. To compactify them, we need to introduce **projective space**

$$\mathbb{P}^n = \{(a_0 : \cdots : a_n) \mid a_i \in k, \text{ and } a_j \neq 0 \text{ for some } j \in \{0, \dots, n\}\}$$

with the condition that $(a_0 : \cdots : a_n) = (\lambda a_0 : \cdots : \lambda a_n)$ for any $\lambda \neq 0$.

A **projective variety** X in \mathbb{P}^n is defined as the set of simultaneous zeros

$$X = \{(a_0 : \cdots : a_n) \in \mathbb{P}^n \mid g_1(a_0, \dots, a_n) = \cdots = g_k(a_0, \dots, a_n) = 0\}$$

of homogenous polynomial equations $g_1, \dots, g_k \in k[x_0, \dots, x_n]$

Projective varieties, II

\mathbb{P}^n and projective varieties are compact and moreover

$$\mathbb{P}^n = \begin{matrix} k^n \\ (a_0 : \cdots : a_{n-1} : 1) \end{matrix} \cup \begin{matrix} \mathbb{P}^{n-1} \\ (b_0 : \cdots : b_{n-1} : 0) \end{matrix}$$

For example, $x_0^2 + x_1^2 + x_2^2 \in k[x_0, x_1, x_2]$
 defines a quadric, which is the
 compactification of the affine variety
 $X = (t_1^2 + t_2^2 + 1 = 0)$ that we saw earlier.



Projective varieties, II

\mathbb{P}^n and projective varieties are compact and moreover

$$\mathbb{P}^n = \begin{matrix} k^n \\ (a_0 : \cdots : a_{n-1} : 1) \end{matrix} \cup \begin{matrix} \mathbb{P}^{n-1} \\ (b_0 : \cdots : b_{n-1} : 0) \end{matrix}$$

For example, $x_0^2 + x_1^2 + x_2^2 \in k[x_0, x_1, x_2]$
 defines a quadric, which is the
 compactification of the affine variety
 $X = (t_1^2 + t_2^2 + 1 = 0)$ that we saw earlier.



Rational maps are now defined as defined as quotients of homogenous polynomials.

Projective varieties, II

\mathbb{P}^n and projective varieties are compact and moreover

$$\mathbb{P}^n = \begin{matrix} k^n & \cup & \mathbb{P}^{n-1} \\ (a_0 : \cdots : a_{n-1} : 1) & (b_0 : \cdots : b_{n-1} : 0) \end{matrix}$$

For example, $x_0^2 + x_1^2 + x_2^2 \in k[x_0, x_1, x_2]$
 defines a quadric, which is the
 compactification of the affine variety
 $X = (t_1^2 + t_2^2 + 1 = 0)$ that we saw earlier.



Rational maps are now defined as defined as quotients of homogenous polynomials.

We will consider from now on only projective varieties.

Equivalence of varieties

We say that two varieties are **isomorphic** if there are morphisms $f : X \rightarrow Y$ and $g : Y \rightarrow X$ that are inverse of each other.

Equivalence of varieties

We say that two varieties are **isomorphic** if there are morphisms $f : X \rightarrow Y$ and $g : Y \rightarrow X$ that are inverse of each other.

Definition

X and Y are **birational equivalent** if there are closed subsets $Z \subset X$ and $W \subset Y$ such that $X \setminus Z$ is isomorphic to $Y \setminus W$.

Equivalence of varieties

We say that two varieties are **isomorphic** if there are morphisms $f : X \rightarrow Y$ and $g : Y \rightarrow X$ that are inverse of each other.

Definition

X and Y are **birational equivalent** if there are closed subsets $Z \subset X$ and $W \subset Y$ such that $X \setminus Z$ is isomorphic to $Y \setminus W$.

X is **birational** to Y if and only if there are rational maps $f : X \dashrightarrow Y$ and $g : Y \dashrightarrow X$ that are inverse of each other.

Equivalence of varieties

We say that two varieties are **isomorphic** if there are morphisms $f : X \rightarrow Y$ and $g : Y \rightarrow X$ that are inverse of each other.


Definition

X and Y are **birational equivalent** if there are closed subsets $Z \subset X$ and $W \subset Y$ such that $X \setminus Z$ is isomorphic to $Y \setminus W$.

X is **birational** to Y if and only if there are rational maps $f : X \dashrightarrow Y$ and $g : Y \dashrightarrow X$ that are inverse of each other.

Being birational is an **equivalence relation**. Hence it partitions algebraic varieties into equivalence classes.

Cone vs. Cilinder



Birational equivalence, II

Birational equivalence is a much more flexible equivalence relation than isomorphism. Yet it retains a lot of informations about the structure of the original variety (e.g., holomorphic differential forms, fundamental group).

Birational equivalence, II

Birational equivalence is a much more flexible equivalence relation than isomorphism. Yet it retains a lot of informations about the structure of the original variety (e.g., holomorphic differential forms, fundamental group).

When we consider varieties up to birational equivalence, once we are given a variety X we are free to switch to a new variety X' birational to X , if X' has better geometric features which may simplify our task.

Birational equivalence, II

Birational equivalence is a much more flexible equivalence relation than isomorphism. Yet it retains a lot of informations about the structure of the original variety (e.g., holomorphic differential forms, fundamental group).

When we consider varieties up to birational equivalence, once we are given a variety X we are free to switch to a new variety X' birational to X , if X' has better geometric features which may simplify our task.

New Goal

Classify projective algebraic varieties up to birational equivalence.

Classifying algebraic varieties

As we want to classify algebraic varieties, we need to understand how exactly we are going to do that.

Classifying algebraic varieties

As we want to classify algebraic varieties, we need to understand how exactly we are going to do that.

It is not hard to prove that X, X' are birational if and only if $\mathbb{C}(X) \simeq \mathbb{C}(X')$.

Classifying algebraic varieties

As we want to classify algebraic varieties, we need to understand how exactly we are going to do that.

It is not hard to prove that X, X' are birational if and only if $\mathbb{C}(X) \simeq \mathbb{C}(X')$.

Question

Let X be a projective algebraic variety. Is there another algebraic variety X' such that $\mathbb{C}(X) \simeq \mathbb{C}(X')$ and the geometry of X' is as simple as possible?

Classifying algebraic varieties

As we want to classify algebraic varieties, we need to understand how exactly we are going to do that.

It is not hard to prove that X, X' are birational if and only if $\mathbb{C}(X) \simeq \mathbb{C}(X')$.

Question

Let X be a projective algebraic variety. Is there another algebraic variety X' such that $\mathbb{C}(X) \simeq \mathbb{C}(X')$ and the geometry of X' is as simple as possible?

What does **simple geometry** mean for X ?

Classifying algebraic varieties

As we want to classify algebraic varieties, we need to understand how exactly we are going to do that.

It is not hard to prove that X, X' are birational if and only if $\mathbb{C}(X) \simeq \mathbb{C}(X')$.

Question

Let X be a projective algebraic variety. Is there another algebraic variety X' such that $\mathbb{C}(X) \simeq \mathbb{C}(X')$ and the geometry of X' is as simple as possible?

What does **simple geometry** mean for X ?

We would like to have some way of measuring these improvements.

Plane Curves

In the projective plane \mathbb{P}^2 , consider the projective variety X_f given by a homogenous polynomial of degree d , $f(X_0, X_1, X_2) = \sum a_{ijk} X_0^i X_1^j X_2^k$, $i + j + k = d$.

Plane Curves

In the projective plane \mathbb{P}^2 , consider the projective variety X_f given by a homogenous polynomial of degree d , $f(X_0, X_1, X_2) = \sum a_{ijk} X_0^i X_1^j X_2^k$, $i + j + k = d$.

X_f is a 1-dimensional projective algebraic variety over the complex numbers. When X_f is smooth, it can be given the structure of a compact complex Riemann surface.

Plane Curves

In the projective plane \mathbb{P}^2 , consider the projective variety X_f given by a homogenous polynomial of degree d , $f(X_0, X_1, X_2) = \sum a_{ijk} X_0^i X_1^j X_2^k$, $i + j + k = d$.

X_f is a 1-dimensional projective algebraic variety over the complex numbers. When X_f is smooth, it can be given the structure of a compact complex Riemann surface.

Question


How does the structure of X_f vary with d ?

Projective curves, II


For smooth projective curves isomorphism and birational equivalence are the same.

Projective curves, II


For smooth projective curves isomorphism and birational equivalence are the same.



genus = 0



genus = 1

genus ≥ 2

$\deg(f)$	$g(X)$	Variety	Universal cover	Curvature
1, 2	0	\mathbb{P}^1	S^2	> 0
3	1	Elliptic curves	\mathbb{R}^2	$= 0$
> 3	≥ 2	Hyperbolic curves	$ x \leq 1 \subset \mathbb{R}^2$	< 0

Higher dimension

In dimension higher than 1 it is no longer true that isomorphism and birational equivalence are the same.

Higher dimension


In dimension higher than 1 it is no longer true that isomorphism and birational equivalence are the same.

The situation is much more complicated, because algebraic varieties cannot be described as easily as in the case of curves.

Higher dimension

In dimension higher than 1 it is no longer true that isomorphism and birational equivalence are the same.

The situation is much more complicated, because algebraic varieties cannot be described as easily as in the case of curves.



Smoothness

There is a notion of smoothness for algebraic varieties.

Smoothness

There is a notion of smoothness for algebraic varieties.

Roughly, an algebraic variety X is smooth at a point $x \in X$, if locally around x , X is close to being isomorphic to \mathbb{C}^n . This is similar to the notion of smoothness in differential geometry. In fact, a smooth algebraic variety has a natural structure of complex manifold (implicit function theorem).

Smoothness

There is a notion of smoothness for algebraic varieties.

Roughly, an algebraic variety X is smooth at a point $x \in X$, if locally around x , X is close to being isomorphic to \mathbb{C}^n . This is similar to the notion of smoothness in differential geometry. In fact, a smooth algebraic variety has a natural structure of complex manifold (implicit function theorem).

Hironaka's desingularization

Over \mathbb{C} , every algebraic variety is birational to a smooth variety.

Smoothness


There is a notion of smoothness for algebraic varieties.

Roughly, an algebraic variety X is smooth at a point $x \in X$, if locally around x , X is close to being isomorphic to \mathbb{C}^n . This is similar to the notion of smoothness in differential geometry. In fact, a smooth algebraic variety has a natural structure of complex manifold (implicit function theorem).

Hironaka's desingularization

Over \mathbb{C} , every algebraic variety is birational to a smooth variety.

Cone vs. Cilinder: desingularization



Building blocks

There are 3 classes of varieties whose features are similar to the case of curves.

Building blocks

There are 3 classes of varieties whose features are similar to the case of curves. These classes can be distinguished by looking at the behavior of holomorphic differential forms of top degree on them (canonical divisor) and a certain higher dimensional analogue of the curvature.

Building blocks

There are 3 classes of varieties whose features are similar to the case of curves. These classes can be distinguished by looking at the behavior of holomorphic differential forms of top degree on them (canonical divisor) and a certain higher dimensional analogue of the curvature.

For a smooth (or mildly singular) projective variety X

$$X \text{ is } \begin{cases} \mathbf{Fano} & \text{if it is positively curved} \\ \mathbf{Calabi-Yau} & \text{if it is flat} \\ \text{a } \mathbf{canonical \ model} & \text{if it is negatively curved} \end{cases}$$

Building blocks

There are 3 classes of varieties whose features are similar to the case of curves. These classes can be distinguished by looking at the behavior of holomorphic differential forms of top degree on them (canonical divisor) and a certain higher dimensional analogue of the curvature.

For a smooth (or mildly singular) projective variety X

$$X \text{ is } \begin{cases} \textbf{Fano} & \text{if it is positively curved} \\ \textbf{Calabi-Yau} & \text{if it is flat} \\ \text{a } \textbf{canonical model} & \text{if it is negatively curved} \end{cases}$$

In \mathbb{P}^n , if $X = \{f = 0\}$ for a homogenous polynomial of degree d (and X is smooth) then

$$X \text{ is } \begin{cases} \textbf{Fano} & \text{if} & \deg(f) \leq n - 1 \\ \textbf{Calabi-Yau} & \text{if} & \deg(f) = n \\ \text{a } \textbf{canonical model} & \text{if} & \deg(f) \geq n + 1 \end{cases}$$

Minimal Model Program

The Minimal Model Program is a conjectural prediction of how to develop a systematic classification of algebraic varieties.

Minimal Model Program

The Minimal Model Program is a conjectural prediction of how to develop a systematic classification of algebraic varieties.

Given a smooth projective variety X then

(1) $X \dashrightarrow X'$ pass to a birational model X' by making the curvature as negative as possible using a combination of pre-determined birational maps.

(2) $X' \longrightarrow Z$ X' is expected to have a morphism to another projective variety whose fibers are in one of the 3 building block. Then repeat this process starting with Z .

The MMP is not only a classification algorithm but it is a much more versatile tool.

Given a question for X , it can be adapted to help us to find a birational variety X' best suitable to the problem.

The MMP is known to work for projective varieties of dimension at most 3 by work of Mori (Fields Medal 1992), Kollár (Shaw Prize 2017) and many others. It is now known also to work in any dimension for varieties which are birational to canonical models by work of Shokurov, Birkar-Cascini-Hacon-McKernan and many others.



Figure: Mori



Figure: Kollár



Figure: Shokurov

Families of varieties

It is important to understand how varieties behave not only as standalone objects, but also in families.

Families of varieties

It is important to understand how varieties behave not only as standalone objects, but also in families. Consider the homogenous polynomials of degree 3 in 3 variables,

$$f(X_0, X_1, X_2) = a_1X_0^3 + a_2X_1^3 + a_3X_2^3 + a_4X_0X_1^2 + a_5X_0X_2^2 + a_6X_1X_0^2 + a_7X_1X_2^2 + a_8X_2X_0^2 + a_9X_2X_1^2 + a_{10}X_0X_1X_2.$$

Families of varieties

It is important to understand how varieties behave not only as standalone objects, but also in families. Consider the homogenous polynomials of degree 3 in 3 variables,

$$f(X_0, X_1, X_2) = a_1X_0^3 + a_2X_1^3 + a_3X_2^3 + a_4X_0X_1^2 + a_5X_0X_2^2 + a_6X_1X_0^2 + \\ + a_7X_1X_2^2 + a_8X_2X_0^2 + a_9X_2X_1^2 + a_{10}X_0X_1X_2.$$

We can parametrize these simply by looking at their coefficients $(a_1 : a_2 : \cdots : a_{10})$.

Families of varieties

It is important to understand how varieties behave not only as standalone objects, but also in families. Consider the homogenous polynomials of degree 3 in 3 variables,

$$f(X_0, X_1, X_2) = a_1X_0^3 + a_2X_1^3 + a_3X_2^3 + a_4X_0X_1^2 + a_5X_0X_2^2 + a_6X_1X_0^2 + a_7X_1X_2^2 + a_8X_2X_0^2 + a_9X_2X_1^2 + a_{10}X_0X_1X_2.$$

We can parametrize these simply by looking at their coefficients $(a_1 : a_2 : \cdots : a_{10})$.

To each such f , there corresponds an algebraic curve $X_f = \{f = 0\}$, which we saw is an elliptic curve when it is smooth.

Boundedness of curves

The space of coefficients $(a_1 : \cdots : a_{10}) = \mathbb{P}^9$ parametrizes these elliptic curves (and their degenerations).

Boundedness of curves

The space of coefficients $(a_1 : \cdots : a_{10}) = \mathbb{P}^9$ parametrizes these elliptic curves (and their degenerations).

Figure: Imagine a beautiful picture of a family of elliptic curves

Boundedness of curves

The space of coefficients $(a_1 : \cdots : a_{10}) = \mathbb{P}^9$ parametrizes these elliptic curves (and their degenerations).

Figure: Imagine a beautiful picture of a family of elliptic curves

When we move the coefficients of f also the structure of X_f varies. Two different polynomials f, f' give non isomorphic $X_f, X_{f'}$, in general.

Boundedness of curves

The space of coefficients $(a_1 : \cdots : a_{10}) = \mathbb{P}^9$ parametrizes these elliptic curves (and their degenerations).

Figure: Imagine a beautiful picture of a family of elliptic curves

When we move the coefficients of f also the structure of X_f varies. Two different polynomials f, f' give non isomorphic $X_f, X_{f'}$, in general.

Moreover, if $W \rightarrow Z$ is a morphism whose fibers are elliptic curves, one should expect to get a map $m: Z \dashrightarrow \mathbb{P}^9$.

Boundedness of curves

The space of coefficients $(a_1 : \cdots : a_{10}) = \mathbb{P}^9$ parametrizes these elliptic curves (and their degenerations).

Figure: Imagine a beautiful picture of a family of elliptic curves

When we move the coefficients of f also the structure of X_f varies. Two different polynomials f, f' give non isomorphic $X_f, X_{f'}$, in general.

Moreover, if $W \rightarrow Z$ is a morphism whose fibers are elliptic curves, one should expect to get a map $m: Z \dashrightarrow \mathbb{P}^9$.

One then could try to classify W by classifying Z and understanding the map $Z \dashrightarrow \mathbb{P}^9$.

Boundedness

As we are interested not only in understanding the behavior of algebraic varieties as standalone objects, but also in families, we need a suitable notion to indicate that a set of varieties can be described by many families.

Boundedness

An infinite number of algebraic varieties can be defined by a finite number of parameters.

How to show boundedness?

Given a class \mathcal{D} of projective algebraic varieties, we need to show that they can all be embedded in a fixed projective space \mathbb{P}^l and the volume of these embeddings is bounded.

State of the art

Question

Are varieties in each of the 3 building block (Fano, Calabi–Yau, canonical models) bounded?

State of the art

Question

Are varieties in each of the 3 building block (Fano, Calabi–Yau, canonical models) bounded?

This is a very difficult problem as in general it is quite complicated to construct embeddings for these varieties with controlled volumes.

State of the art

Question

Are varieties in each of the 3 building block (Fano, Calabi–Yau, canonical models) bounded?

This is a very difficult problem as in general it is quite complicated to construct embeddings for these varieties with controlled volumes.

Canonical models [Hacon-McKernan-Xu, 2009-15]:

N -dimensional canonical models with mild singularities with fixed volume v are bounded.

Fano varieties [Birkar, 2016]:

N -dimensional Fano varieties with bounded singularities are bounded.

These results were one of 2 main achievements that earned Hacon-McKernan the 2017 Breakthrough Prize and Birkar the 2018 Fields Medal.



Figure: McKernan and Hacon



Figure: Birkar

Calabi-Yau varieties and boundedness

Question

What happens for Calabi-Yau varieties?

If $n = 1$, X is an elliptic curve.

Calabi-Yau varieties and boundedness

Question

What happens for Calabi-Yau varieties?

If $n = 1$, X is an elliptic curve.

If $n = 2$, X is a K3 surface. K3 surfaces share the same underlying topological space but there are ∞ -many families of those.

Calabi-Yau varieties and boundedness

Question

What happens for Calabi-Yau varieties?

If $n = 1$, X is an elliptic curve.

If $n = 2$, X is a K3 surface. K3 surfaces share the same underlying topological space but there are ∞ -many families of those.

Conjecture

Fix a positive integer $n \geq 3$. Are n -dimensional projective Calabi-Yau varieties bounded?

Calabi-Yau varieties and boundedness

Question

What happens for Calabi-Yau varieties?

If $n = 1$, X is an elliptic curve.

If $n = 2$, X is a K3 surface. K3 surfaces share the same underlying topological space but there are ∞ -many families of those.

Conjecture

Fix a positive integer $n \geq 3$. Are n -dimensional projective Calabi-Yau varieties bounded?

This is a very obscure problem. Only known result:

Elliptic CY 3-folds [Gross, 1994]:

Elliptic Calabi-Yau threefolds $f: X \rightarrow Y$ are bounded up to birational equivalence, assuming that f is non-isotrivial.

My work

In a recent project, joint with G. Di Cerbo, we focus on Calabi-Yau varieties with an elliptic fibration. They have been intensively studied in mathematics and they seem to be the most relevant in physics (string theory and F-theory).

My work

In a recent project, joint with G. Di Cerbo, we focus on Calabi-Yau varieties with an elliptic fibration. They have been intensively studied in mathematics and they seem to be the most relevant in physics (string theory and F-theory).

Theorem

Let $n \leq 5$ be an integer. Then there are finitely many families of elliptic Calabi-Yau varieties $f: X \rightarrow Y$, up to birational isomorphism, such that $\dim(X) = n$, X is not a product and there exists a section $s: Y \rightarrow X$.

My work

In a recent project, joint with G. Di Cerbo, we focus on Calabi-Yau varieties with an elliptic fibration. They have been intensively studied in mathematics and they seem to be the most relevant in physics (string theory and F-theory).

Theorem

Let $n \leq 5$ be an integer. Then there are finitely many families of elliptic Calabi-Yau varieties $f: X \rightarrow Y$, up to birational isomorphism, such that $\dim(X) = n$, X is not a product and there exists a section $s: Y \rightarrow X$.

A section $s: Y \rightarrow X$ is simply a map such that $f \circ s = \text{Id}_Y$. In particular, it gives a point of reference on every fiber.

Idea of the proof

We split the proof of the Theorem in 2 parts.

Idea of the proof

We split the proof of the Theorem in 2 parts.

$$\begin{array}{c}
 Y \dashrightarrow Y' \\
 \downarrow \\
 Y_1 \\
 \downarrow \\
 Y_2 \\
 \downarrow \\
 Y_3
 \end{array}$$

(1) We first show that bases of elliptic CYs are bounded. Using the MMP, they decompose birationally into a tower of morphisms. We show that at each step in the tower boundedness holds and it propagates upwards.

$$\begin{array}{c}
 s(Y') = S \longrightarrow X' \\
 \downarrow \\
 Y'
 \end{array}$$

Idea of the proof

We split the proof of the Theorem in 2 parts.

$$\begin{array}{c}
 Y \dashrightarrow Y' \\
 \downarrow \\
 Y_1 \\
 \downarrow \\
 Y_2 \\
 \downarrow \\
 Y_3
 \end{array}$$

(1) We first show that bases of elliptic CYs are bounded. Using the MMP, they decompose birationally into a tower of morphisms. We show that at each step in the tower boundedness holds and it propagates upwards.

$$\begin{array}{ccc}
 s(Y') = S & \longrightarrow & X' \\
 & & \downarrow \\
 & & Y'
 \end{array}$$

(2) Since the elliptic fibration has a section, we can deduce boundedness of the total space from boundedness of the bases.

Future directions:

The previous theorem can still be improved in few directions.

Future directions:

The previous theorem can still be improved in few directions.

- First of all, we would like to prove the same statement in any dimension. One approach is to prove the existence of a parameter space for the fibers of the morphism appearing the tower. Once such space is constructed, the result should follow.

Future directions:

The previous theorem can still be improved in few directions.

- First of all, we would like to prove the same statement in any dimension. One approach is to prove the existence of a parameter space for the fibers of the morphism appearing the tower. Once such space is constructed, the result should follow.
- It would be interesting to remove the assumption on the existence of the section. The result of M. Gross in dimension 3 does not have this assumption. In higher dimension, there are subtle problems related to the type of singular fibers that appear in the fibration and how to deal with those.

Future directions:

The previous theorem can still be improved in few directions.

- First of all, we would like to prove the same statement in any dimension. One approach is to prove the existence of a parameter space for the fibers of the morphism appearing the tower. Once such space is constructed, the result should follow.
- It would be interesting to remove the assumption on the existence of the section. The result of M. Gross in dimension 3 does not have this assumption. In higher dimension, there are subtle problems related to the type of singular fibers that appear in the fibration and how to deal with those.
- Finally, it should be enough to assume that the base of the fibration is covered by many copies of \mathbb{P}^1 . At the moment we can prove this only in dimension 4 and we are working with Birkar to remove this obstruction.

Thank you!