## Group 1: Tennis

Olivia Beck, Emma Lewis, Ryan Volkert

December 12, 2019


## **Project**

- Project based off a paper by Paul K. Newton and Joseph B. Keller called Probability of Winning at Tennis 1. Theory and Data
- Paper discussed how the probability of winning a game, a set, and a match in tennis are computed based on each player's probability of winning a point on a serve
- Based on other papers the conclusion was made to treat serves in tennis as independent identically distributed (iid) random variables
- ▶ It is noted that points in tennis are not iid for most purposes this is not a bad assumption as the divergence from iid is small
- Both two out of three and three out of five set matches were considered allowing for 13-point tiebreaker in each set if needed

#### The Game of Tennis

- ▶ Point: Smallest unit of measurement (Love-15-30-40-game)
- ► Game: A game is one when a player reaches 4 points with at least a 2 point advantage
- ► Set: A set consists of 6 games and is won by the player who reaches 6 games first
- Advantage Set: If a game score of 6-6 is reached and advantage set rules are used, a player can only win a set with a 2 game lead
- ▶ Matches: Best of 3 set (for women) or 5 sets (for men)
- ► Tie-break game: If a game score of 6-6 is reached and tie-break set rules are used. In a tie-break game, a player/team must reach 7 points with a two point advantage to win

## Game of Tennis



#### Values

#### Things we need to know:

- ► Empirical probability of winning a rally on serve
- Empirical probability of winning a game on serve

#### **Values**

#### Things we need to know:

- Empirical probability of winning a rally on serve
- Empirical probability of winning a game on serve

#### Things we need to calculate:

- Theoretical probability of winning a game on serve
- Theoretical probability of winning a set
- Theoretical probability of winning the tie-breaker
- Theoretical probability of winning a match

## Theoretical Probabilities: Rally and Game

$$P(A \text{ Winning a Rally } | \text{ A Served}) = p_A^R = \frac{\text{Points Won On Serve}}{\text{Points Served}}$$

# Theoretical Probabilities: Rally and Game

$$P(A \text{ Winning a Rally } | A \text{ Served}) = p_A^R = \frac{Points \text{ Won On Serve}}{Points \text{ Served}}$$

P(A Winning a Game | A Served)  
= 
$$p_A^G = (p_A^R)^4 [1 + 4q_A^R + 10(q_A^R)^2] + 20(p_A^R q_A^R)^3 (p_A^R)^2 [1 - 2p_A^R q_A^R]^{-1}$$

Where Probability A loses a rally given A served:

$$q_A^R = 1 - \frac{\text{Points Won On Serve}}{\text{Points Served}}$$

### Theoretical Probabilities: Set

$$=p_A^S=\sum_{i=0}^4p_A^S(6,j)+p_A^S(7,5)+p_A^S(6,6)p_A^T$$

### Theoretical Probabilities: Set

$$= p_A^S = \sum_{j=0}^4 p_A^S(6,j) + p_A^S(7,5) + p_A^S(6,6)p_A^T$$

Where,  $p_A^S(i,j)$  is defined recursivly as:

$$p_A^S(0,0) = 1$$
,  $p_A^S(i,j) = 0$  if i<0 or j<0.

- if i+j -1 is even:  $p_A^S(i,j) = p_A^S(i-1,j)p_A^G + p_A^S(i,j-1)q_A^G$ 
  - ▶ omit i-1 term if j=6 and i<6
  - ▶ omit j-1 term if i=6 and j<6
- ▶ if i+j -1 is odd:  $p_A^S(i,j) = p_A^S(i-1,j)q_B^G + p_A^S(i,j-1)p_B^G$ 
  - ▶ omit i-1 term if j=6 and i<6
  - ▶ omit j-1 term if i=6 and j<6</p>

### Theoretical Probabilities: Tie Breaker

P(A Winning the Tie Breaker | A Served Initially)

$$= p_A^T = \sum_{j=0}^5 p_A^T(7,j) + p_A^T(6,6) \sum_{n=1}^\infty p_A^T(n+2,n)$$

### Theoretical Probabilities: Tie Breaker

P(A Winning the Tie Breaker | A Served Initially)

$$= p_A^T = \sum_{j=0}^5 p_A^T(7,j) + p_A^T(6,6) \sum_{n=1}^\infty p_A^T(n+2,n)$$

Where,  $p_{\Delta}^{T}(i,j)$  is defined recursivly as:

$$p_A^T(0,0) = 1$$
,  $p_A^T(i,j) = 0$  if i<0 or j<0.

- ▶ if  $i + j 1 \mod 4 \equiv 0$  or 3:
  - $p_A^S(i,j) = p_A^T(i-1,j)p_A^R + p_A^T(i,j-1)q_A^R$ 
    - ▶ omit i-1 term if j=7 and i<7
    - $\,\blacktriangleright\,$  omit j-1 term if i=7 and j<7
- if  $i + j 1 \mod 4 \equiv 1$  or 2:  $p_A^T(i, j) = p_A^T(i - 1, j) q_B^R + p_A^T(i, j - 1) p_B^R$ 
  - $\blacktriangleright$  omit i-1 term if j=7 and i<7
  - ▶ omit j-1 term if i=7 and j<7

#### Theoretical Probabilities: Match

Women's (Best of 3):

 $P(A \text{ Winning the Match} \mid A \text{ Served First})$ 

$$=p_A^M=(p_A^S)^2+2(p_A^S)^2p_B^S$$

#### Theoretical Probabilities: Match

Women's (Best of 3):

$$P(A \text{ Winning the Match} \mid A \text{ Served First})$$

$$= p_A^M = (p_A^S)^2 + 2(p_A^S)^2 p_B^S$$

Men's (Best of 5):

$$= p_A^M = (p_A^S)^3 + 3(p_A^S)^3 p_B^S + 6(p_A^S)^3 (p_B^S)^2$$

Where:

- $ightharpoonup p_A^S =$ Probabilty A wins a set given A served
- $p_B^S$  = Probabilty B wins a set given B served

## Data being used

#### Data used in paper:

- Men and Women semifinalists in the 2002 U.S Open tournament
- Men and Women semifinalists in the 2002 Wimbledon tournament

#### Data used in our simulation:

Men and Women semifinalists in the 2019 Wibledon tournament

#### Results of Game

Table 1: Data for the Womens Semifinalists in the 2002 U.S. Open Tournament

	P(Win a Rally)	Empirical P(Win a Game)	Paper P(Win a Game)	Our P(Win a Game)
S. Williams	0.69	0.71	0.89	0.89
V. Williams	0.63	0.80	0.79	0.79
L. Davenport	0.65	0.85	0.83	0.83
A. Mauresmo	0.63	0.77	0.79	0.79

#### Results of Game


Table 1: Data for the Womens Semifinalists in the 2002 U.S. Open Tournament

	P(Win a Rally)	Empirical P(Win a Game)	Paper P(Win a Game)	Our P(Win a Game)
S. Williams	0.69	0.71	0.89	0.89
V. Williams	0.63	0.80	0.79	0.79
L. Davenport	0.65	0.85	0.83	0.83
A. Mauresmo	0.63	0.77	0.79	0.79

Table 2: Data for the Mens Semifinalists in the 2002 U.S. Open Tournament

	P(Win a Rally)	Empirical P(Win a Game)	Paper P(Win a Game)	Our P(Win a Game)
P. Sampras	0.73	0.95	0.93	0.93
A. Agassi	0.66	0.87	0.85	0.85
L. Hewitt	0.67	0.85	0.86	0.86
S. Schalken	0.68	0.90	0.88	0.88


# 2002 U.S Open: Women



## 2002 U.S Open: Women




# 2002 U.S Open: Men



## 2002 U.S Open: Men




## 2019 Wimbledon: Women



### 2019 Wimbledon: Women



### 2019 Wimbledon: Men



### 2019 Wimbledon: Men

