
burn_4


Map 1 of 5: Landsat 8/9 before and after the burn | prescribed burn 27 Oct 2026 | reported 11 ac



Satellite estimate from a recorded point, date and acres (free data); not field-verified. Prescribed burns under canopy often read as unburned even where surface fuels burned.
Landsat 8/9 surface reflectance (30 m resampled to 20 m). Coordinates WGS84 decimal degrees.

burn_4

Map 2 of 5: Sentinel-2 before and after the burn | prescribed burn 27 Oct 2026 | reported 11 ac



Reported: 11 ac Mapped: NOT DETECTED

Images: 21 before, 4 after (searched 60 days before and 45 days after). Each panel is the per-pixel median of the 3 clearest looks closest to the burn; dates above are those looks. Flat grey = no clear view.

How the outline is drawn:
Sentinel-2 (20 m) burn index drop that is next to the recorded point and dated to the burn.


False colour (SWIR-NIR-red): healthy vegetation green, fresh burn scar dark red / brown / purple.



Satellite estimate from a recorded point, date and acres (free data); not field-verified. Prescribed burns under canopy often read as unburned even where surface fuels burned.
Sentinel-2 surface reflectance (20 m). Coordinates WGS84 decimal degrees.

burn_4

Map 3 of 5: Sentinel-2 + Landsat combined | prescribed burn 27 Oct 2026 | reported 11 ac



Combined method

- Mapped burned area
- 1 km search radius around point
- Recorded lat/long

Reported: 11 ac Mapped: NOT DETECTED
Images: 21 + 4 Sentinel-2, 4 + 0 Landsat (before + after)
First clear image after burn: 3 days


How the outline is drawn:
Both satellites in one time series (more clear looks).
Each image is compared with land far from the point,
so seasonal change cancels out. No embeddings.



Satellite estimate from a recorded point, date and acres (free data); not field-verified. Prescribed burns under canopy often read as unburned even where surface fuels burned.
Combined Sentinel-2 + Landsat burn index, point + date match. Coordinates WGS84 decimal degrees.

burn_4

Map 4 of 5: VIIRS active-fire detections (NASA FIRMS) | prescribed burn 27 Oct 2026 | reported 11 ac



VIIRS active fire

- VIIRS fire detection (375 m pixel footprint)
- 1 km search radius around point
- Recorded lat/long

Window: 26 Oct - 29 Oct 2026
Detections within 1 km of the point (+ half a pixel): 12
All detections loaded: 17 detections within ~30 km
Dates: 28 Oct, 29 Oct


VIIRS (Suomi-NPP / NOAA-20 / NOAA-21) sees heat from active flames when a satellite passes (about twice a day per satellite). It confirms fire was present, not how far it spread; small, cool or short burns are often missed.



Satellite estimate from a recorded point, date and acres (free data); not field-verified. Prescribed burns under canopy often read as unburned even where surface fuels burned.
VIIRS 375 m active-fire product, NASA FIRMS. Coordinates WGS84 decimal degrees.

burn_4

Map 5 of 5: best effort (adds embeddings) | prescribed burn 27 Oct 2026 | reported 11 ac



Burn severity (satellite index)

- Unburned / very low (dNBR < 0.10): 0 ac
- Low (0.10 - 0.27): 1 ac
- Moderate-low (0.27 - 0.44): 0 ac
- Moderate-high (0.44 - 0.66): 0 ac
- High (> 0.66): 0 ac
- Mapped burn perimeter
- 1 km search radius around point
- Recorded lat/long

Reported: 11 ac Mapped: NOT DETECTED
Images: 26 Sentinel-2 + 4 Landsat
First clear image after burn: 3 days
Confidence: LOW
 - not detected (< 1 ha mapped)

How the outline is drawn:
As the combined method, but each pixel is compared only with look-alike land (same vegetation type, from TESSERA satellite embeddings of the year before), which removes more green-up / dry-down noise.



Satellite estimate from a recorded point, date and acres (free data); not field-verified. Prescribed burns under canopy often read as unburned even where surface fuels burned.
Severity = green-up-corrected dNBR, Key & Benson (2006) classes, not calibrated to field (CBI) plots. Coordinates WGS84 decimal degrees.