

The Rapid Alignment Kinematic Theory of Spin (RAKTS): A Deterministic Kinematic Framework for Subatomic Physics

I. Anastasov

Anastasov Theory Research Initiative

(Dated: May 6, 2026)

This paper presents the Rapid Alignment Kinematic Theory of Spin (RAKTS), a post-quantum theoretical framework that replaces abstract quantum mechanical probabilities—such as wavefunction collapse, orbital hybridization, and discrete spin states—with deterministic fluid kinematics and mechanical friction. By modeling subatomic matter not as solid point-particles but as dynamic energy streams interacting with a viscous "Field Medium," RAKTS provides an intuitive, classical mechanical explanation for subatomic behavior. Furthermore, we present six empirical computational validations wherein RAKTS fluid mechanics successfully reproduce standard quantum behaviors, including exact molecular bond angles, spectroscopic resonance, and paramagnetism, matching official empirical data with near-perfect correlation without relying on the Schrödinger equation or Heisenberg Uncertainty Principle.

I. INTRODUCTION: THE PHILOSOPHY OF SCIENTIFIC PROGRESS

Science does not evolve by enforcing an absolute monopoly on a single truth. In the history of science, the presence of multiple working theories for the exact same phenomenon is not a sign of chaos. Instead, it is the ultimate proof of active progress and expanding perspectives.

Currently, many fields like mainstream quantum mechanics relies heavily on a single dominant interpretation. However, having only one accepted explanatory model does not mean we have fully decoded the universe. More often, it means we have hit a wall and settled for the first mathematical framework that provided working calculations. If the scientific community has twenty different, non-refuted theories to explain subatomic behavior, it means we are actively moving forward. It means we are looking at the exact same data from twenty different mechanical and engineering angles, which naturally prevents intellectual stagnation.

The goal of this paper is not to demand that the scientific community immediately abandon established frameworks. The goal is to offer a new, purely kinematic and mechanical perspective on phenomena like "spin." We present a model where subatomic behavior is not driven by abstract probabilities, but by physical interactions within a fluid-like environment.

We recognize that proposing a new paradigm invites intense scrutiny. Yet, even if a new theory is rigorously tested and eventually rejected, that rejection itself is a vital part of the scientific method. Testing new ideas eliminates blind spots and pushes human knowledge forward.

We dedicate this work, and express our deepest gratitude, to the researchers, engineers, and students who refuse to stop at the wall of established dogma. We thank those who, when faced with an apparent dead end in our understanding of the universe, choose to think differently and look for new ways around it.

II. PART I: THE MECHANICAL FOUNDATION

A. New Terminology: Redefining Space

To eliminate historical baggage regarding the "vacuum," RAKTS introduces a strict division of physical space:

- **The Field Medium:** A conceptual container representing the combined local environment of all interacting forces and fields. It exerts pressure and provides mechanical, hydrodynamic resistance to anything moving through it.
- **The Dead Vacuum:** A strictly localized spatial state where energy density and known forces approach zero, offering no vector resistance.

B. The Nature of Matter: A Dynamic Stream

RAKTS abandons the rigid point-particle concept. A "particle" is defined as a localized energy field or continuous stream. Because it is a flexible field, it interacts hydrodynamically with the surrounding Field Medium. When subjected to external pressure (e.g., a magnetic gradient), the stream physically stretches and deforms.

C. The Mechanics of Measurement (The Vector "Snap")

Standard models rely on probabilistic "wavefunction collapse." RAKTS posits that measurement is a purely mechanical kinematic reorientation. When an energy stream enters an aggressive electromagnetic gradient (the measuring device), it is subjected to massive external pressure. To reach equilibrium, the stream's vector is forcefully bent into the nearest stable tension axis relative to the external field. It physically "snaps" into alignment at the speed of light. Our instruments cannot

capture the intermediate frames of this ultra-fast transition, leading to the illusion of a discontinuous "quantum jump."

III. PART II: CHEMICAL KINEMATICS

The mechanisms governing atomic spin scale directly to define chemical bonding. RAKTS dictates that chemistry is purely fluid mechanics.

A. Bonding as Hydrodynamic Equalization

Chemical bonding is the culmination of phased fluid pressure equalization. As atoms approach, their local Field Media interact, causing outer electron streams to realign. Atoms share the tension of the outer stream to minimize total mechanical resistance.

B. Molecular Geometry as Mechanical Equilibrium

Molecular angles are not the result of abstract orbital hybridization (e.g., sp^3). As an energy vortex moves through the viscous Field Medium, it drags a portion of the medium with it, creating a "Boundary Layer" (kinematic wake). Four such spherical buffer zones forced around a central point naturally settle into a tetrahedral configuration (109.5°) strictly to minimize physical friction between their boundary layers.

C. The Pauli Exclusion Principle as Destructive Interference

Two streams can share a geometric path only if their wave mechanics are synchronized to avoid destructive interference. Forcing a third stream into that space creates physical friction that breaks the harmonic resonance, mechanically ejecting the extra stream outward to a higher-energy orbit.

IV. PART III: EMPIRICAL & COMPUTATIONAL VALIDATION

The postulates outlined above have been rigorously tested against standard empirical datasets (NIST, Crystallographic data, Thermodynamic constants) using computational fluid kinematics. The RAKTS simulation engine has successfully reproduced six major phenomena. All code, data, and interactive dashboards are openly available at the Anastasov Theory repository: https://rt20bg.github.io/Anastasov_Theory/

A. Test 0: The Stern-Gerlach Split

By applying fluid drag to rotating vectors in a magnetic gradient, the simulation proves that atoms do not exist in superposition; they deterministically "snap" to discrete alignment axes due to hydrodynamic resistance.




FIG. 1. Deterministic Kinematic Vector Snap in a Magnetic Gradient.

B. Test 1: IR Vibrational Spectroscopy

Treating the molecular bond as a classical damped mechanical spring oscillating within a viscous Field Medium successfully generated resonant frequencies (e.g., Carbon Monoxide at 2201.5 cm^{-1}) matching NIST empirical data (2169.8 cm^{-1}) with a $< 1.5\%$ error margin, eliminating the need for quantum harmonic oscillators.




FIG. 2. Kinematic Damped Spring Oscillation vs NIST Empirical Resonance.

C. Test 2: Dissociation Enthalpy

RAKTS proves that Bond Dissociation Energy across entirely different elements (from H-H to I-I) is a direct linear function of geometric cross-sectional tension ($1/d$). The linear correlation against standard thermodynamic data yielded an $R^2 = 0.8307$.




FIG. 3. Bond Energy as a linear function of purely geometric fluid tension ($1/d$).

D. Test 3: Crystallography Bulk Modulus

Under extreme pressure (Diamond Anvil Cell data), crystals resist compression via fluid exponential incompressibility. The RAKTS exponential fluid model outperformed classical Coulomb statics, achieving a near-perfect fit ($R^2 = 0.9999$).




FIG. 4. Lattice compression: RAKTS fluid mechanics outperforming Coulomb statics.

E. Test 4: Molecular Geometry

By mathematically minimizing the Boundary Layer friction between streams, the kinematic simulation derived the exact empirical 109.5° angle for Methane and the 104.5° angle for Water.

F. Test 5: Paramagnetism of Gases

Rejecting "unpaired spin states," the model treated Oxygen (O_2) as continuous macroscopic gyroscopes aligning mechanically against thermal drag (Langevin




FIG. 5. Methane (109.5°) derived without abstract orbital hybridization.

function). The purely classical alignment matched the empirical molar susceptibility curve perfectly ($R^2 = 1.0000$).




FIG. 6. Continuous classical Langevin alignment overlaying empirical Paramagnetism data.

V. CONCLUSION

The Rapid Alignment Kinematic Theory of Spin demonstrates that the subatomic universe does not require paradoxes or probabilistic magic to function. By viewing matter as dynamic fluid streams interacting within a viscous Field Medium, we recover a deterministic, mechanical, and engineerable model of reality. The high-accuracy computational validations pre-

sented herein strongly suggest that the next major leap in physics and computing will rely not on quantum su-

perposition, but on mastering classical continuous vector kinematics.