

Chapter 1

Introduction

Contact tracing is a non-pharmaceutical intervention that aims to halt the spread of infectious disease by identifying and quarantining individuals that have been in close physical proximity with the infected [3, 5].

In response to the pandemic of coronavirus disease 2019 (COVID-19) [4, 6, 7],

Effective contact tracing is inherently difficult because of the complex interactions between epidemiology, public health, ethics, politics, and sociocultural norms [3].

A common design element across all of the aforementioned contact tracing methodologies is that they only consider direct interactions between users. While there are privacy benefits to this approach, a major limitation is that they cannot utilize information about indirect contact to more effectively reduce the spread of disease. ShareTrace addresses this limitation by constructing a factor graph and estimating infection risk via a message-passing algo-

rithm. As such, this work labels the ShareTrace algorithm as *risk propagation*. The first work on ShareTrace was a white paper that focused on the motivation, design, and engineering details. Exclusive to Ayday et al. [2] is a discussion on privacy, network roaming, protocol interoperability, and the usage of geolocation data. Furthermore, it includes detail on the system model and data flow. The second work on ShareTrace [1] formalizes the algorithmic details in a centralized setting and demonstrates its improved efficacy, compared to the framework developed by Apple and Google [**AppleGoogle**].

Higher-order contact tracing has demonstrated increased efficacy [5]

Bibliography

- [1] Erman Ayday, Youngjin Yoo, and Anisa Halimi. “ShareTrace: An Iterative Message Passing Algorithm for Efficient and Effective Disease Risk Assessment on an Interaction Graph”. In: *Proc. 12th ACM Con. Bioinformatics, Comput. Biology, Health Inform.* BCB 2021. 2021 (cit. on p. 2).
- [2] Erman Ayday et al. *ShareTrace: A Smart Privacy-Preserving Contact Tracing Solution by Architectural Design During an Epidemic*. Tech. rep. Case Western Reserve University, 2020. URL: <https://github.com/cwru-xlab/sharetrace-papers/blob/main/sharetrace-whitepaper.pdf> (cit. on p. 2).
- [3] Allan M. Brandt. “The History of Contact Tracing and the Future of Public Health”. In: *American Journal of Public Health* 112.8 (2022), pp. 1097–1099. DOI: 10.2105/AJPH.2022.306949 (cit. on p. 1).
- [4] Alexander E. Gorbalenya et al. “The species severe acute respiratory syndrome-related coronavirus: classifying 2019-nCoV and naming it SARS-CoV-2”. In: *Nature Microbiology* 5.4 (2020), pp. 536–544. DOI: 10.1038/s41564-020-0695-z (cit. on p. 1).

- [5] Francisco Pozo-Martin et al. “Comparative effectiveness of contact tracing interventions in the context of the COVID-19 pandemic: a systematic review”. In: *European Journal of Epidemiology* 38.3 (2023), pp. 243–266. DOI: 10.1007/s10654-023-00963-z (cit. on pp. 1, 2).
- [6] Sudhvir Singh et al. “How an outbreak became a pandemic: a chronological analysis of crucial junctures and international obligations in the early months of the COVID-19 pandemic”. In: *Lancet* 398.10316 (2021), pp. 2109–2124 (cit. on p. 1).
- [7] Na Zhu et al. “A Novel Coronavirus from Patients with Pneumonia in China, 2019”. In: *New England Journal of Medicine* 382.8 (2020), pp. 727–733. DOI: 10.1056/NEJMoa2001017 (cit. on p. 1).