


# A new proof of Viazovska's modular form inequalities for sphere packing in dimension 8

Dan Romik  
UC Davis

Computer Algebra Workshop  
Institut Henri Poincaré

December 6, 2023



# Talk outline

# Talk outline

- ① Background: sphere packings in  $\mathbb{R}^d$

# Talk outline

- ① Background: sphere packings in  $\mathbb{R}^d$
- ② Viazovska's solution of the sphere packing problem in dimension 8

# Talk outline

- ① Background: sphere packings in  $\mathbb{R}^d$
- ② Viazovska's solution of the sphere packing problem in dimension 8
- ③ Viazovska's modular form inequalities

# Talk outline

- ① Background: sphere packings in  $\mathbb{R}^d$
- ② Viazovska's solution of the sphere packing problem in dimension 8
- ③ Viazovska's modular form inequalities
- ④ A new proof

# Some useful references

Source material for this talk:

# Some useful references

Source material for this talk:

- The paper “The sphere packing problem in dimension 8” by Maryna Viazovska (*Ann. Math.* (2017), 991–1015).

# Some useful references

Source material for this talk:


- The paper “The sphere packing problem in dimension 8” by Maryna Viazovska (*Ann. Math.* (2017), 991–1015).
- My paper “On Viazovska’s modular form inequalities” (PNAS, 2023).

# Some useful references

Source material for this talk:

- The paper “The sphere packing problem in dimension 8” by Maryna Viazovska (*Ann. Math.* (2017), 991–1015).
- My paper “On Viazovska’s modular form inequalities” (PNAS, 2023).
- Chapter 6 + Appendix of my book “Topics in Complex Analysis”

[https://www.math.ucdavis.edu/  
~romik/topics-in-complex-analysis/](https://www.math.ucdavis.edu/~romik/topics-in-complex-analysis/)



## Background: sphere packings in $\mathbb{R}^d$

## Background: sphere packings in $\mathbb{R}^d$

**The sphere packing problem in  $\mathbb{R}^d$**  asks: what is the densest way to pack unit spheres in  $d$ -dimensional space?

## Background: sphere packings in $\mathbb{R}^d$

**The sphere packing problem in  $\mathbb{R}^d$**  asks: what is the densest way to pack unit spheres in  $d$ -dimensional space?

- Trivial case:  $d = 1$ .

## Background: sphere packings in $\mathbb{R}^d$

**The sphere packing problem in  $\mathbb{R}^d$**  asks: what is the densest way to pack unit spheres in  $d$ -dimensional space?

- Trivial case:  $d = 1$ .
- (Relatively) easy case:  $d = 2$ .

## Background: sphere packings in $\mathbb{R}^d$

**The sphere packing problem in  $\mathbb{R}^d$**  asks: what is the densest way to pack unit spheres in  $d$ -dimensional space?

- Trivial case:  $d = 1$ .
- (Relatively) easy case:  $d = 2$ . The optimal *circle packing* is the **hexagonal lattice packing**, with packing density  $\frac{\pi}{2\sqrt{3}}$ .  
Proved by Gauss (1831), Thue (1890), Tóth (1941).

## Background: sphere packings in $\mathbb{R}^d$

**The sphere packing problem in  $\mathbb{R}^d$**  asks: what is the densest way to pack unit spheres in  $d$ -dimensional space?

- Trivial case:  $d = 1$ .
- (Relatively) easy case:  $d = 2$ . The optimal *circle packing* is the **hexagonal lattice packing**, with packing density  $\frac{\pi}{2\sqrt{3}}$ .  
Proved by Gauss (1831), Thue (1890), Tóth (1941).
- Famous case:  $d = 3$ .

## Background: sphere packings in $\mathbb{R}^d$

The **sphere packing problem** in  $\mathbb{R}^d$  asks: what is the densest way to pack unit spheres in  $d$ -dimensional space?

- Trivial case:  $d = 1$ .
- (Relatively) easy case:  $d = 2$ . The optimal *circle packing* is the **hexagonal lattice packing**, with packing density  $\frac{\pi}{2\sqrt{3}}$ .  
Proved by Gauss (1831), Thue (1890), Tóth (1941).
- Famous case:  $d = 3$ . Kepler's conjecture from 1611 stated that the optimal density for sphere packing is  $\frac{\pi}{3\sqrt{2}}$ , achieved by the **cubic close packing** and the **hexagonal close packing**. Proved by Thomas Hales in 1998.

## Background: sphere packings in $\mathbb{R}^d$

The **sphere packing problem** in  $\mathbb{R}^d$  asks: what is the densest way to pack unit spheres in  $d$ -dimensional space?

- Trivial case:  $d = 1$ .
- (Relatively) easy case:  $d = 2$ . The optimal *circle packing* is the **hexagonal lattice packing**, with packing density  $\frac{\pi}{2\sqrt{3}}$ . Proved by Gauss (1831), Thue (1890), Tóth (1941).
- Famous case:  $d = 3$ . Kepler's conjecture from 1611 stated that the optimal density for sphere packing is  $\frac{\pi}{3\sqrt{2}}$ , achieved by the **cubic close packing** and the **hexagonal close packing**. Proved by Thomas Hales in 1998.
- The case  $d = 8$ .

## Background: sphere packings in $\mathbb{R}^d$

The **sphere packing problem** in  $\mathbb{R}^d$  asks: what is the densest way to pack unit spheres in  $d$ -dimensional space?

- Trivial case:  $d = 1$ .
- (Relatively) easy case:  $d = 2$ . The optimal *circle packing* is the **hexagonal lattice packing**, with packing density  $\frac{\pi}{2\sqrt{3}}$ .  
Proved by Gauss (1831), Thue (1890), Tóth (1941).
- Famous case:  $d = 3$ . Kepler's conjecture from 1611 stated that the optimal density for sphere packing is  $\frac{\pi}{3\sqrt{2}}$ , achieved by the **cubic close packing** and the **hexagonal close packing**. Proved by Thomas Hales in 1998.
- The case  $d = 8$ . Maryna Viazovska proved in 2016 that for  $d = 8$ , the densest packing is the  **$E_8$  lattice packing**, with packing density  $\frac{\pi^4}{384}$ .

## Background: sphere packings in $\mathbb{R}^d$

The **sphere packing problem** in  $\mathbb{R}^d$  asks: what is the densest way to pack unit spheres in  $d$ -dimensional space?

- Trivial case:  $d = 1$ .
- (Relatively) easy case:  $d = 2$ . The optimal *circle packing* is the **hexagonal lattice packing**, with packing density  $\frac{\pi}{2\sqrt{3}}$ .  
Proved by Gauss (1831), Thue (1890), Tóth (1941).
- Famous case:  $d = 3$ . Kepler's conjecture from 1611 stated that the optimal density for sphere packing is  $\frac{\pi}{3\sqrt{2}}$ , achieved by the **cubic close packing** and the **hexagonal close packing**. Proved by Thomas Hales in 1998.
- The case  $d = 8$ . Maryna Viazovska proved in 2016 that for  $d = 8$ , the densest packing is the  **$E_8$  lattice packing**, with packing density  $\frac{\pi^4}{384}$ .
- The case  $d = 24$ .

## Background: sphere packings in $\mathbb{R}^d$





**The sphere packing problem in  $\mathbb{R}^d$**  asks: what is the densest way to pack unit spheres in  $d$ -dimensional space?

- Trivial case:  $d = 1$ .
- (Relatively) easy case:  $d = 2$ . The optimal *circle packing* is the **hexagonal lattice packing**, with packing density  $\frac{\pi}{2\sqrt{3}}$ .  
Proved by Gauss (1831), Thue (1890), Tóth (1941).
- Famous case:  $d = 3$ . Kepler's conjecture from 1611 stated that the optimal density for sphere packing is  $\frac{\pi}{3\sqrt{2}}$ , achieved by the **cubic close packing** and the **hexagonal close packing**. Proved by Thomas Hales in 1998.
- The case  $d = 8$ . Maryna Viazovska proved in 2016 that for  $d = 8$ , the densest packing is the  **$E_8$  lattice packing**, with packing density  $\frac{\pi^4}{384}$ .
- The case  $d = 24$ . Viazovska with Cohn, Kumar, Miller, and Radchenko then proved that for  $d = 24$ , the densest packing is the **Leech lattice packing**, with packing density  $\frac{\pi^{12}}{12!}$ .

## Background: sphere packings in $\mathbb{R}^d$ (continued)

In other dimensions the problem remains open.

## Background: sphere packings in $\mathbb{R}^d$ (continued)



The optimal lattices for sphere packing in dimensions 2, 3, 8

# Viazovska's proof in dimension 8

## Viazovska's proof in dimension 8

- It had previously been conjectured that the  $E_8$  lattice packing, with packing density  $\frac{\pi^4}{384}$ , is optimal. This gives a lower bound on the optimal packing density; Viazovska proved a matching upper bound, solving the problem.

## Viazovska's proof in dimension 8

- It had previously been conjectured that the  $E_8$  lattice packing, with packing density  $\frac{\pi^4}{384}$ , is optimal. This gives a lower bound on the optimal packing density; Viazovska proved a matching upper bound, solving the problem.
- Viazovska made use of a remarkable theorem from 2001, the **Cohn-Elkies linear programming bounds**. It reduced the problem to finding a **magic function**, an analytic object with certain properties.

## Viazovska's proof in dimension 8

- It had previously been conjectured that the  $E_8$  lattice packing, with packing density  $\frac{\pi^4}{384}$ , is optimal. This gives a lower bound on the optimal packing density; Viazovska proved a matching upper bound, solving the problem.
- Viazovska made use of a remarkable theorem from 2001, the **Cohn-Elkies linear programming bounds**. It reduced the problem to finding a **magic function**, an analytic object with certain properties.
- Viazovska's proof is complex-analytic. She used modular forms to construct the magic function for dimension 8. An extension of the method works for dimension 24.

## Viazovska's proof in dimension 8

- It had previously been conjectured that the  $E_8$  lattice packing, with packing density  $\frac{\pi^4}{384}$ , is optimal. This gives a lower bound on the optimal packing density; Viazovska proved a matching upper bound, solving the problem.
- Viazovska made use of a remarkable theorem from 2001, the **Cohn-Elkies linear programming bounds**. It reduced the problem to finding a **magic function**, an analytic object with certain properties.
- Viazovska's proof is complex-analytic. She used modular forms to construct the magic function for dimension 8. An extension of the method works for dimension 24.
- One component of the proof makes extensive use of computer calculations.

# The Cohn-Elkies linear programming bounds

# The Cohn-Elkies linear programming bounds

## Theorem (Cohn-Elkies, 2001)

Let  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  be a function and  $\rho > 0$  a number. Assume that the following conditions are satisfied:

# The Cohn-Elkies linear programming bounds

## Theorem (Cohn-Elkies, 2001)

Let  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  be a function and  $\rho > 0$  a number. Assume that the following conditions are satisfied:

- ①  $f$  is a Schwartz function

# The Cohn-Elkies linear programming bounds

## Theorem (Cohn-Elkies, 2001)

Let  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  be a function and  $\rho > 0$  a number. Assume that the following conditions are satisfied:

- ①  $f$  is a Schwartz function
- ②  $f(0) = \widehat{f}(0) > 0$

# The Cohn-Elkies linear programming bounds

## Theorem (Cohn-Elkies, 2001)

Let  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  be a function and  $\rho > 0$  a number. Assume that the following conditions are satisfied:

- ①  $f$  is a Schwartz function
- ②  $f(0) = \widehat{f}(0) > 0$       ( $\widehat{f}$  = the Fourier transform of  $f$ )

# The Cohn-Elkies linear programming bounds

## Theorem (Cohn-Elkies, 2001)

Let  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  be a function and  $\rho > 0$  a number. Assume that the following conditions are satisfied:

- ①  $f$  is a Schwartz function
- ②  $f(0) = \widehat{f}(0) > 0$       ( $\widehat{f}$  = the Fourier transform of  $f$ )
- ③  $f(x) \leq 0$  for all  $x \in \mathbb{R}^d$  such that  $\|x\| \geq \rho$

# The Cohn-Elkies linear programming bounds

## Theorem (Cohn-Elkies, 2001)

Let  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  be a function and  $\rho > 0$  a number. Assume that the following conditions are satisfied:

- ①  $f$  is a Schwartz function
- ②  $f(0) = \widehat{f}(0) > 0$       ( $\widehat{f}$  = the Fourier transform of  $f$ )
- ③  $f(x) \leq 0$  for all  $x \in \mathbb{R}^d$  such that  $\|x\| \geq \rho$
- ④  $\widehat{f}(x) \geq 0$  for all  $x \in \mathbb{R}^d$

# The Cohn-Elkies linear programming bounds

## Theorem (Cohn-Elkies, 2001)

Let  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  be a function and  $\rho > 0$  a number. Assume that the following conditions are satisfied:

- ①  $f$  is a Schwartz function
- ②  $f(0) = \widehat{f}(0) > 0$  ( $\widehat{f}$  = the Fourier transform of  $f$ )
- ③  $f(x) \leq 0$  for all  $x \in \mathbb{R}^d$  such that  $\|x\| \geq \rho$
- ④  $\widehat{f}(x) \geq 0$  for all  $x \in \mathbb{R}^d$

Then the optimal packing density  $\delta_d$  in  $\mathbb{R}^d$  satisfies

$$\delta_d \leq \text{vol}(B_{\rho/2}(0))$$

# The Cohn-Elkies linear programming bounds

## Theorem (Cohn-Elkies, 2001)

Let  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  be a function and  $\rho > 0$  a number. Assume that the following conditions are satisfied:

- ①  $f$  is a Schwartz function
- ②  $f(0) = \widehat{f}(0) > 0$  ( $\widehat{f}$  = the Fourier transform of  $f$ )
- ③  $f(x) \leq 0$  for all  $x \in \mathbb{R}^d$  such that  $\|x\| \geq \rho$
- ④  $\widehat{f}(x) \geq 0$  for all  $x \in \mathbb{R}^d$

Then the optimal packing density  $\delta_d$  in  $\mathbb{R}^d$  satisfies

$$\delta_d \leq \text{vol}(B_{\rho/2}(0)) = \frac{\rho^d}{2^d} \times [\text{vol. of unit ball}]$$

# The Cohn-Elkies linear programming bounds

## Theorem (Cohn-Elkies, 2001)

Let  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  be a function and  $\rho > 0$  a number. Assume that the following conditions are satisfied:

- ①  $f$  is a Schwartz function
- ②  $f(0) = \widehat{f}(0) > 0$  ( $\widehat{f}$  = the Fourier transform of  $f$ )
- ③  $f(x) \leq 0$  for all  $x \in \mathbb{R}^d$  such that  $\|x\| \geq \rho$
- ④  $\widehat{f}(x) \geq 0$  for all  $x \in \mathbb{R}^d$

Then the optimal packing density  $\delta_d$  in  $\mathbb{R}^d$  satisfies

$$\delta_d \leq \text{vol}(B_{\rho/2}(0)) = \frac{\rho^d}{2^d} \times [\text{vol. of unit ball}]$$

- The proof is an application of the **Poisson summation formula** from harmonic analysis; see the appendix of my book.

# The Cohn-Elkies linear programming bounds

## Theorem (Cohn-Elkies, 2001)

Let  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  be a function and  $\rho > 0$  a number. Assume that the following conditions are satisfied:

- ①  $f$  is a Schwartz function
- ②  $f(0) = \widehat{f}(0) > 0$  ( $\widehat{f}$  = the Fourier transform of  $f$ )
- ③  $f(x) \leq 0$  for all  $x \in \mathbb{R}^d$  such that  $\|x\| \geq \rho$
- ④  $\widehat{f}(x) \geq 0$  for all  $x \in \mathbb{R}^d$

Then the optimal packing density  $\delta_d$  in  $\mathbb{R}^d$  satisfies

$$\delta_d \leq \text{vol}(B_{\rho/2}(0)) = \frac{\rho^d}{2^d} \times [\text{vol. of unit ball}]$$

- The proof is an application of the **Poisson summation formula** from harmonic analysis; see the appendix of my book.
- For the case  $d = 8$ , the sharp bound  $\frac{\pi^4}{384}$  is obtained when  $\rho = \sqrt{2}$ . A function satisfying the conditions of the theorem for that  $\rho$  is called a **magic function**.

# Applying the Cohn-Elkies bounds in practice

## Applying the Cohn-Elkies bounds in practice

Cohn and Elkies applied their bound to numerically optimized bounding functions  $f$ , obtaining the best known (at the time) upper bounds for the sphere packing density in dimensions 4–36.

# Applying the Cohn-Elkies bounds in practice

Cohn and Elkies applied their bound to numerically optimized bounding functions  $f$ , obtaining the best known (at the time) upper bounds for the sphere packing density in dimensions 4–36.





Image source: Henry Cohn, A conceptual breakthrough in sphere packing (Notices of AMS, 2017)


# Applying the Cohn-Elkies bounds in practice (continued)

In dimensions 2, 8 and 24, their bounds came extremely close to matching the known lower bounds.




# Applying the Cohn-Elkies bounds in practice (continued)

In dimensions 2, 8 and 24, their bounds came extremely close to matching the known lower bounds.



# Applying the Cohn-Elkies bounds in practice (continued)

In dimensions 2, 8 and 24, their bounds came extremely close to matching the known lower bounds.



They conjectured that in those dimensions there exists a “magic function”  $f$  certifying a *sharp* bound.

# Viazovska's magic function

Viazovska's function  $\varphi : \mathbb{R}^8 \rightarrow \mathbb{R}$  is defined by

# Viazovska's magic function

Viazovska's function  $\varphi : \mathbb{R}^8 \rightarrow \mathbb{R}$  is defined by (the analytic continuation of)

$$\begin{aligned}\varphi(x) = & -4 \sin^2 \left( \frac{\pi \|x\|^2}{2} \right) \\ & \times \int_0^\infty e^{-\pi t \|x\|^2} \left[ 108 \frac{(itE'_4(it) + 4E_4(it))^2}{E_4(it)^3 - E_6(it)^2} \right. \\ & \quad \left. + 128 \left( \frac{\theta_3(it)^4 + \theta_4(it)^4}{\theta_2(it)^8} + \frac{\theta_4(it)^4 - \theta_2(it)^4}{\theta_3(it)^8} \right) \right] dt,\end{aligned}$$

# Viazovska's magic function

Viazovska's function  $\varphi : \mathbb{R}^8 \rightarrow \mathbb{R}$  is defined by (the analytic continuation of)

$$\begin{aligned}\varphi(x) = & -4 \sin^2 \left( \frac{\pi \|x\|^2}{2} \right) \\ & \times \int_0^\infty e^{-\pi t \|x\|^2} \left[ 108 \frac{(itE'_4(it) + 4E_4(it))^2}{E_4(it)^3 - E_6(it)^2} \right. \\ & \quad \left. + 128 \left( \frac{\theta_3(it)^4 + \theta_4(it)^4}{\theta_2(it)^8} + \frac{\theta_4(it)^4 - \theta_2(it)^4}{\theta_3(it)^8} \right) \right] dt,\end{aligned}$$

where  $E_4, E_6$  are the **Eisenstein series** and  $\theta_2, \theta_3, \theta_4$  are the **Jacobi theta null functions**, defined by

$$E_4(z) = 1 + 240 \sum_{n=1}^{\infty} \sigma_3(n) q^{2n}, \quad \theta_2(z) = \sum_{n=-\infty}^{\infty} q^{(n+1/2)^2},$$

$$E_6(z) = 1 - 504 \sum_{n=1}^{\infty} \sigma_5(n) q^{2n}, \quad \theta_3(z) = \sum_{n=-\infty}^{\infty} q^{n^2},$$

$$\theta_4(z) = \sum_{n=-\infty}^{\infty} (-1)^n q^{n^2},$$

(with the standard notation  $q = e^{\pi iz}$ ,  $\sigma_\alpha(n) = \sum_{d|n} d^\alpha$ ).

# Viazovska's magic function (continued)

## Theorem (Viazovska)

$\varphi$  is a magic function for dimension 8. More precisely, it has the following properties:

# Viazovska's magic function (continued)

## Theorem (Viazovska)

$\varphi$  is a magic function for dimension 8. More precisely, it has the following properties:

- ①  $\varphi$  is a Schwartz function

# Viazovska's magic function (continued)

## Theorem (Viazovska)

$\varphi$  is a magic function for dimension 8. More precisely, it has the following properties:

- ①  $\varphi$  is a Schwartz function
- ②  $\varphi(0) = \widehat{\varphi}(0) = 240\pi$

# Viazovska's magic function (continued)

## Theorem (Viazovska)

$\varphi$  is a magic function for dimension 8. More precisely, it has the following properties:

- ①  $\varphi$  is a Schwartz function
- ②  $\varphi(0) = \widehat{\varphi}(0) = 240\pi$       ( $\widehat{\varphi}$  = the Fourier transform of  $\varphi$ )

# Viazovska's magic function (continued)

## Theorem (Viazovska)

$\varphi$  is a magic function for dimension 8. More precisely, it has the following properties:

- ①  $\varphi$  is a Schwartz function
- ②  $\varphi(0) = \widehat{\varphi}(0) = 240\pi$       ( $\widehat{\varphi}$  = the Fourier transform of  $\varphi$ )
- ③  $\widehat{\varphi}(x) \geq 0$  for all  $x \in \mathbb{R}^8$

# Viazovska's magic function (continued)

## Theorem (Viazovska)

$\varphi$  is a magic function for dimension 8. More precisely, it has the following properties:

- ①  $\varphi$  is a Schwartz function
- ②  $\varphi(0) = \widehat{\varphi}(0) = 240\pi$       ( $\widehat{\varphi}$  = the Fourier transform of  $\varphi$ )
- ③  $\widehat{\varphi}(x) \geq 0$  for all  $x \in \mathbb{R}^8$
- ④  $\varphi(x) \leq 0$  for all  $x \in \mathbb{R}^8$  with  $\|x\| \geq \sqrt{2}$

# Viazovska's magic function (continued)

## Theorem (Viazovska)

$\varphi$  is a magic function for dimension 8. More precisely, it has the following properties:

- ①  $\varphi$  is a Schwartz function
- ②  $\varphi(0) = \widehat{\varphi}(0) = 240\pi$       ( $\widehat{\varphi}$  = the Fourier transform of  $\varphi$ )
- ③  $\widehat{\varphi}(x) \geq 0$  for all  $x \in \mathbb{R}^8$
- ④  $\varphi(x) \leq 0$  for all  $x \in \mathbb{R}^8$  with  $\|x\| \geq \sqrt{2}$

Using the Cohn-Elkies linear programming bound, the above properties imply that  $\varphi$  certifies an upper bound of  $\frac{\pi^4}{384}$  for sphere packing density in  $\mathbb{R}^8$ . This matches the packing density of the  $E_8$  lattice packing.

# Viazovska's magic function (continued)

## Theorem (Viazovska)

$\varphi$  is a magic function for dimension 8. More precisely, it has the following properties:

- ①  $\varphi$  is a Schwartz function
- ②  $\varphi(0) = \widehat{\varphi}(0) = 240\pi$       ( $\widehat{\varphi}$  = the Fourier transform of  $\varphi$ )
- ③  $\widehat{\varphi}(x) \geq 0$  for all  $x \in \mathbb{R}^8$
- ④  $\varphi(x) \leq 0$  for all  $x \in \mathbb{R}^8$  with  $\|x\| \geq \sqrt{2}$

Using the Cohn-Elkies linear programming bound, the above properties imply that  $\varphi$  certifies an upper bound of  $\frac{\pi^4}{384}$  for sphere packing density in  $\mathbb{R}^8$ . This matches the packing density of the  $E_8$  lattice packing.

It remains to prove the claimed properties. This is not trivial.

# Viazovska's magic function (continued)

## Theorem (Viazovska)

$\varphi$  is a magic function for dimension 8. More precisely, it has the following properties:

- ①  $\varphi$  is a Schwartz function
- ②  $\varphi(0) = \widehat{\varphi}(0) = 240\pi$       ( $\widehat{\varphi}$  = the Fourier transform of  $\varphi$ )
- ③  $\widehat{\varphi}(x) \geq 0$  for all  $x \in \mathbb{R}^8$
- ④  $\varphi(x) \leq 0$  for all  $x \in \mathbb{R}^8$  with  $\|x\| \geq \sqrt{2}$

Using the Cohn-Elkies linear programming bound, the above properties imply that  $\varphi$  certifies an upper bound of  $\frac{\pi^4}{384}$  for sphere packing density in  $\mathbb{R}^8$ . This matches the packing density of the  $E_8$  lattice packing.

It remains to prove the claimed properties. This is not trivial.  
(Related, and much more nontrivial: the reasoning that led to the strange formula for  $\varphi$ .)

## The modular forms in the definition of $\varphi$

The problem boils down to understanding the properties of the modular forms in the definition of  $\varphi$ . Let  $\mathbb{H}$  denote the upper half plane. Define functions  $U : \mathbb{H} \rightarrow \mathbb{C}$ ,  $V : \mathbb{H} \rightarrow \mathbb{C}$  by

$$U(z) = 108 \frac{(zE'_4(z) + 4E_4(z))^2}{E_4(z)^3 - E_6(z)^2}$$
$$V(z) = 128 \left( \frac{\theta_3(z)^4 + \theta_4(z)^4}{\theta_2(z)^8} + \frac{\theta_4(z)^4 - \theta_2(z)^4}{\theta_3(z)^8} \right).$$

## The modular forms in the definition of $\varphi$

The problem boils down to understanding the properties of the modular forms in the definition of  $\varphi$ . Let  $\mathbb{H}$  denote the upper half plane. Define functions  $U : \mathbb{H} \rightarrow \mathbb{C}$ ,  $V : \mathbb{H} \rightarrow \mathbb{C}$  by

$$U(z) = 108 \frac{(zE'_4(z) + 4E_4(z))^2}{E_4(z)^3 - E_6(z)^2}$$
$$V(z) = 128 \left( \frac{\theta_3(z)^4 + \theta_4(z)^4}{\theta_2(z)^8} + \frac{\theta_4(z)^4 - \theta_2(z)^4}{\theta_3(z)^8} \right).$$

Define functions  $\varphi_{\pm} : \mathbb{R}^8 \rightarrow \mathbb{R}$  by (the analytic continuation of)

$$\varphi_+(x) = -4 \sin^2 \left( \frac{\pi \|x\|^2}{2} \right) \int_0^\infty e^{-\pi t \|x\|^2} U(it) dt$$

$$\varphi_-(x) = -4 \sin^2 \left( \frac{\pi \|x\|^2}{2} \right) \int_0^\infty e^{-\pi t \|x\|^2} V(it) dt,$$

# The modular forms in the definition of $\varphi$

The problem boils down to understanding the properties of the modular forms in the definition of  $\varphi$ . Let  $\mathbb{H}$  denote the upper half plane. Define functions  $U : \mathbb{H} \rightarrow \mathbb{C}$ ,  $V : \mathbb{H} \rightarrow \mathbb{C}$  by

$$U(z) = 108 \frac{(zE'_4(z) + 4E_4(z))^2}{E_4(z)^3 - E_6(z)^2}$$
$$V(z) = 128 \left( \frac{\theta_3(z)^4 + \theta_4(z)^4}{\theta_2(z)^8} + \frac{\theta_4(z)^4 - \theta_2(z)^4}{\theta_3(z)^8} \right).$$

Define functions  $\varphi_{\pm} : \mathbb{R}^8 \rightarrow \mathbb{R}$  by (the analytic continuation of)

$$\varphi_+(x) = -4 \sin^2 \left( \frac{\pi \|x\|^2}{2} \right) \int_0^\infty e^{-\pi t \|x\|^2} U(it) dt$$

$$\varphi_-(x) = -4 \sin^2 \left( \frac{\pi \|x\|^2}{2} \right) \int_0^\infty e^{-\pi t \|x\|^2} V(it) dt,$$

so that  $\varphi = \varphi_+ + \varphi_-$ .

## Viazovska's modular form inequalities

The definitions of  $U(z)$ ,  $V(z)$  were carefully chosen to satisfy several conditions, including, crucially,

$$\widehat{\varphi_+} = \varphi_+, \quad \widehat{\varphi_-} = -\varphi_-.$$

(That is,  $\varphi_{\pm}$  are eigenfunctions of the Fourier transform with resp. eigenvalues  $\pm 1$ .)

## Viazovska's modular form inequalities

The definitions of  $U(z)$ ,  $V(z)$  were carefully chosen to satisfy several conditions, including, crucially,

$$\widehat{\varphi_+} = \varphi_+, \quad \widehat{\varphi_-} = -\varphi_-.$$

(That is,  $\varphi_{\pm}$  are eigenfunctions of the Fourier transform with resp. eigenvalues  $\pm 1$ .) Thus, we have the relations

$$\varphi = \varphi_+ + \varphi_-$$

## Viazovska's modular form inequalities

The definitions of  $U(z)$ ,  $V(z)$  were carefully chosen to satisfy several conditions, including, crucially,

$$\widehat{\varphi_+} = \varphi_+, \quad \widehat{\varphi_-} = -\varphi_-.$$

(That is,  $\varphi_{\pm}$  are eigenfunctions of the Fourier transform with resp. eigenvalues  $\pm 1$ .) Thus, we have the relations

$$\varphi = \varphi_+ + \varphi_- = -4 \sin^2 \left( \frac{\pi \|x\|^2}{2} \right) \int_0^\infty e^{-\pi t \|x\|^2} (U(it) + V(it)) dt,$$

## Viazovska's modular form inequalities

The definitions of  $U(z)$ ,  $V(z)$  were carefully chosen to satisfy several conditions, including, crucially,

$$\widehat{\varphi_+} = \varphi_+, \quad \widehat{\varphi_-} = -\varphi_-.$$

(That is,  $\varphi_{\pm}$  are eigenfunctions of the Fourier transform with resp. eigenvalues  $\pm 1$ .) Thus, we have the relations

$$\varphi = \varphi_+ + \varphi_- = -4 \sin^2\left(\frac{\pi \|x\|^2}{2}\right) \int_0^\infty e^{-\pi t \|x\|^2} (U(it) + V(it)) dt,$$

$$\widehat{\varphi} = \varphi_+ - \varphi_-$$

## Viazovska's modular form inequalities

The definitions of  $U(z)$ ,  $V(z)$  were carefully chosen to satisfy several conditions, including, crucially,

$$\widehat{\varphi_+} = \varphi_+, \quad \widehat{\varphi_-} = -\varphi_-.$$

(That is,  $\varphi_{\pm}$  are eigenfunctions of the Fourier transform with resp. eigenvalues  $\pm 1$ .) Thus, we have the relations

$$\varphi = \varphi_+ + \varphi_- = -4 \sin^2 \left( \frac{\pi \|x\|^2}{2} \right) \int_0^\infty e^{-\pi t \|x\|^2} (U(it) + V(it)) dt,$$

$$\widehat{\varphi} = \varphi_+ - \varphi_- = -4 \sin^2 \left( \frac{\pi \|x\|^2}{2} \right) \int_0^\infty e^{-\pi t \|x\|^2} (U(it) - V(it)) dt.$$

## Viazovska's modular form inequalities

The definitions of  $U(z)$ ,  $V(z)$  were carefully chosen to satisfy several conditions, including, crucially,

$$\widehat{\varphi_+} = \varphi_+, \quad \widehat{\varphi_-} = -\varphi_-.$$

(That is,  $\varphi_{\pm}$  are eigenfunctions of the Fourier transform with resp. eigenvalues  $\pm 1$ .) Thus, we have the relations

$$\varphi = \varphi_+ + \varphi_- = -4 \sin^2 \left( \frac{\pi \|x\|^2}{2} \right) \int_0^\infty e^{-\pi t \|x\|^2} (U(it) + V(it)) dt,$$

$$\widehat{\varphi} = \varphi_+ - \varphi_- = -4 \sin^2 \left( \frac{\pi \|x\|^2}{2} \right) \int_0^\infty e^{-\pi t \|x\|^2} (U(it) - V(it)) dt.$$

The inequalities for  $\varphi$  and  $\widehat{\varphi}$  will therefore follow from the following result:

# Viazovska's modular form inequalities

The definitions of  $U(z)$ ,  $V(z)$  were carefully chosen to satisfy several conditions, including, crucially,

$$\widehat{\varphi_+} = \varphi_+, \quad \widehat{\varphi_-} = -\varphi_-.$$

(That is,  $\varphi_{\pm}$  are eigenfunctions of the Fourier transform with resp. eigenvalues  $\pm 1$ .) Thus, we have the relations

$$\varphi = \varphi_+ + \varphi_- = -4 \sin^2 \left( \frac{\pi \|x\|^2}{2} \right) \int_0^\infty e^{-\pi t \|x\|^2} (U(it) + V(it)) dt,$$

$$\widehat{\varphi} = \varphi_+ - \varphi_- = -4 \sin^2 \left( \frac{\pi \|x\|^2}{2} \right) \int_0^\infty e^{-\pi t \|x\|^2} (U(it) - V(it)) dt.$$

The inequalities for  $\varphi$  and  $\widehat{\varphi}$  will therefore follow from the following result:

## Theorem (Viazovska)

The functions  $U, V$  satisfy the inequalities

$$U(it) + V(it) \geq 0 \quad (t > 0) \tag{V1}$$

$$U(it) - V(it) \leq 0 \quad (t > 0) \tag{V2}$$

# Viazovska's modular form inequalities (continued)

Theorem (Viazovska)

The functions  $U, V$  satisfy the inequalities

$$U(it) + V(it) \geq 0 \quad (t > 0) \quad (\text{V1})$$

$$U(it) - V(it) \leq 0 \quad (t > 0) \quad (\text{V2})$$

# Viazovska's modular form inequalities (continued)

## Theorem (Viazovska)

The functions  $U, V$  satisfy the inequalities

$$U(it) + V(it) \geq 0 \quad (t > 0) \tag{V1}$$

$$U(it) - V(it) \leq 0 \quad (t > 0) \tag{V2}$$

- The inequalities (V1)–(V2) seem unnatural, because they relate modular forms that belong to different modular form spaces. This makes it difficult to think of a conceptual reason why they should be true.

# Viazovska's modular form inequalities (continued)

## Theorem (Viazovska)

The functions  $U, V$  satisfy the inequalities

$$U(it) + V(it) \geq 0 \quad (t > 0) \tag{V1}$$

$$U(it) - V(it) \leq 0 \quad (t > 0) \tag{V2}$$

- The inequalities (V1)–(V2) seem unnatural, because they relate modular forms that belong to different modular form spaces. This makes it difficult to think of a conceptual reason why they should be true.
- Viazovska proved them using (rigorously supported) numerical calculations.

# Viazovska's modular form inequalities (continued)


## Theorem (Viazovska)

The functions  $U, V$  satisfy the inequalities

$$U(it) + V(it) \geq 0 \quad (t > 0) \tag{V1}$$

$$U(it) - V(it) \leq 0 \quad (t > 0) \tag{V2}$$

- The inequalities (V1)–(V2) seem unnatural, because they relate modular forms that belong to different modular form spaces. This makes it difficult to think of a conceptual reason why they should be true.
- Viazovska proved them using (rigorously supported) numerical calculations.



## A new proof

I will now show a new proof of (V1)–(V2) that does not rely on computer calculations.

## A new proof

I will now show a new proof of (V1)–(V2) that does not rely on computer calculations. Recall a few facts about modular forms:

## A new proof

I will now show a new proof of (V1)–(V2) that does not rely on computer calculations. Recall a few facts about modular forms:

**Reminder (1).**  $E_4, E_6, \theta_2, \theta_3, \theta_4$  are **modular forms** satisfying the modular transformation properties

$$E_4(z+1) = E_4(z), \quad E_4(-1/z) = z^4 E_4(z),$$

$$E_6(z+1) = E_6(z), \quad E_6(-1/z) = z^6 E_6(z),$$

$$\theta_2(z+1)^4 = -\theta_2(z)^4, \quad \theta_2(-1/z)^4 = -z^2 \theta_4(z)^4,$$

$$\theta_3(z+1)^4 = \theta_4(z)^4, \quad \theta_3(-1/z)^4 = -z^2 \theta_3(z)^4,$$

$$\theta_4(z+1)^4 = \theta_3(z)^4, \quad \theta_4(-1/z)^4 = -z^2 \theta_2(z)^4.$$

# A new proof

I will now show a new proof of (V1)–(V2) that does not rely on computer calculations. Recall a few facts about modular forms:

**Reminder (1).**  $E_4, E_6, \theta_2, \theta_3, \theta_4$  are **modular forms** satisfying the modular transformation properties

$$E_4(z+1) = E_4(z), \quad E_4(-1/z) = z^4 E_4(z),$$

$$E_6(z+1) = E_6(z), \quad E_6(-1/z) = z^6 E_6(z),$$

$$\theta_2(z+1)^4 = -\theta_2(z)^4, \quad \theta_2(-1/z)^4 = -z^2 \theta_4(z)^4,$$

$$\theta_3(z+1)^4 = \theta_4(z)^4, \quad \theta_3(-1/z)^4 = -z^2 \theta_3(z)^4,$$

$$\theta_4(z+1)^4 = \theta_3(z)^4, \quad \theta_4(-1/z)^4 = -z^2 \theta_2(z)^4.$$

**Reminder (2).**  $E_4^3 - E_6^2$  is a scaling of the **modular discriminant**:

$$E_4^3 - E_6^2 = \frac{1728}{(2\pi)^{12}} \Delta(z)$$

# A new proof

I will now show a new proof of (V1)–(V2) that does not rely on computer calculations. Recall a few facts about modular forms:

**Reminder (1).**  $E_4, E_6, \theta_2, \theta_3, \theta_4$  are **modular forms** satisfying the modular transformation properties

$$E_4(z+1) = E_4(z), \quad E_4(-1/z) = z^4 E_4(z),$$

$$E_6(z+1) = E_6(z), \quad E_6(-1/z) = z^6 E_6(z),$$

$$\theta_2(z+1)^4 = -\theta_2(z)^4, \quad \theta_2(-1/z)^4 = -z^2 \theta_4(z)^4,$$

$$\theta_3(z+1)^4 = \theta_4(z)^4, \quad \theta_3(-1/z)^4 = -z^2 \theta_3(z)^4,$$

$$\theta_4(z+1)^4 = \theta_3(z)^4, \quad \theta_4(-1/z)^4 = -z^2 \theta_2(z)^4.$$

**Reminder (2).**  $E_4^3 - E_6^2$  is a scaling of the **modular discriminant**:

$$E_4^3 - E_6^2 = \frac{1728}{(2\pi)^{12}} \Delta(z) = 1728q^2 \prod_{n=1}^{\infty} (1 - q^{2n})^{24}$$

# A new proof

I will now show a new proof of (V1)–(V2) that does not rely on computer calculations. Recall a few facts about modular forms:

**Reminder (1).**  $E_4, E_6, \theta_2, \theta_3, \theta_4$  are **modular forms** satisfying the modular transformation properties

$$E_4(z+1) = E_4(z), \quad E_4(-1/z) = z^4 E_4(z),$$

$$E_6(z+1) = E_6(z), \quad E_6(-1/z) = z^6 E_6(z),$$

$$\theta_2(z+1)^4 = -\theta_2(z)^4, \quad \theta_2(-1/z)^4 = -z^2 \theta_4(z)^4,$$

$$\theta_3(z+1)^4 = \theta_4(z)^4, \quad \theta_3(-1/z)^4 = -z^2 \theta_3(z)^4,$$

$$\theta_4(z+1)^4 = \theta_3(z)^4, \quad \theta_4(-1/z)^4 = -z^2 \theta_2(z)^4.$$

**Reminder (2).**  $E_4^3 - E_6^2$  is a scaling of the **modular discriminant**:

$$E_4^3 - E_6^2 = \frac{1728}{(2\pi)^{12}} \Delta(z) = 1728q^2 \prod_{n=1}^{\infty} (1 - q^{2n})^{24} = \frac{27}{4} (\theta_2 \theta_3 \theta_4)^8.$$

## A new proof

I will now show a new proof of (V1)–(V2) that does not rely on computer calculations. Recall a few facts about modular forms:

**Reminder (1).**  $E_4, E_6, \theta_2, \theta_3, \theta_4$  are **modular forms** satisfying the modular transformation properties

$$E_4(z+1) = E_4(z), \quad E_4(-1/z) = z^4 E_4(z),$$

$$E_6(z+1) = E_6(z), \quad E_6(-1/z) = z^6 E_6(z),$$

$$\theta_2(z+1)^4 = -\theta_2(z)^4, \quad \theta_2(-1/z)^4 = -z^2 \theta_4(z)^4,$$

$$\theta_3(z+1)^4 = \theta_4(z)^4, \quad \theta_3(-1/z)^4 = -z^2 \theta_3(z)^4,$$

$$\theta_4(z+1)^4 = \theta_3(z)^4, \quad \theta_4(-1/z)^4 = -z^2 \theta_2(z)^4.$$

**Reminder (2).**  $E_4^3 - E_6^2$  is a scaling of the **modular discriminant**:

$$E_4^3 - E_6^2 = \frac{1728}{(2\pi)^{12}} \Delta(z) = 1728q^2 \prod_{n=1}^{\infty} (1 - q^{2n})^{24} = \frac{27}{4} (\theta_2 \theta_3 \theta_4)^8.$$

**Reminder (3).** The **modular lambda function** is

$$\lambda = \frac{\theta_2^4}{\theta_3^4} = 1 - \frac{\theta_4^4}{\theta_3^4}.$$

# A new proof

I will now show a new proof of (V1)–(V2) that does not rely on computer calculations. Recall a few facts about modular forms:

**Reminder (1).**  $E_4, E_6, \theta_2, \theta_3, \theta_4$  are **modular forms** satisfying the modular transformation properties

$$E_4(z+1) = E_4(z), \quad E_4(-1/z) = z^4 E_4(z),$$

$$E_6(z+1) = E_6(z), \quad E_6(-1/z) = z^6 E_6(z),$$

$$\theta_2(z+1)^4 = -\theta_2(z)^4, \quad \theta_2(-1/z)^4 = -z^2 \theta_4(z)^4,$$

$$\theta_3(z+1)^4 = \theta_4(z)^4, \quad \theta_3(-1/z)^4 = -z^2 \theta_3(z)^4,$$

$$\theta_4(z+1)^4 = \theta_3(z)^4, \quad \theta_4(-1/z)^4 = -z^2 \theta_2(z)^4.$$

**Reminder (2).**  $E_4^3 - E_6^2$  is a scaling of the **modular discriminant**:

$$E_4^3 - E_6^2 = \frac{1728}{(2\pi)^{12}} \Delta(z) = 1728q^2 \prod_{n=1}^{\infty} (1 - q^{2n})^{24} = \frac{27}{4} (\theta_2 \theta_3 \theta_4)^8.$$

**Reminder (3).** The **modular lambda function** is

$$\lambda = \frac{\theta_2^4}{\theta_3^4} = 1 - \frac{\theta_4^4}{\theta_3^4}. \text{ For } t > 0, \lambda(it) \in (0, 1).$$

## A new proof of (V1)–(V2), part I: proof of (V1)

## A new proof of (V1)–(V2), part I: proof of (V1)

First, observe that  $U(it) \geq 0$  for all  $t > 0$  since, by inspection of the relevant Fourier series, we have  $itE'_4(it) + 4E_4(it) \in \mathbb{R}$ , and separately we have  $E_4(z)^3 - E_6(z)^2 > 0$  for  $t > 0$  by the infinite product formula from the previous slide.

## A new proof of (V1)–(V2), part I: proof of (V1)

First, observe that  $U(it) \geq 0$  for all  $t > 0$  since, by inspection of the relevant Fourier series, we have  $itE'_4(it) + 4E_4(it) \in \mathbb{R}$ , and separately we have  $E_4(z)^3 - E_6(z)^2 > 0$  for  $t > 0$  by the infinite product formula from the previous slide.

Similarly, it also holds that  $V(it) \geq 0$  for  $t > 0$ . To see this, rewrite  $V(z)$  in terms of  $\theta_3$  and the modular lambda function as

## A new proof of (V1)–(V2), part I: proof of (V1)

First, observe that  $U(it) \geq 0$  for all  $t > 0$  since, by inspection of the relevant Fourier series, we have  $itE'_4(it) + 4E_4(it) \in \mathbb{R}$ , and separately we have  $E_4(z)^3 - E_6(z)^2 > 0$  for  $t > 0$  by the infinite product formula from the previous slide.

Similarly, it also holds that  $V(it) \geq 0$  for  $t > 0$ . To see this, rewrite  $V(z)$  in terms of  $\theta_3$  and the modular lambda function as

$$V = 128 \left( \frac{\theta_3^4 + \theta_4^4}{\theta_2^8} + \frac{\theta_4^4 - \theta_2^4}{\theta_3^8} \right) = \dots = \frac{128}{\theta_3^4} \frac{(1 - \lambda)(2 + \lambda + 2\lambda^2)}{\lambda^2}.$$

## A new proof of (V1)–(V2), part I: proof of (V1)

First, observe that  $U(it) \geq 0$  for all  $t > 0$  since, by inspection of the relevant Fourier series, we have  $itE'_4(it) + 4E_4(it) \in \mathbb{R}$ , and separately we have  $E_4(z)^3 - E_6(z)^2 > 0$  for  $t > 0$  by the infinite product formula from the previous slide.

Similarly, it also holds that  $V(it) \geq 0$  for  $t > 0$ . To see this, rewrite  $V(z)$  in terms of  $\theta_3$  and the modular lambda function as

$$V = 128 \left( \frac{\theta_3^4 + \theta_4^4}{\theta_2^8} + \frac{\theta_4^4 - \theta_2^4}{\theta_3^8} \right) = \dots = \frac{128}{\theta_3^4} \frac{(1-\lambda)(2+\lambda+2\lambda^2)}{\lambda^2}.$$

Then use the facts that  $\theta_3(it) > 0$  (trivially), that  $\lambda(it) \in (0, 1)$  for  $t > 0$ , and that the map  $x \mapsto \frac{(1-x)(2+x+2x^2)}{x^2}$  takes positive values for  $x \in (0, 1)$ .

## A new proof of (V1)–(V2), part II: proof of (V2)

# A new proof of (V1)–(V2), part II: proof of (V2)

## **Step 1: A bit of cleanup**

# A new proof of (V1)–(V2), part II: proof of (V2)

## Step 1: A bit of cleanup

Define functions

$$F(z) = \frac{1}{108}(E_4^3 - E_6^2)U(z) = (E'_4)^2 z^2 + 8E_4 E'_4 z + 16E_4^2,$$

$$\tilde{F}(z) = \frac{1}{108}(E_4^3 - E_6^2)z^2 U(-1/z) = (E'_4)^2,$$

$$G(z) = \frac{1}{108}(E_4^3 - E_6^2)V(z) = 8\theta_4^8(\theta_3^{12} + \theta_4^4\theta_3^8 + \theta_2^8\theta_4^4 - \theta_2^{12}),$$

$$\tilde{G}(z) = \frac{1}{108}(E_4^3 - E_6^2)z^2 V(-1/z) = -8\theta_2^8(\theta_3^{12} + \theta_2^4\theta_3^8 + \theta_2^4\theta_4^8 - \theta_4^{12})$$

# A new proof of (V1)–(V2), part II: proof of (V2)

## Step 1: A bit of cleanup

Define functions

$$F(z) = \frac{1}{108}(E_4^3 - E_6^2)U(z) = (E'_4)^2 z^2 + 8E_4 E'_4 z + 16E_4^2,$$

$$\tilde{F}(z) = \frac{1}{108}(E_4^3 - E_6^2)z^2 U(-1/z) = (E'_4)^2,$$

$$G(z) = \frac{1}{108}(E_4^3 - E_6^2)V(z) = 8\theta_4^8(\theta_3^{12} + \theta_4^4\theta_3^8 + \theta_2^8\theta_4^4 - \theta_2^{12}),$$

$$\tilde{G}(z) = \frac{1}{108}(E_4^3 - E_6^2)z^2 V(-1/z) = -8\theta_2^8(\theta_3^{12} + \theta_2^4\theta_3^8 + \theta_2^4\theta_4^8 - \theta_4^{12})$$

(making use of the modular transformation properties).

# A new proof of (V1)–(V2), part II: proof of (V2)

## Step 1: A bit of cleanup

Define functions

$$F(z) = \frac{1}{108}(E_4^3 - E_6^2)U(z) = (E'_4)^2 z^2 + 8E_4 E'_4 z + 16E_4^2,$$

$$\tilde{F}(z) = \frac{1}{108}(E_4^3 - E_6^2)z^2 U(-1/z) = (E'_4)^2,$$

$$G(z) = \frac{1}{108}(E_4^3 - E_6^2)V(z) = 8\theta_4^8(\theta_3^{12} + \theta_4^4\theta_3^8 + \theta_2^8\theta_4^4 - \theta_2^{12}),$$

$$\tilde{G}(z) = \frac{1}{108}(E_4^3 - E_6^2)z^2 V(-1/z) = -8\theta_2^8(\theta_3^{12} + \theta_2^4\theta_3^8 + \theta_2^4\theta_4^8 - \theta_4^{12})$$

(making use of the modular transformation properties).

Trivially, the inequality (V2) is equivalent to the pair of inequalities

$$-\tilde{F}(it) < -\tilde{G}(it) \quad (t \geq 1), \tag{V2-I}$$

$$F(it) < G(it) \quad (t \geq 1). \tag{V2-II}$$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Step 2: Understanding the behavior at  $t = 1$**

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

### Step 2: Understanding the behavior at $t = 1$

Theorem (Gauss, Ramanujan, folklore)

We have the explicit evaluations

$$\begin{aligned} E_4(i) &= \frac{3\Gamma(1/4)^8}{64\pi^6}, & \theta_2(i) &= \frac{\Gamma(1/4)}{(2\pi)^{3/4}}, \\ E'_4(i) &= \frac{3\Gamma(1/4)^8}{32\pi^6}i, & \theta_3(i) &= \frac{\Gamma(1/4)}{\sqrt{2}\pi^{3/4}}, \\ && \theta_4(i) &= \frac{\Gamma(1/4)}{(2\pi)^{3/4}}, \end{aligned}$$

(where  $\Gamma(\cdot)$  denotes the Euler gamma function).

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

### Step 2: Understanding the behavior at $t = 1$

Theorem (Gauss, Ramanujan, folklore)

We have the explicit evaluations

$$\begin{aligned} E_4(i) &= \frac{3\Gamma(1/4)^8}{64\pi^6}, & \theta_2(i) &= \frac{\Gamma(1/4)}{(2\pi)^{3/4}}, \\ E'_4(i) &= \frac{3\Gamma(1/4)^8}{32\pi^6}i, & \theta_3(i) &= \frac{\Gamma(1/4)}{\sqrt{2}\pi^{3/4}}, \\ && \theta_4(i) &= \frac{\Gamma(1/4)}{(2\pi)^{3/4}}, \end{aligned}$$

(where  $\Gamma(\cdot)$  denotes the Euler gamma function).

See p. 257 of my book for a proof sketch and references.

# A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

## **Step 3: Leveraging monotonicity**

# A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

## Step 3: Leveraging monotonicity

**Proof of (V2-I).** Observe that

$$\begin{aligned}-\tilde{F}(z) &= 230400\pi^2 q^4 + 8294400\pi^2 q^6 + 113356800\pi^2 q^8 \\ &\quad + 831283200\pi^2 q^{10} + 4337971200\pi^2 q^{12} + \dots,\end{aligned}$$

$$\begin{aligned}-\tilde{G}(z) &= 163840q^3 + 16121856q^5 + 333250560q^7 + \\ &\quad + 3199467520q^9 + 19472547840q^{11} + \dots.\end{aligned}$$

# A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

## Step 3: Leveraging monotonicity

**Proof of (V2-I).** Observe that

$$\begin{aligned}-\tilde{F}(z) &= 230400\pi^2 q^4 + 8294400\pi^2 q^6 + 113356800\pi^2 q^8 \\ &\quad + 831283200\pi^2 q^{10} + 4337971200\pi^2 q^{12} + \dots,\end{aligned}$$

$$\begin{aligned}-\tilde{G}(z) &= 163840q^3 + 16121856q^5 + 333250560q^7 + \\ &\quad + 3199467520q^9 + 19472547840q^{11} + \dots\end{aligned}$$

Note that  $q^4 = e^{-4\pi t} \ll e^{-3\pi t} = q^3$  for  $t$  large, so the inequality  $-\tilde{F}(it) < -\tilde{G}(it)$  holds asymptotically.

# A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

## Step 3: Leveraging monotonicity

**Proof of (V2-I).** Observe that

$$\begin{aligned}-\tilde{F}(z) &= 230400\pi^2 q^4 + 8294400\pi^2 q^6 + 113356800\pi^2 q^8 \\ &\quad + 831283200\pi^2 q^{10} + 4337971200\pi^2 q^{12} + \dots,\end{aligned}$$

$$\begin{aligned}-\tilde{G}(z) &= 163840q^3 + 16121856q^5 + 333250560q^7 + \\ &\quad + 3199467520q^9 + 19472547840q^{11} + \dots\end{aligned}$$

Note that  $q^4 = e^{-4\pi t} \ll e^{-3\pi t} = q^3$  for  $t$  large, so the inequality  $-\tilde{F}(it) < -\tilde{G}(it)$  holds asymptotically. To prove the stronger claim that it holds for  $t \geq 1$ , note that the Fourier coefficients in both series are positive.\*

# A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

## Step 3: Leveraging monotonicity

**Proof of (V2-I).** Observe that

$$\begin{aligned}-\tilde{F}(z) &= 230400\pi^2 q^4 + 8294400\pi^2 q^6 + 113356800\pi^2 q^8 \\ &\quad + 831283200\pi^2 q^{10} + 4337971200\pi^2 q^{12} + \dots,\end{aligned}$$

$$\begin{aligned}-\tilde{G}(z) &= 163840q^3 + 16121856q^5 + 333250560q^7 + \\ &\quad + 3199467520q^9 + 19472547840q^{11} + \dots\end{aligned}$$

Note that  $q^4 = e^{-4\pi t} \ll e^{-3\pi t} = q^3$  for  $t$  large, so the inequality  $-\tilde{F}(it) < -\tilde{G}(it)$  holds asymptotically. To prove the stronger claim that it holds for  $t \geq 1$ , note that the Fourier coefficients in both series are positive.\*

\* This is easy to prove from the definitions.

# A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

## Step 3: Leveraging monotonicity

**Proof of (V2-I).** Observe that

$$\begin{aligned}-\tilde{F}(z) &= 230400\pi^2 q^4 + 8294400\pi^2 q^6 + 113356800\pi^2 q^8 \\ &\quad + 831283200\pi^2 q^{10} + 4337971200\pi^2 q^{12} + \dots,\end{aligned}$$

$$\begin{aligned}-\tilde{G}(z) &= 163840q^3 + 16121856q^5 + 333250560q^7 + \\ &\quad + 3199467520q^9 + 19472547840q^{11} + \dots\end{aligned}$$

Note that  $q^4 = e^{-4\pi t} \ll e^{-3\pi t} = q^3$  for  $t$  large, so the inequality  $-\tilde{F}(it) < -\tilde{G}(it)$  holds asymptotically. To prove the stronger claim that it holds for  $t \geq 1$ , note that the Fourier coefficients in both series are positive.\* In particular, the function  $t \mapsto -q^{-3}\tilde{F}(it)$  is a decreasing function of  $t$ , so that for  $t \geq 1$ ,

\* This is easy to prove from the definitions.

# A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

## Step 3: Leveraging monotonicity

**Proof of (V2-I).** Observe that

$$\begin{aligned}-\tilde{F}(z) &= 230400\pi^2 q^4 + 8294400\pi^2 q^6 + 113356800\pi^2 q^8 \\ &\quad + 831283200\pi^2 q^{10} + 4337971200\pi^2 q^{12} + \dots,\end{aligned}$$

$$\begin{aligned}-\tilde{G}(z) &= 163840q^3 + 16121856q^5 + 333250560q^7 + \\ &\quad + 3199467520q^9 + 19472547840q^{11} + \dots\end{aligned}$$

Note that  $q^4 = e^{-4\pi t} \ll e^{-3\pi t} = q^3$  for  $t$  large, so the inequality  $-\tilde{F}(it) < -\tilde{G}(it)$  holds asymptotically. To prove the stronger claim that it holds for  $t \geq 1$ , note that the Fourier coefficients in both series are positive.\* In particular, the function  $t \mapsto -q^{-3}\tilde{F}(it)$  is a decreasing function of  $t$ , so that for  $t \geq 1$ ,

$$-e^{3\pi t}\tilde{F}(it)$$

\* This is easy to prove from the definitions.

# A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

## Step 3: Leveraging monotonicity

**Proof of (V2-I).** Observe that

$$\begin{aligned}-\tilde{F}(z) &= 230400\pi^2 q^4 + 8294400\pi^2 q^6 + 113356800\pi^2 q^8 \\ &\quad + 831283200\pi^2 q^{10} + 4337971200\pi^2 q^{12} + \dots,\end{aligned}$$

$$\begin{aligned}-\tilde{G}(z) &= 163840q^3 + 16121856q^5 + 333250560q^7 + \\ &\quad + 3199467520q^9 + 19472547840q^{11} + \dots\end{aligned}$$

Note that  $q^4 = e^{-4\pi t} \ll e^{-3\pi t} = q^3$  for  $t$  large, so the inequality  $-\tilde{F}(it) < -\tilde{G}(it)$  holds asymptotically. To prove the stronger claim that it holds for  $t \geq 1$ , note that the Fourier coefficients in both series are positive.\* In particular, the function  $t \mapsto -q^{-3}\tilde{F}(it)$  is a decreasing function of  $t$ , so that for  $t \geq 1$ ,

$$-e^{3\pi t}\tilde{F}(it) \leq e^{3\pi}\tilde{F}(i)$$

\* This is easy to prove from the definitions.

# A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

## Step 3: Leveraging monotonicity

**Proof of (V2-I).** Observe that

$$\begin{aligned}-\tilde{F}(z) &= 230400\pi^2 q^4 + 8294400\pi^2 q^6 + 113356800\pi^2 q^8 \\ &\quad + 831283200\pi^2 q^{10} + 4337971200\pi^2 q^{12} + \dots,\end{aligned}$$

$$\begin{aligned}-\tilde{G}(z) &= 163840q^3 + 16121856q^5 + 333250560q^7 + \\ &\quad + 3199467520q^9 + 19472547840q^{11} + \dots\end{aligned}$$

Note that  $q^4 = e^{-4\pi t} \ll e^{-3\pi t} = q^3$  for  $t$  large, so the inequality  $-\tilde{F}(it) < -\tilde{G}(it)$  holds asymptotically. To prove the stronger claim that it holds for  $t \geq 1$ , note that the Fourier coefficients in both series are positive.\* In particular, the function  $t \mapsto -q^{-3}\tilde{F}(it)$  is a decreasing function of  $t$ , so that for  $t \geq 1$ ,

$$-e^{3\pi t}\tilde{F}(it) \leq e^{3\pi}\tilde{F}(i) = -e^{3\pi}E'_4(i)^2$$

\* This is easy to prove from the definitions.

# A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

## Step 3: Leveraging monotonicity

**Proof of (V2-I).** Observe that

$$\begin{aligned}-\tilde{F}(z) &= 230400\pi^2 q^4 + 8294400\pi^2 q^6 + 113356800\pi^2 q^8 \\ &\quad + 831283200\pi^2 q^{10} + 4337971200\pi^2 q^{12} + \dots,\end{aligned}$$

$$\begin{aligned}-\tilde{G}(z) &= 163840q^3 + 16121856q^5 + 333250560q^7 + \\ &\quad + 3199467520q^9 + 19472547840q^{11} + \dots\end{aligned}$$

Note that  $q^4 = e^{-4\pi t} \ll e^{-3\pi t} = q^3$  for  $t$  large, so the inequality  $-\tilde{F}(it) < -\tilde{G}(it)$  holds asymptotically. To prove the stronger claim that it holds for  $t \geq 1$ , note that the Fourier coefficients in both series are positive.\* In particular, the function  $t \mapsto -q^{-3}\tilde{F}(it)$  is a decreasing function of  $t$ , so that for  $t \geq 1$ ,

$$-e^{3\pi t}\tilde{F}(it) \leq e^{3\pi}\tilde{F}(i) = -e^{3\pi}E'_4(i)^2 = e^{3\pi} \frac{9\Gamma(1/4)^{16}}{1024\pi^{12}}$$

\* This is easy to prove from the definitions.

# A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

## Step 3: Leveraging monotonicity

**Proof of (V2-I).** Observe that

$$\begin{aligned}-\tilde{F}(z) &= 230400\pi^2 q^4 + 8294400\pi^2 q^6 + 113356800\pi^2 q^8 \\ &\quad + 831283200\pi^2 q^{10} + 4337971200\pi^2 q^{12} + \dots,\end{aligned}$$

$$\begin{aligned}-\tilde{G}(z) &= 163840q^3 + 16121856q^5 + 333250560q^7 + \\ &\quad + 3199467520q^9 + 19472547840q^{11} + \dots\end{aligned}$$

Note that  $q^4 = e^{-4\pi t} \ll e^{-3\pi t} = q^3$  for  $t$  large, so the inequality  $-\tilde{F}(it) < -\tilde{G}(it)$  holds asymptotically. To prove the stronger claim that it holds for  $t \geq 1$ , note that the Fourier coefficients in both series are positive.\* In particular, the function  $t \mapsto -q^{-3}\tilde{F}(it)$  is a decreasing function of  $t$ , so that for  $t \geq 1$ ,

$$-e^{3\pi t}\tilde{F}(it) \leq e^{3\pi}\tilde{F}(i) = -e^{3\pi}E'_4(i)^2 = e^{3\pi} \frac{9\Gamma(1/4)^{16}}{1024\pi^{12}} \approx 105043.78.$$

\* This is easy to prove from the definitions.

# A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

## Step 3: Leveraging monotonicity

**Proof of (V2-I).** Observe that

$$\begin{aligned}-\tilde{F}(z) &= 230400\pi^2 q^4 + 8294400\pi^2 q^6 + 113356800\pi^2 q^8 \\ &\quad + 831283200\pi^2 q^{10} + 4337971200\pi^2 q^{12} + \dots,\end{aligned}$$

$$\begin{aligned}-\tilde{G}(z) &= 163840q^3 + 16121856q^5 + 333250560q^7 + \\ &\quad + 3199467520q^9 + 19472547840q^{11} + \dots\end{aligned}$$

Note that  $q^4 = e^{-4\pi t} \ll e^{-3\pi t} = q^3$  for  $t$  large, so the inequality  $-\tilde{F}(it) < -\tilde{G}(it)$  holds asymptotically. To prove the stronger claim that it holds for  $t \geq 1$ , note that the Fourier coefficients in both series are positive.\* In particular, the function  $t \mapsto -q^{-3}\tilde{F}(it)$  is a decreasing function of  $t$ , so that for  $t \geq 1$ ,

$$-e^{3\pi t}\tilde{F}(it) \leq e^{3\pi}\tilde{F}(i) = -e^{3\pi}E'_4(i)^2 = e^{3\pi} \frac{9\Gamma(1/4)^{16}}{1024\pi^{12}} \approx 105043.78.$$

This in turn is  $< 163840$ , which is a lower bound for  $-e^{3\pi t}\tilde{G}(it)$ .


\* This is easy to prove from the definitions.

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

Summarizing this argument:

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)


Summarizing this argument:




Plots of  $-\tilde{F}(it)$ ,  $-\tilde{G}(it)$

# A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

Summarizing this argument:



Plots of  $-\tilde{F}(it)$ ,  $-\tilde{G}(it)$



Plots of  $-e^{3\pi t}\tilde{F}(it)$ ,  $-e^{3\pi t}\tilde{G}(it)$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Proof of (V2-II).** Imitating the approach for (V2-I), note that

$$\begin{aligned} F(it) &= 16 + (-3840\pi t + 7680)q^2 \\ &\quad + (230400\pi^2 t^2 - 990720\pi t + 990720)q^4 \\ &\quad + (8294400\pi^2 t^2 - 25205760\pi t + 16803840)q^6 + \dots, \end{aligned}$$

$$\begin{aligned} G(it) &= 16 + 1920q^2 - 81920q^3 + 1077120q^4 - 8060928q^5 \\ &\quad + 41725440q^6 - 166625280q^7 + 553054080q^8 + \dots, \end{aligned}$$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Proof of (V2-II).** Imitating the approach for (V2-I), note that

$$\begin{aligned} F(it) &= 16 + (-3840\pi t + 7680)q^2 \\ &\quad + (230400\pi^2 t^2 - 990720\pi t + 990720)q^4 \\ &\quad + (8294400\pi^2 t^2 - 25205760\pi t + 16803840)q^6 + \dots, \end{aligned}$$

$$\begin{aligned} G(it) &= 16 + 1920q^2 - 81920q^3 + 1077120q^4 - 8060928q^5 \\ &\quad + 41725440q^6 - 166625280q^7 + 553054080q^8 + \dots, \end{aligned}$$

Define renormalized functions

$$K(z) = -\frac{F(z) - 16}{q^2} = -q^{-2}(E'_4)^2 z^2 - 8q^{-2} E'_4 E_4 z - 16q^{-2}(E_4^2 - 1),$$

$$L(z) = -\frac{G(z) - 16}{q^2} = -8q^{-2} [\theta_4^8(\theta_3^{12} + \theta_4^4\theta_3^8 + \theta_2^8\theta_4^4 - \theta_2^{12}) - 2],$$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Proof of (V2-II).** Imitating the approach for (V2-I), note that

$$\begin{aligned} F(it) &= 16 + (-3840\pi t + 7680)q^2 \\ &\quad + (230400\pi^2 t^2 - 990720\pi t + 990720)q^4 \\ &\quad + (8294400\pi^2 t^2 - 25205760\pi t + 16803840)q^6 + \dots, \end{aligned}$$

$$\begin{aligned} G(it) &= 16 + 1920q^2 - 81920q^3 + 1077120q^4 - 8060928q^5 \\ &\quad + 41725440q^6 - 166625280q^7 + 553054080q^8 + \dots, \end{aligned}$$

Define renormalized functions

$$K(z) = -\frac{F(z) - 16}{q^2} = -q^{-2}(E'_4)^2 z^2 - 8q^{-2} E'_4 E_4 z - 16q^{-2}(E_4^2 - 1),$$

$$L(z) = -\frac{G(z) - 16}{q^2} = -8q^{-2} [\theta_4^8(\theta_3^{12} + \theta_4^4\theta_3^8 + \theta_2^8\theta_4^4 - \theta_2^{12}) - 2],$$

The inequality (V2-II) is thus equivalent to the inequality

$$K(it) > L(it) \quad (t \geq 1).$$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

As in the earlier proof, we will bound each of  $K(it)$  and  $L(it)$  separately, obtaining the inequality (V2-II) from the combination of the following two lemmas:

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

As in the earlier proof, we will bound each of  $K(it)$  and  $L(it)$  separately, obtaining the inequality (V2-II) from the combination of the following two lemmas:

### Lemma (1)

$$L(it) \leq 2297 \text{ for } t \geq 1.$$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

As in the earlier proof, we will bound each of  $K(it)$  and  $L(it)$  separately, obtaining the inequality (V2-II) from the combination of the following two lemmas:

**Lemma (1)**

$$L(it) \leq 2297 \text{ for } t \geq 1.$$

**Lemma (2)**

$$K(it) \geq 3747 \text{ for } t \geq 1.$$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Proof of Lemma (1).** Again the idea is to leverage monotonicity.

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Proof of Lemma (1).** Again the idea is to leverage monotonicity.

Define

$$\begin{aligned} H(z) &= \frac{L(z+1) - L(z)}{2} = \dots = 4q^{-2} (\theta_2^8(\theta_3^{12} - \theta_4^{12}) + \theta_2^{12}(\theta_3^8 + \theta_4^8)) \\ &= 81920q + 8060928q^3 + 166625280q^5 + 1599733760q^7 + \dots \end{aligned}$$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Proof of Lemma (1).** Again the idea is to leverage monotonicity.

Define

$$\begin{aligned} H(z) &= \frac{L(z+1) - L(z)}{2} = \dots = 4q^{-2} (\theta_2^8(\theta_3^{12} - \theta_4^{12}) + \theta_2^{12}(\theta_3^8 + \theta_4^8)) \\ &= 81920q + 8060928q^3 + 166625280q^5 + 1599733760q^7 + \dots \end{aligned}$$

Then for  $t \geq 1$ ,

$$\begin{aligned} L(it) &= -1920 + 81920q - 1077120q^2 + 8060928q^3 \\ &\quad - 41725440q^4 + 166625280q^5 - 553054080q^6 + \dots \end{aligned}$$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Proof of Lemma (1).** Again the idea is to leverage monotonicity.

Define

$$\begin{aligned} H(z) &= \frac{L(z+1) - L(z)}{2} = \dots = 4q^{-2} (\theta_2^8(\theta_3^{12} - \theta_4^{12}) + \theta_2^{12}(\theta_3^8 + \theta_4^8)) \\ &= 81920q + 8060928q^3 + 166625280q^5 + 1599733760q^7 + \dots \end{aligned}$$

Then for  $t \geq 1$ ,

$$\begin{aligned} L(it) &= -1920 + 81920q - 1077120q^2 + 8060928q^3 \\ &\quad - 41725440q^4 + 166625280q^5 - 553054080q^6 + \dots \\ &\leq -1920 + 81920q + 8060928q^3 + 166625280q^5 + \dots \end{aligned}$$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Proof of Lemma (1).** Again the idea is to leverage monotonicity.

Define

$$\begin{aligned} H(z) &= \frac{L(z+1) - L(z)}{2} = \dots = 4q^{-2} (\theta_2^8(\theta_3^{12} - \theta_4^{12}) + \theta_2^{12}(\theta_3^8 + \theta_4^8)) \\ &= 81920q + 8060928q^3 + 166625280q^5 + 1599733760q^7 + \dots \end{aligned}$$

Then for  $t \geq 1$ ,

$$\begin{aligned} L(it) &= -1920 + 81920q - 1077120q^2 + 8060928q^3 \\ &\quad - 41725440q^4 + 166625280q^5 - 553054080q^6 + \dots \end{aligned}$$

↙ (assuming alternating coefficients — need to justify)

$$\leq -1920 + 81920q + 8060928q^3 + 166625280q^5 + \dots$$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Proof of Lemma (1).** Again the idea is to leverage monotonicity.

Define

$$\begin{aligned} H(z) &= \frac{L(z+1) - L(z)}{2} = \dots = 4q^{-2} (\theta_2^8(\theta_3^{12} - \theta_4^{12}) + \theta_2^{12}(\theta_3^8 + \theta_4^8)) \\ &= 81920q + 8060928q^3 + 166625280q^5 + 1599733760q^7 + \dots \end{aligned}$$

Then for  $t \geq 1$ ,

$$\begin{aligned} L(it) &= -1920 + 81920q - 1077120q^2 + 8060928q^3 \\ &\quad - 41725440q^4 + 166625280q^5 - 553054080q^6 + \dots \\ &\quad \downarrow \text{ (assuming alternating coefficients — need to justify)} \\ &\leq -1920 + 81920q + 8060928q^3 + 166625280q^5 + \dots \\ &= -1920 + \frac{L(it+1) - L(it)}{2} \end{aligned}$$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Proof of Lemma (1).** Again the idea is to leverage monotonicity.

Define

$$\begin{aligned} H(z) &= \frac{L(z+1) - L(z)}{2} = \dots = 4q^{-2} (\theta_2^8(\theta_3^{12} - \theta_4^{12}) + \theta_2^{12}(\theta_3^8 + \theta_4^8)) \\ &= 81920q + 8060928q^3 + 166625280q^5 + 1599733760q^7 + \dots \end{aligned}$$

Then for  $t \geq 1$ ,

$$\begin{aligned} L(it) &= -1920 + 81920q - 1077120q^2 + 8060928q^3 \\ &\quad - 41725440q^4 + 166625280q^5 - 553054080q^6 + \dots \\ &\quad \downarrow \text{(assuming alternating coefficients — need to justify)} \\ &\leq -1920 + 81920q + 8060928q^3 + 166625280q^5 + \dots \\ &= -1920 + \frac{L(it+1) - L(it)}{2} = -1920 + H(it) \end{aligned}$$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Proof of Lemma (1).** Again the idea is to leverage monotonicity.

Define

$$\begin{aligned} H(z) &= \frac{L(z+1) - L(z)}{2} = \dots = 4q^{-2} (\theta_2^8(\theta_3^{12} - \theta_4^{12}) + \theta_2^{12}(\theta_3^8 + \theta_4^8)) \\ &= 81920q + 8060928q^3 + 166625280q^5 + 1599733760q^7 + \dots \end{aligned}$$

Then for  $t \geq 1$ ,

$$\begin{aligned} L(it) &= -1920 + 81920q - 1077120q^2 + 8060928q^3 \\ &\quad - 41725440q^4 + 166625280q^5 - 553054080q^6 + \dots \\ &\quad \downarrow \text{—— (assuming alternating coefficients — need to justify)} \\ &\leq -1920 + 81920q + 8060928q^3 + 166625280q^5 + \dots \\ &= -1920 + \frac{L(it+1) - L(it)}{2} = -1920 + H(it) \\ &\leq -1920 + H(i) \end{aligned}$$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Proof of Lemma (1).** Again the idea is to leverage monotonicity.

Define

$$\begin{aligned} H(z) &= \frac{L(z+1) - L(z)}{2} = \dots = 4q^{-2} (\theta_2^8(\theta_3^{12} - \theta_4^{12}) + \theta_2^{12}(\theta_3^8 + \theta_4^8)) \\ &= 81920q + 8060928q^3 + 166625280q^5 + 1599733760q^7 + \dots \end{aligned}$$

Then for  $t \geq 1$ ,

$$\begin{aligned} L(it) &= -1920 + 81920q - 1077120q^2 + 8060928q^3 \\ &\quad - 41725440q^4 + 166625280q^5 - 553054080q^6 + \dots \end{aligned}$$

$\downarrow$  (assuming alternating coefficients — need to justify)

$$\leq -1920 + 81920q + 8060928q^3 + 166625280q^5 + \dots$$

$$= -1920 + \frac{L(it+1) - L(it)}{2} = -1920 + H(it)$$

$$\leq -1920 + H(i) = \dots$$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Proof of Lemma (1).** Again the idea is to leverage monotonicity.

Define

$$\begin{aligned} H(z) &= \frac{L(z+1) - L(z)}{2} = \dots = 4q^{-2} (\theta_2^8(\theta_3^{12} - \theta_4^{12}) + \theta_2^{12}(\theta_3^8 + \theta_4^8)) \\ &= 81920q + 8060928q^3 + 166625280q^5 + 1599733760q^7 + \dots \end{aligned}$$

Then for  $t \geq 1$ ,

$$\begin{aligned} L(it) &= -1920 + 81920q - 1077120q^2 + 8060928q^3 \\ &\quad - 41725440q^4 + 166625280q^5 - 553054080q^6 + \dots \\ &\quad \downarrow \text{ (assuming alternating coefficients — need to justify)} \\ &\leq -1920 + 81920q + 8060928q^3 + 166625280q^5 + \dots \\ &= -1920 + \frac{L(it+1) - L(it)}{2} = -1920 + H(it) \\ &\leq -1920 + H(i) = \dots = -1920 + 3e^{2\pi} \frac{\Gamma(1/4)^{20}}{2048\pi^{15}} \end{aligned}$$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Proof of Lemma (1).** Again the idea is to leverage monotonicity.

Define

$$\begin{aligned} H(z) &= \frac{L(z+1) - L(z)}{2} = \dots = 4q^{-2} (\theta_2^8(\theta_3^{12} - \theta_4^{12}) + \theta_2^{12}(\theta_3^8 + \theta_4^8)) \\ &= 81920q + 8060928q^3 + 166625280q^5 + 1599733760q^7 + \dots \end{aligned}$$

Then for  $t \geq 1$ ,

$$\begin{aligned} L(it) &= -1920 + 81920q - 1077120q^2 + 8060928q^3 \\ &\quad - 41725440q^4 + 166625280q^5 - 553054080q^6 + \dots \\ &\quad \downarrow \text{ (assuming alternating coefficients — need to justify)} \\ &\leq -1920 + 81920q + 8060928q^3 + 166625280q^5 + \dots \\ &= -1920 + \frac{L(it+1) - L(it)}{2} = -1920 + H(it) \\ &\leq -1920 + H(i) = \dots = -1920 + 3e^{2\pi} \frac{\Gamma(1/4)^{20}}{2048\pi^{15}} \approx 2296.16 \end{aligned}$$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Proof of Lemma (1).** Again the idea is to leverage monotonicity.

Define

$$\begin{aligned} H(z) &= \frac{L(z+1) - L(z)}{2} = \dots = 4q^{-2} (\theta_2^8(\theta_3^{12} - \theta_4^{12}) + \theta_2^{12}(\theta_3^8 + \theta_4^8)) \\ &= 81920q + 8060928q^3 + 166625280q^5 + 1599733760q^7 + \dots \end{aligned}$$

Then for  $t \geq 1$ ,

$$\begin{aligned} L(it) &= -1920 + 81920q - 1077120q^2 + 8060928q^3 \\ &\quad - 41725440q^4 + 166625280q^5 - 553054080q^6 + \dots \\ &\quad \downarrow \text{ (assuming alternating coefficients — need to justify)} \\ &\leq -1920 + 81920q + 8060928q^3 + 166625280q^5 + \dots \\ &= -1920 + \frac{L(it+1) - L(it)}{2} = -1920 + H(it) \\ &\leq -1920 + H(i) = \dots = -1920 + 3e^{2\pi} \frac{\Gamma(1/4)^{20}}{2048\pi^{15}} \approx 2296.16 \\ &\leq 2297, \quad \text{which is what we wanted.} \end{aligned}$$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Justification of the assumption about alternating coefficients:** define

$$W(z) = \theta_3^{12}\theta_2^8 + \theta_3^8\theta_2^{12} + \theta_3^{12}\theta_4^8 + \theta_3^8\theta_4^{12}.$$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Justification of the assumption about alternating coefficients:** define

$$W(z) = \theta_3^{12}\theta_2^8 + \theta_3^8\theta_2^{12} + \theta_3^{12}\theta_4^8 + \theta_3^8\theta_4^{12}.$$

By simple algebra,  $-L(z+1) = 8q^{-2}(W(z) - 2)$ , so the claim is equivalent to the statement that the Fourier expansion of  $W(z)$  has nonnegative coefficients.\*

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Justification of the assumption about alternating coefficients:** define

$$W(z) = \theta_3^{12}\theta_2^8 + \theta_3^8\theta_2^{12} + \theta_3^{12}\theta_4^8 + \theta_3^8\theta_4^{12}.$$

By simple algebra,  $-L(z+1) = 8q^{-2}(W(z) - 2)$ , so the claim is equivalent to the statement that the Fourier expansion of  $W(z)$  has nonnegative coefficients.\* This follows from the identity\*\*

$$W = \frac{1}{16}(6X^5 + 15X^4Y + 10X^3Y^2 + Y^5),$$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Justification of the assumption about alternating coefficients:** define

$$W(z) = \theta_3^{12}\theta_2^8 + \theta_3^8\theta_2^{12} + \theta_3^{12}\theta_4^8 + \theta_3^8\theta_4^{12}.$$

By simple algebra,  $-L(z+1) = 8q^{-2}(W(z) - 2)$ , so the claim is equivalent to the statement that the Fourier expansion of  $W(z)$  has nonnegative coefficients.\* This follows from the identity\*\*

$$W = \frac{1}{16}(6X^5 + 15X^4Y + 10X^3Y^2 + Y^5),$$

where  $X := \theta_2^4$  and  $Y := 2\theta_3^4 - \theta_2^4 = \theta_3(z)^4 + \theta_3(z+1)^4$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Justification of the assumption about alternating coefficients:** define

$$W(z) = \theta_3^{12}\theta_2^8 + \theta_3^8\theta_2^{12} + \theta_3^{12}\theta_4^8 + \theta_3^8\theta_4^{12}.$$

By simple algebra,  $-L(z+1) = 8q^{-2}(W(z) - 2)$ , so the claim is equivalent to the statement that the Fourier expansion of  $W(z)$  has nonnegative coefficients.\* This follows from the identity\*\*

$$W = \frac{1}{16}(6X^5 + 15X^4Y + 10X^3Y^2 + Y^5),$$

where  $X := \theta_2^4$  and  $Y := 2\theta_3^4 - \theta_2^4 = \theta_3(z)^4 + \theta_3(z+1)^4$  are both easily seen to have nonnegative Fourier coefficients.

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Justification of the assumption about alternating coefficients:** define

$$W(z) = \theta_3^{12}\theta_2^8 + \theta_3^8\theta_2^{12} + \theta_3^{12}\theta_4^8 + \theta_3^8\theta_4^{12}.$$

By simple algebra,  $-L(z+1) = 8q^{-2}(W(z) - 2)$ , so the claim is equivalent to the statement that the Fourier expansion of  $W(z)$  has nonnegative coefficients.\* This follows from the identity\*\*

$$W = \frac{1}{16}(6X^5 + 15X^4Y + 10X^3Y^2 + Y^5),$$

where  $X := \theta_2^4$  and  $Y := 2\theta_3^4 - \theta_2^4 = \theta_3(z)^4 + \theta_3(z+1)^4$  are both easily seen to have nonnegative Fourier coefficients.

\* This nonnegativity result was first proved by Slipper (2018), with a more complicated proof. See also <https://mathoverflow.net/q/441749/78525>.

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Justification of the assumption about alternating coefficients:** define

$$W(z) = \theta_3^{12}\theta_2^8 + \theta_3^8\theta_2^{12} + \theta_3^{12}\theta_4^8 + \theta_3^8\theta_4^{12}.$$

By simple algebra,  $-L(z+1) = 8q^{-2}(W(z) - 2)$ , so the claim is equivalent to the statement that the Fourier expansion of  $W(z)$  has nonnegative coefficients.\* This follows from the identity\*\*

$$W = \frac{1}{16}(6X^5 + 15X^4Y + 10X^3Y^2 + Y^5),$$

where  $X := \theta_2^4$  and  $Y := 2\theta_3^4 - \theta_2^4 = \theta_3(z)^4 + \theta_3(z+1)^4$  are both easily seen to have nonnegative Fourier coefficients.

\* This nonnegativity result was first proved by Slipper (2018), with a more complicated proof. See also <https://mathoverflow.net/q/441749/78525>.

\*\* I discovered this identity using computer algebra + a linear program solver.

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Proof of Lemma (2).**

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Proof of Lemma (2).** The asymptotic expansion of  $K(it)$  is

$$\begin{aligned} K(it) = & (3840\pi t - 7680) + (-230400\pi^2 t^2 + 990720\pi t - 990720)q^2 \\ & + (-8294400\pi^2 t^2 + 25205760\pi t - 16803840)q^4 + \dots \end{aligned}$$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Proof of Lemma (2).** The asymptotic expansion of  $K(it)$  is

$$\begin{aligned} K(it) = & (3840\pi t - 7680) + (-230400\pi^2 t^2 + 990720\pi t - 990720)q^2 \\ & + (-8294400\pi^2 t^2 + 25205760\pi t - 16803840)q^4 + \dots \end{aligned}$$

With this motivation in mind, define

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Proof of Lemma (2).** The asymptotic expansion of  $K(it)$  is

$$\begin{aligned} K(it) = & (3840\pi t - 7680) + (-230400\pi^2 t^2 + 990720\pi t - 990720)q^2 \\ & + (-8294400\pi^2 t^2 + 25205760\pi t - 16803840)q^4 + \dots \end{aligned}$$

With this motivation in mind, define

$$K_1(t) = 3840\pi t + (-230400\pi^2 t^2 + 990720\pi t - 990720)q^2,$$

$$\begin{aligned} K_2(t) = & q^{-2} E'_4(it)^2 t^2 - 16q^{-2}(E_4(it)^2 - 1) \\ & + (230400\pi^2 t^2 + 990720)q^2, \end{aligned}$$

$$K_3(t) = -8iq^{-2} E'_4(it) E_4(it) t - (3840\pi t + 990720\pi t q^2),$$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Proof of Lemma (2).** The asymptotic expansion of  $K(it)$  is

$$\begin{aligned} K(it) = & (3840\pi t - 7680) + (-230400\pi^2 t^2 + 990720\pi t - 990720)q^2 \\ & + (-8294400\pi^2 t^2 + 25205760\pi t - 16803840)q^4 + \dots \end{aligned}$$

With this motivation in mind, define

$$K_1(t) = 3840\pi t + (-230400\pi^2 t^2 + 990720\pi t - 990720)q^2,$$

$$\begin{aligned} K_2(t) = & q^{-2} E'_4(it)^2 t^2 - 16q^{-2}(E_4(it)^2 - 1) \\ & + (230400\pi^2 t^2 + 990720)q^2, \end{aligned}$$

$$K_3(t) = -8iq^{-2} E'_4(it) E_4(it) t - (3840\pi t + 990720\pi t q^2),$$

so that we have

$$K(it) = K_1(t) + K_2(t) + K_3(t).$$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Proof of Lemma (2).** The asymptotic expansion of  $K(it)$  is

$$\begin{aligned} K(it) = & (3840\pi t - 7680) + (-230400\pi^2 t^2 + 990720\pi t - 990720)q^2 \\ & + (-8294400\pi^2 t^2 + 25205760\pi t - 16803840)q^4 + \dots \end{aligned}$$

With this motivation in mind, define

$$K_1(t) = 3840\pi t + (-230400\pi^2 t^2 + 990720\pi t - 990720)q^2,$$

$$\begin{aligned} K_2(t) = & q^{-2} E'_4(it)^2 t^2 - 16q^{-2}(E_4(it)^2 - 1) \\ & + (230400\pi^2 t^2 + 990720)q^2, \end{aligned}$$

$$K_3(t) = -8iq^{-2} E'_4(it) E_4(it) t - (3840\pi t + 990720\pi t q^2),$$

so that we have

$$K(it) = K_1(t) + K_2(t) + K_3(t).$$

The following claims are easy to check:

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Proof of Lemma (2).** The asymptotic expansion of  $K(it)$  is

$$\begin{aligned} K(it) = & (3840\pi t - 7680) + (-230400\pi^2 t^2 + 990720\pi t - 990720)q^2 \\ & + (-8294400\pi^2 t^2 + 25205760\pi t - 16803840)q^4 + \dots \end{aligned}$$

With this motivation in mind, define

$$K_1(t) = 3840\pi t + (-230400\pi^2 t^2 + 990720\pi t - 990720)q^2,$$

$$\begin{aligned} K_2(t) = & q^{-2} E'_4(it)^2 t^2 - 16q^{-2}(E_4(it)^2 - 1) \\ & + (230400\pi^2 t^2 + 990720)q^2, \end{aligned}$$

$$K_3(t) = -8iq^{-2} E'_4(it) E_4(it) t - (3840\pi t + 990720\pi t q^2),$$

so that we have

$$K(it) = K_1(t) + K_2(t) + K_3(t).$$

The following claims are easy to check:

- ① The function  $K_1(t)$  is monotone increasing on  $[1, \infty)$ ,

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Proof of Lemma (2).** The asymptotic expansion of  $K(it)$  is

$$\begin{aligned} K(it) = & (3840\pi t - 7680) + (-230400\pi^2 t^2 + 990720\pi t - 990720)q^2 \\ & + (-8294400\pi^2 t^2 + 25205760\pi t - 16803840)q^4 + \dots \end{aligned}$$

With this motivation in mind, define

$$K_1(t) = 3840\pi t + (-230400\pi^2 t^2 + 990720\pi t - 990720)q^2,$$

$$\begin{aligned} K_2(t) = & q^{-2} E'_4(it)^2 t^2 - 16q^{-2}(E_4(it)^2 - 1) \\ & + (230400\pi^2 t^2 + 990720)q^2, \end{aligned}$$

$$K_3(t) = -8iq^{-2} E'_4(it) E_4(it) t - (3840\pi t + 990720\pi t q^2),$$

so that we have

$$K(it) = K_1(t) + K_2(t) + K_3(t).$$

The following claims are easy to check:

- ① The function  $K_1(t)$  is monotone increasing on  $[1, \infty)$ ,
- ② The function  $K_2(t)$  is monotone increasing on  $[1, \infty)$ .

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

**Proof of Lemma (2).** The asymptotic expansion of  $K(it)$  is

$$\begin{aligned} K(it) = & (3840\pi t - 7680) + (-230400\pi^2 t^2 + 990720\pi t - 990720)q^2 \\ & + (-8294400\pi^2 t^2 + 25205760\pi t - 16803840)q^4 + \dots \end{aligned}$$

With this motivation in mind, define

$$K_1(t) = 3840\pi t + (-230400\pi^2 t^2 + 990720\pi t - 990720)q^2,$$

$$\begin{aligned} K_2(t) = & q^{-2} E'_4(it)^2 t^2 - 16q^{-2}(E_4(it)^2 - 1) \\ & + (230400\pi^2 t^2 + 990720)q^2, \end{aligned}$$

$$K_3(t) = -8iq^{-2} E'_4(it) E_4(it) t - (3840\pi t + 990720\pi t q^2),$$

so that we have

$$K(it) = K_1(t) + K_2(t) + K_3(t).$$

The following claims are easy to check:

- ① The function  $K_1(t)$  is monotone increasing on  $[1, \infty)$ ,
- ② The function  $K_2(t)$  is monotone increasing on  $[1, \infty)$ .
- ③  $K_3(t) \geq 0$  for all  $t > 0$ .

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

Therefore, assuming  $t \geq 1$ ,

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

Therefore, assuming  $t \geq 1$ ,

$$K(it)$$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

Therefore, assuming  $t \geq 1$ ,

$$K(it) = K_1(t) + K_2(t) + K_3(t)$$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

Therefore, assuming  $t \geq 1$ ,

$$K(it) = K_1(t) + K_2(t) + K_3(t) \geq K_1(t) + K_2(t)$$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

Therefore, assuming  $t \geq 1$ ,

$$K(it) = K_1(t) + K_2(t) + K_3(t) \geq K_1(t) + K_2(t) \geq K_1(1) + K_2(1)$$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

Therefore, assuming  $t \geq 1$ ,

$$\begin{aligned} K(it) &= K_1(t) + K_2(t) + K_3(t) \geq K_1(t) + K_2(t) \geq K_1(1) + K_2(1) \\ &= -e^{2\pi} (-E'_4(i)^2 + 16E_4(i)^2 - 16) + 3840\pi + 990720\pi e^{-2\pi} \end{aligned}$$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

Therefore, assuming  $t \geq 1$ ,

$$\begin{aligned} K(it) &= K_1(t) + K_2(t) + K_3(t) \geq K_1(t) + K_2(t) \geq K_1(1) + K_2(1) \\ &= -e^{2\pi} (-E'_4(i)^2 + 16E_4(i)^2 - 16) + 3840\pi + 990720\pi e^{-2\pi} \\ &= -e^{2\pi} \left( \frac{45\Gamma(1/4)^{16}}{1024\pi^{12}} - 16 \right) + 3840\pi + 990720\pi e^{-2\pi} \end{aligned}$$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

Therefore, assuming  $t \geq 1$ ,

$$\begin{aligned} K(it) &= K_1(t) + K_2(t) + K_3(t) \geq K_1(t) + K_2(t) \geq K_1(1) + K_2(1) \\ &= -e^{2\pi} (-E'_4(i)^2 + 16E_4(i)^2 - 16) + 3840\pi + 990720\pi e^{-2\pi} \\ &= -e^{2\pi} \left( \frac{45\Gamma(1/4)^{16}}{1024\pi^{12}} - 16 \right) + 3840\pi + 990720\pi e^{-2\pi} \\ &\approx 3747.1, \end{aligned}$$

## A new proof of (V1)–(V2), part II: proof of (V2) (cont'd)

Therefore, assuming  $t \geq 1$ ,

$$\begin{aligned} K(it) &= K_1(t) + K_2(t) + K_3(t) \geq K_1(t) + K_2(t) \geq K_1(1) + K_2(1) \\ &= -e^{2\pi} (-E'_4(i)^2 + 16E_4(i)^2 - 16) + 3840\pi + 990720\pi e^{-2\pi} \\ &= -e^{2\pi} \left( \frac{45\Gamma(1/4)^{16}}{1024\pi^{12}} - 16 \right) + 3840\pi + 990720\pi e^{-2\pi} \\ &\approx 3747.1, \end{aligned}$$

as claimed.

## Final thoughts and open problems

## Final thoughts and open problems

- Viazovska's magic function  $\varphi$  is a mathematical object of remarkable beauty and mystery.

## Final thoughts and open problems

- Viazovska's magic function  $\varphi$  is a mathematical object of remarkable beauty and mystery.
- We now understand it a bit better than before. But there seems more to understand still.
- Open problems:

## Final thoughts and open problems

- Viazovska's magic function  $\varphi$  is a mathematical object of remarkable beauty and mystery.
- We now understand it a bit better than before. But there seems more to understand still.
- Open problems:
  - Find a human proof of the analogous inequalities for the case of dimension 24.

## Final thoughts and open problems

- Viazovska's magic function  $\varphi$  is a mathematical object of remarkable beauty and mystery.
- We now understand it a bit better than before. But there seems more to understand still.
- Open problems:
  - Find a human proof of the analogous inequalities for the case of dimension 24.
  - Prove the analogous inequalities for dimensions that are multiples of 4.

## Final thoughts and open problems

- Viazovska's magic function  $\varphi$  is a mathematical object of remarkable beauty and mystery.
- We now understand it a bit better than before. But there seems more to understand still.
- Open problems:
  - Find a human proof of the analogous inequalities for the case of dimension 24.
  - Prove the analogous inequalities for dimensions that are multiples of 4. (Might require computer assistance?)

## Final thoughts and open problems

- Viazovska's magic function  $\varphi$  is a mathematical object of remarkable beauty and mystery.
- We now understand it a bit better than before. But there seems more to understand still.
- Open problems:
  - Find a human proof of the analogous inequalities for the case of dimension 24.
  - Prove the analogous inequalities for dimensions that are multiples of 4. (Might require computer assistance?)
- A philosophical lesson:

## Final thoughts and open problems

- Viazovska's magic function  $\varphi$  is a mathematical object of remarkable beauty and mystery.
- We now understand it a bit better than before. But there seems more to understand still.
- Open problems:
  - Find a human proof of the analogous inequalities for the case of dimension 24.
  - Prove the analogous inequalities for dimensions that are multiples of 4. (Might require computer assistance?)
- A philosophical lesson:

*Human analysis and insights can still triumph over computer-assisted mathematics*

## Final thoughts and open problems

- Viazovska's magic function  $\varphi$  is a mathematical object of remarkable beauty and mystery.
- We now understand it a bit better than before. But there seems more to understand still.
- Open problems:
  - Find a human proof of the analogous inequalities for the case of dimension 24.
  - Prove the analogous inequalities for dimensions that are multiples of 4. (Might require computer assistance?)
- A philosophical lesson:

*Human analysis and insights can still triumph over computer-assisted mathematics*

*(...in 2023).*

# Final thoughts and open problems

- Viazovska's magic function  $\varphi$  is a mathematical object of remarkable beauty and mystery.
- We now understand it a bit better than before. But there seems more to understand still.
- Open problems:
  - Find a human proof of the analogous inequalities for the case of dimension 24.
  - Prove the analogous inequalities for dimensions that are multiples of 4. (Might require computer assistance?)
- A philosophical lesson:

*Human analysis and insights can still triumph over computer-assisted mathematics  
(...in 2023).*

That's all — thank you!