


# Physics 30 1997 January

## Numerical Response

1. A horizontal force of 207 N acts on a 7.80 kg bowling ball for 0.520 s. The change in the ball's speed is \_\_\_\_\_ m/s.





(Record your **three-digit answer** in the numerical-response section on the answer sheet.)

*Use the following information to answer the next two questions.*




1. The speed of the block at point Z is
- A. 4.4 m/s
  - B. 4.9 m/s
  - C. 8.9 m/s
  - D. 14 m/s
2. Assuming that the potential energy of the block is zero at ground level, then the total mechanical energy of the block at point Y is
- A. 4.9 J
  - B. 9.8 J
  - C. 15 J
  - D. 20 J

3. A single stationary railway car is bumped by a five-car train moving at 9.3 km/h. The six cars move off together after the collision. Assuming that the masses of all the railway cars are the same, then the speed of the new six-car train immediately after impact is
- A. 7.8 km/h  
B. 8.5 km/h  
C. 9.3 km/h  
D. 11 km/h
4. When the electron and the proton in a hydrogen atom are  $5.3 \times 10^{-11}$  m apart, the magnitude of the electrostatic force on the electron is
- A.  $4.3 \times 10^{-20}$  N  
B.  $4.3 \times 10^{-18}$  N  
C.  $8.2 \times 10^{-12}$  N  
D.  $8.2 \times 10^{-8}$  N
5. Which graph **best** represents the magnitude of the electrostatic force,  $F$ , as a function of the distance,  $r$ , between two point charges?




6. The field resulting from a positive point charge and a negative point charge is best represented by


A.




B.



C.



D.




*Use the following information to answer the next question.*

Two parallel metal plates are 1.5 cm apart and are maintained at a potential difference of  $2.5 \times 10^2$  V.

7. The magnitude of the electrical force on an alpha particle when the alpha particle is between the plates is
- A.  $5.3 \times 10^{-15}$  N
  - B.  $2.7 \times 10^{-15}$  N
  - C.  $5.3 \times 10^{-17}$  N
  - D.  $2.7 \times 10^{-17}$  N

Use the following information to answer the next three questions.

**Electrostatics**



Two identical conducting balls,  $b_1$  and  $b_2$ , each of mass 25.0 g, are hanging on 50.0 cm-long insulating threads. They become equally charged and come to rest with angles of deviation of  $45.0^\circ$  from the vertical.

8. The electrostatic force between the charged balls can **best** be described as
- A. an attraction due to dissimilar charges
  - B. a repulsion due to dissimilar charges
  - C. an attraction due to similar charges
  - D. a repulsion due to similar charges
9. What is the tension in the thread that is supporting one of the balls?
- A. 0.173 N
  - B. 0.245 N
  - C. 0.347 N
  - D. 9.81 N
10. If the charge on  $b_1$  is tripled and the charge on  $b_2$  is reduced to one-third of its original amount, the angles of deviation from centre would
- A. increase for  $b_1$  and decrease for  $b_2$
  - B. remain the same for both  $b_1$  and  $b_2$
  - C. increase for both  $b_1$  and  $b_2$
  - D. decrease for both  $b_1$  and  $b_2$

*Use the following information to answer the next three questions.*

The headlight of a car operates with an input power of 75.0 W and draws a current of 6.25 A.

**Numerical Response**

2. The voltage supplied to the headlight is \_\_\_\_\_ V.

(Record your **three-digit answer** in the numerical-response section on the answer sheet.)

**Numerical Response**

3. The number of electrons passing through the headlight every minute, expressed in scientific notation, is **a.b** x 10<sup>cd</sup> electrons. The values of **a**, **b**, **c**, and **d** are \_\_\_\_\_, \_\_\_\_\_, \_\_\_\_\_, and \_\_\_\_\_.


(Record all **four digits** in the numerical-response section on the answer sheet.)

**Numerical Response**

4. The energy used by the headlight during 1.00 h of operation, expressed in scientific notation, is **b** x 10<sup>w</sup> J. The value of **b** is \_\_\_\_\_.

(Record your **three-digit answer** in the numerical-response section on the answer sheet.)

Use the following information to answer the next three questions.



**Numerical Response**

5. The equivalent resistance for the circuit is \_\_\_\_\_  $\Omega$ .

(Record your **three-digit answer** in the numerical-response section on the answer sheet.)

Use your recorded answer for **Numerical Response 5** to solve **Numerical Response 6**.

**Numerical Response**

6. The total current supplied by the 46.0 V power supply to the circuit is \_\_\_\_\_ A.

(Record your **three-digit answer** in the numerical-response section on the answer sheet.)

11. The voltage drop across the  $20.0 \Omega$  resistor in the circuit is

- A. 30.6 V
- B. 24.5 V
- C. 16.0 V
- D. 1.53 V

Use the following information to answer the next three questions.

Lightning storms are one of the most spectacular phenomena in nature and play a key role in maintaining the electrical balance of Earth. One lightning strike occurred over a potential difference of 200 MV and transferred 12 C of charge to the ground in 0.010 s.

12. The energy released by the lightning strike in this time interval was
- A.  $3.2 \times 10^{-11} \text{ J}$
  - B.  $2.4 \times 10^3 \text{ J}$
  - C.  $2.4 \times 10^9 \text{ J}$
  - D.  $2.4 \times 10^{11} \text{ J}$

Use your recorded answer for **Multiple Choice 12** to solve **Numerical Response 7**.

**Numerical Response**

7. The power released in this lightning strike, expressed in scientific notation, is **a.b**  $\times 10^{\text{cd}}$  W. The values of **a**, **b**, **c**, and **d** are \_\_\_\_\_, \_\_\_\_\_, \_\_\_\_\_, and \_\_\_\_\_.

(Record all **four digits** in the numerical-response section on the answer sheet.)

13. People inside a car are protected from the electric fields associated with lightning. Many parts of stereo components are contained in metal boxes. These two examples demonstrate that the electric field inside a closed metal container is
- A. zero
  - B. opposite to the field outside
  - C. equal to the field outside
  - D. half the field outside
- \_\_\_\_\_
14. Which of the following unit combinations is **not** equivalent to an ampere?
- A. Watt/volt
  - B. Volt/ohm
  - C. Watt/ohm
  - D. Coulomb/second

15. Energy is used to move a charge of 3.00 C through a circuit with a resistance of  $1.00 \times 10^2 \Omega$  in 1.00 s. If the same amount of energy is used to throw a 1.00 kg ball vertically upward, the maximum height of the ball would be
- A.  $1.09 \times 10^{-2} \text{ m}$
  - B.  $3.27 \times 10^{-2} \text{ m}$
  - C.  $3.06 \times 10^1 \text{ m}$
  - D.  $9.17 \times 10^1 \text{ m}$
16. To boil three cups of water each day for one year using an old model of microwave oven costs \$1.10. A new 750 W microwave oven boils one cup of water in 100 s. If the cost of energy is \$0.0100/MJ, how much money will a consumer **save** by using the new microwave oven to boil three cups of water each day for one year? Assume in each case that the water is at the same starting temperature.
- A. \$0.28
  - B. \$0.82
  - C. \$1.38
  - D. \$2.80

*Use the following information to answer the next question,*


Unit Combinations	
I.	J/C
II.	N/(A·m)
III.	T
IV.	(N·s)/(C·m)

17. Which unit combinations could be used correctly for a magnetic field?
- A. I, III, and IV
  - B. II and III only
  - C. II and IV only
  - D. II, III, and IV



18. An electric field of strength  $1.5 \times 10^4 \text{ N/C}$  is perpendicular to a magnetic field of strength  $3.0 \times 10^{-3} \text{ T}$ . An electron moves perpendicular to both fields and is undeflected as it passes through the fields. The speed of the electron is
- A.  $2.0 \times 10^{-7} \text{ m/s}$
  - B.  $2.0 \times 10^{-1} \text{ m/s}$
  - C.  $5.0 \times 10^6 \text{ m/s}$
  - D.  $5.0 \times 10^7 \text{ m/s}$
19. An ideal transformer steps down 25 000 V to 120 V for use in a house. Several appliances draw a total of 2 000 W from the 120 V side of the transformer. What is the current in the 25 000 V line?
- A.  $6.00 \times 10^{-2} \text{ A}$
  - B.  $8.00 \times 10^{-2} \text{ A}$
  - C. 12.5 A
  - D. 16.7 A
20. When two parallel conducting wires repel each other, the currents in the wires are
- A. in opposite directions
  - B. in the same direction
  - C. oscillating in phase
  - D. oppositely charged


Use the following information to answer the next question.




21. When the string is plucked, a small current is produced in the coil of wire because
- A. a potential difference is produced in the string
  - B. there is a current in the string that can be amplified
  - C. there is a charge buildup on the string
  - D. the string behaves as a magnet moving toward and away from the coil
22. French high-speed trains operate using power lines that have an effective voltage of 25.0 kV and a frequency of 50.0 Hz. The maximum or peak voltage of the power lines is
- A. 12.5 kV
  - B. 17.7 kV
  - C. 35.4 kV
  - D. 50.0 kV

23. Which of the following diagrams best illustrates the magnetic field near a wire that carries an electron current out of the plane of the paper?


A.




B.



C.



D.



### Numerical Response

8. An electromagnetic wave has a frequency of  $2.00 \times 10^8$  Hz. The speed of the wave in a vacuum, expressed in scientific notation, is  $b \times 10^x$  m/s. The value of  $b$  is \_\_\_\_\_.

(Record your **three-digit answer** in the numerical-response section on the answer sheet.)

24. In a vacuum, the period of oscillation of a microwave with a wavelength of 2.5 cm is

- A.  $8.3 \times 10^{-11}$  s
- B.  $8.3 \times 10^{-9}$  s
- C.  $1.2 \times 10^8$  s
- D.  $1.2 \times 10^{10}$  s

### Numerical Response

9. The minimum potential difference through which an electron must be accelerated to produce an X-ray of energy  $1.62 \times 10^4$  eV, expressed in scientific notation, is  $b \times 10^w$  V. The value of  $b$  is \_\_\_\_\_.

(Record your **three-digit answer** in the numerical-response section on the answer sheet.)

### Numerical Response

10. A term used in aviation is **radar mile**, which is the time it takes a radar pulse to travel to a target 1.00 mile away and return (1.00 mile = 1.625 km). The radar mile, expressed in scientific notation, is  $b \times 10^{-w}$  s. The value of  $b$  is \_\_\_\_\_.

(Record your **three-digit answer** in the numerical-response section on the answer sheet.)


*Use the following information to answer the next question.*

A technological application of quantum theory is the development of “electric eyes,” which can be used in automatic door openers or burglar alarms. A light beam shines across a door opening and causes the production of a current in a circuit. When the beam is broken, the current stops and a mechanism is triggered to open a door or sound an alarm.

25. The operation of an electric eye is an application of
- A. the Compton effect
  - B. the wave nature of matter
  - C. the photoelectric effect
  - D. Maxwell's electromagnetic wave theory

26. To determine the speed of charged particles in a cathode-ray tube, Thomson balanced the forces produced by
- A. an electromagnetic field and a gravitational field
  - B. a magnetic field and a gravitational field
  - C. an electric field and a gravitational field
  - D. an electric field and a magnetic field
27. A student performs a photoelectric experiment in which a photoelectric current is observed for all colours of visible light. The student wants to investigate what effect varying the intensity and colour of the incident light has on the photoelectric current and kinetic energy of the photoelectrons. If the brightness of the light is decreased and the colour is changed from yellow to blue, the photoelectric
- A. current and photoelectron energy both decrease
  - B. current and photoelectron energy both increase
  - C. current decreases and the photoelectron energy increases
  - D. current increases and the photoelectron energy decreases
28. When a blue laser beam is incident upon the surface of the metal of a photoelectric cell, there is no photoemission. A second beam of radiation causes photoelectrons to be emitted. The second beam may consist of
- A. ultraviolet radiation
  - B. infrared radiation
  - C. red laser radiation
  - D. microwave radiation
29. A photon of energy 1.13 eV is emitted by a hydrogen atom when the electron “jumps” from
- A.  $n=6$  to  $n=3$
  - B.  $n=3$  to  $n=6$
  - C.  $n=5$  to  $n=2$
  - D.  $n=2$  to  $n=5$

Use the following information to answer the next two questions.




30. What is the frequency of light emitted from the laser when the electron in the chromium atom goes from state  $E_1$  to state  $E_0$ ?
- A.  $9.7 \times 10^{13}$  Hz
  - B.  $4.3 \times 10^{14}$  Hz
  - C.  $5.3 \times 10^{14}$  Hz
  - D.  $1.1 \times 10^{15}$  Hz

### Numerical Response

11. Flashes of light pump electrons in the Cr atoms from the ground state,  $E_0$ , to state  $E_2$ . The wavelength of these flashes of light, expressed in scientific notation, is  $b \times 10^{-w}$  m. The value of  $b$  is \_\_\_\_\_.

(Record your **two-digit answer** in the numerical-response section on the answer sheet.)

Use the following information to answer the next three questions.



31. The maximum kinetic energy of the electrons ejected from the photocathode is
- A.  $6.7 \times 10^{-20} \text{ J}$
  - B.  $3.4 \times 10^{-19} \text{ J}$
  - C.  $4.0 \times 10^{-19} \text{ J}$
  - D.  $7.4 \times 10^{-19} \text{ J}$
32. The electrons leaving the photocathode are attracted by the 0.20 kV electrode. The maximum speed they attain is
- A.  $8.6 \times 10^5 \text{ m/s}$
  - B.  $8.4 \times 10^6 \text{ m/s}$
  - C.  $7.0 \times 10^{13} \text{ m/s}$
  - D.  $2.1 \times 10^{15} \text{ m/s}$

Use the following additional information to answer the next question.

**Inner Workings of the Photomultiplier Tube**

An electron striking the 0.20 kV electrode will use its energy to eject multiple secondary electrons from the electrode. The secondary electrons accelerate toward the next electrode, and the process continues along successive increases in voltage. The energy required to release a single electron is 40.0 eV.

33. Assume all of the kinetic energy of an electron striking the 0.20 kV electrode is used to eject secondary electrons. The number of electrons released from the 0.20 kV electrode is
- A. 5
  - B. 10
  - C. 100
  - D. 200
- 

**Numerical Response**

12. In the ground state of a hydrogen atom, the radius of the electron orbit is  $5.3 \times 10^{-11}$  m. According to the Bohr model, the radius of the electron orbit corresponding to the third energy level, expressed in scientific notation, is **a.b**  $\times 10^{-cd}$  m. The values of **a**, **b**, **c**, and **d** are \_\_\_\_\_, \_\_\_\_\_, \_\_\_\_\_, and \_\_\_\_\_.

(Record all **four digits** in the numerical-response section on the answer sheet.)

34. The half-life of radium-226 is  $1.6 \times 10^3$  years. How long will it take for 20.0 mg of radium-226 to decay to 2.50 mg?
- A.  $1.3 \times 10^3$  years
  - B.  $1.6 \times 10^3$  years
  - C.  $3.2 \times 10^3$  years
  - D.  $4.8 \times 10^3$  years



*Use the following information to answer the next two questions.*

Recently H. J. Rose and G. A. Jones of Oxford University predicted that the decay of radium-223 would emit alpha particles ( $^4\text{He}$ ) as well as a few double alphas ( $^8\text{Be}$ ) and even triple alphas ( $^{12}\text{C}$ ).

The experiment was run for 600 days. The results of the experiment showed the detection of  $2.2 \times 10^{10}$  alpha particles, but the double and triple alphas were not detected. However, unexpectedly, nineteen carbon-14 nuclei were detected. This experiment has led scientists to continue to look for other examples of radioactive decay in which medium-sized nuclei such as carbon-14 are produced.

35. Which of the following initial products was observed in the greatest abundance?
- A. Radon-219
  - B. Polonium-215
  - C. Lead-211
  - D. Lead-209
36. Which of the following products was **not** predicted by Rose and Jones?
- A. Radon-219
  - B. Polonium-215
  - C. Lead-211
  - D. Lead-209
- 
37. In a nuclear reaction, the mass of the products was determined to be considerably less than the mass of the reactants. A correct explanation of this is that
- A. the reaction was a beta-decay
  - B. a large amount of energy was released in the reaction
  - C. the mass of the alpha and beta particles was not accounted for
  - D. a large amount of energy was required to cause the reaction to occur

## Written Response

Use the following information to answer written- response question 1.

**A Spring Ram Stapler**

The diagram illustrates the operation of a spring ram stapler in three stages:

- No compression:** The spring is at its natural length, labeled  $x = 0$ . A ram is positioned below the spring.
- Maximum compression:** The spring is compressed to a maximum value of  $x_1 = 3.50 \times 10^{-2} \text{ m}$ . The ram is positioned below the spring. A staple is shown below the ram. A dashed line indicates the position when the ram is released.
- Partial compression:** The spring is compressed to a partial value of  $x_2 = 1.00 \times 10^{-2} \text{ m}$ . The ram is positioned below the spring. A staple is shown below the ram. A dashed line indicates the position when the ram hits the staple.

Heavy-duty stapling guns use powerful springs in combination with a small metal rod (called a ram) to produce the impact necessary to move staples or nails into materials such as wood, wallboard, or even concrete.

A particular stapler gun has a ram with mass  $0.200 \text{ kg}$  and a spring with a spring constant of  $35\,000 \text{ N/m}$ . When the handle of the gun is squeezed, the spring is compressed to a maximum value of  $3.50 \times 10^{-2} \text{ m}$ . When the ram makes contact with the staple, the spring is still compressed  $1.00 \times 10^{-2} \text{ m}$ . Assume that  $3.00\%$  of the ram's kinetic energy is transferred to the  $2.00 \text{ g}$  staple when the ram hits the staple. The potential energy of a spring is  $\frac{1}{2} kx^2$ .

1. Describe and calculate the energy transformations involved in the operation of the spring ram stapler. Use conservation laws, physics concepts, and related equations to support your answer. Ignore the mass of the spring and the effects of gravitational potential energy on the system.


(8 marks)

Use the following information to answer written-response question 2.

2. A student used the apparatus shown below to measure the radius of curvature of the path of electrons as they pass through a magnetic field that is perpendicular to their path. This experimental design has the voltage as the manipulated variable, the speed calculated from the voltage, and the radius as the responding variable.

**Accelerating  
Potential**


Difference (V)	Speed ( $10^6$ m/s)	Radius ( $10^{-2}$ m)
20.0	2.65	7.2
40.0	3.75	9.1
60.0	4.59	11.0
80.0	5.30	12.8
100.0	5.93	14.1
120.0	6.49	16.3



x indicates magnetic field into the page

- a. Plot the graph of radius as a function of speed, and construct a best-fit line.

(3)



- b. Using the slope or other appropriate averaging technique, determine the strength of the magnetic field.

(3)

- c. Derive the equation that would allow the student to calculate the speed of the electrons from their accelerating potential.

(2)