

Your MMM Is Broken: Identification of Nonlinear and Time-varying Effects in Marketing Mix Models


Ryan Dew

*The Wharton School
University of Pennsylvania*

Joint work with Nicolas Padilla (LBS) and Anya Shchetkina (Wharton)


The Cookie-free Future

- For the past 10+ years, individual-level trackability has been central to measuring marketing efficacy




The Cookie-free Future

- For the past 10+ years, individual-level trackability has been central to measuring marketing efficacy
- Now? Gone!




The Cookie-free Future

- For the past 10+ years, individual-level trackability has been central to measuring marketing efficacy
- Now? Gone!




The Cookie-free Future

- For the past 10+ years, individual-level trackability has been central to measuring marketing efficacy
- Now? Gone!
- Result: huge resurgence in interest in models based on **aggregate data**



THE RETURN OF THE MARKETING MIX MODEL



Statistics > Applications

[Submitted on 7 Jun 2021 ([v1](#)), last revised 5 Sep 2021 (this version, v3)]

Bayesian Time Varying Coefficient Model with Applications to Marketing Mix Modeling

Edwin Ng, Zhishi Wang, Athena Dai

Both Bayesian and varying coefficient models are very useful tools in practice as they can be used to model parameter heterogeneity in a generalizable way. Motivated by the need of enhancing Marketing Mix Modeling at Uber, we propose a Bayesian Time Varying Coefficient model, equipped with a hierarchical Bayesian structure. This model is different from other time varying coefficient models in the sense that the coefficients are weighted over a set of local latent variables following certain probabilistic distributions. Stochastic Variational Inference is used to approximate the posteriors of latent variables and dynamic coefficients. The proposed model also helps address many challenges faced by traditional MMM approaches. We used simulations as well as real world marketing datasets to demonstrate our model superior performance in terms of both accuracy and interpretability.

[Submitted on 7 Jun 2021 ([v1](#)), last revised 5 Sep 2021 (this version)]

Bayesian Time Varying Coefficients

Edwin Ng, Zhishi Wang, Athena Dai

Both Bayesian and varying coefficient models are very generalizable way. Motivated by the need of enhancing model, equipped with a hierarchical Bayesian structure coefficients are weighted over a set of local latent variables to approximate the posteriors of latent variables and dynamic coefficients. The proposed model also helps address many challenges faced by traditional MMM approaches. We used simulations as well as real world marketing datasets to demonstrate our model superior performance in terms of both accuracy and interpretability.



Orbit

[Submitted on 7 Ju



Robyn

Documentation

Getting Started

Case Studies

Resources

Bayesian

Edwin Ng, Zhi

Both Bayesian and frequentist approaches have their own merits and generalizability. Robyn is built on a Bayesian framework, which provides a more principled model, equipped with prior distributions that can incorporate domain knowledge and coefficients are interpreted as posterior distributions. Robyn also provides methods to approximate the posterior distributions of the parameters, which are useful for traditional Marketing Mix Modeling (MMM) analysis. Robyn is designed to be user friendly in terms of both the interface and the underlying statistical models.

Robyn

Robyn is an experimental, AI/ML-powered and open sourced Marketing Mix Modeling (MMM) package from Meta Marketing Science.

Getting Started

Statistics > Applications

[Submitted on 7 Ju

Bayesian

Edwin Ng, Zhi

Both Bayesian and frequentist methods have their own strengths and weaknesses. Robyn is a generalizable, open-source Python package that provides a unified interface for fitting Bayesian models, equipped with a variety of priors and samplers. It also provides a way to approximate coefficients and uncertainty intervals for parameters in a model. Robyn is designed to approximate coefficients and uncertainty intervals for parameters in a model. Robyn is designed to be used in both traditional MCMC and variational inference frameworks. Robyn is designed to be used in both traditional MCMC and variational inference frameworks. Robyn is designed to be used in both traditional MCMC and variational inference frameworks.

Robyn

Robyn is an experimental, AI-powered, open-source Python package for Marketing
Modeling (MMM) package

Getting Started



Case Studies

Resources



PyMC-Marketing

Statistics > Applications

[Submitted on 7 Ju

Bayesian

Edwin Ng, Zhi

Both Bayesian and frequentist approaches have their own strengths and weaknesses. Bayesian methods are more generalizable and can incorporate prior knowledge, while frequentist methods are more robust to model misspecification. Bayesian methods also provide a natural way to handle uncertainty and make predictions based on the posterior distribution of parameters. In contrast, frequentist methods often rely on asymptotic approximations and can be less transparent about the assumptions underlying their results.



Robyn

Documentation

Getting Started

Case Studies

Resources



arXiv

Meridian

Search

English ▾

Home

Modeling

Empower your team with best-in-class marketing mix models and drive better business outcome

Meridian is an open-source MMM built by Google that provides innovative solutions to key measurement challenges

Apply for early access

Meridian is currently offered in limited availability

Marketing

Marketing Mix Models

- Very long history in marketing

Borden (1964), Palda (1965), Bultez and Naert (1979), Little (1979), Winer (1979), ..., Hanssens et al. (2003), ...

Marketing Mix Models

- Very long history in marketing

Borden (1964), Palda (1965), Bultez and Naert (1979), Little (1979), Winer (1979), ..., Hanssens et al. (2003), ...

- At its heart: just regression

$$y_t = \sum_j \beta_j x_{jt} + \epsilon_t \quad -\text{or}- \quad \log y_t = \sum_j \beta_j \log x_{jt} + \epsilon_t$$

Marketing Mix Models

- Very long history in marketing

Borden (1964), Palda (1965), Bultez and Naert (1979), Little (1979), Winer (1979), ..., Hanssens et al. (2003), ...

- At its heart: just regression

$$y_t = \sum_j \beta_j x_{jt} + \epsilon_t \quad -\text{or}- \quad \log y_t = \sum_j \beta_j \log x_{jt} + \epsilon_t$$

- ...with many well-documented **issues and extensions**

Marketing Mix Models

- Very long history in marketing

Borden (1964), Palda (1965), Bultez and Naert (1979), Little (1979), Winer (1979), ..., Hanssens et al. (2003), ...

- At its heart: just regression

$$y_t = \sum_j \beta_j x_{jt} + \epsilon_t \quad -\text{or}- \quad \log y_t = \sum_j \beta_j \log x_{jt} + \epsilon_t$$

- ...with many well-documented **issues and extensions**

- Long-term effects, often captured through **AdStock**: $S(x_t, L, \alpha) = \frac{\sum_{\ell=0}^{L-1} \alpha^\ell x_{t-\ell}}{\sum_{\ell=0}^{L-1} \alpha^\ell}$

Marketing Mix Models

- Very long history in marketing

Borden (1964), Palda (1965), Bultez and Naert (1979), Little (1979), Winer (1979), ..., Hanssens et al. (2003), ...

- At its heart: just regression

$$y_t = \sum_j \beta_j x_{jt} + \epsilon_t \quad -\text{or}- \quad \log y_t = \sum_j \beta_j \log x_{jt} + \epsilon_t$$

- ...with many well-documented **issues and extensions**

- Long-term effects, often captured through **AdStock**: $S(x_t, L, \alpha) = \frac{\sum_{\ell=0}^{L-1} \alpha^\ell x_{t-\ell}}{\sum_{\ell=0}^{L-1} \alpha^\ell}$
- **Nonlinear** effects: $y_t = \sum_j f_j(x_{jt}) + \epsilon_t$

Marketing Mix Models

- Very long history in marketing

Borden (1964), Palda (1965), Bultez and Naert (1979), Little (1979), Winer (1979), ..., Hanssens et al. (2003), ...

- At its heart: just regression

$$y_t = \sum_j \beta_j x_{jt} + \epsilon_t \quad -\text{or}- \quad \log y_t = \sum_j \beta_j \log x_{jt} + \epsilon_t$$

- ...with many well-documented **issues and extensions**

- Long-term effects, often captured through **AdStock**: $S(x_t, L, \alpha) = \frac{\sum_{\ell=0}^{L-1} \alpha^\ell x_{t-\ell}}{\sum_{\ell=0}^{L-1} \alpha^\ell}$
- **Nonlinear** effects: $y_t = \sum_j f_j(x_{jt}) + \epsilon_t$
- **Time-varying** effects: $y_t = \sum_j \beta_{jt} x_{jt} + \epsilon_t$

Marketing Mix Models

- Very long history in marketing

Borden (1964), Palda (1965), Bultez and Naert (1979), Little (1979), Winer (1979), ..., Hanssens et al. (2003), ...

- At its heart: just regression

$$y_t = \sum_j \beta_j x_{jt} + \epsilon_t \quad -\text{or}- \quad \log y_t = \sum_j \beta_j \log x_{jt} + \epsilon_t$$

- ...with many well-documented **issues and extensions**

- Long-term effects, often captured through **AdStock**: $S(x_t, L, \alpha) = \frac{\sum_{\ell=0}^{L-1} \alpha^\ell x_{t-\ell}}{\sum_{\ell=0}^{L-1} \alpha^\ell}$
- **Nonlinear** effects: $y_t = \sum_j f_j(x_{jt}) + \epsilon_t$
- **Time-varying** effects: $y_t = \sum_j \beta_{jt} x_{jt} + \epsilon_t$
- Endogeneity 

Marketing Mix Models

- Very long history in marketing

Borden (1964), Palda (1965), Bultez and Naert (1979), Little (1979), Winer (1979), ..., Hanssens et al. (2003), ...

- At its heart: just regression

$$y_t = \sum_j \beta_j x_{jt} + \epsilon_t \quad -\text{or}- \quad \log y_t = \sum_j \beta_j \log x_{jt} + \epsilon_t$$

- ...with many well-documented **issues and extensions**

- Long-term effects, often captured through **AdStock**: $S(x_t, L, \alpha) = \frac{\sum_{\ell=0}^{L-1} \alpha^\ell x_{t-\ell}}{\sum_{\ell=0}^{L-1} \alpha^\ell}$
- **Nonlinear** effects: $y_t = \sum_j f_j(x_{jt}) + \epsilon_t$
- **Time-varying** effects: $y_t = \sum_j \beta_{jt} x_{jt} + \epsilon_t$
- Endogeneity 

Note: Terminology

Throughout the talk,
dynamic = time-varying

What's new? Firepower



What's new? Firepower



Meridian

Wang et al. (2017), Jin et al. (2017), Sun et al. (2017), Zhang et al. (2023)

$$\begin{aligned}y_{g,t} = & \mu_t + \tau_g + \sum_{c=1}^C \gamma_{g,c} z_{g,t,c} \\& + \sum_{m=1}^M \beta_{g,m} HillAdstock \left(\left\{ x_{g,t-s,m} \right\}_{s=0}^L ; \alpha_m, ec_m, slope_m \right) \\& + \sum_{n=1}^N \beta_{g,n}^{(rf)} Adstock \left(\left\{ r_{g,t-s,n} \cdot Hill \left(f_{g,t-s,n}; ec_n^{(rf)}, slope_n^{(rf)} \right) \right\}_{s=0}^L ; \alpha_n^{(rf)} \right) \\& + \epsilon_{g,t}\end{aligned}$$

What's new? Firepower



Meridian


Wang et al. (2017), Jin et al. (2017), Sun et al. (2017), Zhang et al. (2023)


$$\begin{aligned}y_{g,t} = & \mu_t + \tau_g + \sum_{c=1}^C \gamma_{g,c} z_{g,t,c} \\& + \sum_{m=1}^M \beta_{g,m} HillAdstock \left(\left\{ x_{g,t-s,m} \right\}_{s=0}^L ; \alpha_m, ec_m, slope_m \right) \\& + \sum_{n=1}^N \beta_{g,n}^{(rf)} Adstock \left(\left\{ r_{g,t-s,n} \cdot Hill \left(f_{g,t-s,n}; ec_n^{(rf)}, slope_n^{(rf)} \right) \right\}_{s=0}^L ; \alpha_n^{(rf)} \right) \\& + \epsilon_{g,t}\end{aligned}$$

Uber's Orbit

Ng et al., (2021)

$$\ln(\hat{y}_t) = l_t + s_t + \sum_{p=1}^P \ln(x_{t,p}) \beta_{t,p}$$
$$\beta_{t,p} = \sum_j w_j(t) \cdot b_{j,p},$$
$$w_j(t) = k(t, t_j) / \sum_{i=1}^J k(t, t_i),$$






**Surely this is nonlinear,
right...?**

True data generating process (DGP):


$$y_t = \beta_t x_t + \epsilon_t$$




**Surely this is nonlinear,
right...?**

True data generating process (DGP):

$$y_t = \beta_t x_t + \epsilon_t$$




**Surely this is nonlinear,
right...?**




True data generating process (DGP):

$$y_t = \beta_t x_t + \epsilon_t$$



**Surely this is nonlinear,
right...?**



**When can
nonlinear and
time-varying
effects be identified
in marketing mix
models?**

When can nonlinear and time-varying effects be identified in marketing mix models?

1. Introduce a **flexible, nonparametric framework** to estimate both types of effects

When can nonlinear and time-varying effects be identified in marketing mix models?

1. Introduce a **flexible, nonparametric framework** to estimate both types of effects
2. Through both theory and simulations, we show that, under realistic conditions, **the two models will be conflated**

When can nonlinear and time-varying effects be identified in marketing mix models?

1. Introduce a **flexible, nonparametric framework** to estimate both types of effects
2. Through both theory and simulations, we show that, under realistic conditions, **the two models will be conflated**
3. Problems are exacerbated under common managerial practices, like **autoregressive decision-making**

When can nonlinear and time-varying effects be identified in marketing mix models?

1. Introduce a **flexible, nonparametric framework** to estimate both types of effects
2. Through both theory and simulations, we show that, under realistic conditions, **the two models will be conflated**
3. Problems are exacerbated under common managerial practices, like **autoregressive decision-making**
4. Similarly fitting models can have **fundamentally different implications** in terms of optimal decision-making

(A little) Math

When can dynamic approximate nonlinear?

When can dynamic approximate nonlinear?

$$y_t = \sum_j \beta_{jt} x_{jt} + \epsilon_t$$

$$y_t = \sum_j f_j(x_{jt}) + \epsilon_t$$

When can dynamic approximate nonlinear?

$$y_t = \sum_j \beta_{jt} x_{jt} + \epsilon_t$$

$$y_t = \sum_j f_j(x_{jt}) + \epsilon_t$$

- First, a simple point: if $\beta_t = y_t/x_t$, the dynamic model will fit perfectly!

When can dynamic approximate nonlinear?

$$y_t = \sum_j \beta_{jt} x_{jt} + \epsilon_t$$

$$y_t = \sum_j f_j(x_{jt}) + \epsilon_t$$


- First, a simple point: if $\beta_t = y_t/x_t$, the dynamic model will fit perfectly!
- A more nuanced answer: in any local region of x , a local linear approximation will work pretty well

When can dynamic approximate nonlinear?

$$y_t = \sum_j \beta_{jt} x_{jt} + \epsilon_t$$

$$y_t = \sum_j f_j(x_{jt}) + \epsilon_t$$

- First, a simple point: if $\beta_t = y_t/x_t$, the dynamic model will fit perfectly!
- A more nuanced answer: in any local region of x , a local linear approximation will work pretty well




When can dynamic approximate nonlinear?

$$y_t = \sum_j \beta_{jt} x_{jt} + \epsilon_t$$

$$y_t = \sum_j f_j(x_{jt}) + \epsilon_t$$

- First, a simple point: if $\beta_t = y_t/x_t$, the dynamic model will fit perfectly!
- A more nuanced answer: in any local region of x , a local linear approximation will work pretty well
- When will β_t be smooth (“forecastable”)? If f is **smooth** and x_t and x_{t+1} are **close together**!



When can nonlinear approximate dynamic?

$$y_t = \sum_j f_j(x_{jt}) + \epsilon_t$$

$$y_t = \sum_j \beta_{jt}x_{jt} + \epsilon_t$$

When can nonlinear approximate dynamic?



$$y_t = \sum_j f_j(x_{jt}) + \epsilon_t$$

$$y_t = \sum_j \beta_{jt}x_{jt} + \epsilon_t$$

- Additional notation: let $\beta_t = \mathcal{B}(t)$, $x_t = \mathcal{X}(t)$

When can nonlinear approximate dynamic?



$$y_t = \sum_j f_j(x_{jt}) + \epsilon_t$$

$$y_t = \sum_j \beta_{jt}x_{jt} + \epsilon_t$$

- Additional notation: let $\beta_t = \mathcal{B}(t)$, $x_t = \mathcal{X}(t)$
- The nonlinear model can fully approximate the dynamic model when $\mathcal{B}(t) = g(\mathcal{X}(t))$ for some function g

When can nonlinear approximate dynamic?



$$y_t = \sum_j f_j(x_{jt}) + \epsilon_t$$

$$y_t = \sum_j \beta_{jt} x_{jt} + \epsilon_t$$

- Additional notation: let $\beta_t = \mathcal{B}(t)$, $x_t = \mathcal{X}(t)$
- The nonlinear model can fully approximate the dynamic model when $\mathcal{B}(t) = g(\mathcal{X}(t))$ for some function g
- Simple case: $\mathcal{X}(t)$ is **monotonic**, in which case the nonlinear model is:

$$f(x) = \mathcal{B}(\mathcal{X}^{-1}(t)) x$$

When can nonlinear approximate dynamic?



$$y_t = \sum_j f_j(x_{jt}) + \epsilon_t$$

$$y_t = \sum_j \beta_{jt} x_{jt} + \epsilon_t$$

- Additional notation: let $\beta_t = \mathcal{B}(t)$, $x_t = \mathcal{X}(t)$
- The nonlinear model can fully approximate the dynamic model when $\mathcal{B}(t) = g(\mathcal{X}(t))$ for some function g
- Simple case: $\mathcal{X}(t)$ is **monotonic**, in which case the nonlinear model is:

$$f(x) = \mathcal{B}(\mathcal{X}^{-1}(t)) x$$

- Less simple case: if x and β both derive from a **common parent process**

When can nonlinear approximate dynamic?



$$y_t = \sum_j f_j(x_{jt}) + \epsilon_t$$





$$y_t = \sum_j \beta_{jt} x_{jt} + \epsilon_t$$

- Additional notation: let $\beta_t = \mathcal{B}(t)$, $x_t = \mathcal{X}(t)$
- The nonlinear model can fully approximate the dynamic model when $\mathcal{B}(t) = g(\mathcal{X}(t))$ for some function g
- Simple case: $\mathcal{X}(t)$ is **monotonic**, in which case the nonlinear model is:

$$f(x) = \mathcal{B}(\mathcal{X}^{-1}(t)) x$$

- Less simple case: if x and β both derive from a **common parent process**



Simulations

When does conflation *actually* happen?

When does conflation *actually* happen?

Four types of simulations:

1. Flexible nonlinear response

$$y_t = f(x_t) + \varepsilon_t$$

2. Dynamic coefficients, linear (or log-linear) model

$$y_t = \beta(t)x_t + \varepsilon_t$$

3. Nonlinear response, parametric hill function

$$y_t = \text{Hill}(x_t) + \varepsilon_t$$

4. Dynamic coefficients inherited from common parent

$$y_t = \beta(t)x(t) + \varepsilon_t, (\beta(t), x(t)) \sim \text{Pa}(t)$$

When does conflation *actually* happen?

Four types of simulations:

1. Flexible nonlinear response

$$y_t = f(x_t) + \varepsilon_t$$

2. Dynamic coefficients, linear (or log-linear) model

$$y_t = \beta(t)x_t + \varepsilon_t$$

3. Nonlinear response, parametric hill function

$$y_t = \text{Hill}(x_t) + \varepsilon_t$$

4. Dynamic coefficients inherited from common parent

$$y_t = \beta(t)x(t) + \varepsilon_t, (\beta(t), x(t)) \sim \text{Pa}(t)$$

When does conflation *actually* happen?

Four types of simulations:

1. Flexible nonlinear response

$$y_t = \boxed{f(x_t)} + \varepsilon_t$$

2. Dynamic coefficients, linear (or log-linear) model

$$y_t = \boxed{\beta(t)x_t} + \varepsilon_t$$

3. Nonlinear response, parametric hill function

$$y_t = \text{Hill}(x_t) + \varepsilon_t$$

4. Dynamic coefficients inherited from common parent

$$y_t = \beta(t)x(t) + \varepsilon_t, (\beta(t), x(t)) \sim \text{Pa}(t)$$

When does conflation *actually* happen?

Four types of simulations:

1. Flexible nonlinear response

$$y_t = \boxed{f(x_t)} + \varepsilon_t$$

2. Dynamic coefficients, linear (or log-linear) model

$$y_t = \boxed{\beta(t)x_t} + \varepsilon_t$$

3. Nonlinear response, parametric hill function

$$y_t = \text{Hill}(x_t) + \varepsilon_t$$

4. Dynamic coefficients inherited from common parent

$$y_t = \beta(t)x(t) + \varepsilon_t, (\beta(t), x(t)) \sim \text{Pa}(t)$$

Gaussian processes

Two important levers:

- Smoothness
- Amplitude

Primer: Gaussian Processes

Gaussian processes: a Bayesian nonparametric approach to modeling unknown functions

Primer: Gaussian Processes

Gaussian processes: a Bayesian nonparametric approach to modeling unknown functions

$$f(x) \sim \mathcal{GP}(m(x), k(x, x'))$$

Primer: Gaussian Processes

Gaussian processes: a Bayesian nonparametric approach to modeling unknown functions

$$f(x) \sim \mathcal{GP}(m(x), k(x, x'))$$



$$f(x_1, \dots, x_N) \sim \mathcal{N}(m(x_1, \dots, x_N), K), \text{ s.t. } K_{ij} = k(x_i, x_j)$$

Primer: Gaussian Processes

Gaussian processes: a Bayesian nonparametric approach to modeling unknown functions

$$f(x) \sim \mathcal{GP}(m(x), k(x, x'))$$

$\uparrow\downarrow$

$$f(x_1, \dots, x_N) \sim \mathcal{N}(m(x_1, \dots, x_N), K), \text{ s.t. } K_{ij} = k(x_i, x_j)$$

$$k(x, x') = \eta^2 \exp \left\{ -\frac{(x - x')^2}{2\rho^2} \right\}$$

Primer: Gaussian Processes

Gaussian processes: a Bayesian nonparametric approach to modeling unknown functions

$$f(x) \sim \mathcal{GP}(m(x), k(x, x'))$$

↑↓

$$f(x_1, \dots, x_N) \sim \mathcal{N}(m(x_1, \dots, x_N), K), \text{ s.t. } K_{ij} = k(x_i, x_j)$$

Amplitude

$$k(x, x') = \eta^2 \exp \left\{ -\frac{(x - x')^2}{2\rho^2} \right\}$$

Lengthscale

Primer: Gaussian Processes

Gaussian processes: a Bayesian nonparametric approach to modeling unknown functions

$$f(x) \sim \mathcal{GP}(m(x), k(x, x'))$$




$$f(x_1, \dots, x_N) \sim \mathcal{N}(m(x_1, \dots, x_N), K), \text{ s.t. } K_{ij} = k(x_i, x_j)$$

Amplitude

$$k(x, x') = \eta^2 \exp \left\{ -\frac{(x - x')^2}{2\rho^2} \right\}$$

Lengthscale



When does conflation *actually* happen?

Four types of simulations:

1. Flexible nonlinear response

$$y_t = \boxed{f(x_t)} + \varepsilon_t$$

2. Dynamic coefficients, linear (or log-linear) model

$$y_t = \boxed{\beta(t)x_t} + \varepsilon_t$$

3. Nonlinear response, parametric hill function

$$y_t = \text{Hill}(x_t) + \varepsilon_t$$

4. Dynamic coefficients inherited from common parent

$$y_t = \beta(t)x(t) + \varepsilon_t, (\beta(t), x(t)) \sim \text{Pa}(t)$$

Gaussian processes

Two important levers:

- Smoothness
- Amplitude

When does conflation *actually* happen?

Four types of simulations:

1. Flexible nonlinear response

$$y_t = f(x_t) + \varepsilon_t$$

2. Dynamic coefficients, linear (or log-linear) model

$$y_t = \beta(t)x_t + \varepsilon_t$$

3. Nonlinear response, parametric hill function

Other manipulated features:

4. Dynamic coefficient
- Autoregressive coefficient in x
 - Noise in x 's autoregressive process
 - Variance of the error term

Gaussian processes

Two important levers:

- Smoothness
- Amplitude


Simulation Results

- For each simulation type, >300 settings, systematically varying the previously described factors, with 100 simulations per setting
- Fit both models (nonlinear and dynamic), measure conflation through validation RMSE

Examples

Nonlinear DGP


Smooth f , low noise



Data

Examples

Data




Examples


Nonlinear DGP

Smooth f , low noise

Data




Implied
 $\beta(t)$




Examples

Data




Implied
 $\beta(t)$




Examples

Data






Implied
 $\beta(t)$






Examples

Data






Implied
 $\beta(t)$






Examples

Data



Implied
 $\beta(t)$



Simulation

- Systematically vary the previously described factors, running 100 simulations per setting, and fit both models (nonlinear and dynamic)
- Any conflation = for at least one simulation of that setting, the wrong model had better validation RMSE
- Major conflation = conflation on over 25% of that setting's simulations

Simulation

- Systematically vary the previously described factors, running 100 simulations per setting, and fit both models (nonlinear and dynamic)
- Any conflation = for at least one simulation of that setting, the wrong model had better validation RMSE
- Major conflation = conflation on over 25% of that setting's simulations

True DGP	No Stock		High Stock	
	Any	Major	Any	Major
Nonlinear	81%	27%	85%	30%
Time-varying	91%	40%	99%	44%
Hill	83%	46%	92%	47%

Diving Deeper

DV: % Conflated Simulations

Variable	Level	Nonlinear DGP		Time-varying DGP	
		Coef	$P(> t)$	Coef	$P(> t)$
Amplitude, f :	Low (1)	-	-	-	-
	Middle (2)	-0.24	0.65	0.23	0.62
	High (5)	-0.65	0.22	0.09	0.85
Smoothness, f :	Low (0.1)	-	-	-	-
	Middle (0.5)	10.36	0.00	7.93	0.00
	High (1)	18.86	0.00	12.21	0.00
AR coef, x :	Low (0)	-	-	-	-
	Middle (0.5)	-0.56	0.29	0.63	0.18
	High (1)	0.42	0.43	13.38	0.00
AR Variance, x :	Low (1)	-	-	-	-
	Middle (5)	0.11	0.84	0.06	0.90
	High (10)	-0.47	0.37	0.36	0.44
Noise, y	Low (0.01)	-	-	-	-
	Middle (0.1)	14.10	0.00	13.05	0.00
	High (0.2)	23.28	0.00	22.71	0.00
AdStock	Low (0)	-	-	-	-
	Middle (0.3)	-0.14	0.79	0.83	0.08
	High (0.8)	1.59	0.00	2.49	0.00

Table 2: Simulation Results
 DV = Percentage Conflation; Intercept omitted for clarity.

Diving Deeper

DV: % Conflated Simulations

Conflation more likely with...

- Noisier data (i.e., ϵ_t)

Variable	Level	Nonlinear DGP		Time-varying DGP	
		Coef	$P(> t)$	Coef	$P(> t)$
Amplitude, f :	Low (1)	-	-	-	-
	Middle (2)	-0.24	0.65	0.23	0.62
	High (5)	-0.65	0.22	0.09	0.85
Smoothness, f :	Low (0.1)	-	-	-	-
	Middle (0.5)	10.36	0.00	7.93	0.00
	High (1)	18.86	0.00	12.21	0.00
AR coef, x :	Low (0)	-	-	-	-
	Middle (0.5)	-0.56	0.29	0.63	0.18
	High (1)	0.42	0.43	13.38	0.00
AR Variance, x :	Low (1)	-	-	-	-
	Middle (5)	0.11	0.84	0.06	0.90
	High (10)	-0.47	0.37	0.36	0.44
Noise, y	Low (0.01)	-	-	-	-
	Middle (0.1)	14.10	0.00	13.05	0.00
	High (0.2)	23.28	0.00	22.71	0.00
AdStock	Low (0)	-	-	-	-
	Middle (0.3)	-0.14	0.79	0.83	0.08
	High (0.8)	1.59	0.00	2.49	0.00

Table 2: Simulation Results
DV = Percentage Conflation; Intercept omitted for clarity.

Diving Deeper

DV: % Conflated Simulations

Conflation more likely with...

- Noisier data (i.e., ϵ_t)
- Smoother functions

Variable	Level	Nonlinear DGP		Time-varying DGP	
		Coef	$P(> t)$	Coef	$P(> t)$
Amplitude, f :	Low (1)	-	-	-	-
	Middle (2)	-0.24	0.65	0.23	0.62
	High (5)	-0.65	0.22	0.09	0.85
Smoothness, f :	Low (0.1)	-	-	-	-
	Middle (0.5)	10.36	0.00	7.93	0.00
	High (1)	18.86	0.00	12.21	0.00
AR coef, x :	Low (0)	-	-	-	-
	Middle (0.5)	-0.56	0.29	0.63	0.18
	High (1)	0.42	0.43	13.38	0.00
AR Variance, x :	Low (1)	-	-	-	-
	Middle (5)	0.11	0.84	0.06	0.90
	High (10)	-0.47	0.37	0.36	0.44
Noise, y	Low (0.01)	-	-	-	-
	Middle (0.1)	14.10	0.00	13.05	0.00
	High (0.2)	23.28	0.00	22.71	0.00
AdStock	Low (0)	-	-	-	-
	Middle (0.3)	-0.14	0.79	0.83	0.08
	High (0.8)	1.59	0.00	2.49	0.00

Table 2: Simulation Results
DV = Percentage Conflation; Intercept omitted for clarity.

Diving Deeper

DV: % Conflated Simulations

Conflation more likely with...

- Noisier data (i.e., ϵ_t)
- Smoother functions
- High degree of autocorrelation

Variable	Level	Nonlinear DGP		Time-varying DGP	
		Coef	$P(> t)$	Coef	$P(> t)$
Amplitude, f :	Low (1)	-	-	-	-
	Middle (2)	-0.24	0.65	0.23	0.62
	High (5)	-0.65	0.22	0.09	0.85
Smoothness, f :	Low (0.1)	-	-	-	-
	Middle (0.5)	10.36	0.00	7.93	0.00
	High (1)	18.86	0.00	12.21	0.00
AR coef, x :	Low (0)	-	-	-	-
	Middle (0.5)	-0.56	0.29	0.63	0.18
	High (1)	0.42	0.43	13.38	0.00
AR Variance, x :	Low (1)	-	-	-	-
	Middle (5)	0.11	0.84	0.06	0.90
	High (10)	-0.47	0.37	0.36	0.44
Noise, y	Low (0.01)	-	-	-	-
	Middle (0.1)	14.10	0.00	13.05	0.00
	High (0.2)	23.28	0.00	22.71	0.00
AdStock	Low (0)	-	-	-	-
	Middle (0.3)	-0.14	0.79	0.83	0.08
	High (0.8)	1.59	0.00	2.49	0.00

Table 2: Simulation Results
DV = Percentage Conflation; Intercept omitted for clarity.

Diving Deeper

DV: % Conflated Simulations

Conflation more likely with...

- Noisier data (i.e., ϵ_t)
- Smoother functions
- High degree of autocorrelation


Decisions are often autocorrelated!

Variable	Level	Nonlinear DGP		Time-varying DGP	
		Coef	$P(> t)$	Coef	$P(> t)$
Amplitude, f :	Low (1)	-	-	-	-
	Middle (2)	-0.24	0.65	0.23	0.62
	High (5)	-0.65	0.22	0.09	0.85
Smoothness, f :	Low (0.1)	-	-	-	-
	Middle (0.5)	10.36	0.00	7.93	0.00
	High (1)	18.86	0.00	12.21	0.00
AR coef, x :	Low (0)	-	-	-	-
	Middle (0.5)	-0.56	0.29	0.63	0.18
	High (1)	0.42	0.43	13.38	0.00
AR Variance, x :	Low (1)	-	-	-	-
	Middle (5)	0.11	0.84	0.06	0.90
	High (10)	-0.47	0.37	0.36	0.44
Noise, y	Low (0.01)	-	-	-	-
	Middle (0.1)	14.10	0.00	13.05	0.00
	High (0.2)	23.28	0.00	22.71	0.00
AdStock	Low (0)	-	-	-	-
	Middle (0.3)	-0.14	0.79	0.83	0.08
	High (0.8)	1.59	0.00	2.49	0.00

Table 2: Simulation Results
DV = Percentage Conflation; Intercept omitted for clarity.


Implications

One last simulation...




(or from roughly \$8,000 to \$14,000)


One last simulation...





(or from roughly \$8,000 to \$14,000)




One last simulation...





(or from roughly \$8,000 to \$14,000)



One last simulation...






(or from roughly \$8,000 to \$14,000)





Real Data: Classic and Modern

- Classic Application 1: Dietary Weight Control (DWC)
(Bass and Clark, 1972)
- Classic Application 2: Lydia Pinkham (LP)
(Palda, 1964)
- Modern applications: MMM data from Nielsen








	Nonlinear	Time-varying	Hill	Log Time-varying
Dietary Weight Control	3.23 [2.22, 5.58]	2.97 [1.72, 5.09]	3.10 [2.59, 4.44]	2.55 [1.50, 4.13]

Spend Optimization



- DWC data: sales volume, but no prices
- Limitation = opportunity: optimal ad spend across a range of prices

Spend Optimization

- DWC data: sales volume, but no prices
- Limitation = opportunity: optimal ad spend across a range of prices




Lydia Pinkham





	Nonlinear	Time-varying	Hill	Log Time-varying
Lydia Pinkham	89.37 [79.41, 104.88]	83.21 [75.44, 95.24]	100.50 [91.68, 114.97]	88.92 [83.23, 98.00]



Lydia Pinkham



Lydia Pinkham



Lydia Pinkham



	Nonlinear	Time-varying	Hill	Log Time-varying
Lydia Pinkham	89.37 [79.41, 104.88]	83.21 [75.44, 95.24]	100.50 [91.68, 114.97]	88.92 [83.23, 98.00]

Spend Optimization with Carryover


- Trickier: spending today affects AdStock for many future periods

Spend Optimization with Carryover

- Trickier: spending today affects AdStock for many future periods
- Follow Google's practice: for L carryover periods, determine optimal spend in a test window, predict revenue L periods out


Spend Optimization with Carryover

- Trickier: spending today affects AdStock for many future periods
- Follow Google's practice: for L carryover periods, determine optimal spend in a test window, predict revenue L periods out



Spend Optimization with Carryover

- Trickier: spending today affects AdStock for many future periods
- Follow Google's practice: for L carryover periods, determine optimal spend in a test window, predict revenue L periods out
- For LP, optimal ad spend:
 - Nonlinear model: **\$90,127**
 - Time-varying model: **\$78,333**




Modern MMM Data

- Again, clear conflation...

	Chocolate	Pet Food	Coffee	Beer
Nonlinear (GP)	368,766 [207,634, 601,163]	48,438 [27,920, 83,249]	722,076 [509,527, 919,820]	443,817 [236,730, 911,732]
Time-varying (GP)	387,623 [207,350, 671,246]	41,825 [24,631, 65,232]	733,269 [490,785, 994,303]	423,085 [207,914, 802,740]


Modern MMM Data

- Case study: Coffee



Modern MMM Data

- Case study: Coffee





So is MMM doomed?

Test 1: Maximal Separation

- A type of “bump up” incrementality test
- Intuition: set spending next period to maximize the difference in predicted values across the two models

Test 1: Maximal Separation

- A type of “bump up” incrementality test
- Intuition: set spending next period to maximize the difference in predicted values across the two models





Test 2: See-saw

- Easier to implement
- Intuition: high-low spending over adjacent time periods

Test 2: See-saw

- Easier to implement
- Intuition: high-low spending over adjacent time periods



Conclusions

Current MMM Practice Might Be Flawed!

- We show that, under many common spending patterns, **time-varying and nonlinear effects cannot be disentangled**, despite having different implications
- This problem is potentially **very widespread**: increasing complexity in models, widespread practice of “model refreshes” to capture changing markets
- Our work both introduces a **framework for estimating** these types of models, and provides **solutions for understanding and preventing** conflation

Thanks!

Feedback or questions:
ryandew@wharton.upenn.edu

Working paper:
www.rtdew.com



You

generate an image of a marketing mix model paradise with some very happy computers and a cameo from the reverend thomas bayes

