

Understanding the utility of archived tag-recapture data for evaluation of movement and mortality estimation

Robert Leaf, Kasea Price, and Amy Schueller (NOAA)

April 16, 2020

Introduction

A high priority research need for improving the assessment of Gulf Menhaden include understanding stock structure, individual growth dynamics, and improving estimates of natural mortality. One of the primary ways that mortality and growth can be understood is through tagging and recapture of individuals (Leaf et al. 2007, Leaf et al. 2008). Similarly, migration and stock delineation have also been described with such tagging studies. Because of the power of tag and recapture studies for understanding stock dynamics, the Gulf Menhaden Stock Assessment report further recommended replicating the tagging work performed by NOAA scientists, which were extensive: Pristas tagged 75,000 *B. patronus* from 1969 to 1972 and Ahrenholz tagged 237,000 *B. patronus* from 1970 to 1985. These studies involved tagging juveniles and adults with internal, individually-numbered ferro-magnetic tags that were recovered on magnets in fish reduction factories. Because of the costs of such studies, it is unlikely that such a large-scale tagging study will ever be performed again. However, leaps in the computational power and statistical modeling approaches in recent decades has made re-analysis of the previously collected data a very worthwhile endeavor. Currently, the data from these tagging studies reside as paper copies kept at the Southeast Fishery Science Center, Beaufort, NC and are not available for digital analysis.

The project started in 1969 (Table 1, 2). The scientists working on the project went out, a few days every couple of months to tag Gulf Menhaden. They used a small version of a purse seine to collect fish to tag. The tags were magnetic and sized appropriately to the length of the fish, “small” and “medium” sized. The smaller tags were used to tag juvenile fish and medium were used for adults. They inserted the tags into the abdominal cavity of each fish and released it. In order to get the tag recovery (or recapture) data, they depended on the menhaden processing companies. The tags were retrieved by sampling the processing conveyor belts which were equipped with magnets running along the top.

Methods

We partnered with NOAA scientist Dr. Amy Schueller at the SEFSC Beaufort Lab, who supported our efforts by providing hard copies of Dean Ahrenholz’s tag-recapture data collected in the 1960’s, 1970’s, and 1980’s. To perform exploratory analysis to understand the feasibility of a comprehensive reanalysis of the tag and recapture data (the overarching theme of the project) we:

- 1.) Acquired and cataloged existing data,
- 2.) Digitized the data,
- 3.) Performed quality assurance and quality control, and
- 4.) Evaluated the utility of the data by comparing the magnitude of tagged and recaptured fish and the data structure with those studies performed by Liljestrand et al.:
 - Liljestrand, E.M., M.J. Wilberg, and A.M. Schueller. 2019. Estimation of movement and mortality of Atlantic menhaden during 1966-1969 using a Bayesian multi-state mark-recovery model. Fisheries Research 210:204-213.

- Liljestrand, E.M., M.J. Wilberg, and A.M. Schueller. 2019. Multi-state dead recovery mark-recovery model performance for estimating movement and mortality rates. *Fisheries Research* 210:214-223.

Results

Acquired and cataloged existing data, digitize the data, perform quality assurance and quality control

K. Price traveled to Beaufort, NC in October of 2019 to sort through and obtain original data on Gulf Menhaden tag and recapture efforts from the 60s, 70s, and 80s. She and A. Schueller sorted through about 20 books (bound volumes of printed material) of data. Three primary sources of data were identified:

1. The tag and recapture histories of adult and juvenile individuals

These data are the record of individual fish captured (in the field) and recaptured (in a processing plant) over the course of the study. These data were scanned to .pdf files and then digitized and saved as Excel documents.

The data are comprised of the following variables:

- Tag_Year, the year in which the menhaden were originally tagged.
- Tag_Month, the month in which the menhaden were originally tagged.
- Tag_Day, the day on which the menhaden were originally tagged.
- Tag_Series is denoted by the letter, this indicates the series of tags being deployed.
- Tag_Series_Number indicates which group in the tag series with which the Gulf Menhaden were tagged. Tag Series and Tag Series Number are paired. For each group of tags, with a tag series and number, (example: U20) has 100 individual tags and are numbered from 0 to 99. These specific tag IDs were entered under the Recov_Tag_ID column.
- Recov_Year, Recov_Month, and Recov_Day are the date that the individual Gulf Menhaden tags were recovered from the reduction plant.
- Recov_Plant is denoted by a number that has been assigned to each Gulf Menhaden reduction plant. This is the plant that individual fish were processed.

We evaluated this data to understand the magnitude of tag and recaptured individuals, by life-stage (Table 1, 2) and also the characteristics of tagged and recaptured fish.

2. Magnet Efficiency Trials

From Liljestrand Master's Thesis: "Magnet efficiency experiments were conducted to test the efficiency of magnets for collecting tags by introducing batches of tagged Menhaden (usually 100 fish) directly into each reduction plant.". These data were conducted by the original experimenters and used by Liljestrand et al. to determine the probability that a tagged fish would be observed, if it was collected.

The magnet efficiency data are comprised of the following variables:

- Test_Year, Test_Month, and Test_Day indicate the date on which the plant test was performed, meaning the date when the 100 random Gulf Menhaden were tagged to test accuracy of magnets.
- Tag_Series_Number indicates which group in the tag series with which the Gulf Menhaden were tagged. For each group of tags, with a tag series and number, (example: U20) has 100 individual tags and are numbered from 0 to 99. These specific tag IDs were entered under the Recov_Tag_ID column.
- Recov_Year, Recov_Month, and Recov_Day pertain to the date on which the individual Gulf Menhaden tags were recovered from the test plant.

- Plant is denoted by a number that has been assigned to each Gulf Menhaden reduction plant. This is the plant where the tests were performed.

3. The tag and recapture histories of adults with fork-length

A small and unique third source of data in the archived data was a subset of recapture data that included fork length of fish, upon collection by the plant. The tag and recapture component of the data was integrated into the tag and recapture histories of adult individuals.

The utility of these data are limited for analysis of growth or to understand how catchability is influenced by the length of the fish. However, they are useful for indicating what the experimenters considered an “adult” fish length to be.

The variables associated with the tag and recapture histories of adults with fork-length

All of the variables are the same as the tag and recapture histories of adults, but include an additional column which is the fork length in FL (mm).

Evaluation the utility of the data

Magnitude of mark and recapture data

In comparison to the magnitude of tagged and recaptured individuals used by Liljestrand et al., the data we have synthesized has fewer number of recaptures (Fig. 3, 4). However, the overall rate of recapture is generally high (Table 1 and 2) - fewer fish were tagged in the Gulf of Mexico, than in the mid-Atlantic. One difference in the modeling approach we propose to use that contrasts that of Liljestrand et al. is that we will not estimate movement parameters - we don't have geographic movement parameters in these data. Omitting these parameters in the modeling effort should improve the ability of the model to derive precise estimates of mortality.

Having positive rates of recapture (~3%) is a necessary characteristic of using mark and recapture models. Another aspect of the tag-recapture work, that is necessary for using contemporary models for parameter estimation, is to have recapture occurrences at many sampling efforts. Figure 2 indicates that both life stages have times ‘at large’ of less than and greater than 500 days.

Evaluation of observation error

Mark recapture modeling attempts to determine (at least) two related parameters - survival and probability of recapture. The second parameter (probability of recapture) is in part determined by observation error. The plant- and year-specific capture probability derived from the trial data, different plants have varying ability for tag detection. These modeled parameters, derived from the data, can be incorporated into the model, as Liljestrand et al. has done.

In conclusion, the magnitude of the data (number of recaptures, magnet trials) and its structure (varying times ‘at large’, stage specific tagging) are likely appropriate for modeling. I would recommend that this be attempted.

Table 1. Summary of adult tag-recapture data from scanned and cataloged data.

Year	Adult Tagged	Adult Recaptured	Adult % Recaptured
1969	27896	3439	12.3%
1970	-	-	-
1971	-	-	-
1972	-	-	-
1973	-	-	-
1974	-	-	-
1976	12001	503	4.19%
1977	8102	325	4.01%
1978	8802	461	5.24%
1979	2089	9	0.43%
1980	6301	242	3.84%
1981	6500	166	2.55%
1982	13201	583	4.42%
1983	8900	299	3.36%
1984	8500	479	5.64%
1985	15500	766	4.94%

Table 2. Summary of juvenile tag-recapture data from scanned and cataloged data.

Year	Juvenile Tagged	Juvenile Recaptured	Juvenile % Recaptured
1969	-	-	-
1970	10313	250	2.42%
1971	6700	119	1.78%
1972	15700	500	3.18%
1973	4400	26	0.59%
1974	10900	157	1.44%
1976	7700	472	6.13%
1977	15299	573	3.75%
1978	9900	200	2.02%
1979	2000	4	0.20%
1980	10200	498	4.88%
1981	11300	345	3.05%
1982	8500	295	3.47%
1983	7900	411	5.20%
1984	13300	536	4.03%
1985	8200	199	2.43%

Table 3. Summary of juvenile tag-recapture data with fork length information.

Tag_Month	Number of Recaptured Fish	Mimimum Length	Maximum Length
4	2545	133	397
5	811	145	338
9	83	196	312




Figure 1: Plant- and year-specific proportion of capture from tests of known number of fish ($n = 100$). Plant number refers to internal designation of an individual plant. The shaded density regions are the aggregated (all year) density for each plant (row) and provided for understanding the expected probability that a tagged fish collected at that plant would be observed upon recapture.)




Figure 2: Frequency histogram of the number of fish, by lifestage, that were tagged and recaptured as a function of the number of days ‘at large’, the time between tagging and recapture.




Figure 3: Time series of the number of Gulf Menhaden, by lifestage, recaptured in the Gulf of Mexico tagging effort, the date of the recaptured fish are binned in each month of the study.




Fig. 2. Predicted (line) and observed (circles) number of recovered tags for each month after June 1966 in by regions. The predicted values are the median of the posterior probability distribution and the error bars indicate 95% credibility intervals.

Figure 4: Figure reprinted from Liljestrand et al. (2019) provided for comparison to the number of Gulf Menhaden, by lifestage, recaptured in the Gulf of Mexico tagging effort.