

These Notes Are Works-In-Progress

Bayesian Statistics for Mathematicians
or, A Measure-Theoretic Approach to Bayesian Statistics

Raymond T. Melton

May 19, 2024

Contents

Preface	iv
1 Introduction	1
2 Measure Spaces and Measurable Sets	3
3 Measurable Functions and Integration	9
4 Function Spaces and Quotient Spaces	22
5 Product Measures and Product Integration	39
5.1 Product Measures	39
5.2 Product Integration	42
5.3 Applications	45
6 Probability, Independence, and Sampling	55
6.1 Probability Spaces and Distributions	55
6.2 Expected Value	58
6.3 Independence	59
6.4 Strong Law of Large Numbers	60
6.5 Sampling	61
7 Conditional Mathematical Expectation	64

7.1	Properties of Expectation Operators	68
7.2	A Monotone Convergence Theorem	71
7.3	Atoms and Independence	72
7.4	Measurable Functions Factor Out	79
8	Conditional Probability and Markov Kernels	83
8.1	Conditional Probability	83
8.2	Markov Kernels	84
8.3	Regular Version	88
8.4	Conditional Distributions	93
8.5	Conditional Densities	104
9	Bayesian Statistics	112
10	Bayesian Statistical Models	118
10.1	Outline	118
10.2	Globe Tossing Model	118
10.3	Gaussian Model of Height	122
10.4	The Linear Model	126
10.5	Logistic Regression	129
11	Conditional Independence	133
12	Expectation Operators as Projections	141
12.1	Expectation as Projection	142
13	Acknowledgements	145
14	Afterword	146

Preface

THESE notes were not initially intended for anyone else to look at. I am working to make them easier on the eyes.
Feel free to contact me. I am willing to share ideas.

1 Introduction

*Many readers will have noticed
the recent trend toward
quotations at the beginnings of
chapters in scientific books.
Often these quotes are sappy,
dippy little things as if the
authors of the book were
struggling for profundity.*

Heterochrony: The Evolution of
Ontogeny
— M. L. MCKINNEY

THIS is not original material. The underlying reason these notes exist is so I could track the progress I had made while learning statistics. I am putting these notes online, thinking they may help another mathematician trying to learn statistics.

In these notes, integration theory is based upon semi-rings as opposed to σ -algebras, to be explained further in the next few paragraphs. Also,

- We do not confuse equivalence classes of functions with the functions themselves.
- We do not make topological assumptions when it comes to regular conditional distributions.
- We make clear which σ -algebras are involved in statistical models.

In the statistics books that I had read, if the author tried to make clear which integral they were using, it was always the Lebesgue integral defined in terms of a measure on a σ -algebra. But the integration theory I had learned was based upon a measure on a semi-ring. What

is the difference?

One main difference: In the case of a measure on a σ -algebra and the consequent Lebesgue integral, say as developed in *Probability Theory*, Loeve [5], the resulting conditional mathematical expectation is a class of functions, where each function in the class is measurable with respect to a σ -subalgebra, but in the case of a measure on a semi-ring, as developed in *Principles of Real Analysis*, Aliprantis & Burkinshaw [2], the consequent conditional mathematical expectation is a class of functions from which we can select a representative function which is measurable with respect to the σ -subalgebra. Not a big difference, but you still need to make the selection. Then again, you always need to make a selection with a regular conditional probability. Sorry about the pun.

Statistical models only started making sense after learning Bayesian statistical models. That is all you will find here. All of the material regarding frequentist statistics has been edited out, like when the cocoon is left behind. For mathematicians, this material is more of a measure-theoretic companion to something comprehensive and explanatory on applied Bayesian statistical models. I strongly recommend Richard McElreath's *Statistical Rethinking* [6].

One more thing. A quote from *Thinking Like Your Editor*, by Rabiner & Fortunato, 2002, where the question is raised, and I really hope I am paraphrasing its content correctly here: Once you've defined your audience, to which level of sophistication do you write?

The answer is that if you want to write serious nonfiction, and if you want your work to be taken seriously, you never dumb down. You always write up. Writing down runs the risk of having your book read like a young adult book. Assume a high reader level of intellectual sophistication, even though you suspect their knowledge of your particular subject may not be high. Fully explain, but do not simplify.

This is not a children's reader. No attempt has been made to dumb down anything.

2 Measure Spaces and Measurable Sets

THIS is an abbreviated review of measure theory, based upon semirings. Roughly, a measure on a semiring is extended to the power-set, and then restricted to the measurable subsets. Resulting measure spaces are complete in the sense that every subset of a set of measure zero is Carathéodory measurable. That is, every null set is measurable. It remains to be seen whether this particular notion of completeness will be sufficient.

Any missing detail might be found in Aliprantis and Burkinshaw, *Principles of Real Analysis* [2]. Except Proposition 2.23. That was found in Aliprantis and Border's book, *Infinite Dimensional Analysis: A Hitchhiker's Guide*, Third Edition, 2006; see 10.31 Lemma therein, page 386.

Definition 2.1. Let X be a nonempty set. A collection $\mathfrak{S}$ of subsets of X is a **semiring** if it has the following three properties:

1. $\emptyset \in \mathfrak{S}$.
2. If A and B are in $\mathfrak{S}$, then $A \cap B \in \mathfrak{S}$.
3. If A and B in $\mathfrak{S}$, there exist disjoint sets $C_1, \dots, C_n$ in $\mathfrak{S}$ such that $A \setminus B = \bigcup_{i=1}^n C_i$.

Definition 2.2. Let $\mathfrak{S}$ be a semiring of subsets of a set X . A set function $\mu : \mathfrak{S} \rightarrow [0, \infty]$ is a **measure** if it has both of the following properties:

1. $\mu(\emptyset) = 0$.
2. μ is σ -additive.

{measure_space}

Definition 2.3. A triple $(X, \mathfrak{S}, \mu)$, where X is a nonempty set, the collection $\mathfrak{S}$ is a semiring of subsets of X , and μ is a measure, is called a **measure space**.

{algebra}

Definition 2.4. A nonempty collection $\mathfrak{A}$ of subsets of a nonempty set X is an **algebra** if it is closed with respect to finite intersections and complements; that is, if it has both the following properties:

1. If A and B are in $\mathfrak{A}$, then $A \cap B \in \mathfrak{A}$.
2. If $A \in \mathfrak{A}$, then $A^c \in \mathfrak{A}$.

Consequently, both $\emptyset \in \mathfrak{A}$ and $X \in \mathfrak{A}$.

A concept in between those of semirings and algebras of sets is that of rings of sets. A nonempty collection of subsets of a set is a **ring** if it is closed with respect to set differences and finite unions. Every algebra of sets is also a ring of sets. Every ring of sets is also a semiring of sets.

Definition 2.5. An algebra of subsets of a set is a σ -**algebra** if it is closed with respect to countable unions, in which case it must also be closed with respect to countable intersections. If $\mathfrak{A}$ is a σ -algebra and $\mathfrak{B} \subseteq \mathfrak{A}$, then $\mathfrak{B}$ is a σ -**subalgebra** of $\mathfrak{A}$ if $\mathfrak{B}$ is a σ -algebra. If $\mathfrak{F}$ is a collection of subsets of a nonempty set, then $\mathfrak{F}$ is included in a smallest σ -algebra, denoted $\sigma(\mathfrak{F})$. It is the intersection of all σ -algebras containing $\mathfrak{F}$, and is called the σ -**algebra generated** by $\mathfrak{F}$.

Definition 2.6. Let $\mathfrak{P}(X)$ be the power set of a set X . A set function $\mu : \mathfrak{P}(X) \rightarrow [0, \infty]$ is an **outer measure** if it has the following three properties:

1. $\mu(\emptyset) = 0$.
2. μ is monotone. That is, if A and B are in $\mathfrak{P}(X)$ with $A \subseteq B$, then $\mu(A) \leq \mu(B)$.
3. μ is σ -subadditive. That is, $\mu(\bigcup_{n=1}^{\infty} A_n) \leq \sum_{n=1}^{\infty} \mu(A_n)$ for every sequence $\{A_n\}$ of subsets of X .

{caratheodory}

Definition 2.7 (Carathéodory). Let μ be an outer measure on the powerset $\mathfrak{P}(X)$ of some set X . A subset B of X is a μ -**measurable set** if

$$\mu(A) = \mu(A \cap B) + \mu(A \cap B^c)$$

for all $A \subseteq X$. Let $\mu\text{-}\mathfrak{M}_{\text{eas}}$ denote the collection of all μ -measurable subsets of X .

Remark 2.8. Measurable catch phrases:

- A set is μ -measurable if it sharply cuts every set.
- A set is μ -measurable if you can always make change with it.

{outer_measure_generated}

Definition 2.9. Given a measure space $(X, \mathfrak{S}, \mu)$, define a set function $\mu^* : \mathfrak{P}(X) \rightarrow [0, \infty]$ by

$$\mu^*(A) = \inf \left\{ \sum_{n=1}^{\infty} \mu(A_n) : \{A_n\} \text{ is a sequence of } \mathfrak{S} \text{ with } A \subseteq \bigcup_{n=1}^{\infty} A_n \right\},$$

where $\inf \emptyset = \infty$. The set function μ^* is the **outer measure generated** by μ .

It will turn out that μ^* , when restricted to the collection of μ^* -measurable sets, is an honest-to-God measure. We might also denote this restriction by μ^* for now. If this reduces clarity at any point, then we will introduce some fairly cumbersome notation, like $\mu|_{\mu\text{-}\mathfrak{M}\text{eas}}$, or $\mu^*|_{\mu^*\text{-}\mathfrak{M}\text{eas}}$.

Proposition 2.10. *The outer measure generated by a measure really is an outer measure.*

{extension_from_semiring}

Proposition 2.11. *Let $(X, \mathfrak{S}, \mu)$ be a measure space. The outer measure μ^* generated by μ is an extension of μ from $\mathfrak{S}$ to the powerset $\mathfrak{P}(X)$ of X . That is, if $A \in \mathfrak{S}$, then $\mu^*(A) = \mu(A)$. Equivalently, the restriction $\mu^*|_{\mathfrak{S}}$ of μ^* to the semiring $\mathfrak{S}$ is equal to μ .*

Notation 2.12. Let μ be an outer measure on the powerset $\mathfrak{P}(X)$ of some set X . A subset A of X is called a μ -**null set** if $\mu(A) = 0$. Any definition or relation on X that holds except possibly on a μ -null set is said to hold μ -**almost everywhere**, or simply μ -a.e. We might also say that the relation holds for μ -**almost all** $x \in X$. For example, if $f : X \rightarrow \mathbb{R}$ and $g : X \rightarrow \mathbb{R}$, and $f(x) = g(x)$ except for $x \in A$ with $\mu(A) = 0$, then we could say that $f = g$ for μ -almost all x in X .

Proposition 2.13. *Let μ be an outer measure on the powerset $\mathfrak{P}(X)$ of some set X . Every μ -null set is μ -measurable. Consequently, since μ is monotone, every subset of a μ -null set is μ -measurable.*

{restricted_outer_measures}

Proposition 2.14. *If μ is an outer measure on the powerset $\mathfrak{P}(X)$ of a set X , then the collection $\mu\text{-}\mathfrak{M}\text{eas}$ of all μ -measurable subsets of X is a σ -algebra, and the triple $(X, \mu\text{-}\mathfrak{M}\text{eas}, \mu|_{\mu\text{-}\mathfrak{M}\text{eas}})$ is a measure space. That is, the outer measure μ , when restricted to the σ -algebra $\mu\text{-}\mathfrak{M}\text{eas}$ of all μ -measurable subsets of X , is σ -additive.*

{subtractive}

Corollary 2.15. *Let $(X, \mathfrak{S}, \mu)$ be a measure space. Let A and B be μ^* -measurable sets with $B \subseteq A$ and with $\mu^*(A) < \infty$. Then $\mu^*(A \cap B^c) = \mu^*(A) - \mu^*(B)$.*

{semiring_measurable}

Proposition 2.16. *If $(X, \mathfrak{S}, \mu)$ is a measure space, then every element of the semiring $\mathfrak{S}$ is a μ^* -measurable subset. That is, $\mathfrak{S} \subseteq \mu^*\text{-}\mathfrak{M}\text{eas}$.*

{sigma_finite}

Definition 2.17. A measure space $(X, \mathfrak{S}, \mu)$ is **finite** if $\mu^*(X) < \infty$. We may also say that the measure μ is finite. A measure space $(X, \mathfrak{S}, \mu)$ is **σ -finite** if there is a disjoint sequence $\{X_n\}$ of subsets of $\mathfrak{S}$ with $\mu(X_n) < \infty$ and $X \subseteq \bigcup_n X_n$. We may also say that measure μ is σ -finite.

Proposition 2.18. *Let $(X, \mathfrak{S}, \mu)$ be a finite measure space. A subset $E \subseteq X$ is μ^* -measurable if and only if*

$$\mu^*(X) = \mu^*(E) + \mu^*(E^c).$$

Proposition 2.19. *Let $(X, \mathfrak{S}, \mu)$ be a measure space, and let $E \in \mu^*\text{-}\mathfrak{M}\text{eas}$. Define a collection of sets, the restriction of $\mathfrak{S}$ to E , by*

$$\mathfrak{S}_E = \{E \cap A : A \in \mathfrak{S}\},$$

and define a function $\nu : \mathfrak{S}_E \rightarrow [0, \infty]$ by $\nu(E \cap A) = \mu^(E \cap A)$. Then the triple $(X, \mathfrak{S}_E, \nu)$ is a measure space, and $\nu^* = \mu^*|_{\mathfrak{P}(E)}$. Furthermore, $\nu^*\text{-}\mathfrak{M}\text{eas} = \{F \subseteq E : F \in \mu^*\text{-}\mathfrak{M}\text{eas}\}$.*

{sufficient_for_measurability}

Proposition 2.20. *Let $(X, \mathfrak{S}, \mu)$ be a measure space, and let $A \subseteq X$. If there is an $E \in \mu^*\text{-}\mathfrak{M}\text{eas}$ with $A \subseteq E$ such that $\mu^*(E) < \infty$ and $\mu^*(E) = \mu^*(A) + \mu^*(E \cap A^c)$, then $A \in \mu^*\text{-}\mathfrak{M}\text{eas}$.*

{extension_measure}

Proposition 2.21. *Let $(X, \mathfrak{S}, \mu)$ be a σ -finite measure space, let $\mathfrak{T}$ be a semiring of subsets of X with $\mathfrak{S} \subseteq \mathfrak{T} \subseteq \mu^*\text{-}\mathfrak{Meas}$, and let $\nu : \mathfrak{T} \rightarrow [0, \infty]$ be a measure on $\mathfrak{T}$. If $\nu = \mu$ on $\mathfrak{S}$, then $\nu = \mu^*$ on $\mathfrak{T}$. Consequently, the measure $\mu^* : \mu^*\text{-}\mathfrak{Meas} \rightarrow [0, \infty]$ is the unique extension of the measure μ from the semiring $\mathfrak{S}$ to the σ -algebra $\mu^*\text{-}\mathfrak{Meas}$.*

$$\begin{array}{ccc} \mu^*\text{-}\mathfrak{Meas} & \xrightarrow{\mu^*} & [0, \infty] \\ \cup & & \\ \mathfrak{T} & \xrightarrow{\nu} & [0, \infty] \\ \cup & & \\ \mathfrak{S} & \xrightarrow{\mu} & [0, \infty] \end{array}$$

Proposition 2.22. *Let $(X, \mathfrak{S}, \mu)$ be a measure space. If $A \subseteq X$, then there is a superset $B \in \sigma(\mathfrak{S})$ of A with $\mu^*(A) = \mu^*(B)$.*

{approximation_sigma_algebra}

Proof. Let $A \subseteq X$. We first look at the case where the outer measure of A is infinite; let $\mu^*(A) = \infty$. We can take $B = X$; we know $X \in \sigma(\mathfrak{S})$ because $\emptyset \in \mathfrak{S}$ and algebras (therefore σ -algebras) are closed with respect to complements. And outer measures are monotone, so $\mu^*(A) \leq \mu^*(X)$. But this implies that $\mu^*(X) = \infty$, so $\mu^*(A) = \mu^*(X)$, completing the proof of this case.

We now look at the complementary case; let $\mu^*(A)$ be finite. For each $k \in \{1, 2, \dots\}$ there must be, according to the definition of $\mu^*(A)$, a sequence $\{A_1^k, A_2^k, \dots\}$ of $\mathfrak{S}$ whose union covers A , where

$$\mu^*(A) \leq \sum_{n=1}^{\infty} \mu(A_n^k) < \mu^*(A) + 1/k.$$

In this case, we can take $B = \bigcap_{k=1}^{\infty} \left(\bigcup_{n=1}^{\infty} A_n^k \right)$, which is in $\sigma(\mathfrak{S})$. Because $A \subseteq \bigcup_{n=1}^{\infty} A_n^k$ for each k , it follows that $A \subseteq B$. Also,

$$\mu^*(A) \leq \mu^*(B) \leq \mu^* \left(\bigcup_{n=1}^{\infty} A_n^k \right) \leq \sum_{n=1}^{\infty} \mu^*(A_n^k) = \sum_{n=1}^{\infty} \mu(A_n^k) < \mu^*(A) + 1/k.$$

2 Measure Spaces and Measurable Sets

Since this holds for each $k \in \{1, 2, \dots\}$, it follows that $\mu^*(A) = \mu^*(B)$, as required. ■

{approximation_sigma_finite}

Proposition 2.23. *Let $(X, \mathfrak{S}, \mu)$ be a σ -finite measure space. If $A \in \mu^*\text{-}\mathcal{M}\text{eas}$, then there is a subset $C \subseteq X$ with $A \cap C = \emptyset$ and such that $\mu^*(C) = 0$ with $A \cup C \in \sigma(\mathfrak{S})$.*

Proof. Let $A \in \mu^*\text{-}\mathcal{M}\text{eas}$. The measure space $(X, \mathfrak{S}, \mu)$ is supposed to be σ -finite, so there is a sequence $\{X_n\}$ of subsets of $\mathfrak{S}$ with $\mu(X_n) < \infty$ and $X \subseteq \bigcup_n X_n$. By Proposition 2.22, for each subset $A \cap X_n$ there is a subset $B_n \in \sigma(\mathfrak{S})$ with $A \cap X_n \subseteq B_n$ and $\mu^*(A \cap X_n) = \mu^*(B_n)$. We take $C = \left(\bigcup_n B_n\right) \cap A^c$ and show that C has the required properties.

Pure set theory shows that $C \subseteq \bigcup_n (B_n \cap (A \cap X_n)^c)$. Since $\mu(X_n) = \mu^*(X_n) < \infty$, it follows that $\mu^*(A \cap X_n) = \mu^*(B_n) < \infty$ and so we can apply Corollary 2.15 which says $\mu^*(B_n \cap (A \cap X_n)^c) = \mu^*(B_n)\mu^*(A \cap X_n)$. But $\mu^*(B_n) = \mu^*(A \cap X_n)$. This means $\mu^*(B_n \cap (A \cap X_n)^c) = 0$ which implies that $\mu^*\left(\bigcup_n (B_n \cap (A \cap X_n)^c)\right) = 0$. By monotonicity, $\mu^*(C) = 0$.

The fact that $C \subseteq A^c$ implies that $A \cap C = \emptyset$. Finally, to see that $A \cup C \in \sigma(\mathfrak{S})$, since we already know that each B_n is in $\sigma(\mathfrak{S})$, we simply need to show that $A \cup C = \bigcup_n B_n$. Again, this is pure set theory. Briefly, since $A \subseteq \bigcup_n (A \cap X_n) \subseteq \bigcup_n B_n$, it follows that $A \cup \left(\bigcup_n B_n\right) = \bigcup_n B_n$, and so

$$\begin{aligned} A \cup C &= A \cup \left(\left(\bigcup_n B_n \right) \cap A^c \right) \\ &= \left(A \cup \bigcup_n B_n \right) \cap (A \cup A^c) \quad \text{just distribute} \\ &= A \cup \bigcup_n B_n \\ &= \bigcup_n B_n, \end{aligned}$$

as required. ■

3 Measurable Functions and Integration

THROUGHOUT this section, let $(X, \mathfrak{S}, \mu)$ denote a measure space, so $\mathfrak{S}$ is at least a semiring, and we continue to let $\mu^*\text{-}\mathcal{M}\text{eas}$ denote the σ -algebra of μ^* -measurable subsets of X .

Notation 3.1. Let X and Y be sets, and let $\mathfrak{P}(X)$ and $\mathfrak{P}(Y)$ denote their respective powersets. For any function $f : X \rightarrow Y$, the set X is the **domain** of f and the set Y is the **codomain**. For any subset $A \subseteq X$, the set

$$f(A) := \{f(x) : x \in A\}$$

is the **image** of A under f , which we could denote $f^\rightarrow(A)$. The subset $f(X) \subseteq Y$ is the **range** of f . And for any subset $B \subseteq Y$, the set

$$f^\leftarrow(B) := \{x \in X : f(x) \in B\}$$

is the **preimage** of B under f . We might just as well have said let $f^\leftarrow$ denote the image of f under a contravariant powerset functor. This means that

$$f^\leftarrow : \mathfrak{P}(Y) \rightarrow \mathfrak{P}(X) : B \mapsto f^\leftarrow(B).$$

The set $f^\leftarrow(B)$ might also be denoted $f^{-1}(B)$, but not here. Any restriction of such a $f^\leftarrow$ which maps some subset of $\mathfrak{P}(Y)$ to some subset of $\mathfrak{P}(X)$ may still be denoted $f^\leftarrow$, in which case the domain and codomain of $f^\leftarrow$ will be made explicit.

For any topological space X , we will let $\mathfrak{Bor}_X$ denote the σ -algebra of subsets of X generated by the topology, namely the **Borel subsets** of the topological space X .

Definition 3.2. A pair $(X, \mathfrak{A})$, where X is a set and $\mathfrak{A}$ is a σ -algebra of subsets of X , is called a **measurable space**. Better terminology might be something flavored Mackey.

{measurable_space}

Definition 3.3. Let $(X, \mathfrak{A})$ and $(Y, \mathfrak{B})$ be measurable spaces. A function $f : X \rightarrow Y$ is called an $(\mathfrak{A}, \mathfrak{B})$ -**measurable function** if a restriction of $f^{\leftarrow}$ to $\mathfrak{B}$ is a function $\mathfrak{B} \rightarrow \mathfrak{A}$; that is, if $f^{\leftarrow}(B) \in \mathfrak{A}$ for each $B \in \mathfrak{B}$. To indicate that $f : X \rightarrow Y$ is $(\mathfrak{A}, \mathfrak{B})$ -measurable, we might use the notation $f : (X, \mathfrak{A}) \rightarrow (Y, \mathfrak{B})$.

{measurable_function}

Remark 3.4. An attempt at a categorical approach to measurable spaces and measurable functions which does not take into account collections of null sets would be considered a first attempt. There is nothing wrong with first attempts. Who knows? You might end up factoring by an ideal.

{function_induces_measure}

Proposition 3.5. Let $(X, \mathfrak{A})$ and $(Y, \mathfrak{B})$ be a measurable spaces. If μ is a measure on $\mathfrak{A}$, and a function $f : X \rightarrow Y$ is $(\mathfrak{A}, \mathfrak{B})$ -measurable, where $f^{\leftarrow} : \mathfrak{B} \rightarrow \mathfrak{A}$, then the set function

$$\mu \circ f^{\leftarrow} : \mathfrak{B} \rightarrow [0, \infty] : B \mapsto \mu(f^{\leftarrow}(B))$$

defines a measure on $\mathfrak{B}$.

{induced_measure}

Definition 3.6. The measure $\mu \circ f^{\leftarrow}$ on the σ -algebra of the codomain of f in Proposition 3.5 is the **measure induced by f** .

Remark 3.7. Should two $(\mathfrak{A}, \mathfrak{B})$ -measurable functions f and g be equal μ^* -almost everywhere, then the induced σ -algebras $f^{\leftarrow}(\mathfrak{B})$ and $g^{\leftarrow}(\mathfrak{B})$ need not be the same. However, the following proposition says that the induced measures $\mu^* \circ f^{\leftarrow}$ and $\mu^* \circ g^{\leftarrow}$ are the same.

{class_induces_measure}

Proposition 3.8. Let $(X, \mathfrak{S}, \mu)$ be a measure space, and let $(Y, \mathfrak{B})$ be a measurable space. If the functions $f : X \rightarrow Y$ and $g : X \rightarrow Y$ are $(\mu^*\text{-}\mathfrak{Meas}, \mathfrak{B})$ -measurable functions with $f = g$ μ^* -almost everywhere on X , then $\mu^*(A \cap f^{\leftarrow}(B)) = \mu^*(A \cap g^{\leftarrow}(B))$ for all $A \in \mu^*\text{-}\mathfrak{Meas}$ and for all $B \in \mathfrak{B}$. So in particular, $\mu^*(f^{\leftarrow}(B)) = \mu^*(g^{\leftarrow}(B))$ for all $B \in \mathfrak{B}$.

Proof. Define $M = \{x \in X : f(x) = g(x)\}$, so that by hypothesis, $\mu^*(M^c) = 0$. Let $A \in \mu^*\text{-}\mathfrak{Meas}$ and $B \in \mathfrak{B}$. If $x \in f^{\leftarrow}(B)$ and $x \notin g^{\leftarrow}(B)$,

then $f(x) \in B$ and $g(x) \in B^c$, implying $f(x) \neq g(x)$ and $x \notin M$, meaning

$$f^{\leftarrow}(B) \cap (g^{\leftarrow}(B))^c \subseteq M^c.$$

This says

$$\mu^*(f^{\leftarrow}(B) \cap (g^{\leftarrow}(B))^c) = 0$$

since $\mu^*(M^c) = 0$. Because $f^{\leftarrow}(B)$ is a disjoint union, as in

$$f^{\leftarrow}(B) = (f^{\leftarrow}(B) \cap g^{\leftarrow}(B)) \sqcup (f^{\leftarrow}(B) \cap (g^{\leftarrow}(B))^c),$$

we can also write $A \cap f^{\leftarrow}(B)$ as a disjoint union:

$$A \cap f^{\leftarrow}(B) = (A \cap (f^{\leftarrow}(B) \cap g^{\leftarrow}(B))) \sqcup (A \cap (f^{\leftarrow}(B) \cap (g^{\leftarrow}(B))^c)).$$

Because $A \cap f^{\leftarrow}(B)$ is μ^* -measurable, it follows by the very definition of what it means to be measurable that

$$\mu^*(A \cap f^{\leftarrow}(B)) = \mu^*(A \cap f^{\leftarrow}(B) \cap g^{\leftarrow}(B)) + \mu^*(A \cap f^{\leftarrow}(B) \cap (g^{\leftarrow}(B))^c).$$

But the fact that $\mu^*(f^{\leftarrow}(B) \cap (g^{\leftarrow}(B))^c) = 0$ implies that $\mu^*(A \cap f^{\leftarrow}(B) \cap (g^{\leftarrow}(B))^c) = 0$, and so

$$\mu^*(A \cap f^{\leftarrow}(B)) = \mu^*(A \cap f^{\leftarrow}(B) \cap g^{\leftarrow}(B)).$$

By a symmetric argument,

$$\mu^*(A \cap g^{\leftarrow}(B)) = \mu^*(A \cap f^{\leftarrow}(B) \cap g^{\leftarrow}(B)).$$

Therefore by transitivity of equality,

$$\mu^*(A \cap f^{\leftarrow}(B)) = \mu^*(A \cap g^{\leftarrow}(B)),$$

as required. ■

Remark 3.9. The following proposition will be used to extend the notion of independence of $\mathcal{L}_p(X, \mathfrak{S}, \mu)$ functions to independence of $L_p(X, \mathfrak{S}, \mu)$ classes, which will appear in Definition 6.12.

Proposition 3.10. *Let $(X, \mathfrak{S}, \mu)$ be a measure space, and let $(Y, \mathfrak{B})$ be a measurable space, and let $f, g, h, k : X \rightarrow Y$ be $(\mu^*\text{-}\mathfrak{M}\text{eas}, \mathfrak{B})$ -measurable functions such that $f = h$ μ^* -almost everywhere on X and $g = k$ μ^* -almost everywhere on X .*

{class_induced_measure}

If $\mu^(A \cap B) = \mu^*(A)\mu^*(B)$ for all $A \in f^{\leftarrow}(\mathfrak{B})$ and all $B \in g^{\leftarrow}(\mathfrak{B})$, then $\mu^*(C \cap D) = \mu^*(C)\mu^*(D)$ for all $C \in h^{\leftarrow}(\mathfrak{B})$ and all $D \in k^{\leftarrow}(\mathfrak{B})$.*

Proof. Let $E, F \in \mathfrak{B}$. Since $f = h$ μ^* -almost everywhere, by Proposition 3.8,

$$\mu^*(f^{\leftarrow}(E) \cap k^{\leftarrow}(F)) = \mu^*(h^{\leftarrow}(E) \cap k^{\leftarrow}(F))$$

and

$$\mu^*(f^{\leftarrow}(E)) = \mu^*(h^{\leftarrow}(E)).$$

Since $g = k$ μ^* -almost everywhere, again by Proposition 3.8,

$$\mu^*(f^{\leftarrow}(E) \cap k^{\leftarrow}(F)) = \mu^*(f^{\leftarrow}(E) \cap g^{\leftarrow}(F))$$

and

$$\mu^*(k^{\leftarrow}(F)) = \mu^*(g^{\leftarrow}(F)).$$

By hypothesis,

$$\mu^*(f^{\leftarrow}(E) \cap g^{\leftarrow}(F)) = \mu^*(f^{\leftarrow}(E))\mu^*(g^{\leftarrow}(F)),$$

and so by transitivity of equality,

$$\mu^*(h^{\leftarrow}(E) \cap k^{\leftarrow}(F)) = \mu^*(h^{\leftarrow}(E))\mu^*(k^{\leftarrow}(F)),$$

as required. ■

{preimage_of_a_null_set}

Proposition 3.11. *Let $(X, \mathfrak{A}, \mu)$ be a measure space, where $\mathfrak{A}$ is a σ -algebra, and let $(Y, \mathfrak{B})$ be a measurable space. Let the function $\phi : X \rightarrow Y$ be $(\mathfrak{A}, \mathfrak{B})$ -measurable. If a subset B of Y is a $(\mu \circ \phi^{\leftarrow})^*$ -null set, then the subset $\phi^{\leftarrow}(B)$ of X is a μ^* -null set.*

Proof. Let B be a subset of Y which is a $(\mu \circ \phi^{\leftarrow})^*$ -null set. Since $\mu \circ \phi^{\leftarrow}$ is a measure, by Proposition 3.5, it follows that the triple $(Y, \mathfrak{B}, \mu \circ \phi^{\leftarrow})$ is a measure space. So by Proposition 2.22, we can take a superset C of B with C in the σ -algebra $\mathfrak{B}$ and having the same outer measure: $(\mu \circ \phi^{\leftarrow})^*(C) = (\mu \circ \phi^{\leftarrow})^*(B) = 0$. Because the outer measure agrees with the measure on the generating semiring (σ -algebra in this case), and because ϕ is assumed measurable, meaning $\phi^{\leftarrow}(C) \in \mathfrak{A}$, the following equalities hold:

$$\begin{aligned} (\mu \circ \phi^{\leftarrow})^*(C) &= (\mu \circ \phi^{\leftarrow})(C) & C \in \mathfrak{B}, \\ &= \mu(\phi^{\leftarrow}(C)) & \text{definition of induced measure,} \\ &= \mu^*(\phi^{\leftarrow}(C)) & \phi^{\leftarrow}(C) \in \mathfrak{A}. \end{aligned}$$

With $B \subseteq C$, it follows that $\phi^{\leftarrow}(B) \subseteq \phi^{\leftarrow}(C)$ and since outer measure is monotone, $\mu^*(\phi^{\leftarrow}(B)) \leq \mu^*(\phi^{\leftarrow}(C))$. The comparison $0 \leq \mu^*(\phi^{\leftarrow}(B)) \leq \mu^*(\phi^{\leftarrow}(C)) = 0$ then shows that $\phi^{\leftarrow}(B)$ is a μ^* -null subset of X , as required. ■

{preimage_of_a_measurable_set}

Proposition 3.12. *Let $(X, \mathfrak{A}, \mu)$ be a measure space, where $\mathfrak{A}$ is a σ -algebra, and let $(Y, \mathfrak{B})$ be a measurable space. Let the function $\phi : X \rightarrow Y$ be $(\mathfrak{A}, \mathfrak{B})$ -measurable, and let B be a $(\mu \circ \phi^{\leftarrow})^*$ -measurable subset of Y with $(\mu \circ \phi^{\leftarrow})^*(B) < \infty$. Then $\phi^{\leftarrow}(B)$ is a μ^* -measurable subset of X , and $\mu^*(\phi^{\leftarrow}(B)) = (\mu \circ \phi^{\leftarrow})^*(B)$.*

Proof. As in the proof of Proposition 3.11, and according to Proposition 2.22, we can take a superset C of B with $C \in \mathfrak{B}$ and having the same outer measure:

$$(\mu \circ \phi^{\leftarrow})^*(C) = (\mu \circ \phi^{\leftarrow})^*(B).$$

With $C \in \mathfrak{B}$, it follows that C is $(\mu \circ \phi^{\leftarrow})^*$ -measurable by Proposition 2.16. Since B is a subset of C , we can write C as the disjoint union

$$C = B \sqcup (C \cap B^c).$$

3 Measurable Functions and Integration

Because both C (by construction) and B (by hypothesis) are $(\mu \circ \phi^\leftarrow)^*$ -measurable, it follows that their set difference $C \cap B^c$ is also $(\mu \circ \phi^\leftarrow)^*$ -measurable. And the outer measure $(\mu \circ \phi^\leftarrow)^*$ is additive on the σ -algebra $(\mu \circ \phi^\leftarrow)^*\text{-}\mathcal{M}\text{eas}$, so

$$(\mu \circ \phi^\leftarrow)^*(C) = (\mu \circ \phi^\leftarrow)^*(B) + (\mu \circ \phi^\leftarrow)^*(C \cap B^c).$$

We have assumed that $(\mu \circ \phi^\leftarrow)^*(B) < \infty$, and by construction $(\mu \circ \phi^\leftarrow)^*(C) = (\mu \circ \phi^\leftarrow)^*(B)$, so $(\mu \circ \phi^\leftarrow)^*(C) < \infty$. It follows that $(\mu \circ \phi^\leftarrow)^*(C \cap B^c) = 0$. This would not follow were $(\mu \circ \phi^\leftarrow)^*(B) = \infty$.

Since $C \cap B^c$ is a $(\mu \circ \phi^\leftarrow)^*$ -null set of Y , it follows by Proposition 3.11 that $\phi^\leftarrow(C \cap B^c)$ is a μ^* -null set of X , and therefore $\mu^*\text{-}\mathcal{M}\text{eas}$. And since $\phi^\leftarrow(C)$ is $\mu^*\text{-}\mathcal{M}\text{eas}$, it follows that the set difference $\phi^\leftarrow(B)$ of $\phi^\leftarrow(C)$ with $\phi^\leftarrow(C \cap B^c)$ is $\mu^*\text{-}\mathcal{M}\text{eas}$. We can also write $\phi^\leftarrow(C)$ as the disjoint union

$$\phi^\leftarrow(C) = \phi^\leftarrow(B) \sqcup \phi^\leftarrow(C \cap B^c).$$

The outer measure μ^* is additive on these $\mu^*\text{-}\mathcal{M}\text{eas}$ sets, so

$$\mu^*(\phi^\leftarrow(C)) = \mu^*(\phi^\leftarrow(B)) + \mu^*(\phi^\leftarrow(C \cap B^c)).$$

But $\phi^\leftarrow(C \cap B^c)$ is a μ^* -null set, so

$$\mu^*(\phi^\leftarrow(C)) = \mu^*(\phi^\leftarrow(B)).$$

Also, as displayed in the proof of Proposition 3.11,

$$(\mu \circ \phi^\leftarrow)^*(C) = \mu^*(\phi^\leftarrow(C)).$$

It follows by transitivity of equality that

$$\mu^*(\phi^\leftarrow(B)) = (\mu \circ \phi^\leftarrow)^*(B),$$

as required. ■

Remark 3.13. One of the difficulties in working with σ -subalgebras is that the restriction of a σ -finite measure to a σ -subalgebra need not be σ -finite. This is a clear indication that our definitions of σ -finite measures and/or σ -subalgebras need more work. A definition that does not respect sub-objects must not yet be right.

{measurable_inclusion}

Proposition 3.14. *Let $(X, \mathfrak{A}, \mu)$ be a finite measure space, where $\mathfrak{A}$ is a σ -algebra. If $\mathfrak{B}$ is a σ -subalgebra of $\mathfrak{A}$, then $\mu|_{\mathfrak{B}}^* \text{-Meas} \subseteq \mu^* \text{-Meas}$, and $\mu|_{\mathfrak{B}}^*(A) = \mu^*(A)$ for all $A \in \mu|_{\mathfrak{B}}^* \text{-Meas}$.*

Proof. Let $\mathfrak{B}$ be a σ -subalgebra of $\mathfrak{A}$, and let $A \in \mu|_{\mathfrak{B}}^* \text{-Meas}$. The function $i : X \rightarrow X : A \mapsto A$, where $i^{\leftarrow} : \mathfrak{B} \rightarrow \mathfrak{A}$, is $(\mathfrak{A}, \mathfrak{B})$ -measurable since $\mathfrak{B}$ is supposed to be a σ -subalgebra of $\mathfrak{A}$. In order to apply Proposition 3.12, we first show that $\mu|_{\mathfrak{B}} = \mu \circ i^{\leftarrow}$ on $\mathfrak{B}$: If $B \in \mathfrak{B}$, then $\mu|_{\mathfrak{B}}(B) = \mu(B) = \mu(i^{\leftarrow}(B)) = (\mu \circ i^{\leftarrow})(B)$. And so by Proposition 2.21, it follows that $\mu|_{\mathfrak{B}}^* = (\mu \circ i^{\leftarrow})^*$ on $\mu|_{\mathfrak{B}}^* \text{-Meas} = (\mu \circ i^{\leftarrow})^* \text{-Meas}$.

Since $A \in \mu|_{\mathfrak{B}}^* \text{-Meas} = (\mu \circ i^{\leftarrow})^* \text{-Meas}$, and since $(X, \mathfrak{A}, \mu)$ is finite, it follows by Proposition 3.12 that $i^{\leftarrow}(A) = A$ is a μ^* -measurable set. This shows that $\mu|_{\mathfrak{B}}^* \text{-Meas} \subseteq \mu^* \text{-Meas}$. It also follows by Proposition 3.12 that $\mu^*(i^{\leftarrow}(A)) = (\mu \circ i^{\leftarrow})^*(A)$, which shows that $\mu^*(A) = \mu|_{\mathfrak{B}}^*(A)$, as required. ■

Remark 3.15. The following proposition says that two functions which are equal almost everywhere are either simultaneously measurable or not. As it should be.

{measurable_almost_everywhere}

Proposition 3.16. *If a function $f : X \rightarrow \mathbb{R}$ is $(\mu^* \text{-Meas}, \text{Bot}_{\mathbb{R}})$ -measurable, and a function g is equal to f μ^* -almost everywhere, then g is also $(\mu^* \text{-Meas}, \text{Bot}_{\mathbb{R}})$ -measurable.*

{measurable_limits}

Theorem 3.17. *Let $\{f_n\}$ be a sequence of $(\mu^* \text{-Meas}, \text{Bot}_{\mathbb{R}})$ -measurable functions. If $f : X \rightarrow \mathbb{R}$ is a function such that $f_n \rightarrow f$ μ^* -almost everywhere, then f is also $(\mu^* \text{-Meas}, \text{Bot}_{\mathbb{R}})$ -measurable.*

Underlying the mechanics of Proposition 3.16 and Theorem 3.17 is the property of completeness. This property describes a tight interplay between a measure and the collection of sets which forms its domain; let's say a measure space is *complete* if every subset of a set of measure zero is measurable. It is almost worth ferreting this out, and then restating the proposition and theorem in these terms. Vaguely, let $(X, \mathfrak{A}, \mu)$ be a complete measure space, where

3 Measurable Functions and Integration

$\mathfrak{A}$ is at least an algebra of sets. Then if $f : X \rightarrow \mathbb{R}$ is an $(\mathfrak{A}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable function, and $g : X \rightarrow \mathbb{R}$ is a function with $g = f$ μ^* -almost everywhere, then g is also an $(\mathfrak{A}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable function. Since the measure space $(X, \mu^*\text{-}\mathfrak{Meas}, \mu^*)$ is a complete measure space, the proposition would follow. But this is a type of interference that generates problems which I do not want to solve here.

Proposition 3.18. *The collection of all $(\mu^*\text{-}\mathfrak{Meas}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable functions forms a vector lattice.*

{simple_function}

Definition 3.19. Let $(X, \mathfrak{A})$ be a measurable space. An $(\mathfrak{A}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable function having finite range is called an $(\mathfrak{A}, \mathfrak{Bor}_{\mathbb{R}})$ -**simple function**.

{measurable_implies_simple_function_limit}

Proposition 3.20. *Let $(X, \mathfrak{A})$ be a measurable space, and let $f : X \rightarrow \mathbb{R}$ be an $(\mathfrak{A}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable function such that $f(x) \geq 0$ for all $x \in X$. There is a sequence $\{\phi_n\}$ of $(\mathfrak{A}, \mathfrak{Bor}_{\mathbb{R}})$ -simple functions such that $\phi_n(x) \geq 0$ and $\phi_n(x) \uparrow f(x)$ for all $x \in X$.*

{measurable_equivalent_simple_function_limit}

Proposition 3.21. *Let $(X, \mathfrak{A})$ be a measurable space. A function $f : X \rightarrow \mathbb{R}$ is $(\mathfrak{A}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable if and only if there is a sequence $\{\phi_n\}$ of $(\mathfrak{A}, \mathfrak{Bor}_{\mathbb{R}})$ -simple functions such that $\lim \phi_n(x) = f(x)$ for all $x \in X$.*

{mixing_measurable_functions}

Theorem 3.22. *Let $(X, \mathfrak{A}, \mu)$ be a σ -finite measure space, where $\mathfrak{A}$ is a σ -algebra. If $f : X \rightarrow \mathbb{R}$ is a $(\mu^*\text{-}\mathfrak{Meas}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable function, then there exists an $(\mathfrak{A}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable function $g : X \rightarrow \mathbb{R}$ with $f = g$ μ^* -almost everywhere.*

Proof. Roughly, we show the result holds for characteristic functions, then for simple functions, then for pointwise limits of simple functions. Assume $f(x) \geq 0$ for all $x \in X$.

Let A be a μ^* -measurable subset of X , and suppose $f = \chi_A$, so f is a $(\mu^*\text{-}\mathfrak{Meas}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable characteristic function. By Proposition 2.23, there is a subset $C \subseteq X$ with $A \cap C = \emptyset$ and such that $\mu^*(C) = 0$ with $A \cup C \in \mathfrak{A}$. Take $g = \chi_{A \cup C}$, so g is an $(\mathfrak{A}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable characteristic function. Then $g - f = \chi_C$, and $\mu^*(C) = 0$, which says that $f = g$ μ^* -almost everywhere.

Now let f be a $(\mu^*\text{-Meas}, \mathfrak{Bot}_{\mathbb{R}})$ -simple function, so by definition we can let the set $\{a_1, \dots, a_n\}$ denote the range of f . Further, let $A_i = f^{-1}(a_i)$, so each A_i is a μ^* -measurable subset of X . We can write $f = \sum_{i=1}^n a_i \chi_{A_i}$. Again, by Proposition 2.23, there are subsets $C_i \subseteq X$ with $A_i \cap C_i = \emptyset$ and such that $\mu^*(C_i) = 0$ with $A_i \cup C_i \in \mathfrak{A}$. Now take $g = \sum_{i=1}^n a_i \chi_{A_i \cup C_i}$, so g is an $(\mathfrak{A}, \mathfrak{Bot}_{\mathbb{R}})$ -simple function. The difference $g - f = \sum_{i=1}^n a_i \chi_{C_i}$. Since $\mu^*(C_i) = 0$, this says that $g - f = 0$ μ^* -almost everywhere; equivalently, $f = g$ μ^* -almost everywhere.

Finally, let $f : X \rightarrow \mathbb{R}$ be any $(\mu^*\text{-Meas}, \mathfrak{Bot}_{\mathbb{R}})$ -measurable function. By Proposition 3.20, there is a sequence $\{\phi_n\}$ of $(\mu^*\text{-Meas}, \mathfrak{Bot}_{\mathbb{R}})$ -simple functions such that $\phi_n(x) \geq 0$ and $\phi_n(x) \uparrow f(x)$ for all $x \in X$. And we have just shown that for each of these ϕ_n , there is an $(\mathfrak{A}, \mathfrak{Bot}_{\mathbb{R}})$ -simple function, call it y_n , such that $\phi_n = y_n$ μ^* -almost everywhere. That is, $\phi_n(x) = y_n(x)$ for all x in the complement of some set, call it A_n , with $\mu^*(A_n) = 0$. The fact that $\mu^*(A_n) = 0$ only says that $A_n \in \mu^*\text{-Meas}$. However, by Proposition 2.22, there is a superset $B_n \in \mathfrak{A}$ with $A_n \subseteq B_n$ and $\mu^*(A_n) = \mu^*(B_n)$. Then $\phi_n(x) = y_n(x)$ for all x in the complement of B_n with $\mu^*(B_n) = 0$ and $B_n \in \mathfrak{A}$. Write $B = \bigcup B_n$, so $B \in \mathfrak{A}$. It follows that $y_n(x) \uparrow f(x)$ for all x in the complement of B . Consequently, $y_n \chi_{B^c} \uparrow f \chi_{B^c}$ on all of X . This time take $g = f \chi_{B^c}$. Then $f = g$ μ^* -almost everywhere. To complete the proof, we need only show that g is $(\mathfrak{A}, \mathfrak{Bot}_{\mathbb{R}})$ -measurable. See that $B \in \mathfrak{A}$ implies $B^c \in \mathfrak{A}$, so χ_{B^c} is $(\mathfrak{A}, \mathfrak{Bot}_{\mathbb{R}})$ -measurable. Each y_n is $(\mathfrak{A}, \mathfrak{Bot}_{\mathbb{R}})$ -measurable, and so the product $y_n \chi_{B^c}$ is an $(\mathfrak{A}, \mathfrak{Bot}_{\mathbb{R}})$ -measurable function and consequently an $(\mathfrak{A}, \mathfrak{Bot}_{\mathbb{R}})$ -simple function. Therefore by Proposition 3.21, the function g is $(\mathfrak{A}, \mathfrak{Bot}_{\mathbb{R}})$ -measurable, as required. $\blacksquare$

{step_function}

Definition 3.23. Let $(X, \mathfrak{A}, \mu)$ be a measure space. Let us say that a $(\mu^*\text{-Meas}, \mathfrak{Bot}_{\mathbb{R}})$ -simple function is a $(\mu^*\text{-Meas}, \mathfrak{Bot}_{\mathbb{R}})$ -**step function** if it can be expressed as a linear combination of characteristic functions of subsets of $\mu^*\text{-Meas}$ of finite μ^* -measure. That is, if ϕ is a $(\mu^*\text{-Meas}, \mathfrak{Bot}_{\mathbb{R}})$ -simple function having range $\{a_1, \dots, a_n\}$, then ϕ is a $(\mu^*\text{-Meas}, \mathfrak{Bot}_{\mathbb{R}})$ -**step function** if $\phi = \sum_{i=1}^n a_i \chi_{A_i}$ with each $A_i \in \mu^*\text{-Meas}$, and with $\mu^*(A_i) < \infty$, where we can suppose the A_i to be pairwise disjoint. Is

3 Measurable Functions and Integration

it clear that with a finite measure space, every $(\mu^*\text{-}\mathcal{M}\text{eas}, \mathfrak{B}\text{o}\mathfrak{r}_{\mathbb{R}})$ -simple function is a $(\mu^*\text{-}\mathcal{M}\text{eas}, \mathfrak{B}\text{o}\mathfrak{r}_{\mathbb{R}})$ -step function?

{sigma_finite_step_functions}

Proposition 3.24. *If $(X, \mathfrak{S}, \mu)$ is a σ -finite measure space, and $f : X \rightarrow \mathbb{R}$ is a $(\mu^*\text{-}\mathcal{M}\text{eas}, \mathfrak{B}\text{o}\mathfrak{r}_{\mathbb{R}})$ -measurable function with $f(x) \geq 0$ on X , then there is a sequence $\{\phi_n\}$ of $(\mu^*\text{-}\mathcal{M}\text{eas}, \mathfrak{B}\text{o}\mathfrak{r}_{\mathbb{R}})$ -step functions such that $\phi_n(x) \geq 0$ and $\phi_n(x) \uparrow f(x)$ for all $x \in X$.*

{integral_step_function}

Definition 3.25. Let $\phi : X \rightarrow \mathbb{R}$ be a $(\mu^*\text{-}\mathcal{M}\text{eas}, \mathfrak{B}\text{o}\mathfrak{r}_{\mathbb{R}})$ -step function such that $\phi = \sum_{i=1}^n a_i \chi_{A_i}$, where the range of ϕ is the set $\{a_1, \dots, a_n\}$, and each $A_i = \phi^{-1}(\{a_i\})$. Then the Lebesgue *integral* of ϕ , denoted by $\int_X \phi d\mu$, or simply $\int \phi$, is defined to be the real number

$$\int_X \phi d\mu = \sum_{i=1}^n a_i \mu^*(A_i).$$

Proposition 3.26. *Let $\phi : X \rightarrow \mathbb{R}$ be a $(\mu^*\text{-}\mathcal{M}\text{eas}, \mathfrak{B}\text{o}\mathfrak{r}_{\mathbb{R}})$ -step function.*

1. *If $\phi \geq 0$ μ^* -almost everywhere, then $\int_X \phi d\mu \geq 0$.*
2. *If $\phi = 0$ μ^* -almost everywhere, then $\int_X \phi d\mu = 0$.*

{continuity_at_zero}

Proposition 3.27. *Let $\{\phi_n\}$ be a sequence of $(\mu^*\text{-}\mathcal{M}\text{eas}, \mathfrak{B}\text{o}\mathfrak{r}_{\mathbb{R}})$ -step functions. If $\phi_n \downarrow 0$ μ^* -almost everywhere, then $\int \phi_n \downarrow 0$.*

{integral_is_unique}

Proposition 3.28. *Let $f : X \rightarrow \mathbb{R}$. If $\{\phi_n\}$ and $\{y_n\}$ are sequences of $(\mu^*\text{-}\mathcal{M}\text{eas}, \mathfrak{B}\text{o}\mathfrak{r}_{\mathbb{R}})$ -step functions with $\phi_n \uparrow f$ μ^* -almost everywhere, and with $y_n \uparrow f$ μ^* -almost everywhere, then $\lim \int_X \phi_n d\mu = \lim \int_X y_n d\mu$.*

{upperfunction}

Definition 3.29. Let $f : X \rightarrow \mathbb{R}$. If there is a sequence $\{\phi_n\}$ of $(\mu^*\text{-}\mathcal{M}\text{eas}, \mathfrak{B}\text{o}\mathfrak{r}_{\mathbb{R}})$ -step functions such that $\phi_n \uparrow f$ μ^* -almost everywhere, and $\lim \int_X \phi_n d\mu < \infty$, then f is called an **upper function**, and the Lebesgue integral of f , denoted by $\int_X f d\mu$, or simply $\int f$, is defined to be the real number

$$\int_X f d\mu = \lim \int_X \phi_n d\mu.$$

We might also call f a $(\mu^*\text{-}\mathcal{M}\text{eas}, \mathfrak{B}\text{o}\mathfrak{r}_{\mathbb{R}})$ -upper function. By Proposition 3.28, the value of the Lebesgue integral of an upper function

is independent of a chosen sequence of step functions $\phi_n \uparrow f$. By Theorem 3.17, every upper function is $(\mu^*\text{-}\mathcal{M}\text{eas}, \mathcal{B}\text{or}_{\mathbb{R}})$ -measurable.

Proposition 3.30. *If f and g are upper functions, then:*

1. *The function $f + g$ is an upper function with $\int (f + g) d\mu = \int f d\mu + \int g d\mu$.*
2. *If $0 \leq a \in \mathbb{R}$, then af is an upper function and $\int (af) d\mu = a \int f d\mu$.*
3. *The functions $f \vee g$ and $f \wedge g$ are upper functions.*

Proposition 3.31. *If f and g are upper functions with $f \leq g$ μ^* -almost everywhere, then $\int f \leq \int g$.*

Proposition 3.32. *If $f : X \rightarrow \mathbb{R}$ is a function with $\{f_n\}$ a sequence of upper functions such that $f_n(x) \uparrow f(x)$ for μ^* -almost all $x \in X$, and if $\lim \int f_n d\mu < \infty$, then f is an upper function and $\lim \int f_n d\mu = \int f d\mu$.*

Proposition 3.33. *Let $\{f_n\}$ be a sequence of upper functions. If $f_n \downarrow 0$ μ^* -almost everywhere, then $\int f_n \downarrow 0$.*

Definition 3.34. A function $f : X \rightarrow \mathbb{R}$ is **integrable** over X with respect to μ if there exist two upper functions g and h such that $f = g - h$ μ^* -almost everywhere. Then the Lebesgue integral of f over X with respect to μ , denoted by $\int_X f d\mu$, or simply $\int f$, is defined to be the real number

$$\int_X f d\mu = \int_X g d\mu - \int_X h d\mu.$$

The value of this integral is independent of the representation of f as a difference of upper functions. A function $f : X \rightarrow \mathbb{R}$ is *integrable over a μ^* -measurable subset A with respect to μ* if $f \cdot \chi_A$ is integrable over X with respect to μ , in which case $\int_A f d\mu$ is defined to be $\int_X f \cdot \chi_A d\mu$.

An extended real function $f : X \rightarrow [-\infty, +\infty]$ **defines an integrable function** over X with respect to μ if the function f “attains” $-\infty$ or $+\infty$, or is undefined, on a set of at most μ^* -measure zero, so it is possible to redefine the values of f on this set in order that the corresponding redefined function $X \rightarrow \mathbb{R}$ is integrable over X with respect to μ . We assume all such functions have been so redefined, and consequently lie

{upper_monotone}

{integrable}

in the domain of the Lebesgue integral, if that makes sense. Extended real functions are hard to avoid when it comes to dealing with signed measures, or even positive measures.

{integrable_and_positive_is_upper}

Proposition 3.35. *Let $f : X \rightarrow \mathbb{R}$ be a function integrable over X with respect to μ . If $f \geq 0$ μ^* -almost everywhere, then f is an upper function.*

{sigma_algebra_measurable}

Proposition 3.36. *Let $(X, \mathfrak{A}, \mu)$ be a σ -finite measure space, where $\mathfrak{A}$ is a σ -algebra. If $f : X \rightarrow \mathbb{R}$ is a function integrable over X with respect to μ , then there is an $(\mathfrak{A}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable function $g : X \rightarrow \mathbb{R}$ with $f = g$ μ^* -almost everywhere.*

Proof. Let $f : X \rightarrow \mathbb{R}$ be integrable over X with respect to μ . By Theorem 3.17, the function f is $(\mu^*\text{-}\mathfrak{Meas}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable, and by Theorem 3.22, the function f is μ^* -almost everywhere equal to an $(\mathfrak{A}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable function, as required. ■

Remark 3.37. The collection of real functions on X that can be written as the difference of upper functions forms a vector lattice, and the integral is a positive linear functional on this vector lattice. Maybe the integral here should have been defined as a positive linear functional with a continuity condition at zero, per Daniell. Of which, the Lebesgue, and so the Riemann, integral is simply a concrete example.

{dominated_by_integrable}

Proposition 3.38. *If $f : X \rightarrow \mathbb{R}$ be an integrable function, and if $g : X \rightarrow \mathbb{R}$ is a $(\mu^*\text{-}\mathfrak{Meas}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable function with $0 \leq g \leq f$ μ^* -almost everywhere, then g is an integrable function.*

{integral_absolute_zero}

Proposition 3.39. *Let $f : X \rightarrow \mathbb{R}$ be a function integrable over X with respect to μ . Then $\int |f| d\mu = 0$ if and only if $f = 0$ μ^* -almost everywhere.*

{levi}

Theorem 3.40 (Levi). *Let $\{f_n\}$ be a sequence of functions integrable with respect to μ . If $f_n(x) \leq f_{n+1}(x)$ for μ^* -almost all $x \in X$, and if $\lim \int f_n d\mu < \infty$, then there is a function f integrable with respect to μ such that $f_n(x) \uparrow f(x)$ for μ^* -almost all $x \in X$. Consequently, $\int f_n d\mu \uparrow \int f d\mu$.*

uniqueness_radon_nikodym}

Proposition 3.41. *Let $(X, \mathfrak{S}, \mu)$ be a measure space, and let $f : X \rightarrow \mathbb{R}$ be integrable over X with respect to μ . If $\int_A f \, d\mu = 0$ for all $A \in \mathfrak{S}$, then $f = 0$ μ^* -almost everywhere.*

4 Function Spaces and Quotient Spaces

IN ORDER to deal with the L_p spaces in a manner that is actually mathematically correct, we will use three theorems from algebra; namely, the induced homomorphism theorem, the correspondence theorem, and the induced quotient homomorphism corollary. These appear respectively as Proposition 4.3, Proposition 4.5, and Proposition 4.18. These propositions are couched in terms of group theory because that is all it takes to get to the heart of the matter. Understand that each of these three theorems has a straightforward extension to modules or vector spaces.

We pay attention to the difference between a function and an equivalence class of functions for at least two reasons. The first reason is that clarity is more important here. The second reason is that we will deal with so-called *versions* of maps defined on σ -algebras which will require selecting representatives from equivalence classes of functions. These will appear in Definition 8.10.

{Lp}

Definition 4.1. Let $(X, \mathfrak{S}, \mu)$ be a measure space, and let $p \in (0, \infty)$. The collection of all $(\mu^*\text{-}\mathfrak{M}eas, \mathfrak{B}or_{\mathbb{R}})$ -measurable functions $f : X \rightarrow \mathbb{R}$ for which $|f|^p$ is integrable forms a semi-normed vector space, denoted by $\mathcal{L}_p(X, \mathfrak{S}, \mu)$. If $f \in \mathcal{L}_p(X, \mathfrak{S}, \mu)$, then $\int |f|^p d\mu = 0$ if and only if $f = 0$ μ^* -almost everywhere. A quotient space defined by identifying those functions in $\mathcal{L}_p(X, \mathfrak{S}, \mu)$ which agree μ^* -almost everywhere forms a normed vector space, denoted by $L_p(X, \mathfrak{S}, \mu)$, where the norm will be defined in terms of the integral. That is, let $\mathcal{N}_p(X, \mathfrak{S}, \mu)$ denote the subspace of functions in $\mathcal{L}_p(X, \mathfrak{S}, \mu)$ which are equal to zero μ^* -almost everywhere, then

$$L_p(X, \mathfrak{S}, \mu) := \mathcal{L}_p(X, \mathfrak{S}, \mu) / \mathcal{N}_p(X, \mathfrak{S}, \mu).$$

For each $f \in \mathcal{L}_p(X, \mathfrak{S}, \mu)$, we will let $\bar{f}^\mu$ denote the equivalence class $f + \mathcal{N}_p(X, \mathfrak{S}, \mu)$ in $L_p(X, \mathfrak{S}, \mu)$. Should $(X, \mathfrak{S}, \mu)$ and $(X, \mathfrak{T}, \nu)$ both be measure spaces, and should the function $f : X \rightarrow \mathbb{R}$ be in both $\mathcal{L}_p(X, \mathfrak{S}, \mu)$ and $\mathcal{L}_p(X, \mathfrak{T}, \nu)$, then the equivalence class $\bar{f}^\mu$ need not be related to the equivalence class $\bar{f}^\nu$. Not even when $\mathfrak{T} \subseteq \mathfrak{S}$ and $\nu = \mu|_{\mathfrak{T}}$, as shown in Example 4.15. Should a discussion or proposition involve only a single measure μ , we might inconsistently drop the superscript and denote $\bar{f}^\mu$ by $\bar{f}$.

Since $\mathcal{N}_1(X, \mathfrak{S}, \mu)$ is in the kernel of the integral $\int$, there is a unique operator $\bar{\int}$ which makes the following diagram commute:

$$\begin{array}{ccc} \mathcal{L}_1(X, \mathfrak{S}, \mu) & \xrightarrow{\int} & \mathbb{R} \\ \text{quotient} \downarrow & \searrow \bar{\int} & \\ L_1(X, \mathfrak{S}, \mu) & & \end{array}$$

It means that for a class $\bar{f}^\mu \in L_p(X, \mathfrak{S}, \mu)$, the value of $\bar{\int}_X \bar{f}^\mu d\mu$ is defined to be equal to the value of $\int_X f d\mu$. Verify that the map $\bar{\int}$ is constant on the cosets of $L_p(X, \mathfrak{S}, \mu)$. The integral symbol ' $\bar{\int}$ ' with the overbar was chosen to somewhat distinguish itself from the plain integral symbol ' $\int$ '. We have a very large number of symbols in mathematics. Do not use the exact same symbol for everything. As usual, define the L_p norm by

$$\|\bar{f}^\mu\|_p := \left(\int |f|^p d\mu \right)^{1/p}.$$

The quotient space $L_p(X, \mathfrak{S}, \mu)$ need not form an algebra with respect to pointwise multiplication. Still, for classes $\bar{f}^\mu$ and $\bar{g}^\mu$ in $L_p(X, \mathfrak{S}, \mu)$, if the function $|fg|^p$ is integrable with respect to the measure μ , then we can define the product $\bar{f}^\mu \bar{g}^\mu$ of classes to be the class $\bar{fg}^\mu$. In this case, verify that if $h \in \bar{f}^\mu$, and $k \in \bar{g}^\mu$, then $hk^\mu = \bar{fg}^\mu$.

Remark 4.2. The extension of the following proposition to modules, or vector spaces, underlies the commutative diagram in Definition 4.1.

Proposition 4.3. *If A and B are groups with $C \triangleleft A$, and ϕ is a homomorphism $A \rightarrow B$ with C contained in the $\text{Ker } \phi$, there is a unique induced homomorphism $\bar{\phi} : A/C \rightarrow B$ mapping the coset $a + C$ to $\phi(a)$, so that the following diagram commutes.*

{induced_homomorphism}

$$\begin{array}{ccc} A & \xrightarrow{\phi} & B \\ Q_C \downarrow & \nearrow \bar{\phi} & \\ A/C & & \end{array}$$

The $\text{Ran } \bar{\phi} = \text{Ran } \phi$ and the $\text{Ker } \bar{\phi} = (\text{Ker } \phi)/C$. The induced homomorphism is an isomorphism exactly when both ϕ is epic and $C = \text{Ker } \phi$.

Remark 4.4. The $L_p(X, \mathfrak{S}, \mu)$ spaces can be comparable; for example, let $(X, \mathfrak{S}, \mu)$ be a finite measure space, then the quotient space $L_2(X, \mathfrak{S}, \mu)$ is a vector subspace of the quotient space $L_1(X, \mathfrak{S}, \mu)$. This is the heart of the correspondence theorem, stated in Proposition 4.5, which extends to vector spaces, and it would be necessary that the kernel $\mathcal{N}_2(X, \mathfrak{S}, \mu)$ be equal to $\mathcal{N}_1(X, \mathfrak{S}, \mu)$ in order for the proposition to apply. The result then that $L_2(X, \mathfrak{S}, \mu)$ is a vector subspace of $L_1(X, \mathfrak{S}, \mu)$ follows from the fact that for a finite measure space, $\mathcal{L}_2(X, \mathfrak{S}, \mu) \subseteq \mathcal{L}_1(X, \mathfrak{S}, \mu)$, which in turn is an application of Hölder's inequality.

{correspondence_theorem}

Proposition 4.5. *If $f : G \twoheadrightarrow H$ is an epimorphism of groups, then the assignment $K \mapsto f(K)$ defines a one-to-one correspondence between the set of all subgroups K of G which contain the $\text{Ker } f$ and the set of all subgroups of H .*

$$\begin{array}{ccccc} \text{Ker } f & \leq & K & \leq & G \\ & & \downarrow & & \downarrow \\ & & f(K) & \leq & H \end{array}$$

Consequently, if N is a normal subgroup of a group G , then every subgroup of G/N is of the form K/N , where K is a subgroup of G that

contains N .

$$\begin{array}{ccccc} N & \leq & K & \leq & G \\ \downarrow & & \downarrow & & \downarrow \\ N/N & \leq & K/N & \leq & G/N \end{array}$$

Proposition 4.6. *If $(X, \mathfrak{S}, \mu)$ is a measure space, then $\mathcal{N}_2(X, \mathfrak{S}, \mu) = \mathcal{N}_1(X, \mathfrak{S}, \mu)$.*

Proof. If $f \in \mathcal{L}_2(X, \mathfrak{S}, \mu)$ and $f = 0$ μ^* -almost everywhere, then $f \in \mathcal{L}_1(X, \mathfrak{S}, \mu)$, and if $f \in \mathcal{L}_1(X, \mathfrak{S}, \mu)$ and $f = 0$ μ^* -almost everywhere, then $f \in \mathcal{L}_2(X, \mathfrak{S}, \mu)$. ■

Proposition 4.7. *If $(X, \mathfrak{S}, \mu)$ is finite, then $L_2(X, \mathfrak{S}, \mu)$ is a vector subspace of the quotient space $L_1(X, \mathfrak{S}, \mu)$.*

$$\begin{array}{ccccc} \mathcal{N}_1(X, \mathfrak{A}, \mu) & \leq & \mathcal{L}_2(X, \mathfrak{A}, \mu) & \leq & \mathcal{L}_1(X, \mathfrak{A}, \mu) \\ \downarrow & & \downarrow & & \downarrow \\ \text{trivial} & \leq & L_2(X, \mathfrak{A}, \mu) & \leq & L_1(X, \mathfrak{A}, \mu) \end{array}$$

Proposition 4.8. *Let $(X, \mathfrak{S}, \mu)$ be a measure space, and let f be an integrable function such that $f > 0$ μ^* -almost everywhere. If A is a μ^* -measurable subset of X , and $\int_A f d\mu = 0$, then $\mu^*(A) = 0$.*

{null_set_test}

Proposition 4.9. *Let $(X, \mathfrak{S}, \mu)$ be a measure space. If the class $\bar{f}^\mu \in L_1(X, \mathfrak{S}, \mu)$, and if $\int_A f d\mu = 0$ for all $A \in \mathfrak{S}$, then $\bar{f}^\mu = \bar{0}^\mu$. Alternately, if $\bar{f}^\mu, \bar{g}^\mu \in L_1(X, \mathfrak{S}, \mu)$, and if $\int_A f d\mu = \int_A g d\mu$ for all $A \in \mathfrak{S}$, then $\bar{f}^\mu = \bar{g}^\mu$.*

{equal_function_test}

Compare the following Proposition 4.10 (where we assume integrability) with Proposition 3.24 (where we do not assume integrability, but we do assume σ -finiteness):

Proposition 3.24. If $(X, \mathfrak{S}, \mu)$ is a σ -finite measure space, and $f : X \rightarrow \mathbb{R}$ is a $(\mu^*\text{-Meas}, \mathfrak{Bot}_{\mathbb{R}})$ -measurable function with $f(x) \geq 0$ on X , then there is a sequence $\{\phi_n\}$ of $(\mu^*\text{-Meas}, \mathfrak{Bot}_{\mathbb{R}})$ step functions such that $\phi_n(x) \geq 0$ and $\phi_n(x) \uparrow f(x)$ for all $x \in X$.

Proposition 4.10. For every $p \in [1, \infty)$, the equivalence classes of $(\mu^*\text{-Meas}, \text{Bot}_{\mathbb{R}})$ -step functions form a vector sublattice of the Banach lattice $L_p(X, \mathfrak{S}, \mu)$, which is norm dense.

{step_functions_dense}

{integral_continuous}

Proposition 4.11. If E is a Banach lattice and F is a normed vector lattice, then every positive operator $E \rightarrow F$ is continuous.

Remark 4.12. The integral $\bar{f}$ on $L_1(X, \mathfrak{S}, \mu)$ is an example of a positive operator on a Banach lattice, and so the integral on $L_1(X, \mathfrak{S}, \mu)$ is necessarily continuous. But with respect to which topologies?

The following proposition describes what is sometimes called a *change of variable*, and perhaps should have been couched as:

$$\begin{array}{ccccc} & & f \circ T & & \\ & \nearrow & & \searrow & \\ (X, \mathfrak{A}, \mu) & \xrightarrow[T^{-1}(B)]{T} & (Y, \mathfrak{B}, \mu \circ T^{-1}) & \xrightarrow[B]{f} & (\mathbb{R}, \text{Bot}_{\mathbb{R}}), \end{array}$$

with

$$\int_{T^{-1}(B)} (f \circ T) d\mu = \int_B f d(\mu \circ T^{-1}).$$

{change_of_variable}

Proposition 4.13 (Change of Variable). Let $(X, \mathfrak{A}, \mu)$ be a measure space, where $\mathfrak{A}$ is a σ -algebra, and let $(Y, \mathfrak{B})$ be a measurable space. Also, let the function $T : X \rightarrow Y$ be $(\mathfrak{A}, \mathfrak{B})$ -measurable, and momentarily let $\nu = \mu \circ T^{-1}$. If $\bar{f}^\nu \in L_1(Y, \mathfrak{B}, \nu)$, then $\bar{f} \circ T^\mu \in L_1(X, \mathfrak{A}, \mu)$ and

$$\int_X (f \circ T) d\mu = \int_Y f d\nu.$$

Proof. This proof relies upon the fact that $\chi_B \circ T = \chi_{T^{-1}(B)}$. We will show that the displayed equality holds in case $f = \chi_B$, where we let the subset B of Y be a $(\mu \circ T^{-1})^*$ -measurable set with $(\mu \circ T^{-1})^*(B) < \infty$, and note that the outer measure $(\mu \circ T^{-1})^*(B)$ should be finite in order that χ_B be integrable with respect to the induced measure $\mu \circ T^{-1}$.

By Proposition 3.12, the set $T^{\leftarrow}(B)$ is μ^* -measurable with

$$\mu^*(T^{\leftarrow}(B)) = (\mu \circ T^{\leftarrow})^*(B),$$

and so

$$\begin{aligned} \int_X \chi_B \circ T \, d\mu &= \int_X \chi_{T^{\leftarrow}(B)} \, d\mu && \chi_B \circ T = \chi_{T^{\leftarrow}(B)} \\ &= \mu^*(T^{\leftarrow}(B)) && \text{by Definition 3.25} \\ &= (\mu \circ T^{\leftarrow})^*(B) && \text{Proposition 3.12} \\ &= \int_Y \chi_B \, d(\mu \circ T^{\leftarrow}) && \text{by Definition 3.25,} \end{aligned}$$

as claimed. ■

{integral_over_subalgebra}

Proposition 4.14. *Let $(X, \mathfrak{A}, \mu)$ be a finite measure space, where $\mathfrak{A}$ is a σ -algebra, and $\mathfrak{B}$ is a σ -subalgebra of $\mathfrak{A}$. If $\bar{f}^{\mu|_{\mathfrak{B}}} \in L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$, then $\bar{f}^{\mu} \in L_1(X, \mathfrak{A}, \mu)$ and*

$$\int_X f \, d\mu|_{\mathfrak{B}} = \int_X f \, d\mu.$$

Consequently, the operator

$$\mathfrak{I} : L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}}) \rightarrow L_1(X, \mathfrak{A}, \mu)$$

defined by $\bar{f}^{\mu|_{\mathfrak{B}}} \mapsto \bar{f}^{\mu}$ has norm 1. Furthermore, the operator $\mathfrak{I}$ is injective.

Proof. Let $\bar{f}^{\mu|_{\mathfrak{B}}} \in L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$. The idea is to apply a change of variable. Define the function $i : X \rightarrow X : x \mapsto x$, where $i^{\leftarrow} : \mathfrak{B} \rightarrow \mathfrak{A} : B \mapsto B$. The function i is $(\mathfrak{A}, \mathfrak{B})$ -measurable because $\mathfrak{B}$ is supposed to be a σ -subalgebra of $\mathfrak{A}$.

We now show that $\int_X f \circ i \, d\mu = \int_X f \, d(\mu \circ i^{\leftarrow})$ by applying Proposition 4.13 to the measure space $(X, \mathfrak{A}, \mu)$ and the measurable space

4 Function Spaces and Quotient Spaces

$(X, \mathfrak{B})$ and the function i . Since i is $(\mathfrak{A}, \mathfrak{B})$ -measurable, and $\bar{f}^{\mu|_{\mathfrak{B}}} \in L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$, it follows by Proposition 4.13 that $\bar{f}^{\mu} \in L_1(X, \mathfrak{A}, \mu)$ and

$$\int_X (f \circ i) d\mu = \int_X f d(\mu \circ i^{\leftarrow}).$$

We next show that $\int_X f d(\mu \circ i^{\leftarrow}) = \int_X f d\mu|_{\mathfrak{B}}$ by applying Proposition 3.14. The assumption that the measure space $(X, \mathfrak{A}, \mu)$ be finite is essential here. The measure $\mu \circ i^{\leftarrow}$ is equal to μ on $\mathfrak{B}$; that is, if $B \in \mathfrak{B}$, then $\mu \circ i^{\leftarrow}(B) = \mu(i^{\leftarrow}(B)) = \mu(B)$. This means that $\mu \circ i^{\leftarrow}$ and μ generate the same outer measure on the σ -algebra $\mu|_{\mathfrak{B}}^* \text{-}\mathfrak{M}_{\text{as}}$. That is, $(\mu \circ i^{\leftarrow})^* = \mu^*$ on $\mu|_{\mathfrak{B}}^* \text{-}\mathfrak{M}_{\text{as}}$. Because the measure space $(X, \mathfrak{A}, \mu)$ is finite, it follows by Proposition 3.14 that $\mu^* = \mu|_{\mathfrak{B}}^*$ on $\mu|_{\mathfrak{B}}^* \text{-}\mathfrak{M}_{\text{as}}$. Since $(\mu \circ i^{\leftarrow})^* = \mu|_{\mathfrak{B}}^*$ on the σ -algebra $\mu|_{\mathfrak{B}}^* \text{-}\mathfrak{M}_{\text{as}}$, it now follows by the definition of the integral that

$$\int_X f d(\mu \circ i^{\leftarrow}) = \int_X f d\mu|_{\mathfrak{B}}.$$

We finally see that

$$\int_X f d\mu = \int_X f \circ i d\mu,$$

and this follows from the fact that $f = f \circ i$ on X .

Therefore, by transitivity of equality,

$$(4.1) \quad \int_X f d\mu|_{\mathfrak{B}} = \int_X f d\mu$$

as required; in other words, the operator $\mathfrak{I}$ preserves the integral.

To see that $\|\mathfrak{I}\| = 1$, witness:

$$\begin{aligned} \|\mathfrak{I}\| &= \sup \{ \|\mathfrak{I} \bar{f}^{\mu|_{\mathfrak{B}}}\| : \|\bar{f}^{\mu|_{\mathfrak{B}}}\| = 1 \} \\ &= \sup \{ \|\bar{f}^{\mu}\| : \|\bar{f}^{\mu|_{\mathfrak{B}}}\| = 1 \} \\ &= \sup \{ \int |f| d\mu : \int |f| d\mu|_{\mathfrak{B}} = 1 \}. \end{aligned}$$

And we have just barely shown in equation (4.1) that the operator $\mathcal{I}$ preserves the integral, so $\int |f| d\mu = \int |f| d\mu_{|\mathfrak{B}}$ for all f in $\mathcal{L}_1(X, \mathfrak{B}, \mu_{|\mathfrak{B}})$. Since an algebra is nonempty, there is at least one f in $\mathcal{L}_1(X, \mathfrak{B}, \mu_{|\mathfrak{B}})$ such that $\int |f| d\mu_{|\mathfrak{B}} = 1$, it follows that $\|\mathcal{I}\| = 1$.

To show that $\mathcal{I}$ is injective, it is sufficient to show that the kernel $\mathcal{I}^\leftarrow(\bar{0}^\mu)$ of $\mathcal{I}$ is trivial, which means showing that $\mathcal{I}^\leftarrow(\bar{0}^\mu) = \{\bar{0}^{\mu_{|\mathfrak{B}}}\}$ in $L_1(X, \mathfrak{B}, \mu_{|\mathfrak{B}})$.

To show the inclusion $\{\bar{0}^{\mu_{|\mathfrak{B}}}\} \subseteq \mathcal{I}^\leftarrow(\bar{0}^\mu)$ means showing that $\mathcal{I}(\bar{0}^{\mu_{|\mathfrak{B}}}) = \bar{0}^\mu$. But this holds by the very definition of $\mathcal{I}$.

To show the inclusion $\mathcal{I}^\leftarrow(\bar{0}^\mu) \subseteq \{\bar{0}^{\mu_{|\mathfrak{B}}}\}$, let $\bar{f}^{\mu_{|\mathfrak{B}}}$ be any element of $\mathcal{I}^\leftarrow(\bar{0}^\mu)$, and we want to see that this implies $\bar{f}^{\mu_{|\mathfrak{B}}} = \bar{0}^{\mu_{|\mathfrak{B}}}$. To assume that $\bar{f}^{\mu_{|\mathfrak{B}}}$ is in $\mathcal{I}^\leftarrow(\bar{0}^\mu)$ means to assume that $\mathcal{I}\bar{f}^{\mu_{|\mathfrak{B}}} = \bar{0}^\mu$, or equivalently that $\bar{f}^\mu = \bar{0}^\mu$. In turn, this means to assume that $f \in \mathcal{N}_1(X, \mathfrak{A}, \mu)$, so that $\int |f| d\mu = 0$. We have just shown in equation (4.1) that the operator $\mathcal{I}$ preserves the integral, so $\int |f| d\mu = 0$ implies that $\int |f| d\mu_{|\mathfrak{B}} = 0$. This says $f \in \mathcal{N}_1(X, \mathfrak{B}, \mu_{|\mathfrak{B}})$, which also says $\bar{f}^{\mu_{|\mathfrak{B}}} = \bar{0}^{\mu_{|\mathfrak{B}}}$, as required. ■

{unequal_kernel}

Example 4.15. Let us show by way of example that $\mathcal{N}_1(X, \mathfrak{B}, \mu_{|\mathfrak{B}})$ need not equal $\mathcal{N}_1(X, \mathfrak{A}, \mu)$. Let $\mathfrak{A} = \{\emptyset, A, A^c, X\}$ with the measure μ being Dirac measure δ_x concentrated at a point $x \in A$, with A non-empty, and with A not equal to X . Let $\mathfrak{B} = \{\emptyset, X\}$.

Verify that the collection of $(\delta_x)^*$ -measurable subsets of X is $\mathfrak{A}$, and that the collection of $(\delta_{x|\mathfrak{B}})^*$ -measurable subsets of X is $\mathfrak{B}$. The set A is not $(\delta_{x|\mathfrak{B}})^*$ -measurable since it fails to cut X sharply with respect to $(\delta_{x|\mathfrak{B}})^*$.

The only function in $\mathcal{N}_1(X, \mathfrak{B}, \delta_{x|\mathfrak{B}})$ is the zero function since $\emptyset$ is the only set of $(\delta_{x|\mathfrak{B}})^*$ -measure zero.

But the set A^c is a set of $(\delta_x)^*$ -measure zero. And since the function χ_{A^c} is $((\delta_x)^*\mathcal{M}_{\text{cas}}, \mathfrak{B}_{\text{or}\mathbb{R}})$ -measurable, and $\chi_{A^c} = 0$ $(\delta_x)^*$ -almost everywhere, it follows that the function χ_{A^c} is in $\mathcal{N}_1(X, \mathfrak{A}, \delta_x)$.

Conclude that $\mathcal{N}_1(X, \mathfrak{B}, \delta_{x|\mathfrak{B}})$ need not equal $\mathcal{N}_1(X, \mathfrak{A}, \delta_x)$.

{L1_subspace_embedding}

Discussion 4.16. What does Proposition 4.14 not say? It does not say that the quotient space $L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$ is a subspace of the quotient space $L_1(X, \mathfrak{A}, \mu)$. This is because the quotient space $L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$ is not in general a subspace of the quotient space $L_1(X, \mathfrak{A}, \mu)$, and this is because the kernel $\mathcal{N}_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$ need not equal the kernel $\mathcal{N}_1(X, \mathfrak{A}, \mu)$, as pointed out in Example 4.15; all possible subspaces of quotient spaces are characterized by the correspondence in Proposition 4.5.

Still, the *function* space $\mathcal{L}_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$ is a subspace of the function space $\mathcal{L}_1(X, \mathfrak{A}, \mu)$, and the inclusion $\mathcal{L}_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}}) \subseteq \mathcal{L}_1(X, \mathfrak{A}, \mu)$ does preserve the semi-norm, so there is an inclusion of subspaces: $\mathcal{N}_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}}) \subseteq \mathcal{N}_1(X, \mathfrak{A}, \mu)$. This implies that there is a unique map making the following diagram commute:

$$\begin{array}{ccc} L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}}) & \xrightarrow{\quad \quad \quad} & L_1(X, \mathfrak{A}, \mu) \\ \text{quotient} \uparrow & & \uparrow \text{quotient} \\ \mathcal{L}_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}}) & \xrightarrow{\text{inclusion}} & \mathcal{L}_1(X, \mathfrak{A}, \mu) \end{array}$$

{subset_vs_subspace}

The distinction between a subset and subspace could be a little less blurred. Rather than the notation

$$\mathcal{L}_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}}) \subseteq \mathcal{L}_1(X, \mathfrak{A}, \mu)$$

something better might be

$$\mathcal{L}_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}}) \leq \mathcal{L}_1(X, \mathfrak{A}, \mu)$$

to suggest some kind of structure-preserving thing.

Remark 4.17. The extension of the following proposition to modules or vector spaces underlies some of the mechanics of the map $L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}}) \rightarrow L_1(X, \mathfrak{A}, \mu)$.

{induced_quotient_homomorphism}

Proposition 4.18. *Whenever A and B are groups with $C \triangleleft A$ and $D \triangleleft B$, and ϕ is a homomorphism $A \rightarrow B$ such that its restriction $\phi|_C$ is a homomorphism $C \rightarrow D$, there is a unique induced homomorphism*

$\widehat{\phi} : A/C \rightarrow B/D$ mapping the coset $a + C$ to the coset $\phi(a) + D$.

$$\begin{array}{ccc} A & \xrightarrow{\phi} & B \\ \nabla & & \nabla \\ C & \xrightarrow{\phi|_C} & D \end{array} \implies \begin{array}{ccc} A/C & \xrightarrow{\widehat{\phi}} & B/D \\ \Psi & & \Psi \\ a + C & \mapsto & \phi(a) + D \end{array}$$

The induced homomorphism $\widehat{\phi}$ is an isomorphism exactly when both $(\text{Ran } \phi) \vee D = B$ and $\phi^{\leftarrow}(D) \leq C$.

Remark 4.19. You may be wondering if there is a more direct way of proving Proposition 4.14, possibly one which more directly uses the mechanics of Proposition 4.18, but which avoids the change of variable. Propositions 4.13 and 4.14 are really just elaborations of Propositions 3.12 and 3.14, and it will be interesting to see how you would avoid these latter two.

{submeasurable_step_function}

Proposition 4.20. Let $(X, \mathfrak{A}, \mu)$ be a finite measure space, where $\mathfrak{A}$ is a σ -algebra, and let $\mathfrak{B}$ be a σ -subalgebra of $\mathfrak{A}$. If $\bar{f}^\mu \in L_1(X, \mathfrak{A}, \mu)$, and if f is $(\mathfrak{B}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable, then there is a sequence $\{\phi_n\}$ of $(\mu^*\text{-Meas}, \mathfrak{Bor}_{\mathbb{R}})$ -step functions with $\phi_n(x) \uparrow f(x)$ for all $x \in X$, where each ϕ_n is a sum of the form

$$\sum_{i=1}^m b_i \chi_{B_i}$$

with each $B_i \in \mathfrak{B}$.

Proof. Let $f \in \mathcal{L}_1(X, \mathfrak{A}, \mu)$, and let f be $(\mathfrak{B}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable. We can suppose that $f \geq 0$. With f being $(\mathfrak{B}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable and nonnegative, it follows by Proposition 3.20 that there is a sequence $\{\phi_n\}$ of $(\mathfrak{B}, \mathfrak{Bor}_{\mathbb{R}})$ -simple functions such that $\phi_n(x) \geq 0$ and $\phi_n(x) \uparrow f(x)$ for all $x \in X$. Each ϕ_n is a sum of the form

$$\sum_{i=1}^m b_i \chi_{B_i}$$

with each $B_i \in \mathfrak{B}$. By hypothesis, $(X, \mathfrak{A}, \mu)$ is a finite measure space which means that each $\mu^*(B_i) < \infty$. And since $\mathfrak{B} \subseteq \mathfrak{A}$ by hypothesis, and $\mathfrak{A} \subseteq \mu^*\text{-}\mathcal{M}\text{eas}$ by Proposition 2.16, it follows that each $B_i \in \mu^*\text{-}\mathcal{M}\text{eas}$, and so each ϕ_n is a $(\mu^*\text{-}\mathcal{M}\text{eas}, \mathfrak{B}\text{or}_{\mathbb{R}})$ -step function, completing the proof. ■

Remark 4.21. The following proposition is a partial converse to Proposition 4.14.

{algebra_to_subalgebra}

Proposition 4.22. *Let $(X, \mathfrak{A}, \mu)$ be a finite measure space, where $\mathfrak{A}$ is a σ -algebra, and let $\mathfrak{B}$ be a σ -subalgebra of $\mathfrak{A}$. If $\bar{f}^\mu \in L_1(X, \mathfrak{A}, \mu)$, and if the function f is $(\mathfrak{B}, \mathfrak{B}\text{or}_{\mathbb{R}})$ -measurable, then $\bar{f}^{\mu|_{\mathfrak{B}}} \in L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$.*

Proof. Let $\bar{f}^\mu \in L_1(X, \mathfrak{A}, \mu)$, with f a $(\mathfrak{B}, \mathfrak{B}\text{or}_{\mathbb{R}})$ -measurable representative. By Proposition 4.20, there is a sequence $\{\phi_n\}$ of $(\mu^*\text{-}\mathcal{M}\text{eas}, \mathfrak{B}\text{or}_{\mathbb{R}})$ -step functions with $\phi_n(x) \uparrow f(x)$ for all $x \in X$, where each ϕ_n is a sum of the form

$$\sum_{i=1}^m b_i \chi_{B_i}$$

with each $B_i \in \mathfrak{B}$. By Proposition 2.16, every element of $\mathfrak{B}$ is a $\mu|_{\mathfrak{B}}^*$ -measurable subset, so $\{\phi_n\}$ is a sequence of $(\mu|_{\mathfrak{B}}^*\text{-}\mathcal{M}\text{eas}, \mathfrak{B}\text{or}_{\mathbb{R}})$ -step functions with $\phi_n(x) \uparrow f(x)$ for all $x \in X$. In order to complete the proof that f is integrable with respect to $\mu|_{\mathfrak{B}}$, it is sufficient to show that $\lim \int_X \phi_n d\mu|_{\mathfrak{B}} < \infty$.

Each ϕ_n is integrable with respect to μ and with respect to $\mu|_{\mathfrak{B}}$, and so by Proposition 4.14,

$$\int_X \phi_n d\mu = \int_X \phi_n d\mu|_{\mathfrak{B}}.$$

By hypothesis, $\int f d\mu < \infty$, and $\int_X f d\mu = \lim \int_X \phi_n d\mu$, so $\lim \int_X \phi_n d\mu < \infty$. Then since $\int_X \phi_n d\mu = \int_X \phi_n d\mu|_{\mathfrak{B}}$, it follows that $\lim \int_X \phi_n d\mu = \lim \int_X \phi_n d\mu|_{\mathfrak{B}} < \infty$, completing the proof. ■

Remark 4.23. Let $(X, \mathfrak{A}, \mu)$ be a measure space, where $\mathfrak{A}$ is a σ -algebra, and let $f^\mu \in L_1(X, \mathfrak{A}, \mu)$. The representative function f is necessarily $(\mu^*\text{-}\mathcal{M}\text{eas}, \mathcal{B}\text{or}_{\mathbb{R}})$ -measurable. By Proposition 2.16, we know that $\mathfrak{A}$ is a σ -subalgebra of $\mu^*\text{-}\mathcal{M}\text{eas}$, implying that any $(\mathfrak{A}, \mathcal{B}\text{or}_{\mathbb{R}})$ -measurable function is also a $(\mu^*\text{-}\mathcal{M}\text{eas}, \mathcal{B}\text{or}_{\mathbb{R}})$ -measurable function. The converse need not hold. That is, the function f need *not* also be $(\mathfrak{A}, \mathcal{B}\text{or}_{\mathbb{R}})$ -measurable. Regardless, each class in $L_1(X, \mathfrak{A}, \mu)$ *does* have an $(\mathfrak{A}, \mathcal{B}\text{or}_{\mathbb{R}})$ -measurable representative, as claimed in the following proposition, which is just a restatement of Proposition 3.22 in terms of the quotient space $L_1(X, \mathfrak{A}, \mu)$.

{a_measurable_representative}

Proposition 4.24. Let $(X, \mathfrak{A}, \mu)$ be a σ -finite measure space, where $\mathfrak{A}$ is a σ -algebra. If $\bar{f}^\mu \in L_1(X, \mathfrak{A}, \mu)$, then there is an $(\mathfrak{A}, \mathcal{B}\text{or}_{\mathbb{R}})$ -measurable function $g \in \mathcal{L}_1(X, \mathfrak{A}, \mu)$ such that $\bar{g}^\mu = \bar{f}^\mu$. That is, each class of $L_1(X, \mathfrak{A}, \mu)$ has an $(\mathfrak{A}, \mathcal{B}\text{or}_{\mathbb{R}})$ -measurable representative.

{absolutely_continuous}

Definition 4.25. Let ν and μ be signed measures on a σ -algebra $\mathfrak{A}$. The signed measure ν is **absolutely continuous** with respect to μ on $\mathfrak{A}$ if $|\mu|(A) = 0$ implies $\nu(A) = 0$ whenever $A \in \mathfrak{A}$. Denote this by $\nu \ll \mu$.

{abscont_properties}

Proposition 4.26. Let $(X, \mathfrak{A})$ be a measurable space, and let μ and ν be measures on $\mathfrak{A}$ with $\nu \ll \mu$.

1. If $A \subseteq X$ and $\mu^*(A) = 0$, then $\nu^*(A) = 0$.
2. If μ is σ -finite, then $\mu^*\text{-}\mathcal{M}\text{eas} \subseteq \nu^*\text{-}\mathcal{M}\text{eas}$.

{indefinite_integral}

Definition 4.27. Let $(X, \mathfrak{A}, \mu)$ be a measure space, where $\mathfrak{A}$ is a σ -algebra. For every class $\bar{f}^\mu \in L_1(X, \mathfrak{A}, \mu)$ there is a finite signed measure $\nu : \mathfrak{A} \rightarrow \mathbb{R}$ on the measurable space $(X, \mathfrak{A})$ defined by

$$\nu(A) = \int_A f d\mu \quad \text{for all } A \in \mathfrak{A}.$$

This finite signed measure ν is absolutely continuous with respect to μ on $\mathfrak{A}$. Call ν the **indefinite integral** of $\bar{f}^\mu$. The idea that an integrable function defines a measure that is absolutely continuous with respect to some measure has the following form of converse.

{radon-nikodym}

Theorem 4.28 (Radon-Nikodym). *Let $(X, \mathfrak{A})$ be a measurable space. If ν is a finite signed measure on $\mathfrak{A}$ that is absolutely continuous with respect to a σ -finite measure μ on $\mathfrak{A}$, then there exists a unique class $\bar{f}^\mu$ in $L_1(X, \mathfrak{A}, \mu)$ such that*

$$\nu(A) = \int_A f d\mu \quad \text{for all } A \in \mathfrak{A}.$$

Furthermore, we may suppose the representative function f is $(\mathfrak{A}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable.

{density_function}

Definition 4.29. The unique class $\bar{f}^\mu \in L_1(X, \mathfrak{A}, \mu)$ in the previous theorem is called the **Radon-Nikodym derivative** of ν with respect to μ , and may be denoted $d\nu/d\mu^\mu$. The $(\mathfrak{A}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable representative function f may be denoted $d\nu/d\mu$, and will be referred to as the **density function** of ν with respect to μ whenever having to say “a Radon-Nikodym derivative representative” becomes tiresome, which happens fairly quickly.

Sometimes, as in the case of Proposition 4.30, the notation $d\nu = f d\mu$ even makes sense.

{radon_nikodym_corollary}

Proposition 4.30. *Let $(X, \mathfrak{A})$ be a measurable space, and let ν be a finite measure on $\mathfrak{A}$ and μ be σ -finite measure on $\mathfrak{A}$ with $\nu \ll \mu$. Also let $\bar{f}^\mu = d\nu/d\mu^\mu \in L_1(X, \mathfrak{A}, \mu)$.*

1. *Let $A = \{x \in X : f(x) > 0\}$. If $B \in \nu^*\text{-}\mathfrak{M}eas$, then $A \cap B \in \mu^*\text{-}\mathfrak{M}eas$.*
2. *If the class $\bar{g}^\nu \in L_1(X, \mathfrak{A}, \nu)$, then the class $gf^\mu \in L_1(X, \mathfrak{A}, \mu)$, and*

$$\int_X g d\nu = \int_X gf d\mu,$$

in which case “ $d\nu = f d\mu$.”

Let us combine a change of variable with the previous proposition.

{combine}

Proposition 4.31. *Let $(X, \mathfrak{A})$ and $(Y, \mathfrak{B})$ be measurable spaces, and let μ be a finite measure on $\mathfrak{A}$. Also, let $f : X \rightarrow Y$ be $(\mathfrak{A}, \mathfrak{B})$ -measurable,*

and let $g : Y \rightarrow \mathbb{R}$ be integrable with respect to $\mu \circ f^\leftarrow$. If λ is a σ -finite measure on $\mathfrak{B}$ such that $\mu \circ f^\leftarrow \ll \lambda$, then, setting $p = d(\mu \circ f^\leftarrow)/d\lambda$,

$$\int_X (g \circ f) d\mu = \int_Y g \cdot p d\lambda.$$

{chain_rule}

Proposition 4.32. Let $(X, \mathfrak{A})$ be a measurable space, and let ν and μ be finite measures on $\mathfrak{A}$, and let λ be a σ -finite measure on $\mathfrak{A}$. If $\nu \ll \mu$ and $\mu \ll \lambda$, then $\nu \ll \lambda$ and

$$\frac{d\nu}{d\lambda} = \frac{d\nu}{d\mu} \frac{d\mu}{d\lambda}.$$

{induced_sigmaalgebra}

Definition 4.33. Let $(Y, \mathfrak{B})$ be a measurable space, and let $f : X \rightarrow Y$ be a function, where $f^\leftarrow : \mathfrak{B} \rightarrow \mathfrak{P}(X)$. Then $f^\leftarrow(\mathfrak{B}) := \{f^\leftarrow(B) : B \in \mathfrak{B}\}$ is a σ -subalgebra of the power set $\mathfrak{P}(X)$; it's the σ -algebra induced by f . The notation $\sigma(f)$ is also used.

The function $f : X \rightarrow Y$ is always $(f^\leftarrow(\mathfrak{B}), \mathfrak{B})$ -measurable by the very definition of what it means for a function to be measurable. Should a σ -subalgebra $\mathfrak{A}$ of $\mathfrak{P}(X)$ be specified, so that $(X, \mathfrak{A})$ is a measurable space, and should $f : X \rightarrow Y$ be $(\mathfrak{A}, \mathfrak{B})$ -measurable, then $f^\leftarrow(\mathfrak{B})$ is a σ -subalgebra of $\mathfrak{A}$. To indicate that $f : X \rightarrow Y$ is $(\mathfrak{A}, \mathfrak{B})$ -measurable, we might use the notation $f : (X, \mathfrak{A}) \rightarrow (Y, \mathfrak{B})$.

Notation 4.34. When it comes to commutative diagrams, you have to decide what it means to commute. In the following, there will be diagrams that commute up to sets of measure zero. For example, in the following diagram

$$\begin{array}{ccc} (X, \mathfrak{A}) & \xrightarrow{g} & (\mathbb{R}, \mathfrak{Bor}_{\mathbb{R}}) \\ & \searrow f & \uparrow h \\ & & (Y, \mathfrak{B}) \end{array}$$

if f and h are functions, and if g is a representative of an $L_1(X, \mathfrak{A}, \mu)$ class, then what does it mean to say that the diagram commutes? If we

mean that it commutes up to subsets of $\mathfrak{A}$ having μ^* -measure zero, or equivalently that $\bar{g}^\mu = \overline{h \circ f^\mu}$, then we will qualify the diagram, saying just that:

$$\begin{array}{ccc} (X, \mathfrak{A}) & \xrightarrow{g} & (\mathbb{R}, \mathfrak{Bor}_{\mathbb{R}}) \\ & \searrow f & \uparrow h \\ & & (Y, \mathfrak{B}) \end{array} \quad \text{meaning } \bar{g}^\mu = \overline{h \circ f^\mu}.$$

Remark 4.35. Although the following proposition looks like the Doob-Dynkin lemma, it is not simply about measurability. It's also about a form of converse to Proposition 4.13.

The double-headed surjective arrow ' $\twoheadrightarrow$ ' indicates that a function is surjective.

{factoring_functions}

Proposition 4.36. Let $(X, \mathfrak{A}, \mu)$ be a finite measure space, where $\mathfrak{A}$ is a σ -algebra, and let $(Y, \mathfrak{B})$ be a measurable space. Also let $f : X \rightarrow Y$ be an $(\mathfrak{A}, \mathfrak{B})$ -measurable function such that $f^{\leftarrow} : \mathfrak{B} \twoheadrightarrow \mathfrak{A}$ is surjective. Let $\bar{g}^\mu \in L_1(X, \mathfrak{A}, \mu)$, then there is a unique class $\bar{h}^\mu \in L_1(Y, \mathfrak{B}, \mu \circ f^{\leftarrow})$ with $\bar{g}^\mu = \overline{h \circ f^\mu}$ and with h being a $(\mathfrak{B}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable function:

$$\begin{array}{ccc} (X, \mathfrak{A}) & \xrightarrow{g} & (\mathbb{R}, \mathfrak{Bor}_{\mathbb{R}}) \\ & \searrow f & \uparrow \text{---} h \\ & & (Y, \mathfrak{B}) \end{array} \quad \text{meaning } \bar{g}^\mu = \overline{h \circ f^\mu}$$

Proof. The idea is to use the Radon-Nikodym Theorem to get the class $\bar{h}^{\mu \circ f^{\leftarrow}} \in L_1(Y, \mathfrak{B}, \mu \circ f^{\leftarrow})$ with $\bar{g}^\mu = \overline{h \circ f^\mu}$.

Let ϕ denote the indefinite integral of g with respect to μ . That is,

$$\phi(A) = \int_A g \, d\mu \quad \text{for all } A \in \mathfrak{A}.$$

The set function $\phi \circ f^{\leftarrow} : \mathfrak{B} \rightarrow [0, \infty]$ defines a finite measure that is σ -additive and absolutely continuous with respect to the finite (and

therefore σ -finite) measure $\mu \circ f^{\leftarrow}$ on $(Y, \mathfrak{B})$. By the Radon-Nikodym theorem (4.28), there is a unique class $\bar{h}^{\mu \circ f^{\leftarrow}} \in L_1(Y, \mathfrak{B}, \mu \circ f^{\leftarrow})$ such that

$$(\phi \circ f^{\leftarrow})(B) = \int_B h d(\mu \circ f^{\leftarrow}) \quad \text{for all } B \in \mathfrak{B},$$

and with h a $(\mathfrak{B}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable function. On the one hand, by definition,

$$(\phi \circ f^{\leftarrow}(B)) = \phi(f^{\leftarrow}(B)) = \int_{f^{\leftarrow}(B)} g d\mu.$$

On the other hand, by Proposition 4.13,

$$\int_B h d(\mu \circ f^{\leftarrow}) = \int_{f^{\leftarrow}(B)} h \circ f d\mu.$$

By transitivity of equality, it follows that

$$\int_{f^{\leftarrow}(B)} g d\mu = \int_{f^{\leftarrow}(B)} h \circ f d\mu \quad \text{for all } B \in \mathfrak{B}.$$

The assumption that $f^{\leftarrow}$ is surjective means that $f^{\leftarrow}(\mathfrak{B}) = \mathfrak{A}$, and so

$$\int_A g d\mu = \int_A h \circ f d\mu \quad \text{for all } A \in \mathfrak{A}.$$

By Proposition 4.9, this implies that $\bar{g}^{\mu} = \overline{h \circ f}^{\mu}$, completing the proof. ■

For comparison with Proposition 4.36, let us look at the so-called Doob-Dynkin lemma.

{doob_dynkin}

Proposition 4.37 (Doob-Dynkin). *Let the space $(Y, \mathfrak{B})$ be measurable, and let $g : X \rightarrow \mathbb{R}$, and $f : X \rightarrow Y$. The function g is $(f^{\leftarrow}(\mathfrak{B}), \mathfrak{Bor}_{\mathbb{R}})$ -measurable if and only if there is a $(\mathfrak{B}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable function $h : Y \rightarrow \mathbb{R}$ such that $g = h \circ f$:*

4 Function Spaces and Quotient Spaces

$$\begin{array}{ccc}
 (X, f^{\leftarrow}(\mathfrak{B})) & \xrightarrow{g} & (\mathbb{R}, \mathfrak{Bor}_{\mathbb{R}}) \\
 & \searrow f & \uparrow \hat{h} \\
 & & (Y, \mathfrak{B})
 \end{array}$$

5 Product Measures and Product Integration

THROUGHOUT this section, let $(X, \mathfrak{S}, \mu)$ and $(Y, \mathfrak{T}, \nu)$ be measure spaces, so $\mathfrak{S}$ and $\mathfrak{T}$ are at least semirings. The powerset of any set A is still denoted $\mathfrak{P}(A)$. And $\mu^*\text{-}\mathfrak{M}\text{eas}$ still denotes the σ -algebra of μ^* -measurable subsets of X . As stated in Definition 2.9, we may let μ^* also denote a restriction $\mu^*|$.

5.1 Product Measures

{product_semiring}

Definition 5.1. If $\mathfrak{S}$ and $\mathfrak{T}$ are semirings of subsets of respective sets X and Y , then the collection of Cartesian products

$$\{A \times B \subseteq X \times Y : A \in \mathfrak{S}, B \in \mathfrak{T}\}$$

forms a semiring of subsets of $X \times Y$, generated in the first component by $\mathfrak{S}$ and in the second component by $\mathfrak{T}$, and will consequently be denoted $\mathfrak{S}\text{emi}(\mathfrak{S}, \mathfrak{T})$. It may be called the **product semiring** of $\mathfrak{S}$ and $\mathfrak{T}$, however, verify that the semiring $\mathfrak{S}\text{emi}(\mathfrak{S}, \mathfrak{T})$ is a subset of $\mathfrak{P}(X \times Y)$, and it is not the same as the Cartesian product $\mathfrak{S} \times \mathfrak{T}$, which is a subset of $\mathfrak{P}(X) \times \mathfrak{P}(Y)$. Semirings are non-empty; they have at least one element.

The idea of a product semiring generalizes in a clear way to the product of more than two factors if we make an identification. The Cartesian product is not associative, so we treat $(A \times B) \times C$ and $A \times (B \times C)$ as equivalent, and may (co)universally write $A \times B \times C$. It means that if $\mathfrak{U}$ were a semiring of subsets of a set Z , then expressions like $\mathfrak{S}\text{emi}(\mathfrak{S}, \mathfrak{T}, \mathfrak{U})$, under the identification, are well-defined.

{product_measure}

Proposition 5.2. *The set function $\mu \times \nu$ defined by*

$$\mu \times \nu : \mathfrak{Semi}(\mathfrak{S}, \mathfrak{T}) \rightarrow [0, \infty] : A \times B \mapsto \mu(A) \cdot \nu(B)$$

is a measure, if the symbols $0 \cdot \infty = 0$ make sense.

{product_measure_space}

Proposition 5.3. *If $(X, \mathfrak{S}, \mu)$ and $(Y, \mathfrak{T}, \nu)$ are measure spaces, then $(X \times Y, \mathfrak{Semi}(\mathfrak{S}, \mathfrak{T}), \mu \times \nu)$ is also a measure space. And if $(X, \mathfrak{S}, \mu)$ and $(Y, \mathfrak{T}, \nu)$ are σ -finite, then $(X \times Y, \mathfrak{Semi}(\mathfrak{S}, \mathfrak{T}), \mu \times \nu)$ is also σ -finite.*

Remark 5.4. By Proposition 2.14, the triples

$$(X, \mu^*-\mathfrak{Meas}, \mu^*) \quad \text{and} \quad (Y, \nu^*-\mathfrak{Meas}, \nu^*)$$

are measure spaces. By Proposition 5.3, the triple

$$(X \times Y, \mathfrak{Semi}(\mu^*-\mathfrak{Meas}, \nu^*-\mathfrak{Meas}), \mu^* \times \nu^*)$$

is also measure space.

{outer_product_measure}

Proposition 5.5. *If $A \in \mu^*-\mathfrak{Meas}$ and $B \in \nu^*-\mathfrak{Meas}$ with $\mu^*(A) < \infty$ and $\nu^*(B) < \infty$, then*

$$\begin{aligned} (\mu \times \nu)^*(A \times B) &= (\mu^* \times \nu^*)(A \times B) \\ &= \mu^*(A) \cdot \nu^*(B). \end{aligned}$$

{semiring_inclusion}

Proposition 5.6. *If $A \in \mu^*-\mathfrak{Meas}$ and $B \in \nu^*-\mathfrak{Meas}$, then $A \times B \in (\mu \times \nu)^*-\mathfrak{Meas}$. That is, $\mathfrak{Semi}(\mu^*-\mathfrak{Meas}, \nu^*-\mathfrak{Meas}) \subseteq (\mu \times \nu)^*-\mathfrak{Meas}$.*

{agreeable_product_measures}

Proposition 5.7. *If the measure spaces $(X, \mathfrak{S}, \mu)$ and $(Y, \mathfrak{T}, \nu)$ are both σ -finite, then the measure $\mu^* \times \nu^*$ agrees with $(\mu \times \nu)^*$ on the semiring $\mathfrak{Semi}(\mu^*-\mathfrak{Meas}, \nu^*-\mathfrak{Meas})$.*

Proof. We will apply Proposition 2.21.

The measure space $(X \times Y, \mathfrak{Semi}(\mathfrak{S}, \mathfrak{T}), \mu \times \nu)$ is σ -finite by Proposition 5.3. Since $\mathfrak{S} \subseteq \mu^*-\mathfrak{Meas}$ and $\mathfrak{T} \subseteq \nu^*-\mathfrak{Meas}$, it follows that

$$\mathfrak{Semi}(\mathfrak{S}, \mathfrak{T}) \subseteq \mathfrak{Semi}(\mu^*-\mathfrak{Meas}, \nu^*-\mathfrak{Meas}),$$

and by Proposition 5.6, it follows that

$$\mathfrak{Semi}(\mu^*\text{-}\mathcal{M}\text{eas}, \nu^*\text{-}\mathcal{M}\text{eas}) \subseteq (\mu \times \nu)^*\text{-}\mathcal{M}\text{eas}.$$

By Proposition 5.3, the set function $\mu^* \times \nu^*$ is a measure on the semiring $\mathfrak{Semi}(\mu^*\text{-}\mathcal{M}\text{eas}, \nu^*\text{-}\mathcal{M}\text{eas})$. Further, the measure $\mu^* \times \nu^*$ agrees with the measure $\mu \times \nu$ on the semiring $\mathfrak{Semi}(\mathfrak{S}, \mathfrak{T})$ for if $A \in \mathfrak{S}$ and $B \in \mathfrak{T}$, then

$$\begin{aligned} (\mu^* \times \nu^*)(A \times B) &= \mu^*(A) \cdot \nu^*(B) && \text{Proposition 5.2} \\ &= \mu(A) \cdot \nu(B) && \text{Proposition 2.11} \\ &= (\mu \times \nu)(A \times B) && \text{Proposition 5.2 again} \end{aligned}$$

Since $\mu^* \times \nu^* = \mu \times \nu$ on $\mathfrak{Semi}(\mathfrak{S}, \mathfrak{T})$, it follows by Proposition 2.21 that $\mu^* \times \nu^* = (\mu \times \nu)^*$ on $\mathfrak{Semi}(\mu^*\text{-}\mathcal{M}\text{eas}, \nu^*\text{-}\mathcal{M}\text{eas})$, completing the proof. ■

{product_sigmaalgebra}

Definition 5.8. Let $\mathfrak{S}$ and $\mathfrak{T}$ be semirings of subsets of respective sets X and Y . The σ -algebra generated by $\mathfrak{Semi}(\mathfrak{S}, \mathfrak{T})$ may be called the **product σ -algebra** of $\mathfrak{S}$ and $\mathfrak{T}$, and will be denoted $\sigma(\mathfrak{S}, \mathfrak{T})$. It is the smallest σ -algebra of $\mathfrak{P}(X \times Y)$ which contains the product semiring $\mathfrak{Semi}(\mathfrak{S}, \mathfrak{T})$. The notation $\mathfrak{S} \otimes \mathfrak{T}$ might also be used, but not here. The notation $\mathfrak{S} \vee \mathfrak{T}$ is compelling.

Remark 5.9. A consequence of the identification of $A \times B \times C$ with $(A \times B) \times C$ and $A \times (B \times C)$ is that

$$\sigma(\mathfrak{A}, \mathfrak{B}, \mathfrak{C}) = \sigma(\sigma(\mathfrak{A}, \mathfrak{B}), \mathfrak{C}) = \sigma(\mathfrak{A}, \sigma(\mathfrak{B}, \mathfrak{C})).$$

{extension_to_sigmaalgebra}

Notation 5.10. Let us apply Proposition 2.21 to the inclusion

$$\mathfrak{Semi}(\mathfrak{S}, \mathfrak{T}) \subseteq \sigma(\mathfrak{S}, \mathfrak{T}) \subseteq (\mu \times \nu)^*\text{-}\mathcal{M}\text{eas}.$$

The proposition says that if a measure on $\sigma(\mathfrak{S}, \mathfrak{T})$ agrees with $\mu \times \nu$ on $\mathfrak{Semi}(\mathfrak{S}, \mathfrak{T})$, then that measure agrees with $(\mu \times \nu)^*$ on $\sigma(\mathfrak{S}, \mathfrak{T})$. We will let $(\mu \times \nu)^*$ denote both the measure $(\mu \times \nu)^*$ on $(\mu \times \nu)^*\text{-}\mathcal{M}\text{eas}$ and its

restriction $(\mu \times \nu)^*|_{\sigma(\mathfrak{S}, \mathfrak{T})}$ to $\sigma(\mathfrak{S}, \mathfrak{T})$, but we will always make the domain clear.

$$\begin{array}{ccc} (\mu \times \nu)^* \mathfrak{Meas} & \xrightarrow{(\mu \times \nu)^*} & [0, \infty] \\ \cup & & \\ \sigma(\mathfrak{S}, \mathfrak{T}) & \xrightarrow{(\mu \times \nu)^*} & [0, \infty] \\ \cup & & \\ \mathfrak{Semi}(\mathfrak{S}, \mathfrak{T}) & \xrightarrow{\mu \times \nu} & [0, \infty] \end{array}$$

5.2 Product Integration

{projections}

Notation 5.11. Let π_1 and π_2 denote the respective **coordinate projections** of $X \times Y$ onto X and Y . That is, $\pi_1(x, y) = x$ and $\pi_2(x, y) = y$.

{sections}

Definition 5.12. Let $E \subseteq X \times Y$, and let $x \in X$ and $y \in Y$. Define sets

$$E_x := \pi_2(E \cap \pi_1^{\leftarrow}\{x\}) = \{y \in Y : (x, y) \in E\} \subseteq Y,$$

and

$$E^y := \pi_1(E \cap \pi_2^{\leftarrow}\{y\}) = \{x \in X : (x, y) \in E\} \subseteq X.$$

{measurable_sections}

The set E_x is the x -**section** of E , and the set E^y is the y -**section** of E .

Proposition 5.13. Let $(X, \mathfrak{A})$ and $(Y, \mathfrak{B})$ be measurable spaces. If $E \subseteq \sigma(\mathfrak{A}, \mathfrak{B})$, then $E_x \in \mathfrak{B}$ for all $x \in X$, and $E^y \in \mathfrak{A}$ for all $y \in Y$.

Notation 5.14. Whenever X , Y , and Z are sets and $f : X \times Y \rightarrow Z$ is a function, we will let f^y denote the map

$$f^y : X \rightarrow Z : x \mapsto f(x, y),$$

and we will let f_x denote the map

$$f_x : Y \rightarrow Z : y \mapsto f(x, y).$$


Also, for each $y \in Y$ let fix^y denote the map

$$\text{fix}^y : X \rightarrow X \times Y : x \mapsto (x, y)$$

and for each $x \in X$ let fix_x denote the map

$$\text{fix}_x : Y \rightarrow X \times Y : y \mapsto (x, y)$$

as illustrated:



Proposition 5.15. Let $(X, \mathfrak{A})$, $(Y, \mathfrak{B})$, and $(Z, \mathfrak{C})$ be measurable spaces. Let $f : X \times Y \rightarrow Z$. If f is $(\sigma(\mathfrak{A}, \mathfrak{B}), \mathfrak{C})$ -measurable, then f^y is $(\mathfrak{A}, \mathfrak{C})$ -measurable for each $y \in Y$, and f_x is $(\mathfrak{B}, \mathfrak{C})$ -measurable for each $x \in X$.

{measurable_out_of_product}

Proposition 5.16. Let $(X, \mathfrak{A})$, $(Y, \mathfrak{B})$, and $(Z, \mathfrak{C})$ be measurable spaces. Let $g : Z \rightarrow X \times Y$. Then the function g is $(\mathfrak{C}, \sigma(\mathfrak{A}, \mathfrak{B}))$ -measurable if and only if both $\pi_1 \circ g : Z \rightarrow X$ is $(\mathfrak{C}, \mathfrak{A})$ -measurable and $\pi_2 \circ g : Z \rightarrow Y$ is $(\mathfrak{C}, \mathfrak{B})$ -measurable.

{measurable_into_product}

Definition 5.17. Let $E \subseteq X \times Y$. Define the **section functions**

{section_functions}

$$\text{Sec}_X E : X \rightarrow \mathfrak{P}(Y) : x \mapsto E_x,$$

and

$$\text{Sec}_Y E : Y \rightarrow \mathfrak{P}(X) : y \mapsto E^y.$$

See that $\nu^* \circ \text{Sec}_X E : X \rightarrow \mathbb{R}$ and $\mu^* \circ \text{Sec}_Y E : Y \rightarrow \mathbb{R}$.

Proposition 5.18. Let $E \in (\mu \times \nu)^* \text{-Meas}$ with $(\mu \times \nu)^*(E) < \infty$. For μ^* -almost all $x \in X$, the set E_x is ν^* -measurable. The function $\nu^* \circ \text{Sec}_X E$ defines an integrable function over X with respect to μ , and

$$(\mu \times \nu)^*(E) = \int_X (\nu^* \circ \text{Sec}_X E) d\mu.$$

Likewise, for ν^* -almost all $y \in Y$, the set E^y is μ^* -measurable. The function $\mu^* \circ \text{Sec}^Y E$ defines an integrable function over Y with respect to ν , and

$$(\mu \times \nu)^*(E) = \int_Y (\mu^* \circ \text{Sec}^Y E) d\nu.$$

{product_integration}

Definition 5.19. Let $f : X \times Y \rightarrow \mathbb{R}$. Say that the **iterated integral** $\int_Y \int_X f d\mu d\nu$ **exists** if, for ν^* -almost all $y \in Y$, the function

$$f^y : X \rightarrow \mathbb{R} : x \mapsto f(x, y)$$

is integrable over X with respect to μ , and the function

$$g : Y \rightarrow \mathbb{R} : y \mapsto \int_X f^y d\mu$$

defines an integrable function over Y with respect to ν . In this case, define

$$\int_Y \int_X f d\mu d\nu := \int_Y \left(\int_X f^y d\mu \right) d\nu = \int_Y g d\nu$$

{fubini}

Theorem 5.20 (Fubini). Let $(X, \mathfrak{S}, \mu)$ and $(Y, \mathfrak{T}, \nu)$ both be measure spaces. If $f : X \times Y \rightarrow \mathbb{R}$ is a $\mu \times \nu$ -integrable function, then both of the iterated integrals $\int_Y \int_X f d\mu d\nu$ and $\int_X \int_Y f d\nu d\mu$ exist, and

$$\int_{X \times Y} f d(\mu \times \nu) = \int_Y \int_X f d\mu d\nu = \int_X \int_Y f d\nu d\mu.$$

{tonelli}

Theorem 5.21 (Tonelli). Let $(X, \mathfrak{S}, \mu)$ and $(Y, \mathfrak{T}, \nu)$ both be σ -finite measure spaces, and let $f : X \times Y \rightarrow \mathbb{R}$ be a $(\mu \times \nu)^*$ -measurable function. If one of the iterated integrals $\int_Y \int_X |f| d\mu d\nu$ or $\int_X \int_Y |f| d\nu d\mu$ exists, then the function f is $\mu \times \nu$ -integrable. Consequently,

$$\int_{X \times Y} f d(\mu \times \nu) = \int_Y \int_X f d\mu d\nu = \int_X \int_Y f d\nu d\mu.$$

Remark 5.22. It has been said that, “It is a difficult problem to determine whether a given function $f : X \times Y \rightarrow \mathbb{R}$ is $\mu \times \nu$ -measurable.” Maybe a slight understatement.

5.3 Applications


Definition 5.23. Let $(X, \mathfrak{A})$ and $(Y, \mathfrak{B})$ be measurable spaces. Let Γ be any measure on $\sigma(\mathfrak{A}, \mathfrak{B})$. Set

$$\alpha = \Gamma \circ \pi_1^{\leftarrow} \quad \text{on } \mathfrak{A}$$

and set

$$\beta = \Gamma \circ \pi_2^{\leftarrow} \quad \text{on } \mathfrak{B}.$$

Call Γ the **joint** measure. Call the induced measures α and β the **marginal** measures, as illustrated:



The marginal measures are determined by the joint, meaning that

$$\alpha(A) = \Gamma(A \times Y) \quad \text{for all } A \in \mathfrak{A}$$

and

$$\beta(B) = \Gamma(X \times B) \quad \text{for all } B \in \mathfrak{B}$$

since $\pi_1^{\leftarrow}(A) = A \times Y$ and $\pi_2^{\leftarrow}(B) = X \times B$.

In case $(X, \mathfrak{A}, \mu)$ and $(Y, \mathfrak{B}, \nu)$ are already measure spaces, we might refer to the measures μ and ν as **reference** measures, with the understanding that the reference measures are usually not the same as the marginal measures.

Remark 5.24. In the remaining propositions of this section, the only assumptions we will make about the joint measure Γ on $\sigma(\mathfrak{A}, \mathfrak{B})$ will be that it is a finite measure, and that it is absolutely continuous with respect to various product measures.

Proposition 5.25. *Let $(X, \mathfrak{A})$ and $(Y, \mathfrak{B})$ be measurable spaces, and let μ and ν be σ -finite measures on respective σ -algebras $\mathfrak{A}$ and $\mathfrak{B}$. Let Γ be a finite measure on $\sigma(\mathfrak{A}, \mathfrak{B})$. If $\Gamma \ll (\mu \times \nu)^*$ on $\sigma(\mathfrak{A}, \mathfrak{B})$, then $\alpha \ll \mu$ on $\mathfrak{A}$ and $\beta \ll \nu$ on $\mathfrak{B}$. In particular,*

$$\frac{d\alpha}{d\mu} : X \rightarrow \mathbb{R} : x \mapsto \int_Y \frac{d\Gamma}{d(\mu \times \nu)^*}(x, y) d\nu(y)$$

and

$$\frac{d\beta}{d\nu} : Y \rightarrow \mathbb{R} : y \mapsto \int_X \frac{d\Gamma}{d(\mu \times \nu)^*}(x, y) d\mu(x).$$

Proof. Set $f = d\Gamma/d(\mu \times \nu)^*$. We will only show that the Radon-Nikodym derivative $d\alpha/d\mu$ is the map $X \rightarrow \mathbb{R} : x \mapsto \int_Y f(x, y) d\nu(y)$. This is just an application of Fubini's theorem.

The function f is supposed to be integrable over $X \times Y$ with respect to $\mu \times \nu$. It follows by Fubini's Theorem 5.20 that the function

$$f_x : Y \rightarrow \mathbb{R} : y \mapsto f(x, y)$$

is integrable over Y with respect to ν for μ^* -almost all $x \in X$, and the function

$$j : X \rightarrow \mathbb{R} : x \mapsto \int_Y f_x d\nu = \int_Y f(x, y) d\nu(y)$$

defines an integrable function over X with respect to μ , with

$$\int_{X \times Y} f d(\mu \times \nu) = \int_X j d\mu = \int_X \left(\int_Y f(x, y) d\nu(y) \right) d\mu(x).$$

For each set $A \in \mathfrak{A}$, we know

$$\alpha(A) = \Gamma \circ \pi_1^+(A) = \Gamma(A \times Y).$$

By hypothesis,

$$\Gamma(A \times Y) = \int_{A \times Y} f d(\mu \times \nu).$$

By Fubini's,

$$\int_{A \times Y} f d(\mu \times \nu) = \int_A j d\mu.$$

By transitivity of equality,

$$\alpha(A) = \int_A j d\mu \quad \text{for all } A \in \mathfrak{A},$$

and this says $\alpha \ll \mu$ on $\mathfrak{A}$ with $d\alpha/d\mu = j$. ■

{reference_implies_mixed_and_marginals}

Proposition 5.26. *Let $(X, \mathfrak{A})$ and $(Y, \mathfrak{B})$ be measurable spaces, and let μ and ν be σ -finite measures on respective σ -algebras $\mathfrak{A}$ and $\mathfrak{B}$. Let Γ be a finite measure on $\sigma(\mathfrak{A}, \mathfrak{B})$. If $\Gamma \ll (\mu \times \nu)^*$ on $\sigma(\mathfrak{A}, \mathfrak{B})$, then $\Gamma \ll (\alpha \times \nu)^*$, and $\Gamma \ll (\mu \times \beta)^*$, and furthermore, $\Gamma \ll (\alpha \times \beta)^*$. In particular,*

$$\frac{d\Gamma}{d(\alpha \times \nu)^*} : X \times Y \rightarrow \mathbb{R} : (x, y) \mapsto \frac{\frac{d\Gamma}{d(\mu \times \nu)^*}(x, y)}{\frac{d\alpha}{d\mu}(x)},$$

$$\frac{d\Gamma}{d(\mu \times \beta)^*} : X \times Y \rightarrow \mathbb{R} : (x, y) \mapsto \frac{\frac{d\Gamma}{d(\mu \times \nu)^*}(x, y)}{\frac{d\beta}{d\nu}(y)},$$

and

$$\frac{d\Gamma}{d(\alpha \times \beta)^*} : X \times Y \rightarrow \mathbb{R} : (x, y) \mapsto \frac{\frac{d\Gamma}{d(\mu \times \nu)^*}(x, y)}{\frac{d\alpha}{d\mu}(x) \cdot \frac{d\beta}{d\nu}(y)}.$$

Proof. Let $\Gamma \ll (\mu \times \nu)^*$. Let $f = d\Gamma/d(\mu \times \nu)^*$ and let

$$j : X \rightarrow \mathbb{R} : x \mapsto \int_Y f_x d\nu$$

and

$$k : Y \rightarrow \mathbb{R} : y \mapsto \int_X f^y d\mu,$$

and define

$$g : X \times Y \rightarrow \mathbb{R} : (x, y) \mapsto \begin{cases} \frac{f(x, y)}{j(x)} & \text{if } j(x) > 0 \\ 1 & \text{if } j(x) = 0 \end{cases}$$

and

$$h : X \times Y \rightarrow \mathbb{R} : (x, y) \mapsto \begin{cases} \frac{f(x, y)}{j(x)k(y)} & \text{if } j(x)k(y) > 0 \\ 1 & \text{if } j(x)k(y) = 0 \end{cases}$$

The claim is that $g = d\Gamma/d(\alpha \times \nu)^*$ and $h = d\Gamma/d(\alpha \times \beta)^*$.

Let us now show that $\Gamma \ll (\alpha \times \nu)^*$. It is sufficient to show that $g = d\Gamma/d(\alpha \times \nu)^*$, meaning

$$\Gamma(C) = \int_C g d(\alpha \times \nu) \quad \text{for all } C \in \sigma(\mathfrak{A}, \mathfrak{B}).$$

In order to apply Tonelli's Theorem 5.21 to show that the function g is integrable with respect to $\alpha \times \nu$, we first show that the iterated integral $\int_X \int_Y g d\nu d\alpha$ exists.

We have assumed $f = d\Gamma/d(\mu \times \nu)^*$, which implies that we have assumed the function f is integrable with respect to $\mu \times \nu$. By Fubini's Theorem 5.20 then, the corresponding iterated integral $\int_X \int_Y f d\nu d\mu$ exists. This means that the function

$$f_x : Y \rightarrow \mathbb{R} : y \mapsto f(x, y)$$

is integrable over Y with respect to ν for μ^* -almost all $x \in X$, and the function $x \mapsto \int_Y f_x d\nu$ defines an integrable function over X with respect to μ . Let us carry this a little further. Since $\Gamma \ll (\mu \times \nu)^*$, it follows by Proposition 5.25 that $\alpha \ll \mu$ on $\mathfrak{A}$. We can then apply Proposition 4.26 which says that if $\mu^*(A) = 0$, then $\alpha^*(A) = 0$ for any $A \in \mathfrak{A}$. It means

the function f_x is integrable over Y with respect to ν for α^* -almost all $x \in X$. Since the function $Y \rightarrow \mathbb{R} : y \mapsto 1/j(x)$ is constant, it follows that the function

$$\frac{f_x}{j(x)} : Y \rightarrow \mathbb{R} : y \mapsto \frac{f(x, y)}{j(x)}$$

is also integrable over Y with respect to ν for α^* -almost all $x \in X$. See that the function

$$X \rightarrow \mathbb{R} : x \mapsto \int_Y \frac{f_x}{j(x)} d\nu$$

defines an integrable function over X with respect to α ; in fact,

$$\int_Y \frac{f_x}{j(x)} d\nu = 1,$$

and constant functions are integrable with respect to finite measures.

Consequently, for $C \in \sigma(\mathfrak{A}, \mathfrak{B})$, the function

$$Y \rightarrow \mathbb{R} : y \mapsto \frac{\chi_C(x, y) f_x(y)}{j(x)}$$

is also integrable over Y with respect to ν for α^* -almost all $x \in X$, and the function

$$X \rightarrow \mathbb{R} : x \mapsto \int_Y \frac{\chi_C(x, y) f_x(y)}{j(x)} d\nu(y)$$

defines an integrable function over X with respect to α . Applying Tonelli's Theorem 5.21 then,

$$\begin{aligned} \int_{X \times Y} \chi_C g d(\alpha \times \nu) &= \int_{X \times Y} \frac{\chi_C(x, y) f_x(y)}{j(x)} d(\alpha \times \nu)(x, y) \\ &= \int_Y \int_X \frac{\chi_C(x, y) f_x(y)}{j(x)} d\alpha(x) d\nu(y). \end{aligned}$$

By Proposition 4.30, we can change measures, after $d\alpha = j d\mu$:

$$\begin{aligned} \int_X \frac{\chi_C(x, y) f_x(y)}{j(x)} d\alpha(x) &= \int_X \frac{\chi_C(x, y) f_x(y)}{j(x)} \cdot j(x) d\mu(x) \\ &= \int_X \chi_C(x, y) f_x(y) d\mu(x), \end{aligned}$$

and so

$$\int_{X \times Y} \chi_C g d(\alpha \times \nu) = \int_Y \int_X \chi_C(x, y) f_x(y) d\mu(x) d\nu(y).$$

It follows that

$$\begin{aligned} \int_C g d(\alpha \times \nu) &= \int_{X \times Y} \chi_C g d(\alpha \times \nu) \\ &= \int_Y \int_X \chi_C(x, y) f_x(y) d\mu(x) d\nu(y) \\ &= \int_C f(x, y) d(\mu \times \nu) \\ &= \Gamma(C), \end{aligned}$$

as required.

Let us sketch the proof that $h = d\Gamma/d(\alpha \times \beta)^*$. We have shown that $\Gamma \ll (\mu \times \nu)^*$ implies $\Gamma \ll (\alpha \times \nu)^*$. Simply use the symmetric proof to show $\Gamma \ll (\alpha \times \nu)^*$ implies $h = d\Gamma/d(\alpha \times \beta)^*$. Use the fact shown earlier that g is integrable with respect to α and so

$$\int_X g(x, y) d\alpha(x) = \int_X \frac{f(x, y)}{j(x)} d\alpha(x) = \int_X \frac{f(x, y)}{j(x)} j(x) d\mu(x) = k(y)$$

to get

$$\int_X \frac{g^y}{k(y)} d\alpha = 1$$

in order to show that the function

$$Y \rightarrow \mathbb{R} : y \mapsto \int_X \frac{g^y}{k(y)} d\alpha$$

defines an integrable function over Y with respect to the finite measure β . And, for future reference:

$$\int_X \frac{g^y}{k(y)} d\alpha = \int_X \frac{\frac{d\Gamma}{d(\mu \times \nu)^*}(x, y)}{\frac{d\alpha}{d\mu}(x) \cdot \frac{d\beta}{d\nu}(y)} d\alpha(x).$$

■

{bayes_rule}

Proposition 5.27. *Let $(X, \mathfrak{A})$ and $(Y, \mathfrak{B})$ be measurable spaces, and let μ and ν be σ -finite measures on respective σ -algebras $\mathfrak{A}$ and $\mathfrak{B}$. Let Γ be a finite measure on $\sigma(\mathfrak{A}, \mathfrak{B})$. If $\Gamma \ll (\mu \times \nu)^*$ on $\sigma(\mathfrak{A}, \mathfrak{B})$, then*

$$\frac{d\Gamma}{d(\mu \times \nu)^*}(x, y) = \frac{\frac{d\Gamma}{d(\alpha \times \nu)^*}(x, y) \cdot \frac{d\alpha}{d\mu}(x)}{\int_X \frac{d\Gamma}{d(\alpha \times \nu)^*}(x, y) \cdot \frac{d\alpha}{d\mu}(x) d\mu(x)}.$$

Proof. This just combines Proposition 5.25 and Proposition 5.26.

It follows from Proposition 5.26 that

$$\frac{d\Gamma}{d(\mu \times \beta)^*}(x, y) = \frac{\frac{d\Gamma}{d(\alpha \times \nu)^*}(x, y) \cdot \frac{d\alpha}{d\mu}(x)}{\frac{d\beta}{d\nu}(y)}.$$

We will simply make a substitution for $d\beta/d\nu$. By Proposition 5.25,

$$\frac{d\beta}{d\nu}(y) = \int_X \frac{d\Gamma}{d(\mu \times \nu)^*}(x, y) d\mu(x)$$

and by Proposition 5.26,

$$\frac{d\Gamma}{d(\mu \times \nu)^*}(x, y) = \frac{d\Gamma}{d(\alpha \times \nu)^*}(x, y) \cdot \frac{d\alpha}{d\mu}(x),$$

so

$$\frac{d\beta}{d\nu}(y) = \int_X \frac{d\Gamma}{d(\alpha \times \nu)^*}(x, y) \cdot \frac{d\alpha}{d\mu}(x) d\mu(x),$$

as required.

■

{mixed_implies_marginal}

Proposition 5.28. *Let $(X, \mathfrak{A})$ and $(Y, \mathfrak{B})$ be measurable spaces, and let μ and ν be σ -finite measures on respective σ -algebras $\mathfrak{A}$ and $\mathfrak{B}$. Let*

Γ be a finite measure on $\sigma(\mathfrak{A}, \mathfrak{B})$. If $\Gamma \ll (\alpha \times \nu)^*$, then $\beta \ll \nu$ and $\Gamma \ll (\alpha \times \beta)^*$. In particular,

$$\frac{d\beta}{d\nu} : Y \mapsto \mathbb{R} : y \mapsto \int_X \frac{d\Gamma}{d(\alpha \times \nu)^*}(x, y) d\alpha(x),$$

and

$$\frac{d\Gamma}{d(\alpha \times \beta)^*} : X \times Y \rightarrow \mathbb{R} : (x, y) \mapsto \frac{\frac{d\Gamma}{d(\alpha \times \nu)^*}(x, y)}{\frac{d\beta}{d\nu}(y)}.$$

Likewise, if $\Gamma \ll (\mu \times \beta)^*$, then $\alpha \ll \mu$ and $\Gamma \ll (\alpha \times \beta)^*$. In particular,

$$\frac{d\alpha}{d\mu} : X \mapsto \mathbb{R} : x \mapsto \int_Y \frac{d\Gamma}{d(\mu \times \beta)^*}(x, y) d\beta(y),$$

and

$$\frac{d\Gamma}{d(\alpha \times \beta)^*} : X \times Y \rightarrow \mathbb{R} : (x, y) \mapsto \frac{\frac{d\Gamma}{d(\mu \times \beta)^*}(x, y)}{\frac{d\alpha}{d\mu}(x)}.$$

Proof. This is nothing but an application of Proposition 5.25 and Proposition 5.26. First set $\mu = \alpha$ and apply them, then set $\nu = \beta$ and apply them again, noting that α and β are both finite and therefore σ -finite. ■

{mixed_implies_miracle}

Proposition 5.29. Let $(X, \mathfrak{A})$ and $(Y, \mathfrak{B})$ be measurable spaces, and let ν be a σ -finite measure on $\mathfrak{B}$. Let Γ be a finite measure on $\sigma(\mathfrak{A}, \mathfrak{B})$. If $\Gamma \ll (\alpha \times \nu)^*$, then

$$\int_B \frac{d\Gamma}{d(\alpha \times \nu)^*}(x, y) d\nu(y) = \int_B \frac{d\Gamma}{d(\alpha \times \beta)^*}(x, y) d\beta(y)$$

for all $B \in \mathfrak{B}$.

Proof. This is just a corollary to the first half of Proposition 5.28 with a change of measure according to $d\beta = (d\beta/d\nu) d\nu$. ■

Proposition 5.30. *Let $(X, \mathfrak{A})$ and $(Y, \mathfrak{B})$ be measurable spaces, and let μ and ν be σ -finite measures on respective σ -algebras $\mathfrak{A}$ and $\mathfrak{B}$. Let Γ be a finite measure on $\sigma(\mathfrak{A}, \mathfrak{B})$. If $\Gamma \ll (\alpha \times \nu)^*$, then*

$$\int_A \frac{d\Gamma}{d(\alpha \times \beta)^*}(x, y) d\alpha(x) = \frac{\int_A \frac{d\Gamma}{d(\alpha \times \nu)^*}(x, y) d\alpha(x)}{\int_X \frac{d\Gamma}{d(\alpha \times \nu)^*}(x, y) d\alpha(x)}$$

for all $A \in \mathfrak{A}$.

Proof. This is the slightest possible re-statement of the first half of Proposition 5.28. ■

Proposition 5.31. *Let $(X, \mathfrak{A})$ and $(Y, \mathfrak{B})$ be measurable spaces, and let Γ be a finite measure on $\sigma(\mathfrak{A}, \mathfrak{B})$. If $\Gamma \ll (\alpha \times \beta)^*$, then*

$$X \rightarrow \mathbb{R} : x \mapsto \int_Y \frac{d\Gamma}{d(\alpha \times \beta)^*}(x, y) d\beta(y) = 1 \quad \alpha^* \text{-a.e. on } X,$$

and

$$Y \rightarrow \mathbb{R} : y \mapsto \int_X \frac{d\Gamma}{d(\alpha \times \beta)^*}(x, y) d\alpha(x) = 1 \quad \beta^* \text{-a.e. on } Y.$$

Proof. We will prove the first displayed equality. Let $\Gamma \ll (\alpha \times \beta)^*$. With Γ finite, it follows that $\alpha := \Gamma \circ \pi_1^+$ on $\mathfrak{A}$ and $\beta := \Gamma \circ \pi_2^+$ on $\mathfrak{B}$ are finite, therefore σ -finite. If we now set $\mu = \alpha$ and set $\nu = \beta$ and apply Proposition 5.25, then we get:

$$\frac{d\alpha}{d\alpha}(x) = \int_Y \frac{d\Gamma}{d(\alpha \times \beta)^*}(x, y) d\beta(y),$$

or equivalently,

$$\alpha(A) = \int_A \left(\int_Y \frac{d\Gamma}{d(\alpha \times \beta)^*}(x, y) d\beta(y) \right) d\alpha(x) \quad \text{for all } A \in \mathfrak{A}.$$

But also

$$\alpha(A) = \int_A 1 d\alpha(x) \quad \text{for all } A \in \mathfrak{A}.$$

By the uniqueness of the Radon-Nikodym derivative, which in this case is a class in $L_1(X, \mathfrak{A}, \alpha)$, it follows that

$$\int_Y \frac{d\Gamma}{d(\alpha \times \beta)^*}(x, y) d\beta(y) = 1 \quad \alpha^*\text{-almost everywhere,}$$

as required. ■

6 Probability, Independence, and Sampling

6.1 Probability Spaces and Distributions

Definition 6.1. A **probability space** is a measure space $(X, \mathfrak{A}, \mu)$ such that $\mathfrak{A}$ is a σ -algebra of subsets of a set X with $\mu(X) = 1$. Any measure ν on a measurable space $(X, \mathfrak{A})$ for which $\nu(X) = 1$ is called a **probability measure**. If $f : X \rightarrow \mathbb{R}^n$ is an $(\mathfrak{A}, \mathfrak{Bor}_{\mathbb{R}^n})$ -measurable function, then f might be called a **random variable**.

{probability_space}

Remark 6.2. When it comes to probability spaces, why change structure from measure spaces $(X, \mathfrak{S}, \mu)$, where $\mathfrak{S}$ is only assumed to be a semiring, to $(X, \mathfrak{A}, \mu)$, where $\mathfrak{A}$ is assumed to be a σ -algebra? One reason may be the use of *density* functions, which represent Radon-Nikodym derivatives. Recall that the Radon-Nikodym Theorem 4.28 applies to measures on a measurable space $(X, \mathfrak{A})$, in which case $\mathfrak{A}$ is supposed to be a σ -algebra. Density functions are described further in Notation 6.4.

{semiring_to_sigmaalgebra}

Even for probability spaces $(X, \mathfrak{A}, \mu)$ and $(Y, \mathfrak{B}, \nu)$, the measure $\mu \times \nu$ on the semiring $\mathfrak{Semi}(\mathfrak{A}, \mathfrak{B})$ need not be a measure on $\sigma(\mathfrak{A}, \mathfrak{B})$, unless possibly $\sigma(\mathfrak{A}, \mathfrak{B})$ happens to equal $\mathfrak{Semi}(\mathfrak{A}, \mathfrak{B})$. But its unique extension $(\mu \times \nu)^*$ restricts to a measure on $\sigma(\mathfrak{A}, \mathfrak{B})$. This was described in Notation 5.10, and is reproduced somewhat here: Let us apply Proposition 2.21 to the inclusion

$$\mathfrak{Semi}(\mathfrak{S}, \mathfrak{T}) \subseteq \sigma(\mathfrak{S}, \mathfrak{T}) \subseteq (\mu \times \nu)^*\text{-Meas}.$$

The proposition says that if a measure agrees with $\mu \times \nu$ on $\mathfrak{Semi}(\mathfrak{S}, \mathfrak{T})$, then that measure agrees with $(\mu \times \nu)^*$ on $\sigma(\mathfrak{S}, \mathfrak{T})$. We will let $(\mu \times \nu)^*$

denote both the measure $(\mu \times \nu)^*$ on $(\mu \times \nu)^*\text{-}\mathcal{M}\text{eas}$ and its restriction $(\mu \times \nu)^*|_{\sigma(\mathfrak{S}, \mathfrak{T})}$ to $\sigma(\mathfrak{S}, \mathfrak{T})$. We will always make the domain clear.

$$\begin{array}{ccc} (\mu \times \nu)^*\text{-}\mathcal{M}\text{eas} & \xrightarrow{(\mu \times \nu)^*} & [0, \infty] \\ \cup & & \\ \sigma(\mathfrak{S}, \mathfrak{T}) & \xrightarrow{(\mu \times \nu)^*} & [0, \infty] \\ \cup & & \\ \mathfrak{Semi}(\mathfrak{S}, \mathfrak{T}) & \xrightarrow{\mu \times \nu} & [0, \infty] \end{array}$$

{probability_distribution}

Definition 6.3. Let $(X, \mathfrak{A}, \mu)$ be a probability space, and let $(Y, \mathfrak{B})$ be a measurable space, so $\mathfrak{B}$ is supposed to be a σ -algebra. If a function $f : X \rightarrow Y$ is $(\mathfrak{A}, \mathfrak{B})$ -measurable, where $f^{\leftarrow} : \mathfrak{B} \rightarrow \mathfrak{A}$, then the induced probability measure $\mu \circ f^{\leftarrow}$ on the σ -algebra $\mathfrak{B}$ of the codomain of f , defined by

$$\mu \circ f^{\leftarrow} : \mathfrak{B} \rightarrow [0, 1] : B \mapsto \mu(f^{\leftarrow}(B)),$$

and illustrated

$$X \xrightarrow{f} Y \quad [0, 1] \xleftarrow{\mu \circ f^{\leftarrow}} \mathfrak{A} \xleftarrow{f^{\leftarrow}} \mathfrak{B}$$

is called the **probability distribution** of f on $\mathfrak{B}$, or simply the *probability distribution* of f . It is a probability measure composed with a preimage: $\mu \circ f^{\leftarrow}$.

{density_functions}

Notation 6.4. Probability distributions are probabilities induced on a σ -algebra of the codomain of a measurable function. The notation

$$f \sim \mu \circ f^{\leftarrow}$$

is used to denote the map from the measurable function to the probability distribution, although seldom so explicitly. For example, suppose an author writes

$$f \sim \text{Normal}(m, s),$$

or maybe something like $f \sim \mathcal{N}(m, s)$. There is an underlying probability space $(X, \mathfrak{A}, \mu)$, where $f : X \rightarrow \mathbb{R}$ is $(\mathfrak{A}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable, as illustrated:

$$X \xrightarrow{f} \mathbb{R} \quad [0, 1] \xleftarrow{\mu} \mathfrak{A} \xleftarrow{f^{\leftarrow}} \mathfrak{Bor}_{\mathbb{R}}$$

$\mu \circ f^{\leftarrow} = \text{Normal}(m, s)$

See that $f \sim \mu \circ f^{\leftarrow}$. That is, $\mu \circ f^{\leftarrow}$ is the probability distribution of f on $\mathfrak{Bor}_{\mathbb{R}}$. By writing $f \sim \text{Normal}(m, s)$, the author has indicated that the measures $\mu \circ f^{\leftarrow}$ and $\text{Normal}(m, s)$ are equal, which means

$$\begin{aligned} \mu \circ f^{\leftarrow}(A) &= \text{Normal}(m, s)(A) \\ &:= \int_A g(t) d\lambda(t) \quad \text{for all } A \in \mathfrak{Bor}_{\mathbb{R}}, \end{aligned}$$

where

$$g : \mathbb{R} \rightarrow \mathbb{R} : t \mapsto \frac{1}{s\sqrt{2\pi}} \exp \left\{ -\frac{1}{2} \left(\frac{t-m}{s} \right)^2 \right\},$$

and λ is Lebesgue measure. The function $g \in \mathcal{L}_1(\mathbb{R}, \mathfrak{Bor}_{\mathbb{R}}, \lambda)$ in this case represents the Radon-Nikodym derivative of the probability measure $\mu \circ f^{\leftarrow}$ with respect to Lebesgue measure λ on $\mathfrak{Bor}_{\mathbb{R}}$. Should an author write

$$\text{Normal}(f \mid m, s),$$

they are likely referring to the function g itself, and might call it the **density** of f .

If an author writes $\int g(t) dt$, then we assume the ‘ dt ’ means with respect to Lebesgue measure. If instead there is a summation sign and they use the term ‘probability function,’ or use the word ‘discrete,’ then they likely mean integration with respect to counting measure. In any case, with a density function it is always the Lebesgue integral with respect to some measure.

Remark 6.5. To be sure:

- A *density* is an integrable function that represents a Radon-Nikodym derivative of a finite measure with respect to a σ -finite measure.
- A *probability distribution* is a probability measure induced on a σ -algebra of the codomain of a measurable function.

There is no intelligible sense in which the terms *density* and *probability distribution* are interchangeable.

6.2 Expected Value

{expected_value}

Definition 6.6. For a class $\bar{f}^\mu$ in $L_1(X, \mathfrak{S}, \mu)$, the **expected value** $E \bar{f}^\mu$ of the class $\bar{f}^\mu$ is $\int f d\mu$. That is, the operator E is the linear functional defined by

$$E : L_1(X, \mathfrak{S}, \mu) \rightarrow \mathbb{R} : \bar{f}^\mu \mapsto \int_X f d\mu.$$

{expected_value_probability}

Remark 6.7. In the case that $(X, \mathfrak{A}, \mu)$ is a probability space, and the probability distribution $\mu \circ f^{\leftarrow}$ of f is absolutely continuous with respect to some σ -finite reference measure λ on $\mathfrak{B}\mathfrak{o}\mathfrak{t}_{\mathbb{R}}$, then, setting $p = d(\mu \circ f^{\leftarrow})/d\lambda$, the expected value of $\bar{f}^\mu$ can be computed by

$$\int_X f d\mu = \int_{\mathbb{R}} t p(t) d\lambda(t),$$

and to see this, let $g : \mathbb{R} \rightarrow \mathbb{R}$ be any function integrable with respect to $\mu \circ f^{\leftarrow}$. By Proposition 4.31,

$$\int_X (g \circ f) d\mu = \int_{\mathbb{R}} g p d\lambda.$$

Taking g to be the identity function $g(t) = t$, so that $g(f(x)) = f(x)$, see that

$$\int_X f d\mu = \int_{\mathbb{R}} t p(t) d\lambda(t).$$

6.3 Independence

Definition 6.8. Let $(X, \mathfrak{S}, \mu)$ be a measure space, and let $\mathfrak{T}$ and $\mathfrak{U}$ be subsets of the semiring $\mathfrak{S}$. Then $\mathfrak{T}$ and $\mathfrak{U}$ are **independent collections** of sets with respect to the measure μ if

$$\mu(A \cap B) = \mu(A) \cdot \mu(B) \quad \text{for all } A \in \mathfrak{T} \text{ and for all } B \in \mathfrak{U}.$$

The ‘ $\mathfrak{V}$ ’ symbol is a fraktur V, and ‘ $\mathfrak{W}$ ’ is fraktur W.

Proposition 6.9. If $\mathfrak{T}$ and $\mathfrak{U}$ are independent collections with $\mathfrak{V} \subseteq \mathfrak{T}$ and $\mathfrak{W} \subseteq \mathfrak{U}$, then $\mathfrak{V}$ and $\mathfrak{W}$ are independent collections.

{independent_functions}

Definition 6.10. Let $f : (X, \mathfrak{A}) \rightarrow (Y, \mathfrak{B})$ and $g : (X, \mathfrak{A}) \rightarrow (Z, \mathfrak{C})$ be measurable functions. Then f and g are **independent functions** with respect to a measure μ on $\mathfrak{A}$ if the σ -algebras $f^{\leftarrow}(\mathfrak{B})$ and $g^{\leftarrow}(\mathfrak{C})$ are independent with respect to μ , where $f^{\leftarrow} : \mathfrak{B} \rightarrow \mathfrak{A}$ and $g^{\leftarrow} : \mathfrak{C} \rightarrow \mathfrak{A}$. It means

$$\mu(f^{\leftarrow}(A) \cap g^{\leftarrow}(B)) = \mu(f^{\leftarrow}(A)) \cdot \mu(g^{\leftarrow}(B)) \quad \text{for all } A \in \mathfrak{B} \text{ and } B \in \mathfrak{C}.$$

{independent_compositions}

Proposition 6.11. If functions $h : (X, \mathfrak{A}) \rightarrow (Y, \mathfrak{B})$ and $k : (X, \mathfrak{A}) \rightarrow (Y, \mathfrak{B})$ are independent, and $g : (Y, \mathfrak{B}) \rightarrow (Z, \mathfrak{C})$ is measurable, then $g \circ h$ and $g \circ k$ are independent.

{independent_classes}

Definition 6.12. Let $(X, \mathfrak{S}, \mu)$ be a measure space, and let $\bar{f}^\mu$ and $\bar{g}^\mu$ be classes in $L_p(X, \mathfrak{S}, \mu)$. Then $\bar{f}^\mu$ and $\bar{g}^\mu$ are **independent classes** with respect to μ^* if the $(\mu^*, \mathfrak{Meas}, \mathfrak{Bot}_{\mathbb{R}})$ -measurable functions f and g are independent with respect to μ^* . Proposition 3.10 says that the independence of classes $\bar{f}^\mu$ and $\bar{g}^\mu$ does not depend on the respective chosen representative functions f and g , meaning if $h \in \bar{f}^\mu$ and $k \in \bar{g}^\mu$, then f and g are independent if and only if h and k are independent.

{expected_value_independent}

Proposition 6.13. If classes $\bar{f}^\mu$ and $\bar{g}^\mu$ in $L_1(X, \mathfrak{S}, \mu)$ are independent with respect to μ^* , and if $\bar{f}^\mu \bar{g}^\mu \in L_1(X, \mathfrak{S}, \mu)$, then

$$\mathbb{E}(\bar{f}^\mu \bar{g}^\mu) = (\mathbb{E} \bar{f}^\mu)(\mathbb{E} \bar{g}^\mu).$$

Proof. This is nothing more imaginative than wading through the definitions of independence, and showing that the equality holds for step functions. Then use the continuity of the integral by recalling that the equivalence classes of step functions are norm dense in $L_1(X, \mathfrak{G}, \mu)$. For example, let $f = a\chi_A + b\chi_B$ and $g = c\chi_C + d\chi_D$ be independent with respect to μ^* , where we suppose A, B, C , and D to be disjoint and μ^* -measurable sets of finite measure. Then

$$\begin{aligned} fg &= (a\chi_A + b\chi_B)(c\chi_C + d\chi_D) \\ &= ac\chi_{A \cap C} + ad\chi_{A \cap D} + bc\chi_{B \cap C} + bd\chi_{B \cap D}, \end{aligned}$$

and so

$$\begin{aligned} \int fg &= ac\mu^*(A \cap C) + ad\mu^*(A \cap D) + bc\mu^*(B \cap C) + bd\mu^*(B \cap D) \\ &= ac\mu^*(A)\mu^*(C) + ad\mu^*(A)\mu^*(D) + bc\mu^*(B)\mu^*(C) + bd\mu^*(B)\mu^*(D) \\ &= (a\mu^*(A) + b\mu^*(B))(c\mu^*(C) + d\mu^*(D)) \\ &= \int f \int g. \end{aligned}$$

■

6.4 Strong Law of Large Numbers

{large_numbers}

Theorem 6.14 (Strong Law of Large Numbers). *Let $(X, \mathfrak{A}, \mu)$ be a probability space, and let $\{f, f_1, f_2, \dots\}$ be a collection of functions in $\mathcal{L}_1(X, \mathfrak{A}, \mu)$. Define the subset*

$$A := \left\{ x \in X : \frac{(f_1 + \dots + f_n)(x)}{n} \rightarrow \int_X f d\mu \right\}.$$

If the f_i are independent and all have the same probability distribution as f , then $\mu(A) = 1$.

6.5 Sampling

There are some terms we will not even try to define for now. Terms like *draw*, or *sample*, or even *random*. Especially *random*.

Proposition 6.15. *Let $(X, \mathfrak{A}, \mu)$ be a probability space, and let $(Y, \mathfrak{B}, \nu)$ be σ -finite. Also let $\{h_i : X \rightarrow Y\}$ be a collection of $(\mathfrak{A}, \mathfrak{B})$ -measurable functions. If the h_i are independent, and if the probability distributions of the h_i all have the same density function f with respect to the measure ν , and if $g \in \mathcal{L}_1(Y, \mathfrak{B}, \mu \circ h_i^{\leftarrow})$, then the subset of X defined by*

{sampling}

$$A := \left\{ x \in X : \frac{1}{n} \sum_{i=1}^n (g \circ h_i)(x) \rightarrow \int_Y g f d\nu \right\}$$

is such that $\mu(A) = 1$.

Proof. Let the h_i be independent, and let the probability distributions $\mu \circ h_i^{\leftarrow}$ of the h_i all have the same density function f with respect to ν on $\mathfrak{B}$, and let $g \in \mathcal{L}_1(Y, \mathfrak{B}, \mu \circ h_i^{\leftarrow})$. We will use the strong law of large numbers to show that the μ -measure of the subset

$$\left\{ x \in X : \frac{1}{n} \sum_{i=1}^n (g \circ h_i)(x) \rightarrow \int_X (g \circ h_1) d\mu \right\}$$

is 1, and to complete the proof we will note that

$$\int_X (g \circ h_1) d\mu = \int_Y g f d\nu.$$

For the probability distributions $\mu \circ h_i^{\leftarrow}$ of the h_i to have the same density function f in $\mathcal{L}_1(Y, \mathfrak{B}, \nu)$ with respect to ν on $\mathfrak{B}$ means

$$(\mu \circ h_i^{\leftarrow})(B) = \int_B f d\nu \quad \text{for all } B \in \mathfrak{B},$$

as somewhat illustrated:

$$X \xrightarrow{h_i} Y \xrightarrow{f} \mathbb{R} \quad [0, 1] \xleftarrow{\mu} \mathfrak{A} \xleftarrow{h_i^{\leftarrow}} \mathfrak{B}$$

$\xleftarrow{\mu \circ h_i^{\leftarrow}}$

The collection $\{g \circ h_i\}$ of functions are independent with respect to μ by Proposition 6.11. By hypothesis $g \in \mathcal{L}_1(Y, \mathfrak{B}, \mu \circ h_i^{\leftarrow})$, and so by Proposition 4.13, each $g \circ h_i \in \mathcal{L}_1(X, \mathfrak{A}, \mu)$. To show that the collection $\{g \circ h_i\}$ of functions all have the same probability distribution as $g \circ h_1$, see first that the $\mu \circ h_i^{\leftarrow} = \mu \circ h_1^{\leftarrow}$ since by hypothesis the $\mu \circ h_i^{\leftarrow}$ all have the same density function with respect to λ :

$$\mu \circ h_i^{\leftarrow}(B) = \int_B f d\lambda = \mu \circ h_1^{\leftarrow}(B) \quad \text{for all } B \in \mathfrak{B}_{\text{or}\mathbb{R}}.$$

See next that the $\{g \circ h_i\}$ all have the same probability distribution as $g \circ h_1$ since

$$\begin{aligned} \mu \circ (g \circ h_i)^{\leftarrow} &= \mu \circ (h_i^{\leftarrow} \circ g^{\leftarrow}) \\ &= (\mu \circ h_i^{\leftarrow}) \circ g^{\leftarrow} \\ &= (\mu \circ h_1^{\leftarrow}) \circ g^{\leftarrow} \\ &= \mu \circ (g \circ h_1)^{\leftarrow}. \end{aligned}$$

Since the collection $\{g \circ h_i\}$ of $\mathcal{L}_1(X, \mathfrak{A}, \mu)$ functions are independent with respect to the measure μ , and they all have the same probability distribution as $g \circ h_1$, it follows by the strong law of large numbers, Theorem 6.14, that the subset

$$A := \left\{ x \in X : \frac{1}{n} \sum_{i=1}^n (g \circ h_i)(x) \rightarrow \int_X (g \circ h_1) d\mu \right\}$$

of X has $\mu(A) = 1$.

Finally, let us note that

$$\int_X (g \circ h_1) d\mu = \int_Y gf d\nu$$

by Proposition 4.31, which completes the proof. ■

7 Conditional Mathematical Expectation

CONDITIONAL mathematical expectation is not difficult to define, but there seems to be some subtlety in the proof of its existence, especially when it comes to applying the Radon-Nikodym theorem.

We are supposing that the $\mathfrak{A}$ in any probability space $(X, \mathfrak{A}, \mu)$ is a σ -algebra, as opposed to merely a semiring. The reason for supposing this is because the Radon-Nikodym theorem applies to measurable spaces, such as $(X, \mathfrak{A})$, in which case $\mathfrak{A}$ is certainly a σ -algebra.

{cme_sigma_algebra}

Definition 7.1. Let $(X, \mathfrak{A}, \mu)$ be a probability space, and let $\mathfrak{B}$ be a σ -subalgebra of $\mathfrak{A}$. For each class $\bar{f}$ in $L_1(X, \mathfrak{A}, \mu)$, the **conditional mathematical expectation** $\mathcal{E}_{\mathfrak{B}} \bar{f}$ of $\bar{f}$ with respect to $\mathfrak{B}$, or simply the *conditional expectation* of $\bar{f}$ with respect to $\mathfrak{B}$, is the unique class in $L_1(X, \mathfrak{A}, \mu)$ having a $(\mathfrak{B}, \mathfrak{B}\sigma\tau_{\mathbb{R}})$ -measurable representative function g such that

$$\int_B f d\mu = \int_B g d\mu \quad \text{for all } B \in \mathfrak{B}.$$

{idempotent}

Remark 7.2. Should a class $\bar{f}$ in $L_1(X, \mathfrak{A}, \mu)$ have a $(\mathfrak{B}, \mathfrak{B}\sigma\tau_{\mathbb{R}})$ -measurable representative, say the function f itself is $(\mathfrak{B}, \mathfrak{B}\sigma\tau_{\mathbb{R}})$ -measurable, then $\mathcal{E}_{\mathfrak{B}} \bar{f} = \bar{f}$ since trivially

$$\int_B f d\mu = \int_B f d\mu \quad \text{for all } B \in \mathfrak{B}.$$

And since by definition the class $\mathcal{E}_{\mathfrak{B}} \bar{f}$ does have a $(\mathfrak{B}, \mathfrak{B}\sigma\tau_{\mathbb{R}})$ -measurable representative function, it follows that $\mathcal{E}_{\mathfrak{B}}(\mathcal{E}_{\mathfrak{B}} \bar{f}) = \mathcal{E}_{\mathfrak{B}} \bar{f}$. Another way of saying this is that $\mathcal{E}_{\mathfrak{B}} \circ \mathcal{E}_{\mathfrak{B}} = \mathcal{E}_{\mathfrak{B}}$ on $L_1(X, \mathfrak{A}, \mu)$, or rather, the operator $\mathcal{E}_{\mathfrak{B}}$ is idempotent on $L_1(X, \mathfrak{A}, \mu)$.

existence_of_conditional}

Theorem 7.3. Let $(X, \mathfrak{A}, \mu)$ be a probability space, with $\bar{f}^\mu \in L_1(X, \mathfrak{A}, \mu)$. For every σ -subalgebra $\mathfrak{B}$ of $\mathfrak{A}$ there exists a unique class in $L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$ having a $(\mathfrak{B}, \mathfrak{B}\sigma\mathbb{R})$ -measurable representative function g such that

$$\int_B f d\mu = \int_B g d\mu|_{\mathfrak{B}} \quad \text{for all } B \in \mathfrak{B}.$$

Proof. The existence of such a class is an application of the Radon-Nikodym theorem, and to see this, let $\nu|_{\mathfrak{B}}$ denote the restriction of the indefinite integral ν of $\bar{f}^\mu$ to the σ -subalgebra $\mathfrak{B}$, which means that $\nu|_{\mathfrak{B}} : \mathfrak{B} \rightarrow \mathbb{R}$ is defined by

$$\nu|_{\mathfrak{B}}(B) = \int_B f d\mu \quad \text{for all } B \in \mathfrak{B}.$$

The set function $\nu|_{\mathfrak{B}}$ is a finite signed measure that is absolutely continuous with respect to the σ -finite measure $\mu|_{\mathfrak{B}}$ on $\mathfrak{B}$, and so by the Radon-Nikodym Theorem 4.28 there is a unique class $\bar{g}^{\mu|_{\mathfrak{B}}}$ in $L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$ such that

$$\nu|_{\mathfrak{B}}(B) = \int_B g d\mu|_{\mathfrak{B}} \quad \text{for all } B \in \mathfrak{B},$$

where the representative function g may be taken to be $(\mathfrak{B}, \mathfrak{B}\sigma\mathbb{R})$ -measurable, which completes the proof. ■

{cme_sigmaalgebra}

Definition 7.4. Let $(X, \mathfrak{A}, \mu)$ be a probability space, and let $\bar{f}^\mu \in L_1(X, \mathfrak{A}, \mu)$. By Theorem 7.3, for each σ -subalgebra $\mathfrak{B}$ of $\mathfrak{A}$ there is a unique class, which may be denoted by $E_{\mathfrak{B}} \bar{f}^\mu$ or $E(\bar{f}^\mu | \mathfrak{B})$, in $L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$ having a $(\mathfrak{B}, \mathfrak{B}\sigma\mathbb{R})$ -measurable representative such that

$$\int_B f d\mu = \int_B E_{\mathfrak{B}} \bar{f}^\mu d\mu|_{\mathfrak{B}} \quad \text{for all } B \in \mathfrak{B}.$$

The class $E_{\mathfrak{B}} \bar{f}^\mu$ is the **conditional expectation** of $\bar{f}^\mu$ with respect to the σ -subalgebra $\mathfrak{B}$. This defines an operator:

$$E_{\mathfrak{B}} : L_1(X, \mathfrak{A}, \mu) \rightarrow L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}}) : \bar{f}^\mu \mapsto E_{\mathfrak{B}} \bar{f}^\mu.$$

The fact that

$$\int_X \bar{f}^\mu d\mu = \int_X f d\mu = \int_X E_{\mathfrak{B}} \bar{f}^\mu d\mu|_{\mathfrak{B}}$$

implies that the operator $E_{\mathfrak{B}}$ has norm 1.

Notation 7.5. Let $(X, \mathfrak{A}, \mu)$ be a probability space. Let $(Y, \mathfrak{B})$ be a measurable space, and let $g : X \rightarrow Y$ be any function, where $g^\leftarrow : \mathfrak{B} \rightarrow \mathfrak{P}(X)$, with the qualification that in case no σ -algebra $\mathfrak{B}$ is supplied, we take $\mathfrak{B}$ to be the powerset $\mathfrak{P}(Y)$ of Y . Let $\mathfrak{A}^{(g)}$ denote the σ -subalgebra of $\mathfrak{B}$ consisting of those sets in $\mathfrak{B}$ whose preimages under g lie in $\mathfrak{A}$:

$$\mathfrak{A}^{(g)} = \{B \in \mathfrak{B} : g^\leftarrow(B) \in \mathfrak{A}\} \subseteq \mathfrak{B}.$$

The σ -subalgebra

$$\mathfrak{G} := g^\leftarrow(\mathfrak{B}) \cap \mathfrak{A}$$

is comprised of the preimages under g of the sets in $\mathfrak{A}^{(g)}$:

$$\mathfrak{G} = \{g^\leftarrow(B) : B \in \mathfrak{A}^{(g)}\} \subseteq \mathfrak{A}.$$

Consequently, the function g is $(\mathfrak{G}, \mathfrak{A}^{(g)})$ -measurable. Furthermore, the function defined by

$$g^\leftarrow : \mathfrak{A}^{(g)} \rightarrow \mathfrak{G} : C \mapsto g^\leftarrow(C)$$

is surjective.

In Definition 7.4, we looked at the conditional mathematical expectation of a class defined with respect to a σ -algebra. But a function can also define a σ -algebra, and we now look at that case.

{cme_function}

Definition 7.6. Let $(X, \mathfrak{A}, \mu)$ be a probability space, and let $\bar{f}^\mu \in L_1(X, \mathfrak{A}, \mu)$. Let $(Y, \mathfrak{B})$ be a measurable space, and let $g : X \rightarrow Y$ be any function, where $g^\leftarrow : \mathfrak{B} \rightarrow \mathfrak{P}(X)$. Set

$$\mathfrak{G} = g^\leftarrow(\mathfrak{B}) \cap \mathfrak{A}.$$

Then the class $E(\bar{f}^\mu | \mathfrak{G})$ in $L_1(X, \mathfrak{G}, \mu|_{\mathfrak{G}})$, which may also be denoted by $E(\bar{f}^\mu | g)$, is the *conditional mathematical expectation of $\bar{f}^\mu$ given the function g* , or simply the *conditional expectation of $\bar{f}^\mu$ given g* . We use this definition of $E(\bar{f}^\mu | g)$ because it is precisely the definition of conditional mathematical expectation as set down in Kolmogorov [4].

In what follows, the fact that $(\mu|_{\mathfrak{G}}) \circ g^{\leftarrow} = (\mu \circ g^{\leftarrow})|_{\mathfrak{A}^{(g)}}$ will be used. By Proposition 4.36, there is a unique class

$$r + \mathcal{N}_1(Y, \mathfrak{A}^{(g)}, (\mu \circ g^{\leftarrow})|_{\mathfrak{A}^{(g)}}) \in L_1(Y, \mathfrak{A}^{(g)}, (\mu \circ g^{\leftarrow})|_{\mathfrak{A}^{(g)}})$$

with

$$\begin{aligned} E(\bar{f}^\mu | g) &= r \circ g + \mathcal{N}_1(X, \mathfrak{G}, \mu|_{\mathfrak{G}}) \\ &= \overline{r \circ g}^{\mu|_{\mathfrak{G}}} \end{aligned}$$

and with the representative r being an $(\mathfrak{A}^{(g)}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable function, as illustrated in the following diagram, letting the function e be any representative of the class $E(\bar{f}^\mu | g)$:

$$\begin{array}{ccc} (X, \mathfrak{A}) & \xrightarrow{f} & (\mathbb{R}, \mathfrak{Bor}_{\mathbb{R}}) \\ & \searrow g & \uparrow r \\ (X, \mathfrak{G}) & \xrightarrow{e} & (\mathbb{R}, \mathfrak{Bor}_{\mathbb{R}}) \end{array} \quad \text{meaning} \quad E(\bar{f}^\mu | g) = \overline{r \circ g}^{\mu|_{\mathfrak{G}}}$$

Call the induced function r the **regression function**.

Discussion 7.7. In a special case of Definition 7.6, if the function $g : X \rightarrow Y$ is known to be $(\mathfrak{A}, \mathfrak{B})$ -measurable, then

$$\mathfrak{A}^{(g)} = \mathfrak{B}$$

and

$$\mathfrak{G} = g^{\leftarrow}(\mathfrak{B}).$$

In this case, the conditional expectation $E(\bar{f}^\mu | g)$ of $\bar{f}^\mu$ given the function g denotes the same class as the conditional expectation $E_{\mathfrak{G}} \bar{f}^\mu$ of $\bar{f}^\mu$ with respect to the σ -algebra $\mathfrak{G}$.

{conditional_expectation_operator}

Definition 7.8. Let $\mathfrak{B}$ be a σ -subalgebra of $\mathfrak{A}$. The following defines a contractive operator:

$$E_{\mathfrak{B}} : L_1(X, \mathfrak{A}, \mu) \rightarrow L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}}).$$

By Proposition 4.14, there is another contractive operator:

$$\mathfrak{I} : L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}}) \rightarrow L_1(X, \mathfrak{A}, \mu) : \bar{f}^{\mu|_{\mathfrak{B}}} \mapsto \bar{f}^\mu.$$

Define the **conditional expectation operator**

$$\mathcal{E}_{\mathfrak{B}} : L_1(X, \mathfrak{A}, \mu) \rightarrow L_1(X, \mathfrak{A}, \mu)$$

to be their composition $\mathfrak{I} \circ E_{\mathfrak{B}}$:

$$\begin{array}{ccc} L_1(X, \mathfrak{A}, \mu) & \xrightarrow{\mathcal{E}_{\mathfrak{B}}} & L_1(X, \mathfrak{A}, \mu) \\ & \searrow E_{\mathfrak{B}} & \uparrow \mathfrak{I} \\ & & L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}}) \end{array}$$

7.1 Properties of Expectation Operators

{L1embedding}

Proposition 7.9. The map $\mathfrak{I}$ preserves the integral, meaning if $\bar{g}^\mu \in L_1(X, \mathfrak{A}, \mu)$, then

$$\int_B \mathcal{E}_{\mathfrak{B}} \bar{g}^\mu d\mu = \int_B E_{\mathfrak{B}} \bar{g}^\mu d\mu|_{\mathfrak{B}} \quad \text{for all } B \in \mathfrak{B}.$$

Remark 7.10. For each class $\bar{g}^\mu$ in $L_1(X, \mathfrak{A}, \mu)$, the class $\mathcal{E}_{\mathfrak{B}} \bar{g}^\mu$ in $L_1(X, \mathfrak{A}, \mu)$ has a $(\mathfrak{B}, \mathfrak{B}\sigma_{\mathbb{R}})$ -measurable representative, and to see this,

say the function $e \in \mathcal{L}_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$ is the $(\mathfrak{B}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable representative of the class $E_{\mathfrak{B}} \bar{g}^{\mu}$, as in Definition 7.4. This means $E_{\mathfrak{B}} \bar{g}^{\mu} = \bar{e}^{\mu|_{\mathfrak{B}}}$. Then the equalities

$$\mathcal{E}_{\mathfrak{B}} \bar{g}^{\mu} = \mathfrak{I}(E_{\mathfrak{B}} \bar{g}^{\mu}) = \mathfrak{I}(\bar{e}^{\mu|_{\mathfrak{B}}}) = \bar{e}^{\mu}$$

show that $\mathcal{E}_{\mathfrak{B}} \bar{g}^{\mu}$ also has the function e , as included in $\mathcal{L}_1(X, \mathfrak{A}, \mu)$, as a $(\mathfrak{B}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable representative. You may recall the inclusion $\mathcal{L}_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}}) \subseteq \mathcal{L}_1(X, \mathfrak{A}, \mu)$ of Discussion 4.16.

{conditional_expectation_operator_linear}

Proposition 7.11. *Let $\mathfrak{B}$ be a σ -subalgebra of $\mathfrak{A}$. The operators*

$$E_{\mathfrak{B}} : L_1(X, \mathfrak{A}, \mu) \rightarrow L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}}) : \bar{f}^{\mu} \mapsto E_{\mathfrak{B}} \bar{f}^{\mu}$$

and

$$\mathcal{E}_{\mathfrak{B}} : L_1(X, \mathfrak{A}, \mu) \rightarrow L_1(X, \mathfrak{A}, \mu) : \bar{f}^{\mu} \mapsto \mathcal{E}_{\mathfrak{B}} \bar{f}^{\mu}$$

are positive, linear, and contractive.

Remark 7.12. The correct and complete proof that

$$\mathbb{E} \bar{f}^{\mu} = \mathbb{E} (\mathcal{E}_{\mathfrak{B}} \bar{f}^{\mu}),$$

as in the following proposition, took some effort, tracing back through at least Proposition 4.14. Let my early and admittedly naive attempts to try to simply throw a monotone convergence theorem at this proposition at least bring a smile to your face. Of course there was more to it.

{equality_of_expected_values}

Proposition 7.13. *If $\mathfrak{B}$ is a σ -subalgebra of $\mathfrak{A}$, then*

$$\int_X f d\mu = \int_X \mathcal{E}_{\mathfrak{B}} \bar{f}^{\mu} d\mu$$

for all classes $\bar{f}^{\mu} \in L_1(X, \mathfrak{A}, \mu)$. Alternately,

$$\mathbb{E} \bar{f}^{\mu} = \mathbb{E} (\mathcal{E}_{\mathfrak{B}} \bar{f}^{\mu}).$$

7 Conditional Mathematical Expectation

Proof. Let $\mathfrak{B}$ be a σ -subalgebra of $\mathfrak{A}$. Let $\bar{f}^\mu \in L_1(X, \mathfrak{A}, \mu)$. By definition of conditional mathematical expectation, we know that

$$\int_B f d\mu = \int_B E_{\mathfrak{B}} \bar{f}^\mu d\mu|_{\mathfrak{B}}$$

for all $B \in \mathfrak{B}$, and by taking $B = X$, it follows that

$$\int_X f d\mu = \int_X E_{\mathfrak{B}} \bar{f}^\mu d\mu|_{\mathfrak{B}}.$$

By Proposition 7.9,

$$\int_X E_{\mathfrak{B}} \bar{f}^\mu d\mu|_{\mathfrak{B}} = \int_X \mathcal{E}_{\mathfrak{B}} \bar{f}^\mu d\mu.$$

It follows by transitivity of equality that

$$\int_X f d\mu = \int_X \mathcal{E}_{\mathfrak{B}} \bar{f}^\mu d\mu,$$

or alternately,

$$\mathbb{E} \bar{f}^\mu = \mathbb{E} (\mathcal{E}_{\mathfrak{B}} \bar{f}^\mu),$$

as required. ■

Remark 7.14. Proposition 7.15 says that the following diagram commutes:

$$\begin{array}{ccc} L_1(X, \mathfrak{A}, \mu) & & \\ \uparrow \mathfrak{I} & \searrow E_{\mathfrak{B}} & \\ L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}}) & \xrightarrow{\text{identity}} & L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}}) \end{array}$$

which is part of a larger commutative diagram:

$$\begin{array}{ccc}
 L_1(X, \mathfrak{A}, \mu) & \xrightarrow{E_{\mathfrak{B}}} & L_1(X, \mathfrak{A}, \mu) \\
 \uparrow \mathfrak{I} & \searrow E_{\mathfrak{B}} & \uparrow \mathfrak{I} \\
 L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}}) & \xrightarrow{\text{identity}} & L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}})
 \end{array}$$

Proposition 7.15. *If $\mathfrak{B}$ is a σ -subalgebra of $\mathfrak{A}$, then*

{L1_subalgebra_fixed}

$$E_{\mathfrak{B}} \mathfrak{I}(\bar{f}^{\mu|_{\mathfrak{B}}}) = \bar{f}^{\mu|_{\mathfrak{B}}}$$

for all classes $\bar{f}^{\mu|_{\mathfrak{B}}} \in L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$.

Proof. From the definition of conditional mathematical expectation: if $\mathfrak{I}(\bar{f}^{\mu|_{\mathfrak{B}}})$ in $L_1(X, \mathfrak{A}, \mu)$, there is a unique class $E_{\mathfrak{B}} \mathfrak{I}(\bar{f}^{\mu|_{\mathfrak{B}}})$ in $L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$ such that

$$\int_B \mathfrak{I}(\bar{f}^{\mu|_{\mathfrak{B}}}) d\mu = \int_B E_{\mathfrak{B}} \mathfrak{I}(\bar{f}^{\mu|_{\mathfrak{B}}}) d\mu|_{\mathfrak{B}} \quad \text{for all } B \in \mathfrak{B}.$$

It follows by Proposition 4.14 that

$$\int_B \mathfrak{I}(\bar{f}^{\mu|_{\mathfrak{B}}}) d\mu = \int_B \bar{f}^{\mu|_{\mathfrak{B}}} d\mu|_{\mathfrak{B}} \quad \text{for all } B \in \mathfrak{B}.$$

By uniqueness, $E_{\mathfrak{B}} \mathfrak{I}(\bar{f}^{\mu|_{\mathfrak{B}}}) = \bar{f}^{\mu|_{\mathfrak{B}}}$, completing the proof. $\blacksquare$

7.2 A Monotone Convergence Theorem

Remark 7.16. If $\bar{f}^{\mu}$ is a class in $L_1(X, \mathfrak{A}, \mu)$, then $\bar{0}^{\mu} \leq \bar{f}^{\mu}$ means that $0 \leq f(x)$ for μ^* -almost all $x \in X$. Equivalently, $\bar{f}^{\mu} \leq \bar{g}^{\mu}$ means that $f(x) \leq g(x)$ for μ^* -almost all $x \in X$. What then does $\bar{f}_n^{\mu} \uparrow \bar{f}^{\mu}$ mean? It means that for each n , $f_n(x) \leq f_{n+1}(x)$ for μ^* -almost all $x \in X$, and that $f_n(x) \rightarrow f(x)$ for μ^* -almost all $x \in X$.

Proposition 7.17. *Let $(X, \mathfrak{A}, \mu)$ be a probability space, and let $\mathfrak{B}$ be a σ -subalgebra of $\mathfrak{A}$. Let $\{\bar{f}^\mu, \bar{f}_1^\mu, \bar{f}_2^\mu, \dots\}$ be a collection of classes in $L_1(X, \mathfrak{A}, \mu)$. If $\bar{0}^\mu \leq \bar{f}_n^\mu \uparrow \bar{f}^\mu$, then $\bar{0}^{\mu|_{\mathfrak{B}}} \leq E_{\mathfrak{B}} \bar{f}_n^\mu \uparrow E_{\mathfrak{B}} \bar{f}^\mu$.*

{conditional_monotone}

Proof. Fairly simple consequence of Levi's Theorem 3.40 applied to $E_{\mathfrak{B}} \bar{f}_n^\mu$, and of the definition of conditional mathematical expectation. Will come back and transcribe my hand-written stuff. At some point, maybe. ■

7.3 Atoms and Independence

Definition 7.18. Let $(X, \mathfrak{S}, \mu)$ be a measure space. A μ^* -measurable set A with $\mu^*(A) > 0$ is called an **atom** of $(X, \mathfrak{S}, \mu)$ if for every μ^* -measurable subset B of A either $\mu^*(B) = 0$, or $\mu^*(A \cap B^c) = 0$.

{atom}

Proposition 7.19. *Let $(X, \mathfrak{S}, \mu)$ be a finite measure space. A μ^* -measurable set A with $\mu^*(A) > 0$ is an atom of $(X, \mathfrak{S}, \mu)$ if for every μ^* -measurable subset B of A either $\mu^*(B) = \mu^*(A)$, or $\mu^*(A \cap B^c) = \mu^*(A)$.*

Remark 7.20. If the measure space $(X, \mathfrak{S}, \mu)$ is not finite, then the condition that either $\mu^*(B) = \mu^*(A)$, or $\mu^*(A \cap B^c) = \mu^*(A)$ is not strong enough to imply the condition that either $\mu^*(B) = 0$, or $\mu^*(A \cap B^c) = 0$.

{countable_partition_of_atoms}

Proposition 7.21. *Let $(X, \mathfrak{A}, \mu)$ be a probability space. Let $\mathfrak{B}$ denote the σ -subalgebra of $\mathfrak{A}$ generated by a countable partition $\{B_i\}$ of X , where each set $B_i \in \mathfrak{A}$. Then every set B_i with $\mu(B_i) > 0$ is an atom of the probability space $(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$.*

Proof. Let us suppose that the set B_1 has $\mu(B_1) > 0$, and let us show that this implies B_1 is an atom of $(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$. We first need to show both that B_1 is a $\mu|_{\mathfrak{B}}^*$ -measurable set, and that $\mu|_{\mathfrak{B}}^*(B_1) > 0$. We then need to show (by Proposition 7.19) that if A is a $\mu|_{\mathfrak{B}}^*$ -measurable subset of B_1 , then either $\mu|_{\mathfrak{B}}^*(A) = \mu|_{\mathfrak{B}}^*(B_1)$, or $\mu|_{\mathfrak{B}}^*(B_1 \cap A^c) = \mu|_{\mathfrak{B}}^*(B_1)$.

First, by Proposition 2.16, we know that $\mathfrak{B} \subseteq \mu|_{\mathfrak{B}}^*\text{-}\mathfrak{Meas}$, and since $B_1 \in \mathfrak{B}$, it follows that B_1 is a $\mu|_{\mathfrak{B}}^*$ -measurable subset of X . Also, by

Proposition 3.14 and Proposition 2.11, we know that $\mu(B_1) = \mu_{|\mathfrak{B}}(B_1) = \mu_{|\mathfrak{B}}^*(B_1)$, and so $P(B_1) > 0$ means that $\mu_{|\mathfrak{B}}^*(B_1) > 0$.

Next, let A be a $\mu_{|\mathfrak{B}}^*$ -measurable subset of B_1 . We need to see that either $\mu_{|\mathfrak{B}}^*(A) = \mu_{|\mathfrak{B}}^*(B_1)$, or $\mu_{|\mathfrak{B}}^*(B_1 \cap A^c) = \mu_{|\mathfrak{B}}^*(B_1)$. Should A be empty, then $B_1 \cap A^c = B_1$, and so $\mu_{|\mathfrak{B}}^*(B_1 \cap A^c) = \mu_{|\mathfrak{B}}^*(B_1)$, rather vacuously. Suppose then that A is non-empty, and let us see that this implies $\mu_{|\mathfrak{B}}^*(A) = \mu_{|\mathfrak{B}}^*(B_1)$. Let us determine $\mu_{|\mathfrak{B}}^*(A)$ by tending to the definition:

$$\mu_{|\mathfrak{B}}^*(A) = \inf \left\{ \sum_{n=1}^{\infty} \mu_{|\mathfrak{B}}(A_n) : \{A_n\} \text{ is a sequence of } \mathfrak{B} \text{ with } A \subseteq \bigcup_{n=1}^{\infty} A_n \right\}.$$

If $\{A_n\}$ is a sequence of $\mathfrak{B}$ with $A \subseteq \bigcup_{n=1}^{\infty} A_n$, then because $\mathfrak{B}$ is generated by a countable collection of disjoint sets B_i , we may assume that each of the A_n is either empty, or equal to one of the B_i . Furthermore, since A is non-empty, at least one of the A_n must equal B_1 in order that $A \subseteq \bigcup_{n=1}^{\infty} A_n$. This means that for any sequence, $\{C_n\}$ say, of $\mathfrak{B}$ where exactly one of the C_n is equal to B_1 and the rest of the C_n are empty, we must have

$$\sum_{n=1}^{\infty} \mu_{|\mathfrak{B}}(C_n) \leq \sum_{n=1}^{\infty} \mu_{|\mathfrak{B}}(A_n)$$

if $\{A_n\}$ is an arbitrary sequence of $\mathfrak{B}$ with $A \subseteq \bigcup_{n=1}^{\infty} A_n$. And $\sum_{n=1}^{\infty} \mu_{|\mathfrak{B}}(C_n) = \mu_{|\mathfrak{B}}(B_1) = \mu_{|\mathfrak{B}}^*(B_1)$. So $\mu_{|\mathfrak{B}}^*(B_1)$ is a lower bound for the set

$$\left\{ \sum_{n=1}^{\infty} \mu_{|\mathfrak{B}}(A_n) : \{A_n\} \text{ is a sequence of } \mathfrak{B} \text{ with } A \subseteq \bigcup_{n=1}^{\infty} A_n \right\}.$$

But $\mu_{|\mathfrak{B}}^*(A)$ is the greatest such lower bound, so $\mu_{|\mathfrak{B}}^*(A) \geq \mu_{|\mathfrak{B}}^*(B_1)$. On the other hand, $A \subseteq B_1$, so $\mu_{|\mathfrak{B}}^*(A) \leq \mu_{|\mathfrak{B}}^*(B_1)$. It follows then that $\mu_{|\mathfrak{B}}^*(A) = \mu_{|\mathfrak{B}}^*(B_1)$, completing the proof. ■

{atom_lemma}

Lemma 7.22. *If $f : X \rightarrow \mathbb{R}$ is an $(\mathfrak{A}, \mathfrak{Bor}_{\mathbb{R}})$ -simple function, then f is constant μ^* -almost everywhere on each atom of the probability space $(X, \mathfrak{A}, \mu)$.*

Proof. Let $f : X \rightarrow \mathbb{R}$ be an $(\mathfrak{A}, \mathfrak{Bor}_{\mathbb{R}})$ -simple function, meaning that f is a $(\mathfrak{A}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable function having finite range. Suppose the range of f to be $\{a_1, \dots, a_n\} \subseteq \mathbb{R}$. If $n = 1$, then f is certainly constant since $f = a_1$ on X , so suppose $n > 1$. Define subsets $A_i \subseteq X$ to be the preimages $f^{-1}(a_i)$. Let A be an atom of $(X, \mathfrak{A}, \mu)$. The subsets $\{A_i\}$ partition X , and so the atom A can be written as the disjoint union $\bigcup_{i=1}^n (A_i \cap A)$, where each $A_i \cap A$ is a μ^* -measurable subset of A . Since μ^* is σ -additive on the μ^* -measurable subset of A (by Proposition 2.14), it follows that $\mu^*(A) = \sum_{i=1}^n \mu^*(A_i \cap A)$. Because A is an atom, it follows (by Proposition 7.19) that for each A_i , either $\mu^*(A_i \cap A) = \mu^*(A)$, or $\mu^*(\bigcup_{j \neq i} (A_j \cap A)) = \mu^*(A)$. Let us argue that $\mu^*(A) = \mu^*(A_i \cap A)$ for exactly one of the A_i , for it will then follow that $f = a_i$ μ^* -almost everywhere on A .

Should $\mu^*(A) = \mu^*(A_i \cap A)$ for at least two of the A_i ; say for A_1 and A_2 , we would then have

$$\mu^*(A) = \sum_{i=1}^n \mu^*(A_i \cap A) \geq \mu^*(A_1 \cap A) + \mu^*(A_2 \cap A) = 2 \cdot \mu^*(A),$$

which could only happen were $\mu^*(A) = 0$. But $\mu^*(A) > 0$, by assumption.

Should $\mu^*(A) \neq \mu^*(A_i \cap A)$ for any of the A_i , then $\mu^*(\bigcup_{j \neq i} (A_j \cap A)) = \mu^*(A)$ for each A_i , and so

$$\begin{aligned} \mu^*(A) &= \mu^*\left((A_i \cap A) \cup \left(\bigcup_{j \neq i} (A_j \cap A)\right)\right) \\ &= \mu^*(A_i \cap A) + \mu^*\left(\bigcup_{j \neq i} (A_j \cap A)\right) \\ &= \mu^*(A_i \cap A) + \mu^*(A), \end{aligned}$$

meaning $\mu^*(A_i \cap A) = 0$ for each i . This would imply that $\mu^*(A) = 0$ because $\mu^*(A) = \sum_{i=1}^n \mu^*(A_i \cap A)$. But $\mu^*(A) > 0$ by assumption.

Conclude that $\mu^*(A) = \mu^*(A_i \cap A)$ for exactly one of the A_i , so $\mu^*(A_j \cap A) = 0$ for $j \neq i$, and consequently $f = a_i$ μ^* -almost everywhere on A , completing the proof. ■

{constant_on_the_atoms}

Proposition 7.23. *If $\mathfrak{B}$ is a σ -subalgebra of $\mathfrak{A}$, and if $f : X \rightarrow \mathbb{R}$ is a $(\mathfrak{B}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable function, then f is constant $\mu|_{\mathfrak{B}}$ -almost everywhere on each atom of the probability space $(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$.*

Proof. Let $\mathfrak{B}$ be a σ -subalgebra of $\mathfrak{A}$, and let $f : X \rightarrow \mathbb{R}$ be $(\mathfrak{B}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable. Also, let B be an atom of $(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$. By Proposition 3.21, there is a sequence $\{f_n\}$ of $(\mathfrak{B}, \mathfrak{Bor}_{\mathbb{R}})$ -simple functions on X such that $\lim f_n(x) = f(x)$ for all $x \in X$. Because the f_n converge to f on X , and because each f_n is constant $\mu|_{\mathfrak{B}}$ -almost everywhere on B (by Lemma 7.22), it follows that f is constant $\mu|_{\mathfrak{B}}$ -almost everywhere on B , as required. ■

Discussion 7.24. If $e \in \mathcal{L}_1(X, \mathfrak{A}, \mu)$, and if $B \in \mathfrak{A}$ so that $\chi_B \in \mathcal{L}_1(X, \mathfrak{A}, \mu)$, and if the product of functions $e\chi_B$ is in $\mathcal{L}_1(X, \mathfrak{A}, \mu)$ so that the class $\overline{e\chi_B}^\mu$ is in $L_1(X, \mathfrak{A}, \mu)$, then the product $\overline{e}^\mu \overline{\chi_B}^\mu$ of classes is defined to be the class $\overline{e\chi_B}^\mu$ in $L_1(X, \mathfrak{A}, \mu)$. Should the function e be any representative of the class $\mathcal{E}_{\mathfrak{B}} \bar{g}^\mu$, then $\overline{e}^\mu = \mathcal{E}_{\mathfrak{B}} \bar{g}^\mu$ and so $\overline{e}^\mu \overline{\chi_B}^\mu = (\mathcal{E}_{\mathfrak{B}} \bar{g}^\mu) \overline{\chi_B}^\mu$ in $L_1(X, \mathfrak{A}, \mu)$.

{value_on_the_atoms}

Proposition 7.25. *Let $\bar{f}^\mu \in L_1(X, \mathfrak{A}, \mu)$. If $\mathfrak{B}$ is a σ -subalgebra of $\mathfrak{A}$, and if B is an atom of the probability space $(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$, then*

$$\left(\mathcal{E}_{\mathfrak{B}} \bar{f}^\mu \right) \overline{\chi_B}^\mu = \left(\frac{1}{\mu(B)} \int_B f d\mu \right) \overline{\chi_B}^\mu \quad \text{in } L_1(X, \mathfrak{A}, \mu).$$

Proof. The idea is to show that

$$\left(\mathcal{E}_{\mathfrak{B}} \bar{f}^\mu \right) \overline{\chi_B}^{\mu|_{\mathfrak{B}}} = \left(\frac{1}{\mu(B)} \int_B f d\mu \right) \overline{\chi_B}^{\mu|_{\mathfrak{B}}} \quad \text{in } L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}}),$$

and then apply the map $\mathfrak{J}$ of Proposition 4.14.

Let $\mathfrak{B}$ be a σ -subalgebra of $\mathfrak{A}$, and let B be an atom of $(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$. We may suppose the function $e \in \mathcal{L}_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$ to be the $(\mathfrak{B}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable representative of the class $E_{\mathfrak{B}} \bar{f}^{\mu} \in L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$. It follows by Proposition 7.23 that the function e is constant $\mu|_{\mathfrak{B}}^*$ -almost everywhere on B . Suppose this constant equals a , meaning that the value of e is equal to a $\mu|_{\mathfrak{B}}^*$ -almost everywhere on B . Constants are easy to integrate over sets:

$$\int_B e d\mu|_{\mathfrak{B}} = a \cdot \mu|_{\mathfrak{B}}(B) = a \cdot \mu(B).$$

On the other hand, by the definition of conditional expectation,

$$\int_B e d\mu|_{\mathfrak{B}} = \int_B E_{\mathfrak{B}} \bar{f}^{\mu} d\mu|_{\mathfrak{B}} = \int_B f d\mu.$$

This says that

$$a \cdot \mu(B) = \int_B f d\mu.$$

By hypothesis, the set B is an atom, and so $\mu(B) > 0$. This means we can divide by $\mu(B)$. Therefore

$$a = \frac{1}{\mu(B)} \int_B f d\mu.$$

Because e is equal to a $\mu|_{\mathfrak{B}}^*$ -almost everywhere on B , it follows that

$$e = \frac{1}{\mu(B)} \int_B f d\mu \quad \mu|_{\mathfrak{B}}^* \text{-almost everywhere on } B,$$

and so

$$e \chi_B = \left(\frac{1}{\mu(B)} \int_B f d\mu \right) \chi_B \quad \mu|_{\mathfrak{B}}^* \text{-almost everywhere on } X.$$

By taking equivalence classes in $L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$, we get

$$\overline{e \chi_B}^{\mu|_{\mathfrak{B}}} = \left(\frac{1}{\mu(B)} \int_B f d\mu \right) \overline{\chi_B}^{\mu|_{\mathfrak{B}}} \quad \text{in } L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}}).$$

Since $\overline{e \chi_B}^{\mu|_{\mathfrak{B}}} = (E_{\mathfrak{B}} \bar{f}^{\mu}) \overline{\chi_B}^{\mu|_{\mathfrak{B}}}$ in $L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$, by substitution

$$\left(E_{\mathfrak{B}} \bar{f}^{\mu} \right) \overline{\chi_B}^{\mu|_{\mathfrak{B}}} = \left(\frac{1}{\mu(B)} \int_B f d\mu \right) \overline{\chi_B}^{\mu|_{\mathfrak{B}}} \quad \text{in } L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}}).$$

Applying the map $\mathfrak{I}$, we get

$$\left(\mathcal{E}_{\mathfrak{B}} \bar{f}^{\mu} \right) \overline{\chi_B}^{\mu} = \left(\frac{1}{\mu(B)} \int_B f d\mu \right) \overline{\chi_B}^{\mu} \quad \text{in } L_1(X, \mathfrak{A}, \mu),$$

completing the proof. ■

{conditional_expectation_countable}

Proposition 7.26. *Let $(X, \mathfrak{A}, \mu)$ be a probability space, and suppose that $\mathfrak{A}$ has a σ -subalgebra $\mathfrak{B}$ generated by a countable partition $\{B_i\}$ of X , where each $B_i \in \mathfrak{A}$ and $\mu(B_i) > 0$. If $\bar{f}^{\mu} \in L_1(X, \mathfrak{A}, \mu)$, then*

$$\mathcal{E}_{\mathfrak{B}} \bar{f}^{\mu} = \sum_{i=1}^{\infty} \left(\frac{1}{\mu(B_i)} \int_{B_i} f d\mu \right) \overline{\chi_{B_i}}^{\mu} \quad \text{in } L_1(X, \mathfrak{A}, \mu).$$

Proof. This simply combines Propositions 7.21 and 7.25. ■

Notation 7.27. Let 1_X denote the constant function $X \rightarrow \mathbb{R} : x \mapsto 1$.

{conditional_is_expected}

Proposition 7.28. *Let $\bar{f}^{\mu} \in L_1(X, \mathfrak{A}, \mu)$. If $\mathfrak{B}$ is a σ -subalgebra of $\mathfrak{A}$, and if the σ -subalgebras $\sigma(f)$ and $\mathfrak{B}$ of the σ -algebra $\mathfrak{A}$ are independent with respect to μ^* , then*

$$\mathcal{E}_{\mathfrak{B}} \bar{f}^{\mu} = (E \bar{f}^{\mu}) \cdot \overline{1_X}^{\mu} \quad \text{in } L_1(X, \mathfrak{A}, \mu).$$

Proof. In order to show that the equality $\mathcal{E}_{\mathfrak{B}} \bar{f}^{\mu} = (E \bar{f}^{\mu}) \cdot \overline{1_X}^{\mu}$ holds in $L_1(X, \mathfrak{A}, \mu)$, it's sufficient to show that $E_{\mathfrak{B}} \bar{f}^{\mu} = (E \bar{f}^{\mu}) \cdot \overline{1_X}^{\mu|_{\mathfrak{B}}}$ holds in $L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$, and in turn, to show this, it's sufficient to show

$$\int_B E_{\mathfrak{B}} \bar{f}^{\mu} d\mu|_{\mathfrak{B}} = \int_B (E \bar{f}^{\mu}) \cdot \overline{1_X}^{\mu|_{\mathfrak{B}}} d\mu|_{\mathfrak{B}} \quad \text{for all } B \in \mathfrak{B}.$$

The independence of the σ -subalgebras $\sigma(f)$ and $\mathfrak{B}$ with respect to μ^* implies that the classes $\bar{f}^\mu$ and $\bar{\chi}_B^\mu$ are independent with respect to μ^* for all $B \in \mathfrak{B}$. And, as an application of Proposition 4.14,

$$\mathbb{E} \bar{\chi}_B^\mu = \int_X \chi_B d\mu = \int_X \chi_B d\mu|_B = \int_B \bar{1}_X^{\mu|_{\mathfrak{B}}} d\mu|_{\mathfrak{B}}.$$

Consequently,

$$\begin{aligned} \int_B \bar{E}_{\mathfrak{B}} \bar{f}^\mu d\mu|_{\mathfrak{B}} &= \int_{\mathfrak{B}} f d\mu && \text{by definition} \\ &= \int_X f \chi_B d\mu \\ &= \mathbb{E} (f \bar{\chi}_B^\mu) \\ &= \mathbb{E} (\bar{f}^\mu \bar{\chi}_B^\mu) \\ &= (\mathbb{E} \bar{f}^\mu) (\mathbb{E} \bar{\chi}_B^\mu) && \text{by Prop. 6.13} \\ &= (\mathbb{E} \bar{f}^\mu) \int_B \bar{1}_X^{\mu|_{\mathfrak{B}}} d\mu|_{\mathfrak{B}} && \text{by substitution} \\ &= \int_B (\mathbb{E} \bar{f}^\mu) \cdot \bar{1}_X^{\mu|_{\mathfrak{B}}} d\mu|_{\mathfrak{B}}, \end{aligned}$$

as required. ■

{the_cheapest_subalgebra}

Discussion 7.29. On any probability space $(X, \mathfrak{A}, \mu)$, the σ -algebra generated by the constant function 1_X is $\{X, \emptyset\}$, which is independent of every σ -subalgebra of $\mathfrak{A}$. It follows by Proposition 7.28 that $\mathcal{E}_{\sigma(1_X)} \bar{f}^\mu = (\mathbb{E} \bar{f}^\mu) \cdot \bar{1}_X^\mu$ for every $\bar{f}^\mu \in L_1(X, \mathfrak{A}, \mu)$. This means that any equation involving conditional mathematical expectation which holds for all σ -subalgebras of $\mathfrak{A}$ must hold for expected values as well. You may argue that this is logically backwards since the conditional mathematical expectation is a generalization of the expected value.

{conditional_equals_expected}

Proposition 7.30. *If classes $\bar{f}^\mu$ and $\bar{g}^\mu$ in $L_1(X, \mathfrak{A}, \mu)$ are independent, then*

$$\mathcal{E}_{\sigma(g)} \bar{f}^\mu = (\mathbb{E} \bar{f}^\mu) \cdot \overline{1_X}^\mu$$

and

$$\mathcal{E}_{\sigma(f)} \bar{g}^\mu = (\mathbb{E} \bar{g}^\mu) \cdot \overline{1_X}^\mu.$$

Proof. These are just specific instances of Proposition 7.28. ■

7.4 Measurable Functions Factor Out

Remark 7.31. A commutative diagram is reproduced here to keep track of what's going on in the following proposition:

$$\begin{array}{ccc} L_1(X, \mathfrak{A}, \mu) & \xrightarrow{\mathcal{E}_{\mathfrak{B}}} & L_1(X, \mathfrak{A}, \mu) \\ \uparrow \mathfrak{I} & \searrow E_{\mathfrak{B}} & \uparrow \mathfrak{I} \\ L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}}) & \xrightarrow{\text{identity}} & L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}}) \end{array}$$

The condition $\bar{f}^{\mu|_{\mathfrak{B}}} \cdot E_{\mathfrak{B}} \bar{g}^\mu$ is in $L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$ is extraneous.

Proposition 7.32. *Let $(X, \mathfrak{A}, \mu)$ be a probability space. Let $\mathfrak{B}$ be a σ -subalgebra of $\mathfrak{A}$, and let the classes $\bar{f}^\mu$ and $\bar{g}^\mu$ be in $L_1(X, \mathfrak{A}, \mu)$. If the function $f : X \rightarrow \mathbb{R}$ is $(\mathfrak{B}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable, and if the class $\bar{f}^\mu \cdot \bar{g}^\mu$ is in $L_1(X, \mathfrak{A}, \mu)$, and if the class $\bar{f}^{\mu|_{\mathfrak{B}}} \cdot E_{\mathfrak{B}} \bar{g}^\mu$ is in $L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$, then*

$$E_{\mathfrak{B}}(\bar{f}^\mu \cdot \bar{g}^\mu) = \bar{f}^{\mu|_{\mathfrak{B}}} \cdot E_{\mathfrak{B}} \bar{g}^\mu,$$

and consequently

$$\mathcal{E}_{\mathfrak{B}}(\bar{f}^\mu \cdot \bar{g}^\mu) = \bar{f}^\mu \cdot \mathcal{E}_{\mathfrak{B}} \bar{g}^\mu.$$

Proof. Let the function $f : X \rightarrow \mathbb{R}$ be $(\mathfrak{B}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable, and let the class $\bar{f}^\mu \cdot \bar{g}^\mu$ be in $L_1(X, \mathfrak{A}, \mu)$, and let the class $\bar{f}^{\mu|_{\mathfrak{B}}} \cdot E_{\mathfrak{B}} \bar{g}^\mu$ be in $L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$.

{measurable_functions_factor_out}

What does it mean to show that

$$E_{\mathfrak{B}}(\bar{f}^\mu \cdot \bar{g}^\mu) = \bar{f}^{\mu|_{\mathfrak{B}}} \cdot E_{\mathfrak{B}} \bar{g}^\mu ?$$

By definition, $E_{\mathfrak{B}}(\bar{f}^\mu \cdot \bar{g}^\mu)$ is the unique class in $L_1(X, \mathfrak{A}, \mu)$ having a $(\mathfrak{B}, \mathfrak{Bor}_R)$ -measurable representative such that

$$\int_B f g d\mu = \int_B E_{\mathfrak{B}}(\bar{f}^\mu \cdot \bar{g}^\mu) d\mu|_{\mathfrak{B}}$$

for all $B \in \mathfrak{B}$. So in order to show that

$$E_{\mathfrak{B}}(\bar{f}^\mu \cdot \bar{g}^\mu) = \bar{f}^{\mu|_{\mathfrak{B}}} \cdot E_{\mathfrak{B}} \bar{g}^\mu,$$

we need to show that

$$\int_B f g d\mu = \int_B \bar{f}^{\mu|_{\mathfrak{B}}} \cdot E_{\mathfrak{B}} \bar{g}^\mu d\mu|_{\mathfrak{B}}$$

for all $B \in \mathfrak{B}$. And this is equivalent to showing, letting $e \in E_{\mathfrak{B}} \bar{g}^\mu$, that

$$\int_B f g d\mu = \int_B f e d\mu|_{\mathfrak{B}}$$

for all $B \in \mathfrak{B}$. We will show that this equality holds when the function f is replaced by suitable step functions, and then take limits.

By hypothesis, $f \in L_1(X, \mathfrak{A}, \mu)$ and f is $(\mathfrak{B}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable so by Proposition 4.20, there is a sequence $\{\phi_n\}$ of $(\mu^* \text{-}\mathfrak{Meas}, \mathfrak{Bor}_{\mathbb{R}})$ -step functions with $\phi_n(x) \uparrow f(x)$ for all $x \in X$, where each ϕ_n is a sum of the form

$$\sum_{i=1}^m c_i \chi_{C_i}$$

with each $C_i \in \mathfrak{B}$.

We can suppose that both f and g are nonnegative. It means we can also suppose that the ϕ_n are also nonnegative. It also means the $\phi_n g$

and fg are upper functions. Since $\phi_n g \uparrow fg$, and since fg is integrable, it follows by Proposition 3.32 that

$$\lim \int_B \phi_n g \, d\mu = \int_B \lim \phi_n g \, d\mu = \int_B fg \, d\mu.$$

Similarly,

$$\lim \int_B \phi_n e \, d\mu|_{\mathfrak{B}} = \int_B \lim \phi_n e \, d\mu|_{\mathfrak{B}} = \int_B fe \, d\mu|_{\mathfrak{B}}.$$

To complete the proof then, it is sufficient to show that

$$\int_B \phi_n g \, d\mu = \int_B \phi_n e \, d\mu|_{\mathfrak{B}},$$

and since each $\phi_n = \sum_{i=1}^m c_i \chi_{C_i}$ with $C_i \in \mathfrak{B}$, in turn it is sufficient to show that

$$\int_B \chi_C g \, d\mu = \int_B \chi_C e \, d\mu|_{\mathfrak{B}},$$

for all $C \in \mathfrak{B}$.

But $E_{\mathfrak{B}} \bar{g}^{\mu}$ is the unique class in $L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$ such that

$$\int_B g \, d\mu = \int_B E_{\mathfrak{B}} \bar{g}^{\mu} \, d\mu|_{\mathfrak{B}}$$

for all $B \in \mathfrak{B}$. This means

$$\int_B g \, d\mu = \int_B e \, d\mu|_{\mathfrak{B}}$$

for all $B \in \mathfrak{B}$, and if $C \in \mathfrak{B}$, then $B \cap C \in \mathfrak{B}$, so

$$\int_{B \cap C} g \, d\mu = \int_{B \cap C} e \, d\mu|_{\mathfrak{B}}.$$

If $C \in \mathfrak{B}$, then C is μ^* -meas, and so

$$\int_{B \cap C} g \, d\mu = \int_B \chi_C g \, d\mu,$$

and C is also $\mu|_{\mathfrak{B}}$ -meas, and similarly

$$\int_{B \cap C} e \, d\mu|_{\mathfrak{B}} = \int_B \chi_C e \, d\mu|_{\mathfrak{B}},$$

it follows by transitivity of equality that

$$\int_B \chi_C g \, d\mu = \int_B \chi_C e \, d\mu|_{\mathfrak{B}},$$

completing the proof.

To see that the equality

$$\mathcal{E}_{\mathfrak{B}}(\bar{f}^{\mu} \cdot \bar{g}^{\mu}) = \bar{f}^{\mu} \cdot \mathcal{E}_{\mathfrak{B}} \bar{g}^{\mu}$$

is a consequence, recall that we are letting $e \in E_{\mathfrak{B}} \bar{g}^{\mu}$. We have shown that

$$E_{\mathfrak{B}}(\bar{f}^{\mu} \cdot \bar{g}^{\mu}) = \bar{f}^{\mu}|_{\mathfrak{B}} \cdot E_{\mathfrak{B}} \bar{g}^{\mu},$$

and we know

$$\begin{aligned} \bar{f}^{\mu}|_{\mathfrak{B}} \cdot E_{\mathfrak{B}} \bar{g}^{\mu} &= \bar{f}^{\mu}|_{\mathfrak{B}} \cdot \bar{e}^{\mu}|_{\mathfrak{B}} \\ &= \overline{f e^{\mu}}|_{\mathfrak{B}} \end{aligned}$$

saying that

$$E_{\mathfrak{B}}(\bar{f}^{\mu} \cdot \bar{g}^{\mu}) = \overline{f e^{\mu}}|_{\mathfrak{B}}.$$

Under the embedding $\mathfrak{I}$,

$$E_{\mathfrak{B}}(\bar{f}^{\mu} \cdot \bar{g}^{\mu}) \mapsto \mathcal{E}_{\mathfrak{B}}(\bar{f}^{\mu} \cdot \bar{g}^{\mu})$$

and

$$\overline{f e^{\mu}}|_{\mathfrak{B}} \mapsto \overline{f e^{\mu}} = \bar{f}^{\mu} \cdot \bar{e}^{\mu},$$

so

$$\mathcal{E}_{\mathfrak{B}}(\bar{f}^{\mu} \cdot \bar{g}^{\mu}) = \bar{f}^{\mu} \cdot \bar{e}^{\mu}.$$

But $E_{\mathfrak{B}}(\bar{g}^{\mu}) = \bar{e}^{\mu}|_{\mathfrak{B}}$, so under the embedding $\mathcal{E}_{\mathfrak{B}}(\bar{g}^{\mu}) = \bar{e}^{\mu}$, and by substitution

$$\mathcal{E}_{\mathfrak{B}}(\bar{f}^{\mu} \cdot \bar{g}^{\mu}) = \bar{f}^{\mu} \cdot \mathcal{E}_{\mathfrak{B}}(\bar{g}^{\mu}),$$

as required. ■

8 Conditional Probability and Markov Kernels

SOME authors confuse Markov kernels and conditional distributions. We will not. However, as shown in Proposition 8.21 below, a Markov kernel satisfying an extra condition corresponds to a regular conditional distribution.


8.1 Conditional Probability

Definition 8.1. Let $(X, \mathfrak{A}, \mu)$ be a probability space, and let $\mathfrak{B}$ be a σ -subalgebra of $\mathfrak{A}$. The set function

{conditional_probability}

$$\mathcal{P}_{\mathfrak{B}} : \mathfrak{A} \rightarrow L_1(X, \mathfrak{A}, \mu) : A \mapsto \mathcal{E}_{\mathfrak{B}} \overline{\chi_A}^{\mu}$$

is the **conditional probability** on $\mathfrak{A}$ with respect to the σ -subalgebra $\mathfrak{B}$, or the *conditional probability on $\mathfrak{A}$ given $\mathfrak{B}$* , the composition being illustrated in the following diagram:



Remark 8.2. Should the σ -subalgebra $\mathfrak{B}$ be generated by a countable partition $\{B_i\}$ of X , where each $B_i \in \mathfrak{A}$ and $\mu(B_i) > 0$, then using

Proposition 7.26, for each $A \in \mathfrak{A}$,

$$\begin{aligned}\mathcal{P}_{\mathfrak{B}}(A) &= \mathcal{E}_{\mathfrak{B}} \overline{\chi_A}^\mu \\ &= \sum_{i=1}^{\infty} \left(\frac{1}{\mu(B_i)} \int_{B_i} \chi_A d\mu \right) \overline{\chi_{B_i}}^\mu \\ &= \sum_{i=1}^{\infty} \left(\frac{\mu(A \cap B_i)}{\mu(B_i)} \right) \overline{\chi_{B_i}}^\mu.\end{aligned}$$

The coefficient

$$\frac{\mu(A \cap B_i)}{\mu(B_i)}$$

reminds us of the symbols in the very elementary definition of conditional probability:

$$P(A | B) = \frac{P(A \cap B)}{P(B)}.$$

Proposition 8.3. *Let $(X, \mathfrak{A}, \mu)$ be a probability space, and let $\mathfrak{B}$ be a σ -subalgebra of $\mathfrak{A}$. The conditional probability $\mathcal{P}_{\mathfrak{B}} : \mathfrak{A} \rightarrow L_1(X, \mathfrak{A}, \mu)$ has the following measure-like properties:*

1. $\mathcal{P}_{\mathfrak{B}}(\emptyset) = \overline{0_X}^\mu$.
2. $\mathcal{P}_{\mathfrak{B}}$ is σ -additive.
3. $\mathcal{P}_{\mathfrak{B}}(X) = \overline{1_X}^\mu$.

8.2 Markov Kernels

Notation 8.4. Let X be a space and let $\mathfrak{B}$ be a σ -algebra. For each set B in the σ -algebra $\mathfrak{B}$, define the function

$$\text{fix}^B : X \rightarrow X \times \mathfrak{B} : x \mapsto (x, B),$$

and for each element x in the space X , define the function

$$\text{fix}_x : \mathfrak{B} \rightarrow X \times \mathfrak{B} : B \mapsto (x, B).$$

{markov_kernel}

Definition 8.5. Given measurable spaces $(X, \mathfrak{A})$ and $(Y, \mathfrak{B})$, a **Markov kernel** is a function

$$k : X \times \mathfrak{B} \rightarrow [0, 1]$$

such that for each set $B \in \mathfrak{B}$, the map

$$k \circ \text{fix}^B : X \rightarrow X \times \mathfrak{B} \rightarrow [0, 1] : x \mapsto (x, B) \mapsto k(x, B)$$

is $(\mathfrak{A}, \mathfrak{Bor}_{[0,1]})$ -measurable, and for each element $x \in X$, the map

$$k \circ \text{fix}_x : \mathfrak{B} \rightarrow X \times \mathfrak{B} \rightarrow [0, 1] : B \mapsto (x, B) \mapsto k(x, B)$$

is a probability measure on $\mathfrak{B}$. The map $k \circ \text{fix}^B$ may be denoted by k^B , and likewise the map $k \circ \text{fix}_x$ may be denoted by k_x . It means that for each $x \in X$ and $B \in \mathfrak{B}$ there is a commutative diagram, only one of which will be illustrated, without quantification:

$$\begin{array}{ccc}
 X & & \\
 \text{fix}^B \downarrow & \searrow k^B \text{ } (\mathfrak{A}, \mathfrak{Bor}_{[0,1]})\text{-measurable} & \\
 X \times \mathfrak{B} & \xrightarrow{k} & [0, 1] \\
 \text{fix}_x \uparrow & \nearrow k_x \text{ probability measure} & \\
 \mathfrak{B} & &
 \end{array}$$

A subscript indicating a domain, such as fix_X^B , could be used in case more than one kernel is involved.

We will say that any map $X \times \mathfrak{B} \rightarrow \mathbb{R}$ or $X \times \mathfrak{B} \rightarrow [0, 1]$ **defines a Markov kernel** if its range is $[0, 1]$ except for possibly those x in some null subset of X and if it can be redefined on that null subset, together with a possible redefinition of the codomain, in such a way that it satisfies the above definition.

A Markov kernel given measurable spaces $(X, \mathfrak{A})$ and $(Y, \mathfrak{B})$ is also called a **transition** from $(X, \mathfrak{A})$ to $(Y, \mathfrak{B})$. We may use the notation

$$k : (X, \mathfrak{A}) \prec (Y, \mathfrak{B})$$

instead of saying that k is a Markov kernel given $(X, \mathfrak{A})$ and $(Y, \mathfrak{B})$.

Example 8.6. Every measurable function induces a Markov kernel, and to see this, let $(X, \mathfrak{A})$ and $(Y, \mathfrak{B})$ be measurable spaces, and let $f : X \rightarrow Y$ be an $(\mathfrak{A}, \mathfrak{B})$ -measurable function. Define a function $k : X \times \mathfrak{B} \rightarrow [0, 1]$ by

$$(x, B) \mapsto \begin{cases} 1 & \text{if } f(x) \in B \\ 0 & \text{if } f(x) \notin B. \end{cases}$$

Two other ways to view the value $k(x, B)$ would be as $k(x, B) = \chi_B(f(x))$, and as $k(x, B) = \delta_{f(x)}(B)$ with $\delta_{f(x)}$ denoting Dirac measure concentrated at $f(x)$. Then for each set $B \in \mathfrak{B}$, the map

$$k \circ \text{fix}_x^B : X \rightarrow [0, 1] : x \mapsto k(x, B) = \chi_B(f(x))$$

can be seen as a composition of the $(\mathfrak{B}, \mathfrak{Bor}_{[0,1]})$ -measurable function χ_B with the $(\mathfrak{A}, \mathfrak{B})$ -measurable function f , and so the composition is $(\mathfrak{A}, \mathfrak{Bor}_{[0,1]})$ -measurable. And for each element $x \in X$, the map

$$k \circ \text{fix}_x : \mathfrak{B} \rightarrow [0, 1] : B \mapsto k(x, B) = \delta_{f(x)}(B)$$

visibly defines a measure on $\mathfrak{B}$, in particular Dirac measure concentrated at $f(x)$. And since $f(x) \in Y$ implies that the Dirac measure of the whole space Y is 1, the map $k \circ \text{fix}_x$ further defines a probability measure on $\mathfrak{B}$. This shows that k is a Markov kernel by the very definition of Markov kernel.

{not_a_product_measure}

Proposition 8.7. Let $(X, \mathfrak{A})$ and $(Y, \mathfrak{B})$ be measurable spaces. Let μ be a probability measure on $\mathfrak{A}$. If $\{k^B : B \in \mathfrak{B}\}$ is a collection of $(\mathfrak{A}, \mathfrak{Bor}_{[0,1]})$ -measurable functions, and $\{k_x : x \in X\}$ is a collection of probability measures on $\mathfrak{B}$, where

$$k^B(x) = k_x(B)$$

for all $B \in \mathfrak{B}$ and for all $x \in X$, then there is a unique probability measure Γ on the σ -algebra $\sigma(\mathfrak{A}, \mathfrak{B})$ of subsets of $X \times Y$ such that

$$\Gamma(A \times B) := \int_A k^B(x) d\mu(x)$$

for all $A \times B \in \mathfrak{Semi}(\mathfrak{A}, \mathfrak{B})$. Furthermore, $\mu(A) = \Gamma(A \times Y)$ for all $A \in \mathfrak{A}$.

Proof. Show that Γ restricted to the product semiring $\mathfrak{Semi}(\mathfrak{A}, \mathfrak{B})$ is countably additive. Then take Γ itself to be the corresponding outer measure restricted to $\sigma(\mathfrak{A}, \mathfrak{B})$. Uniqueness follows by Proposition 2.21. Since k_x is a probability measure, it follows that $k_x(Y) = 1$. That is, the measure of the whole space is 1. And by hypothesis, $k^Y(x) = k_x(Y) = 1$ so

$$\Gamma(A \times Y) = \int_A k^Y(x) d\mu(x) = \int_A 1_X d\mu = \mu(A),$$


as required. ■

{induced_markov_kernel}

Proposition 8.8. Let $(X, \mathfrak{A})$ and $(Y, \mathfrak{B})$ be measurable spaces, and let $\pi_1 : X \times Y \rightarrow X$ denote the canonical projection onto the first component, with $\pi_1^+ : \mathfrak{A} \rightarrow \pi_1^+(\mathfrak{A})$. Let $k : X \times \mathfrak{B} \rightarrow [0, 1]$ be a Markov kernel given measurable spaces $(X, \mathfrak{A})$ and $(Y, \mathfrak{B})$. If a function $m : (X \times Y) \times \mathfrak{B} \rightarrow [0, 1]$ is defined in terms of the Markov kernel k by

$$m : (X \times Y) \times \mathfrak{B} \rightarrow [0, 1] : ((x, y), B) \mapsto k(x, B),$$

then m is a Markov kernel given measurable spaces $(X \times Y, \pi_1^+(\mathfrak{A}))$ and $(Y, \mathfrak{B})$, as illustrated:



Conversely, let $m : (X \times Y) \times \mathfrak{B} \rightarrow [0, 1]$ be a Markov kernel given measurable spaces $(X \times Y, \pi_1^+(\mathfrak{A}))$ and $(Y, \mathfrak{B})$. If a function $k : X \times \mathfrak{B} \rightarrow [0, 1]$ is defined in terms of the Markov kernel m by

$$k : X \times \mathfrak{B} \rightarrow [0, 1] : (x, B) \mapsto m((x, y), B),$$

for any $y \in Y$ such that $(x, y) \in X \times Y$, then k is a Markov kernel given measurable spaces $(X, \mathfrak{A})$ and $(Y, \mathfrak{B})$.

Proof. Let $m : (X \times Y) \times \mathfrak{B} \rightarrow [0, 1]$ be defined by

$$m : (X \times Y) \times \mathfrak{B} \rightarrow [0, 1] : ((x, y), B) \mapsto k(x, B).$$

We want to show that m is a Markov kernel given measurable spaces $(X \times Y, \pi_1^+(\mathfrak{A}))$ and $(Y, \mathfrak{B})$.

$$m \circ \text{fix}^B(x, y) = m((x, y), B) = k(x, B) = k \circ \text{fix}_X^B(x) = k \circ \text{fix}_X^B \circ \pi_1(x, y),$$

which is a composition of measurable functions. Also by definition,

$$m \circ \text{fix}_{(x, y)}(B) = m((x, y), B) = k(x, B) = k \circ \text{fix}_x(B),$$

so $m \circ \text{fix}_{(x, y)} = k \circ \text{fix}_x$, which is a probability measure. ■

8.3 Regular Version

{version}

Definition 8.9. Let $(X, \mathfrak{A}, \mu)$ be a measure space. Let $\mathcal{R}$ be a function $L_1(X, \mathfrak{A}, \mu) \rightarrow \mathcal{L}_1(X, \mathfrak{A}, \mu)$ which selects a representative from each equivalence class. If S is a set, and $\mathcal{P} : S \rightarrow L_1(X, \mathfrak{A}, \mu)$ is a function, then the collection $(\mathcal{R} \circ \mathcal{P})(S)$ of $\mathcal{L}_1(X, \mathfrak{A}, \mu)$ functions is called a **version** of $\mathcal{P}$.

$$\begin{array}{ccc} L_1(X, \mathfrak{A}, \mu) & \xrightarrow{\mathcal{R}} & \mathcal{L}_1(X, \mathfrak{A}, \mu) \\ \mathcal{P} \uparrow & \nearrow \mathcal{R} \circ \mathcal{P} & \\ S & & \end{array}$$

{regular_version}

Definition 8.10. Let the collection $(\mathcal{R} \circ \mathcal{P}_{\mathfrak{B}})(\mathfrak{A})$ of $\mathcal{L}_1(X, \mathfrak{A}, \mu)$ functions be a version of the conditional probability $\mathcal{P}_{\mathfrak{B}}$ on $\mathfrak{A}$ given $\mathfrak{B}$:

$$\begin{array}{ccc} L_1(X, \mathfrak{A}, \mu) & \xrightarrow{\mathcal{R}} & \mathcal{L}_1(X, \mathfrak{A}, \mu) \\ \mathcal{P}_{\mathfrak{B}} \uparrow & \nearrow \mathcal{R} \circ \mathcal{P}_{\mathfrak{B}} & \\ \mathfrak{A} & & \end{array}$$

If the selection function $\mathcal{R} : L_1(X, \mathfrak{A}, \mu) \rightarrow \mathcal{L}_1(X, \mathfrak{A}, \mu)$ has the additional property that each map

$$\mathfrak{A} \rightarrow [0, 1] : A \mapsto (\mathcal{R} \circ \mathcal{P}_{\mathfrak{B}}(A))(x)$$

is a probability measure on $\mathfrak{A}$ except for possibly all x in a $\mu_{|\mathfrak{B}}^*$ -null set, and for each $A \in \mathfrak{A}$, the map

$$X \rightarrow [0, 1] : x \mapsto (\mathcal{R} \circ \mathcal{P}_{\mathfrak{B}}(A))(x)$$

is $(\mathfrak{B}, \mathfrak{Bor}_{[0,1]})$ -measurable, then the collection $\mathcal{R} \circ \mathcal{P}_{\mathfrak{B}}(\mathfrak{A})$ of functions, or any such collection of $\mathcal{L}_1(X, \mathfrak{A}, \mu)$ functions without specifying $\mathcal{R}$, is called a **regular version** of the conditional probability $\mathcal{P}_{\mathfrak{B}}$ on $\mathfrak{A}$ given $\mathfrak{B}$. If all of these $\mathcal{L}_1(X, \mathfrak{A}, \mu)$ functions are modified on some fixed $\mu_{|\mathfrak{B}}^*$ -null set, then the new collection of functions is still considered to be a regular version of $\mathcal{P}_{\mathfrak{B}}$ on $\mathfrak{A}$ given $\mathfrak{B}$. It means that any regular version of $\mathcal{P}_{\mathfrak{B}}$ on $\mathfrak{A}$ given $\mathfrak{B}$ could be redefined in such a way that the map $A \mapsto (\mathcal{R} \circ \mathcal{P}_{\mathfrak{B}}(A))(x)$ is a probability measure on all of $\mathfrak{A}$: should $A \mapsto (\mathcal{R} \circ \mathcal{P}_{\mathfrak{B}}(A))(x)$ be a probability measure on $\mathfrak{A}$ except for all x in a $\mu_{|\mathfrak{B}}^*$ -null set N , then choose $y \in X$ with $y \notin N$, and set

$$(\mathcal{R} \circ \mathcal{P}_{\mathfrak{B}}(A))(x) = (\mathcal{R} \circ \mathcal{P}_{\mathfrak{B}}(A))(y)$$

for all $x \in N$. Unless otherwise stated, all regular conditional probabilities will be considered so selected.

We will call a conditional probability $\mathcal{P}_{\mathfrak{B}}$ on $\mathfrak{A}$ given $\mathfrak{B}$ a **regular conditional probability** if there exists a regular version of $\mathcal{P}_{\mathfrak{B}}$.

{quote_loeve}

Remark 8.11. To quote a paragraph of Loeve [5] verbatim:

A c.pr. $P^{\mathfrak{B}}$ is said to be *regular* if, for every $A \in \mathfrak{A}$, it is possible to select $P^{\mathfrak{B}}A$ within its class of equivalence in such a manner that the $P_x^{\mathfrak{B}}$ are pr.'s on $\mathfrak{A}$ except for points x belonging to a $P_{\mathfrak{B}}$ -null event N . A regular pr.f. $P^{\mathfrak{B}}$ can be said to be defined up to an equivalence, in the sense that if all the functions $P^{\mathfrak{B}}A$ are modified arbitrarily on an arbitrary but fixed $P_{\mathfrak{B}}$ -null event, the new c.pr. is still regular.

In particular, a regular c.pr. $P^{\mathfrak{B}}$ can be selected within its equivalence class so that $P_x^{\mathfrak{B}}$ is a pr. on $\mathfrak{A}$ for *every* $x \in X$. For example, for every x belonging to the exceptional $P^{\mathfrak{B}}$ -null event N set $P_x^{\mathfrak{B}} = P_N$ where P_N is a pr. on $\mathfrak{A}$. Unless otherwise stated, regular c.pr.'s will be so selected.

Loeve continues on saying that a *regular* conditional probability is just a conditional probability which induces a particular Markov kernel; we leave it as the statement of Definition 8.10 and Example 8.13.

{regular_version_equivalence}

Discussion 8.12. Let us compare the conditional probability $P^{\mathfrak{B}}$ quoted from Loeve in Remark 8.11 with the conditional probability $\mathcal{P}^{\mathfrak{B}}$ in Definition 8.10. The conditional probability $P^{\mathfrak{B}}$ quoted from Loeve in Remark 8.11 refers to the restriction of the operator

$$E_{\mathfrak{B}} : L_1(X, \mathfrak{A}, \mu) \rightarrow L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$$

to the family of characteristic functions of events in $\mathfrak{A}$, which we illustrate as:

$$\begin{array}{ccc} L_1(X, \mathfrak{A}, \mu) & \xrightarrow{E_{\mathfrak{B}}} & L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}}) \\ \uparrow Q \circ \chi & \nearrow P^{\mathfrak{B}} & \\ \mathfrak{A} & & \end{array}$$

The conditional probability $\mathcal{P}^{\mathfrak{B}}$ in Definition 8.1 is defined as the composition $\mathcal{E}_{\mathfrak{B}} \circ Q \circ \chi$ in the following diagram:

$$\begin{array}{ccc} L_1(X, \mathfrak{A}, \mu) & \xrightarrow{\mathcal{E}_{\mathfrak{B}}} & L_1(X, \mathfrak{A}, \mu) \\ \uparrow Q \circ \chi & \nearrow \mathcal{P}^{\mathfrak{B}} & \\ \mathfrak{A} & & \end{array}$$

How do the regular versions of these conditional probabilities compare? They are equivalent in this sense. There exists a regular version of $P^{\mathfrak{B}}$ if and only if there exists a regular version of $\mathcal{P}^{\mathfrak{B}}$. Briefly, say $\mathcal{I}$

and $\mathcal{R}$ are the selection functions which select representatives from equivalence classes as illustrated in the following diagrams:

$$\begin{array}{ccc} L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}}) & \xrightarrow{\mathcal{S}} & \mathcal{L}_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}}) \\ \uparrow P^{\mathfrak{B}} & \nearrow \mathcal{S} \circ P^{\mathfrak{B}} & \\ \mathfrak{A} & & \end{array} \quad \begin{array}{ccc} L_1(X, \mathfrak{A}, \mu) & \xrightarrow{\mathcal{R}} & \mathcal{L}_1(X, \mathfrak{A}, \mu) \\ \uparrow \mathcal{P}_{\mathfrak{B}} & \nearrow \mathcal{R} \circ \mathcal{P}_{\mathfrak{B}} & \\ \mathfrak{A} & & \end{array}$$

By Proposition 4.14, if the function $\mathcal{S}(P^{\mathfrak{B}}(A))$ is in $\mathcal{L}_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$, then $\mathcal{S}(P^{\mathfrak{B}}(A))$ is also in $\mathcal{L}_1(X, \mathfrak{A}, \mu)$, and we can take

$$\mathcal{R}(\mathcal{P}_{\mathfrak{B}}(A)) = \mathcal{S}(P^{\mathfrak{B}}(A)).$$

And by Proposition 4.22, if the function $\mathcal{R}(\mathcal{P}_{\mathfrak{B}}(A))$ is in $\mathcal{L}_1(X, \mathfrak{A}, \mu)$, and if $\mathcal{R}(\mathcal{P}_{\mathfrak{B}}(A))$ is $(\mathfrak{B}, \text{Bot}_{[0,1]})$ -measurable, then $\mathcal{R}(\mathcal{P}_{\mathfrak{B}}(A))$ is in $\mathcal{L}_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$ and so we can take

$$\mathcal{S}(P^{\mathfrak{B}}(A)) = \mathcal{R}(\mathcal{P}_{\mathfrak{B}}(A)).$$

The version in both cases is the same family of functions, it just depends on whether they are considered as residing in $\mathcal{L}_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$ or $\mathcal{L}_1(X, \mathfrak{A}, \mu)$. Verify that if $\mathcal{S}(P^{\mathfrak{B}}(A)) = \mathcal{R}(\mathcal{P}_{\mathfrak{B}}(A))$ and either of

$$A \mapsto \mathcal{R}(\mathcal{P}_{\mathfrak{B}}(A))(x)$$

or

$$A \mapsto \mathcal{S}(P^{\mathfrak{B}}(A))(x)$$

is a probability measure on $\mathfrak{A}$, then so is the other.

Example 8.13. Every regular conditional probability induces a Markov kernel; let $(X, \mathfrak{A}, \mu)$ be a probability space, and let $\mathfrak{B}$ be a σ -subalgebra of $\mathfrak{A}$. Let the collection $(\mathcal{R} \circ \mathcal{P}_{\mathfrak{B}})(\mathfrak{A})$ of $\mathcal{L}_1(X, \mathfrak{A}, \mu)$ functions be a regular version of the conditional probability $\mathcal{P}_{\mathfrak{B}}$ on $\mathfrak{A}$ given $\mathfrak{B}$, which means by definition that the selection function $\mathcal{R} : L_1(X, \mathfrak{A}, \mu) \rightarrow \mathcal{L}_1(X, \mathfrak{A}, \mu)$ has the additional property that for each $A \in \mathfrak{A}$, the map

$$X \rightarrow [0, 1] : x \mapsto [(\mathcal{R} \circ \mathcal{P}_{\mathfrak{B}})(A)](x)$$

{regular_conditional_probability_induces_markov_k

is $(\mathfrak{B}, \mathfrak{Bor}_{[0,1]})$ -measurable, and for each $x \in X$ the map

$$\mathfrak{A} \rightarrow [0, 1] : A \mapsto [(\mathcal{R} \circ \mathcal{P}_{\mathfrak{B}})(A)](x)$$

is a probability measure on $\mathfrak{A}$. Let us define a Markov kernel

$$k : X \times \mathfrak{A} \rightarrow [0, 1]$$

so that $k \circ \text{fix}^A = (\mathcal{R} \circ \mathcal{P}_{\mathfrak{B}})(A)$; that is, define

$$k : X \times \mathfrak{A} : (x, A) \mapsto [(\mathcal{R} \circ \mathcal{P}_{\mathfrak{B}})(A)](x),$$

which we will show is a Markov kernel given measurable spaces $(X, \mathfrak{B})$ and $(X, \mathfrak{A})$.

For each set $A \in \mathfrak{A}$, the map

$$k \circ \text{fix}^A : X \rightarrow [0, 1] : x \mapsto k(x, A) = [(\mathcal{R} \circ \mathcal{P}_{\mathfrak{B}})(A)](x)$$

is $(\mathfrak{B}, \mathfrak{Bor}_{[0,1]})$ -measurable by hypothesis, and for each element $x \in X$, the map

$$k \circ \text{fix}_x : \mathfrak{A} \rightarrow [0, 1] : A \mapsto k(x, A) = [(\mathcal{R} \circ \mathcal{P}_{\mathfrak{B}})(A)](x)$$

is a probability measure on $\mathfrak{A}$ by hypothesis, as illustrated:

$$\begin{array}{ccc}
 L_1(X, \mathfrak{A}, \mu) & & X \\
 \uparrow \mathcal{P}_{\mathfrak{B}} & & \downarrow \text{fix}^A \\
 \mathfrak{A} & \xrightarrow{\quad k \quad} & [0, 1] \\
 & \uparrow \text{fix}_x & \uparrow \text{probability measure} \\
 & \mathfrak{A} &
 \end{array}$$

This shows that k is a Markov kernel given measurable spaces $(X, \mathfrak{B})$ and $(X, \mathfrak{A})$, according to definition.

Remark 8.14. Instead of using some arbitrary letter k to denote a Markov kernel, notation like $\mu^{\mathfrak{B}}$ seems arguably better. I get the strangest feeling that this kind of notation will be used in the next couple of updates of these notes.

8.4 Conditional Distributions

Definition 8.15. Let $(X, \mathfrak{A}, \mu)$ be a probability space, and let $\mathfrak{B}$ be a σ -subalgebra of $\mathfrak{A}$. Also let $(Y, \mathfrak{C})$ be a measurable space, and let $f : X \rightarrow Y$ be an $(\mathfrak{A}, \mathfrak{C})$ -measurable function, with $f^{\leftarrow} : \mathfrak{C} \rightarrow \mathfrak{A}$. Just as the function f induces the probability distribution $\mu \circ f^{\leftarrow}$ of f on $\mathfrak{C}$, so does the function f together with the σ -subalgebra $\mathfrak{B}$ induce the **conditional probability distribution** $\mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow}$ of f on $\mathfrak{C}$ given $\mathfrak{B}$:

$$\mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow} : \mathfrak{C} \rightarrow L_1(X, \mathfrak{A}, \mu) : C \mapsto \overline{\mathcal{E}_{\mathfrak{B}} \chi_{f^{\leftarrow}(C)}}^{\mu},$$

or simply the *conditional distribution* of f on $\mathfrak{C}$ given $\mathfrak{B}$, as illustrated:

$$\begin{array}{ccc} L_1(X, \mathfrak{A}, \mu) & \xrightarrow{\mathcal{E}_{\mathfrak{B}}} & L_1(X, \mathfrak{A}, \mu) \\ \uparrow Q \circ \chi & \nearrow \mathcal{P}_{\mathfrak{B}} & \uparrow \mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow} \\ \mathfrak{A} & \xleftarrow{f^{\leftarrow}} & \mathfrak{C} \end{array}$$

Remark 8.16. Distributions are functions induced on a σ -algebra of subsets of the codomain of a measurable function, whether the distribution is a probability distribution or a conditional distribution:

- A probability distribution is a probability measure composed with a preimage.
- A conditional distribution is a conditional probability composed with a preimage.

It is just that simple.

Discussion 8.17. You might reasonably try to avoid explicit mention of the map $f : (X, \mathfrak{A}) \rightarrow (Y, \mathfrak{B})$ in the definition of conditional distribution, perhaps by replacing the σ -algebra $\mathfrak{C}$ with $f^{\leftarrow}(\mathfrak{C})$, and replacing $f^{\leftarrow}$

{distributions}

{avoid_function}

with $\text{inc}^{\leftarrow} := \text{inclusion}^{\leftarrow}$

$$\begin{array}{ccc}
 L_1(X, \mathfrak{A}, \mu) & \xrightarrow{\mathcal{E}_{\mathfrak{B}}} & L_1(X, \mathfrak{A}, \mu) \\
 \uparrow Q \circ \chi & \nearrow \mathcal{P}_{\mathfrak{B}} & \uparrow \mathcal{P}_{\mathfrak{B}} \circ \text{inc}^{\leftarrow} \\
 \mathfrak{A} & \xleftarrow{\text{inc}^{\leftarrow}} & f^{\leftarrow}(\mathfrak{C})
 \end{array}$$

But one way or another, with a distribution, there is a preimage involved. There is no way around it. An $f^{\leftarrow}$ is in there somewhere.

{conditional_probability_of_a_set}

Proposition 8.18. *Let $(X, \mathfrak{A}, \mu)$ be a probability space, and let $\mathfrak{B}$ be a σ -subalgebra of $\mathfrak{A}$. Also let $(Y, \mathfrak{C})$ be a measurable space, and let $f : X \rightarrow Y$ be an $(\mathfrak{A}, \mathfrak{C})$ -measurable function, with $f^{\leftarrow} : \mathfrak{C} \rightarrow \mathfrak{A}$. If $B \in \mathfrak{B}$ and $C \in \mathfrak{C}$, then*

$$\int_B (\mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow})(C) d\mu = \mu(B \cap f^{\leftarrow}(C)).$$

Proof.

$$\begin{aligned}
 \int_B (\mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow})(C) d\mu &= \int_B \mathcal{E}_{\mathfrak{B}} \overline{\chi_{f^{\leftarrow}(C)}}^{\mu} d\mu \\
 &= \int_B E_{\mathfrak{B}} \overline{\chi_{f^{\leftarrow}(C)}}^{\mu} d\mu|_{\mathfrak{B}} \\
 &= \int_B \chi_{f^{\leftarrow}(C)} d\mu \\
 &= \mu(B \cap f^{\leftarrow}(C)).
 \end{aligned}$$

■

{regular_conditional_distribution}

Definition 8.19. The definition of a regular conditional probability *distribution* is nothing other than Definition 8.10 of a regular conditional probability combined with a preimage.

Let the collection $(\mathcal{R} \circ \mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow})(\mathfrak{C})$ of $\mathcal{L}_1(X, \mathfrak{A}, \mu)$ functions be a version of the conditional distribution $\mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow}$ of f on $\mathfrak{C}$ given $\mathfrak{B}$. If the selection function $\mathcal{R} : L_1(X, \mathfrak{A}, \mu) \rightarrow \mathcal{L}_1(X, \mathfrak{A}, \mu)$ has the additional property that the map

$$\mathfrak{C} \rightarrow [0, 1] : C \mapsto (\mathcal{R} \circ \mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow}(C))(x)$$

is a probability measure on the σ -algebra $\mathfrak{C}$ except for possibly all x in a $\mu|_{\mathfrak{B}}^*$ -null set, and for each $C \in \mathfrak{C}$, the map

$$X \rightarrow [0, 1] : x \mapsto (\mathcal{R} \circ \mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow}(C))(x)$$

is $(\mathfrak{B}, \mathfrak{Bor}_{[0,1]})$ -measurable, then the collection $\mathcal{R} \circ \mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow}(\mathfrak{C})$ of functions, or any such collection of $\mathcal{L}_1(X, \mathfrak{A}, \mu)$ functions without specifying $\mathcal{R}$, is called a regular version of the conditional distribution $\mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow}$ of f on $\mathfrak{C}$ given $\mathfrak{B}$, as illustrated:

$$\begin{array}{ccccc} L_1(X, \mathfrak{A}, \mu) & \xrightarrow{\mathcal{E}_{\mathfrak{B}}} & L_1(X, \mathfrak{A}, \mu) & \xrightarrow{\mathcal{R}} & \mathcal{L}_1(X, \mathfrak{A}, \mu) \\ \uparrow Q \circ \chi & \nearrow \mathcal{P}_{\mathfrak{B}} & \uparrow & \nearrow \mathcal{R} \circ \mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow} & \\ \mathfrak{A} & \xleftarrow{f^{\leftarrow}} & \mathfrak{C} & & \end{array}$$

If all of the functions in the collection $(\mathcal{R} \circ \mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow})(\mathfrak{C})$ are modified on a fixed $\mu|_{\mathfrak{B}}^*$ -null set, then the new collection of functions is still considered to be a regular version of $\mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow}$ of f on $\mathfrak{C}$ given $\mathfrak{B}$. It means that any regular version of $\mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow}$ of f on $\mathfrak{C}$ given $\mathfrak{B}$ could be redefined in such a way that the map $C \mapsto (\mathcal{R} \circ \mathcal{P}_{\mathfrak{C}} \circ f^{\leftarrow}(C))(x)$ is a probability measure on all of $\mathfrak{A}$: should $C \mapsto (\mathcal{R} \circ \mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow}(C))(x)$ be a probability measure on $\mathfrak{C}$ except for all x in a $\mu|_{\mathfrak{B}}^*$ -null set N , then choose $y \in X$ with $y \notin N$, and set

$$(\mathcal{R} \circ \mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow}(C))(x) = (\mathcal{R} \circ \mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow}(C))(y)$$


for all $x \in N$. Unless otherwise stated, all regular conditional distributions will be considered so selected.

We will call a conditional distribution $\mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow}$ of f on $\mathfrak{C}$ given $\mathfrak{B}$ a **regular conditional probability distribution**, or simply a *regular conditional distribution*, if there exists a regular version of the conditional distribution $\mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow}$. Should the function f here happen to equal the identity function, so that $\mathfrak{C} = \mathfrak{A}$, then this definition agrees with the Definition 8.10 of a regular conditional probability, as it should be.

Remark 8.20. No conditional distribution could possibly be a probability measure since the codomain of a conditional distribution is an L_1 space rather than the interval $[0, 1]$. However, a regular conditional distribution is equivalent to a particular Markov kernel defining a whole family of probability measures, as shown in Proposition 8.21.

{equivalent_definition_regular_distribution}

Proposition 8.21. Let $(X, \mathfrak{A}, \mu)$ be a probability space, and let $\mathfrak{B}$ be a σ -subalgebra of $\mathfrak{A}$. Also let $(Y, \mathfrak{C})$ be a measurable space, and let $f : X \rightarrow Y$ be an $(\mathfrak{A}, \mathfrak{C})$ -measurable function, with $f^{\leftarrow} : \mathfrak{C} \rightarrow \mathfrak{A}$. If, given $(X, \mathfrak{B})$ and $(Y, \mathfrak{C})$, there is a Markov kernel $k : X \times \mathfrak{C} \rightarrow [0, 1]$ with $k^C \in \mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow}(C)$ for each $C \in \mathfrak{C}$, then the collection $\{k^C : C \in \mathfrak{C}\}$ of $(\mathfrak{B}, \mathfrak{B} \circ \text{pr}_{[0,1]})$ -measurable functions is a regular version of the conditional distribution $\mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow}$ of f on $\mathfrak{C}$ given $\mathfrak{B}$, as illustrated:



Conversely, if the conditional distribution $\mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow}$ of f on $\mathfrak{C}$ given $\mathfrak{B}$ is regular, then given $(X, \mathfrak{B})$ and $(Y, \mathfrak{C})$ there is a Markov kernel $k : X \times \mathfrak{C} \rightarrow [0, 1]$ with $k^C \in \mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow}(C)$ for each $C \in \mathfrak{C}$.

Proof. Let $k : X \times \mathfrak{C} \rightarrow [0, 1]$ be a Markov kernel given $(X, \mathfrak{B})$ and $(Y, \mathfrak{C})$, with each $k^C \in \mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow}(C)$. To show that the collection $\{k^C : C \in \mathfrak{C}\}$ is a regular version of the conditional distribution $\mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow}$ of f on $\mathfrak{C}$

given $\mathfrak{B}$, see that we can use the hypothesis that $k^C \in \mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow}(C)$ to specify the selection function:

$$\mathcal{R} : L_1(X, \mathfrak{A}, \mu) \rightarrow \mathcal{L}_1(X, \mathfrak{A}, \mu) : \mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow}(C) \mapsto k^C.$$

By the hypothesis that $k : X \times \mathfrak{C} \rightarrow [0, 1]$ is a Markov kernel given $(X, \mathfrak{B})$ and $(Y, \mathfrak{C})$, this selection function has the additional properties that for each $C \in \mathfrak{C}$, the map

$$X \rightarrow [0, 1] : x \mapsto [(\mathcal{R} \circ \mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow})(C)](x) = k^C(x)$$

is $(\mathfrak{B}, \mathfrak{Bor}_{[0,1]})$ -measurable, and for each $x \in X$ the map

$$\mathfrak{C} \rightarrow [0, 1] : C \mapsto [(\mathcal{R} \circ \mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow})(C)](x) = k \circ \text{fix}_x(C)$$

is a probability measure on $\mathfrak{C}$, showing that $\{k^C : C \in \mathfrak{C}\}$ is a regular version of the conditional distribution $\mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow}$ of f on $\mathfrak{C}$ given $\mathfrak{B}$ by the very definition.

Now let $\mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow}$ be a regular conditional distribution of f on $\mathfrak{C}$ given $\mathfrak{B}$, which means we are supposing there is a selection function $\mathcal{R} : L_1(X, \mathfrak{A}, \mu) \rightarrow \mathcal{L}_1(X, \mathfrak{A}, \mu)$ with the additional properties that for each $C \in \mathfrak{C}$, the map

$$X \rightarrow [0, 1] : x \mapsto [(\mathcal{R} \circ \mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow})(C)](x)$$

is $(\mathfrak{B}, \mathfrak{Bor}_{[0,1]})$ -measurable, and for each $x \in X$ the map

$$\mathfrak{C} \rightarrow [0, 1] : C \mapsto [(\mathcal{R} \circ \mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow})(C)](x)$$

is a probability measure on $\mathfrak{C}$. It means that we can define the Markov kernel $k : X \times \mathfrak{C} \rightarrow [0, 1]$ given $(X, \mathfrak{B})$ and $(Y, \mathfrak{C})$ by

$$(x, C) \mapsto [(\mathcal{R} \circ \mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow})(C)](x).$$

By definition, the selection function has the property that

$$\mathcal{R}((\mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow})(C)) \in (\mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow})(C)$$

and since $k^C = \mathcal{R}((\mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow})(C))$, it follows that $k^C \in \mathcal{P}_{\mathfrak{B}} \circ f^{\leftarrow}(C)$ for each $C \in \mathfrak{C}$, as required. ■

Definition 8.22. Proposition 8.21 says that to each regular conditional distribution there corresponds a particular Markov kernel. We will call that kernel the **corresponding Markov kernel**.

{induced_marginals}


Proposition 8.23. Let λ be a measure on a σ -algebra $\mathfrak{D}$, and let $(f, g) : U \rightarrow X \times Y$ be a $(\mathfrak{D}, \sigma(\mathfrak{A}, \mathfrak{B}))$ -measurable function. Let the joint $\Gamma = \lambda \circ (f, g)^{\leftarrow}$, and let the marginal $\alpha = \Gamma \circ \pi_1^{\leftarrow}$ and the marginal $\beta = \Gamma \circ \pi_2^{\leftarrow}$. Then

$$\alpha = \lambda \circ f^{\leftarrow}$$

and

$$\beta = \lambda \circ g^{\leftarrow}$$

as illustrated:



Proof. This is just “set theory.” For example, while showing $\Gamma \circ \pi_1^{\leftarrow} = \lambda \circ f^{\leftarrow}$, you might use the fact that $\pi_1^{\leftarrow}(S) = S \times Y$ for any $S \subseteq X$. ■

{marginal_implies_markov}

Proposition 8.24. If $\Gamma \ll (\alpha \times \beta)^*$ on $\sigma(\mathfrak{A}, \mathfrak{B})$, then the map

$$Y \times \mathfrak{A} \rightarrow \mathbb{R} : (y, A) \mapsto \int_A \frac{d\Gamma}{d(\alpha \times \beta)^*}(x, y) d\alpha(x)$$

defines a Markov kernel $(Y, \mathfrak{B}) \prec (X, \mathfrak{A})$, and symmetrically, the map


$$X \times \mathfrak{A} \rightarrow \mathbb{R} : (x, B) \mapsto \int_B \frac{d\Gamma}{d(\alpha \times \beta)^*}(x, y) d\beta(y)$$

defines a Markov kernel $(X, \mathfrak{A}) \prec (Y, \mathfrak{B})$.

Proof. We will only show the first part, namely that we get a Markov kernel $(Y, \mathfrak{B}) \prec (X, \mathfrak{A})$. By hypothesis, $\Gamma \ll (\alpha \times \beta)^*$, and since the measure Γ is finite, and the measure $(\alpha \times \beta)^*$ is also finite, therefore σ -finite, we can take the Radon-Nikodym derivative $d\Gamma/d(\alpha \times \beta)^*$, and define:

$$j : Y \times \mathfrak{A} \rightarrow \mathbb{R} : (y, A) \mapsto \int_A \frac{d\Gamma}{d(\alpha \times \beta)^*}(x, y) d\alpha(x),$$

which is illustrated in a fashion similar to a Markov kernel:



At this stage, we can only say that the codomain of j is $\mathbb{R}$. We will show this map can be redefined on a subset of Y of β^* -measure zero, resulting in a Markov kernel.

By Proposition 5.31, $j^X(y) = 1$ for β^* -almost all $y \in Y$; say for all $y \in Y$ except for those y in a β^* -null subset N of Y . The set inclusion $A \subseteq X$ implies the inequality $j^A \leq j^X$, and so $0 \leq j_y \leq 1$ for all $y \notin N$. It means that we can redefine the codomain of each such j_y to be the interval $[0, 1]$, and it implies that each such j_y is a probability measure. Fix $z \in Y$ with $z \notin N$, so j_z is a probability measure. For each $y \in N$, redefine $j_y(A) := j_z(A)$, so now all the j_y are probability measures. In summary,

$$j_y : \mathfrak{A} \rightarrow [0, 1] : A \mapsto \begin{cases} \int_A \frac{d\Gamma}{d(\alpha \times \beta)^*}(x, y) d\alpha(x) & \text{if } y \notin N \\ j_z(A) & \text{for that fixed } z \quad \text{if } y \in N \end{cases}$$

The representative function $d\Gamma/(\alpha \times \beta)^*$ is integrable with respect to the measure $\alpha \times \beta$, and this means by Fubini's 5.20 that the iterated integrals exist and in particular for each $A \in \mathfrak{A}$, the map j^A defines

an integrable function over Y with respect to the measure β . By Proposition 3.36, each j^A can be taken $(\mathfrak{B}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable. We can redefine the codomain of each j^A to be $[0, 1]$ rather than all of $\mathbb{R}$ by modifying each function j^A on that β^* -null set N . Redefine $j^A(y) := j^A(z)$ if $y \in N$, where the $z \in Y$ with $z \notin N$ is the same fixed element from the previous paragraph. We know that $j^A(z) = j_z(A)$ by their definitions, and $j_z(A) \in [0, 1]$ implies $j^A(z) \in [0, 1]$, so the codomain of each j^A can be taken $[0, 1]$. And each such function j^A is $(\mathfrak{B}, \mathfrak{Bor}_{[0,1]})$ -measurable. In summary,

$$j^A : Y \rightarrow [0, 1] : y \mapsto \begin{cases} \int_A \frac{d\Gamma}{d(\alpha \times \beta)^*}(x, y) d\alpha(x) & \text{if } y \notin N \\ j^A(z) & \text{for that fixed } z \quad \text{if } y \in N \end{cases}$$

This gives us a Markov kernel $(Y, \mathfrak{B}) \prec (X, \mathfrak{A})$:

$$\begin{array}{ccc} Y & & \\ \text{fix}^A \downarrow & \searrow j^A & \\ Y \times \mathfrak{A} & \xrightarrow{j} & [0, 1] \\ \text{fix}_y \uparrow & \nearrow j_y & \\ \mathfrak{A} & & \end{array}$$

as required. ■

{markov_composition_measurable}

Proposition 8.25. *If $j : Y \times \mathfrak{A} \rightarrow [0, 1]$ is a Markov kernel $(Y, \mathfrak{B}) \prec (X, \mathfrak{A})$, and if $g : (U, \mathfrak{G}) \rightarrow (Y, \mathfrak{B})$ is measurable, then*

$$k : U \times \mathfrak{A} \rightarrow [0, 1] : (u, A) \mapsto j(g(u), A)$$


is a Markov kernel $(U, \mathfrak{G}) \prec (X, \mathfrak{A})$.

{marginal_implies_regular}

Proposition 8.26. *Let $(U, \mathfrak{D}, \lambda)$ be a probability space, with functions $f : (U, \mathfrak{D}) \rightarrow (X, \mathfrak{A})$ and $g : (U, \mathfrak{D}) \rightarrow (Y, \mathfrak{B})$ measurable. Set $\mathfrak{G} = g^{\leftarrow}(\mathfrak{B})$. Let Γ denote the joint measure $\lambda \circ (f, g)^{\leftarrow}$, and let α denote the marginal measure $\Gamma \circ \pi_1^{\leftarrow}$, and let β denote the marginal measure $\Gamma \circ \pi_2^{\leftarrow}$.*

If $\Gamma \ll (\alpha \times \beta)^*$ on $\sigma(\mathfrak{A}, \mathfrak{B})$, then the conditional distribution $\mathcal{P}_{\mathfrak{G}} \circ f^{\leftarrow}$ of f on $\mathfrak{A}$ given $\mathfrak{G}$ is regular, and symmetrically, setting $\mathfrak{F} = f^{\leftarrow}(\mathfrak{A})$, the conditional distribution $\mathcal{P}_{\mathfrak{F}} \circ g^{\leftarrow}$ of g on $\mathfrak{B}$ given $\mathfrak{F}$ is regular.

Proof. Let $\Gamma \ll (\alpha \times \beta)^*$. We have set $\Gamma = \lambda \circ (f, g)^{\leftarrow}$:




We will define a map $k : U \times \mathfrak{A} \rightarrow [0, 1]$, then show it is a Markov kernel, and then apply Proposition 8.21.

By Proposition 8.24, the map

$$j : Y \times \mathfrak{A} \rightarrow \mathbb{R} : (y, A) \mapsto \int_A \frac{d\Gamma}{d(\alpha \times \beta)^*}(x, y) d\alpha(x)$$

defines the Markov kernel:



By hypothesis, the function $g : U \rightarrow Y$ is $(\mathfrak{D}, \mathfrak{B})$ -measurable, and therefore $(\mathfrak{G}, \mathfrak{B})$ -measurable since $\mathfrak{G} = g^{\leftarrow}(\mathfrak{B})$. By Proposition 8.25, the map

$$k : U \times \mathfrak{A} \rightarrow [0, 1] : (u, A) \mapsto j(g(u), A)$$

is a Markov kernel $(U, \mathfrak{G}) \prec (Y, \mathfrak{A})$. Verify that $k^A \in \mathcal{P}_{\mathfrak{G}} \circ f^{\leftarrow}(A)$ for each $A \in \mathfrak{A}$, so the collection of k^A is a version of $\mathcal{P}_{\mathfrak{G}} \circ f^{\leftarrow}$. It follows by Proposition 8.21 that $\mathcal{P}_{\mathfrak{G}} \circ f^{\leftarrow}$ is regular, as required. ■


{mixed_implies_bayes}

Proposition 8.27. *Let $(U, \mathfrak{D}, \lambda)$ be a probability space, with functions $f : (U, \mathfrak{D}) \rightarrow (X, \mathfrak{A})$ and $g : (U, \mathfrak{D}) \rightarrow (Y, \mathfrak{B})$ measurable. Set $\mathfrak{G} = g^{\leftarrow}(\mathfrak{B})$. Let Γ denote the joint measure $\lambda \circ (f, g)^{\leftarrow}$, and let α denote the marginal measure $\Gamma \circ \pi_1^{\leftarrow}$, and let β denote the marginal measure $\Gamma \circ \pi_2^{\leftarrow}$. Let ν be a σ -finite measure on $\mathfrak{B}$.*


Let $\Gamma \ll (\alpha \times \nu)^$ on $\sigma(\mathfrak{A}, \mathfrak{B})$, and let k denote the Markov kernel corresponding to the regular conditional distribution $\mathcal{P}_{\mathfrak{G}} \circ f^{\leftarrow}$ of f on $\mathfrak{A}$ given $\mathfrak{G}$. If $j^A : Y \rightarrow [0, 1]$ is any collection of $(\mathfrak{B}, \mathfrak{Bor}_{[0,1]})$ -measurable functions such that $k^A = j^A \circ g$, then for ν^* -almost all $y \in Y$,*

$$j^A(y) = \int_A \frac{d\Gamma}{d(\alpha \times \beta)^*}(x, y) d\alpha(x).$$

Proof. With $\Gamma \ll (\alpha \times \nu)^*$ on $\sigma(\mathfrak{A}, \mathfrak{B})$, it follows by Proposition 5.28 that $\Gamma \ll (\alpha \times \beta)^*$, and so by Proposition 8.26, the conditional distribution $\mathcal{P}_{\mathfrak{G}} \circ f^{\leftarrow}$ of f on $\mathfrak{A}$ given $\mathfrak{G}$ is regular. We are letting k denote the Markov kernel corresponding to $\mathcal{P}_{\mathfrak{G}} \circ f^{\leftarrow}$, illustrated as:



Let $j^A : Y \rightarrow [0, 1]$ be any collection of $(\mathfrak{B}, \mathfrak{Bor}_{[0,1]})$ -measurable functions such that $k^A = j^A \circ g$, as illustrated:



With $\Gamma \ll (\alpha \times \nu)^*$, we can apply Proposition 5.25 to get $\beta \ll \nu$ on $\mathfrak{B}$. And in turn, we can apply the Radon-Nikodym corollary, namely Proposition 4.30 to get

$$\int_B j^A \frac{d\beta}{d\nu} d\nu = \int_B j^A d\beta.$$

Applying a change of variable according to Proposition 4.13,

$$\int_B j^A d\beta = \int_{g^{\leftarrow}(B)} k^A d\lambda.$$

With $k^A \in \mathcal{P}_{\mathfrak{G}} \circ f^{\leftarrow}(A)$, it follows that

$$\int_{g^{\leftarrow}(B)} k^A d\lambda = \int_{g^{\leftarrow}(B)} (\mathcal{P}_{\mathfrak{G}} \circ f^{\leftarrow})(A) d\lambda.$$

By Proposition 8.18,

$$\int_{g^{\leftarrow}(B)} (\mathcal{P}_{\mathfrak{G}} \circ f^{\leftarrow})(A) d\lambda = \lambda(g^{\leftarrow}(B) \cap f^{\leftarrow}(A)).$$

Since $\Gamma = \lambda \circ (f, g)^{\leftarrow}$,

$$\lambda(g^{\leftarrow}(B) \cap f^{\leftarrow}(A)) = \Gamma(A \times B).$$

By transitivity of equality, we have shown so far that

$$\Gamma(A \times B) = \int_B j^A \frac{d\beta}{d\nu} d\nu.$$

Again, with $\Gamma \ll (\alpha \times \nu)^*$, we can apply Fubini's Theorem 5.20 to get

$$\Gamma(A \times B) = \int_B \int_A \frac{d\Gamma}{d(\alpha \times \nu)^*} d\alpha d\nu.$$

This says that

$$\int_B \left(j^A \frac{d\beta}{d\nu} \right) d\nu = \int_B \left(\int_A \frac{d\Gamma}{d(\alpha \times \nu)^*} d\alpha \right) d\nu$$

for all $B \in \mathfrak{B}$. By Proposition 3.41,

$$j^A(y) \frac{d\beta}{d\nu}(y) = \int_A \frac{d\Gamma}{d(\alpha \times \nu)^*}(x, y) d\alpha(x)$$

for ν^* -almost all $y \in Y$, and by Proposition 5.25,

$$\frac{d\beta}{d\nu}(y) = \int_X \frac{d\Gamma}{d(\alpha \times \nu)^*}(x, y) d\alpha(x),$$

so

$$j^A(y) = \frac{\int_A \frac{d\Gamma}{d(\alpha \times \nu)^*}(x, y) d\alpha(x)}{\int_X \frac{d\Gamma}{d(\alpha \times \nu)^*}(x, y) d\alpha(x)}$$

for ν^* -almost all $y \in Y$. By Proposition 5.30,

$$\int_A \frac{d\Gamma}{d(\alpha \times \beta)^*}(x, y) d\alpha(x) = \frac{\int_A \frac{d\Gamma}{d(\alpha \times \nu)^*}(x, y) d\alpha(x)}{\int_X \frac{d\Gamma}{d(\alpha \times \nu)^*}(x, y) d\alpha(x)}$$

so

$$j^A(y) = \int_A \frac{d\Gamma}{d(\alpha \times \beta)^*}(x, y) d\alpha(x)$$

for ν^* -almost all $y \in Y$, as required. ■

8.5 Conditional Densities

Remark 8.28. Proposition 8.29 and Proposition 8.31 are supporting lemmas for the conditional independence equivalences given in Proposition 11.6.

{joint_decomposition}

Proposition 8.29. *Let $(U, \mathfrak{D}, \lambda)$ be a probability space, and let the functions $f : (U, \mathfrak{D}) \rightarrow (X, \mathfrak{A})$ and $g : (U, \mathfrak{D}) \rightarrow (Y, \mathfrak{B})$ be measurable. Set $\mathfrak{G} = g^{\leftarrow}(\mathfrak{B})$. Let Γ denote the joint measure $\lambda \circ (f, g)^{\leftarrow}$, and let β denote the marginal measure $\lambda \circ g^{\leftarrow}$. Let the conditional distribution*

$$\mathcal{P}_{\mathfrak{G}} \circ f^{\leftarrow} : \mathfrak{A} \rightarrow L_1(U, \mathfrak{D}, \lambda)$$

of f on $\mathfrak{A}$ given $\mathfrak{G}$ be regular, and let k denote the corresponding Markov kernel given $(U, \mathfrak{G})$ and $(X, \mathfrak{A})$ such that $k^A \in \mathcal{P}_{\mathfrak{G}} \circ f^{\leftarrow}(A)$ for all $A \in \mathfrak{A}$.

If a function $j : X \times Y \rightarrow \mathbb{R}$ is integrable over $X \times Y$ with respect to the joint measure Γ , then


$$\int_{X \times Y} j \, d\Gamma = \int_Y \int_X j(x, z) \, dk_u(x) \, d\beta(y),$$

where $g(u) = y$.

Proof. We have set $\mathfrak{G} = g^{\leftarrow}(\mathfrak{B})$ because subscripts do not suffer symbols like $g^{\leftarrow}(\mathfrak{B})$ or $\sigma(g)$ well. Let the conditional distribution

$$\mathcal{P}_{\mathfrak{G}} \circ f^{\leftarrow} : \mathfrak{A} \rightarrow L_1(U, \mathfrak{D}, \lambda)$$

of f on $\mathfrak{A}$ given $\mathfrak{G}$ be regular, and we let k denote the corresponding Markov kernel given $(U, \mathfrak{G})$ and $(X, \mathfrak{A})$, as illustrated:



For each set $A \in \mathfrak{A}$, the function k^A factors as some function a composed with g , that is, $k^A = a \circ g$, which was the heart of Proposition 4.36:

$$\begin{array}{ccc} (U, \mathfrak{G}) & \xrightarrow{k^A} & ([0, 1], \mathfrak{Bor}_{[0,1]}) \\ & \searrow g & \uparrow a \\ & & (Y, \mathfrak{B}) \end{array}$$

The evaluation $k^A(u) = a(g(u))$ may be restated as $k^A(u) = a(y)$ if $g(u) = y$. Since $k^A(u) = k_u(A)$, we can write

$$a(y) = k_u(A) \text{ for } y = g(u).$$

It is sufficient to show that the required equality holds for characteristic functions of the form $j = \chi_{A \times B}$ with $A \in \mathfrak{A}$ and $B \in \mathfrak{B}$. Set $j = \chi_{A \times B}$, so

$$\int_{X \times Y} j \, d\Gamma = \lambda(f^{\leftarrow}(A) \cap g^{\leftarrow}(B)).$$

By Proposition 8.18,

$$\lambda(f^{\leftarrow}(A) \cap g^{\leftarrow}(B)) = \int_{g^{\leftarrow}(B)} \bar{\mathcal{P}}_{\mathfrak{G}} \circ f^{\leftarrow}(A) \, d\lambda.$$

The hypothesis that the conditional distribution

$$\bar{\mathcal{P}}_{\mathfrak{G}} \circ f^{\leftarrow} : \mathfrak{A} \rightarrow L_1(U, \mathfrak{D}, \lambda)$$

of f on $\mathfrak{A}$ given $\mathfrak{G}$ is regular means that $k^A \in \bar{\mathcal{P}}_{\mathfrak{G}} \circ f^{\leftarrow}(A)$, so

$$\int_{g^{\leftarrow}(B)} \bar{\mathcal{P}}_{\mathfrak{G}} \circ f^{\leftarrow}(A) \, d\lambda = \int_{g^{\leftarrow}(B)} k^A \, d\lambda,$$

and with $k^A = a \circ g$,

$$\int_{g^{\leftarrow}(B)} k^A \, d\lambda = \int_{g^{\leftarrow}(B)} (a \circ g) \, d\lambda.$$

Use the fact that

$$\begin{aligned}\chi_{g^{\leftarrow}(B)} \cdot (a \circ g) &= (\chi_B \circ g) \cdot (a \circ g) \\ &= (\chi_B \cdot a) \circ g\end{aligned}$$

and write

$$\int_{g^{\leftarrow}(B)} (a \circ g) d\lambda = \int_U (\chi_B \cdot a) \circ g d\lambda.$$

Apply a change of variable:

$$\begin{aligned}\int_U (\chi_B \cdot a) \circ g d\lambda &= \int_Y (\chi_B \cdot a) d\beta \\ &= \int_Y \chi_B(y) \cdot a(y) d\beta(y).\end{aligned}$$

We know that

$$a(y) = k_u(A) = \int_X \chi_A(x) dk_u(x)$$

for $y = g(u)$. By substitution for $a(y)$:

$$\int_Y \chi_B(y) \cdot a(y) d\beta(y) = \int_Y \chi_B(y) \left(\int_X \chi_A(x) dk_u(x) \right) d\beta(y).$$

By linearity of the integral, we can rewrite:

$$\int_Y \chi_B(y) \left(\int_X \chi_A(x) dk_u(x) \right) d\beta(y) = \int_Y \int_X \chi_A(x) \chi_B(y) dk_u(x) d\beta(y).$$

And since $\chi_A(x) \chi_B(y) = \chi_{A \times B}(x, y)$:

$$\begin{aligned}\int_Y \int_X \chi_A(x) \chi_B(y) dk_u(x) d\beta(y) &= \int_Y \int_X \chi_{A \times B}(x, y) dk_u(x) d\beta(y) \\ &= \int_Y \int_X j(x, y) dk_u(x) d\beta(y).\end{aligned}$$

By transitivity of equality then,

$$\int_{X \times Y} j \, d\Gamma = \int_Y \int_X j(x, y) \, dk_u(x) \, d\beta(y),$$

as required. ■

Remark 8.30. Just as in Remark 6.7 where the expected value could be computed as an integral with respect to a probability distribution, the regression function can be computed with respect to a probability measure corresponding to a conditional distribution.

Recall from Definition 7.6 how the regression function r was defined. Let $(U, \mathfrak{D}, \lambda)$ be a probability space. Let functions $f : U \rightarrow X$ and $g : U \rightarrow Y$ be measurable. Let $h : Y \rightarrow \mathbb{R}$. Setting $\mathfrak{F} = f^{\leftarrow}(\mathfrak{A})$, let

$$E(\overline{h \circ g}^\lambda \mid f) = \overline{r \circ f}^{\lambda|_{\mathfrak{F}}}$$

with the function r being an $(\mathfrak{A}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable function, as illustrated in the following diagram, letting the function e denote any representative of the class $E(\overline{h \circ g}^\lambda \mid f)$:

$$\begin{array}{ccc} (U, \mathfrak{D}) & \xrightarrow{h \circ g} & (\mathbb{R}, \mathfrak{Bor}_{\mathbb{R}}) \\ & \searrow f & \uparrow r \\ (U, \mathfrak{F}) & \xrightarrow{e} & (\mathbb{R}, \mathfrak{Bor}_{\mathbb{R}}) \end{array} \quad \text{meaning} \quad E(\overline{h \circ g}^\lambda \mid f) = \overline{r \circ f}^{\lambda|_{\mathfrak{F}}}$$

$(X, \mathfrak{A})$

We are calling the induced function r the *regression function*. Under certain conditions, this function can be computed as

$$r(x) = \int_U (h \circ g) \, dk_u \quad \text{where } f(u) = x$$

for $\lambda \circ f^{\leftarrow}$ almost all $x \in X$. In this case, k is the Markov kernel corresponding to the conditional probability distribution $\mathcal{P}_{\mathfrak{F}} \circ g^{\leftarrow}$. Very

roughly, this might involve two changes of variable, and an application of Proposition 8.29. The changes of variable look like

$$\int_{f^{\leftarrow}(C)} (r \circ f) d\lambda = \int_C r d(\lambda \circ f^{\leftarrow})$$

and

$$\int_{f^{\leftarrow}(C)} (h \circ g) d\lambda = \int_{C \times Y} (h \circ \pi_2) d(\lambda \circ (f, g)^{\leftarrow}).$$

It seems the only condition is the function $h \circ \pi_2$ should be integrable with respect to the joint measure $\lambda \circ (f, g)^{\leftarrow}$.

If the idea that the regression function can be computed this way turns out to be useful, then we will come back and fill in the details.


{conditional_density}

Proposition 8.31. *Let $(U, \mathfrak{D}, \lambda)$ be a probability space, and let functions $f : (U, \mathfrak{D}) \rightarrow (X, \mathfrak{A})$ and $g : (U, \mathfrak{D}) \rightarrow (Y, \mathfrak{B})$ be measurable. Set $\mathfrak{G} = g^{\leftarrow}(\mathfrak{B})$. Let Γ denote the joint measure $\lambda \circ (f, g)^{\leftarrow}$, and let β denote the marginal measure $\lambda \circ g^{\leftarrow}$. Let μ be a σ -finite measure on $\mathfrak{A}$.*

If the joint measure $\Gamma \ll (\mu \times \beta)^$, where k denotes the corresponding Markov kernel given $(U, \mathfrak{G})$ and $(X, \mathfrak{A})$ such that $k^A \in \mathcal{P}_{\mathfrak{G}} \circ f^{\leftarrow}(A)$ for all $A \in \mathfrak{A}$, then for β^* -almost all $y \in Y$, each measure k_u on $\mathfrak{A}$ such that $g(u) = y$ has the following density function with respect to the measure μ :*

$$\frac{dk_u}{d\mu} : X \rightarrow \mathbb{R} : x \mapsto \frac{d\Gamma}{d(\mu \times \beta)^*}(x, y).$$

Proof. Let $\Gamma \ll (\mu \times \beta)^*$, with a reminder that μ is a reference measure on $\mathfrak{A}$, and β is a marginal measure on $\mathfrak{B}$. We are letting k denote the corresponding Markov kernel given $(U, \mathfrak{G})$ and $(X, \mathfrak{A})$, as illustrated:



It is sufficient to show that for β^* -almost all $y \in Y$, if $A \in \mathfrak{A}$, and $y = g(u)$, then

$$k_u(A) = \int_A \frac{d\Gamma}{d(\mu \times \beta)^*}(x, y) d\mu(x).$$

Let $A \in \mathfrak{A}$ and $B \in \mathfrak{B}$. Then $\chi_{A \times B}$ is integrable with respect to the joint measure Γ . Since $\Gamma \ll (\mu \times \beta)^*$, it follows by Proposition 4.30 that

$$\int_{X \times Y} \chi_{A \times B} d\Gamma = \int_{X \times Y} \chi_{A \times B} \frac{d\Gamma}{d(\mu \times \beta)^*} d(\mu \times \beta).$$

By Fubini's Theorem 5.20,

$$\int_{X \times Y} \chi_{A \times B} \frac{d\Gamma}{d(\mu \times \beta)^*} d(\mu \times \beta) = \int_Y \int_X \chi_{A \times B} \frac{d\Gamma}{d(\mu \times \beta)^*} d\mu d\beta.$$

But also, by Proposition 8.29,

$$\int_{X \times Y} \chi_{A \times B} d\Gamma = \int_Y \int_X \chi_{A \times B}(x, y) dk_u(x) d\beta(y),$$

where $g(u) = y$. By transitivity of equality, this shows that

$$\int_Y \int_X \chi_{A \times B} \frac{d\Gamma}{d(\mu \times \beta)^*} d\mu d\beta = \int_Y \int_X \chi_{A \times B}(x, y) dk_u(x) d\beta(y),$$

where $g(u) = y$. We can rewrite this as integration over sets:

$$\int_B \int_A \frac{d\Gamma}{d(\mu \times \beta)^*} d\mu d\beta = \int_B \int_A 1_X(x) dk_u(x) d\beta(y),$$

where $g(u) = y$. Since this is true for all $B \in \mathfrak{B}$, it follows by Proposition 4.9 that

$$\int_A \frac{d\Gamma}{d(\mu \times \beta)^*}(x, y) d\mu(x) = \int_A 1_X dk_u$$

for β^* -almost all $y \in Y$, and where $g(u) = y$. See that

$$\int_A 1_X dk_u = k_u(A).$$

This means we have shown that

$$k_u(A) = \int_A \frac{d\Gamma}{d(\mu \times \beta)^*}(x, y) d\mu(x)$$

for β^* -almost all $y \in Y$, and where $g(u) = y$, as required. ■

9 Bayesian Statistics

{bayesian_statistics}

Fix the Markov kernels.


STATISTICS is a mathematical analysis of guesswork. This kind of guesswork is called inductive inference. Its purpose is to identify a probability distribution or a conditional distribution. The particular kind of guesswork we will use is considered *Bayesian* since it involves a parameter that induces a probability distribution. A parameter here is just a measurable function.

There are no topological restrictions imposed on the objects in this chapter. No Borel space conditions, no separable conditions. We rely instead upon measure-theoretic conditions.

This chapter is intended to provide the smallest possible coherent glimpse of Bayesian statistics.

Suppose $(U, \mathfrak{D})$ and $(X, \mathfrak{A})$ are measurable spaces. Let $(U, \mathfrak{D}, \lambda)$ be a probability space, and let $\mathfrak{G}$ be a σ -subalgebra of $\mathfrak{D}$. The conditional probability $\mathcal{P}_{\mathfrak{G}} : \mathfrak{D} \rightarrow L_1(U, \mathfrak{D}, \lambda)$ on $\mathfrak{D}$ given $\mathfrak{G}$ could not possibly be a probability measure since the codomain of the conditional probability $\mathcal{P}_{\mathfrak{G}}$ is the quotient space $L_1(U, \mathfrak{D}, \lambda)$ rather than the interval $[0, 1]$. For the same reason, no conditional distribution $\mathcal{P}_{\mathfrak{G}} \circ f^{\leftarrow} : \mathfrak{A} \rightarrow L_1(U, \mathfrak{D}, \lambda)$ of f on $\mathfrak{A}$ given $\mathfrak{G}$ could possibly be a probability measure, with $f : U \rightarrow X$ here being any $(\mathfrak{D}, \mathfrak{A})$ -measurable function. However, a *regular* conditional distribution corresponds to a Markov kernel k defining a

whole family of probability measures on the σ -algebra $\mathfrak{A}$:



By some schools of thought, just a family of probability measures itself, indexed by a parameter space, is considered to be a statistical experiment.

Let

$$(X \times Y, \sigma(\mathfrak{A}, \mathfrak{B}), \Gamma)$$


be a probability space. The measure Γ is called the *joint*. Call the measurable space $(X, \mathfrak{A})$ the *attribute space*, or *parameter space*, and call the measurable space $(Y, \mathfrak{B})$ the *sample space*. Let π_1 and π_2 denote the respective coordinate projections of $X \times Y$ onto X and Y . Also let $\pi_1^{\leftarrow}$ and $\pi_2^{\leftarrow}$ denote the respective induced maps

$$\pi_1^{\leftarrow} : \mathfrak{A} \rightarrow \sigma(\mathfrak{A}, \mathfrak{B}) : A \mapsto A \times Y$$

and

$$\pi_2^{\leftarrow} : \mathfrak{B} \rightarrow \sigma(\mathfrak{A}, \mathfrak{B}) : B \mapsto X \times B,$$

as illustrated:



The projection π_1 is $(\sigma(\mathfrak{A}, \mathfrak{B}), \mathfrak{A})$ -measurable, and induces the marginal probability distribution $\alpha := \Gamma \circ \pi_1^\leftarrow$ of π_1 on the parameter space σ -algebra $\mathfrak{A}$, which is called the *prior probability distribution*. The projection π_2 is $(\sigma(\mathfrak{A}, \mathfrak{B}), \mathfrak{B})$ -measurable, and induces the marginal probability distribution $\beta := \Gamma \circ \pi_2^\leftarrow$ of π_2 on the sample space σ -algebra $\mathfrak{B}$, which is called the *predictive probability distribution*. Consequently:

$$\alpha(A) = (\Gamma \circ \pi_1^\leftarrow)(A) = \Gamma(A \times Y) \quad \text{for all } A \in \mathfrak{A},$$

and

$$\beta(B) = (\Gamma \circ \pi_2^\leftarrow)(B) = \Gamma(X \times B) \quad \text{for all } B \in \mathfrak{B}.$$

Call the conditional distribution $\mathcal{P}_{\pi_1^\leftarrow(\mathfrak{A})} \circ \pi_2^\leftarrow$ of π_2 on $\mathfrak{B}$ given $\pi_1^\leftarrow(\mathfrak{A})$ the **sampling conditional distribution**, as illustrated:

$$\begin{array}{ccc} L_1(X \times Y, \sigma(\mathfrak{A}, \mathfrak{B}), \Gamma) & \xrightarrow{\mathcal{E}_{\pi_1^\leftarrow(\mathfrak{A})}} & L_1(X \times Y, \sigma(\mathfrak{A}, \mathfrak{B}), \Gamma) \\ \uparrow Q \circ \chi & \nearrow \mathcal{P}_{\pi_1^\leftarrow(\mathfrak{A})} & \uparrow \mathcal{P}_{\pi_1^\leftarrow(\mathfrak{A})} \circ \pi_2^\leftarrow \\ \sigma(\mathfrak{A}, \mathfrak{B}) & \xleftarrow{\pi_2^\leftarrow} & \mathfrak{B} \end{array}$$

Call the conditional distribution $\mathcal{P}_{\pi_2^\leftarrow(\mathfrak{B})} \circ \pi_1^\leftarrow$ of π_1 on $\mathfrak{A}$ given $\pi_2^\leftarrow(\mathfrak{B})$ the **posterior conditional distribution**, as illustrated:

$$\begin{array}{ccc} L_1(X \times Y, \sigma(\mathfrak{A}, \mathfrak{B}), \Gamma) & \xrightarrow{\mathcal{E}_{\pi_2^\leftarrow(\mathfrak{B})}} & L_1(X \times Y, \sigma(\mathfrak{A}, \mathfrak{B}), \Gamma) \\ \uparrow Q \circ \chi & \nearrow \mathcal{P}_{\pi_2^\leftarrow(\mathfrak{B})} & \uparrow \mathcal{P}_{\pi_2^\leftarrow(\mathfrak{B})} \circ \pi_1^\leftarrow \\ \sigma(\mathfrak{A}, \mathfrak{B}) & \xleftarrow{\pi_1^\leftarrow} & \mathfrak{A} \end{array}$$

The sampling and posterior conditional distributions could not possibly be probability measures, and this should be clear since their

codomains are L_1 spaces rather than the interval $[0, 1]$. However, were they *regular* conditional distributions, then they would induce whole families of probability measures on the parameter and sample space σ -algebras by way of their corresponding Markov kernels. What measure-theoretic conditions might be sufficient to imply that these conditional distributions are regular?

One simple condition is when $\Gamma \ll (\alpha \times \beta)^*$. Then both the sampling conditional distribution and the posterior conditional distribution are regular, which is the heart of Proposition 8.26.

Here, in turn, is a measure-theoretic condition sufficient to imply that $\Gamma \ll (\alpha \times \beta)^*$. Suppose $\{P_x : x \in X\}$ is a family of probability measures on the sample space σ -algebra $\mathfrak{B}$, called the **sampling probabilities**, and suppose that the σ -algebra $\mathfrak{A}$ is such that for each $B \in \mathfrak{B}$, the function defined by

$$P^B : X \rightarrow [0, 1] : x \mapsto P_x(B)$$

is $(\mathfrak{A}, \mathfrak{Bor}_{[0,1]})$ -measurable. Also suppose the joint probability measure Γ is defined in terms of a given *prior* probability measure γ on the parameter space σ -algebra $\mathfrak{A}$ by

$$\Gamma(A \times B) := \int_A P^B(x) d\gamma(x) \quad \text{for all } B \in \mathfrak{B} \text{ and all } A \in \mathfrak{A}.$$

Then, by Proposition 8.7, the given *prior* probability measure γ must equal the marginal, or *prior*, probability distribution α . If also ν is a σ -finite measure on $\mathfrak{B}$ such that each $P_x \ll \nu$, and if $f : X \times Y \rightarrow \mathbb{R}$ is a $(\sigma(\mathfrak{A}, \mathfrak{B}), \mathfrak{Bor}_{\mathbb{R}})$ -measurable function such that each $f_x : Y \rightarrow \mathbb{R}$ is equal to the Radon-Nikodym derivative:

$$(9.1) \quad \{\text{as_density}\} \quad f_x = dP_x/d\nu,$$

then $\Gamma \ll (\alpha \times \nu)^*$ with $f = d\Gamma/d(\alpha \times \nu)^*$, and so by Proposition 5.28, $\Gamma \ll (\alpha \times \beta)^*$. Further, if the function $g : X \times Y \rightarrow \mathbb{R}$ is defined by setting

$$g(x, y) = \frac{f(x, y)}{\int_X f(x, y) d\alpha(x)},$$

then $g = d\Gamma/d(\alpha \times \beta)^*$ by Proposition 5.28.

The family $\{P_x : x \in X\}$ of probability measures on the sample space σ -algebra $\mathfrak{B}$ must satisfy

$$P_x(B) = \int_B g(x, y) d\beta(y) \quad \text{for all } B \in \mathfrak{B},$$

by Proposition 5.29, and the family $\{\alpha_y : y \in Y\}$ of probability measures on the parameter space σ -algebra $\mathfrak{A}$, called the **posterior probabilities**, defined for those $y \in Y$ such that f^y is integrable with respect to α , is defined by

$$(9.2) \quad \alpha_y(A) = \int_A g(x, y) d\alpha(x) \quad \text{for all } A \in \mathfrak{A},$$

or equivalently by

$$\frac{d\alpha_y}{d\alpha} : X \rightarrow \mathbb{R} : x \mapsto \frac{f(x, y)}{\int_X f(x, y) d\alpha(x)}.$$


The equality (9.2) may also be seen as Bayes' Theorem in the case where there is a σ -finite measure λ on $\mathfrak{A}$ with $\alpha \ll \lambda$, say with density function $h = d\alpha/d\lambda \in \mathcal{L}_1(X, \mathfrak{A}, \lambda)$. In this case,

$$\alpha_y(A) = \frac{\int_A f(x, y) h(x) d\lambda(x)}{\int_X f(x, y) h(x) d\lambda(x)} \quad \text{for all } A \in \mathfrak{A},$$

or equivalently,

$$\begin{aligned} \frac{d\alpha_y}{d\lambda} : X \rightarrow \mathbb{R} : x \mapsto & \frac{f(x, y) h(x)}{\int_X f(x, y) h(x) d\lambda(x)} \\ & = \frac{f^y(x) h(x)}{\int_X f^y(x) h(x) d\lambda(x)}, \end{aligned}$$

where $f(x, y)$, or in particular f^y , might be referred to as “the likelihood,” illustrated in the following diagram:



The collection $\{f^y : y \in Y\}$ of likelihood functions $X \rightarrow \mathbb{R}$ is indexed by the sample space Y , which is one way that a particular sample y may be used in applications; the sample itself determines the likelihood function by indexing into the collection. It is interesting how the sample, or **observation**, or **data**, can be seen as selecting a likelihood function from a collection, and warrants further thought. On the other hand, you always need information to make an intelligent selection. Perhaps that is all there is to it.

10 Bayesian Statistical Models

{bayesian_statistical_models_examples}

10.1 Outline

Here is a partial list of steps that will be applied to each of the examples that follow:

- Establish an indexed, or parametrized, family $\{P_x : x \in X\}$ of sampling probabilities on the sample space σ -algebra $\mathfrak{B}$. Hopefully, each P_x is absolutely continuous with respect to a σ -finite reference measure on $\mathfrak{B}$.
- Verify that some function $f : X \times Y \rightarrow \mathbb{R}$ has f_x as the density function for P_x with respect to the reference measure.
- Define a prior probability measure on the parameter space σ -algebra $\mathfrak{A}$. Say that h is a density function for the prior with respect to a σ -finite reference measure on $\mathfrak{A}$, temporarily call this measure ν .
- Then understand that the map

$$X \rightarrow \mathbb{R} : x \mapsto \frac{f(x, y)h(x)}{\int_X f(x, y)h(x) d\nu(x)} = \frac{f^y(x)h(x)}{\int_X f^y(x)h(x) d\nu(x)}$$

is a density function for the posterior probability measure of the parameter given the data, with respect to ν on $\mathfrak{A}$.

10.2 Globe Tossing Model

{globe_tossing}

This example of an experiment is highly paraphrased from Richard McElreath's *Statistical Rethinking, Second Edition* [6]. The experiment

is tossing a globe N times into the air, catching it, and wherever an index finger happens to touch the surface of the globe, record *water* should this surface corresponds to water, and record *land* should it correspond to land. This N -tuple is the sample. Deduce from this experiment the probability distribution of the proportion p of the surface covered with water. This proportion p is the parameter. Quoting from *Statistical Rethinking* [6], page 78, verbatim:

4.2.1. Re-describing the globe tossing model. It's good to work with examples. Recall the proportion of water problem from previous chapters. The model in that case was always:

$$\begin{aligned} W &\sim \text{Binomial}(N, p) \\ p &\sim \text{Uniform}(0, 1) \end{aligned}$$

where W was the observed count of water, N was the total number of tosses, and p was the proportion of water on the globe. Read the above statement as:


The count W is distributed binomially with sample size N and probability p .

The prior for p is assumed to be uniform between zero and one.

And in a display following this, find:

$$\Pr(p|w, n) = \frac{\text{Binomial}(w|n, p) \text{Uniform}(p|0, 1)}{\int \text{Binomial}(w|n, p) \text{Uniform}(p|0, 1) dp}$$


Let U denote the domain of the measurable functions p and W , as illustrated:



so that the underlying space U is the Cartesian product of the interval $[0, 1]$, representing all possible proportions of water to land, with the

collection of all N -tuples of *water's* and *land's*. The function W is supposed to count the number of *water's* in an N -tuple of *water's* and *land's*. Let $\mathfrak{D}$ denote a σ -algebra of subsets of U , to be qualified momentarily. Then the parameter space X and sample space Y are the respective codomains, where $X = [0, 1]$ with σ -algebra $\mathfrak{A} = \mathfrak{Bor}_{[0,1]}$, and Y is the set of integers $\{0, \dots, N\}$ with σ -algebra $\mathfrak{B} = \mathfrak{P}(Y)$.

We suppose $(U, \mathfrak{D}, Q)$ is a probability space such that the following diagram commutes:




We can take $\mathfrak{D}$ to be the σ -algebra generated by the product of $[0, 1]$ with the collection of N -tuples of *water's* and *land's*. The probability distributions can be specified, although the specification of the measure Q appears induced, and will not be stressed here:

$$\begin{aligned} W &\sim \text{Binomial}(N, x) = Q \circ W^{\leftarrow} \\ p &\sim \text{Uniform}(0, 1) = Q \circ p^{\leftarrow} \end{aligned}$$

Define $f : X \times Y \rightarrow \mathbb{R}$ by

$$\begin{aligned} f(x, w) &= \frac{N!}{w!(N-w)!} x^w (1-x)^{N-w} \\ &=: \text{Binomial}(w|N, x) \end{aligned}$$

so the following diagram commutes:



Let κ denote counting measure on the sample space σ -algebra $\mathfrak{B}$, and define a family $\{P_x : x \in X\}$ of probability measures on $\mathfrak{B}$ by

$$P_x(B) = \int_{w \in B} f_x(w) d\kappa(w) = \sum_{w \in B} f_x(w) \quad B \in \mathfrak{B}.$$

Let us verify one of the requirements that a measure must satisfy in order to be a probability measure; namely that $P_x(Y) = 1$ for any $x \in X$. This verification is based on the binomial theorem:

$$(a + b)^N = \sum_{w=0}^N \frac{N!}{w!(N-w)!} a^w b^{N-w},$$

taking $a = x$ and $b = 1 - x$.

By definition, each of the sampling probabilities P_x is absolutely continuous with respect to counting measure κ , with $f_x = dP_x/d\kappa$. That is, the function f_x represents the Radon-Nikodym derivative of P_x with respect to κ .

Let α be the probability measure on the parameter space σ -algebra $\mathfrak{A} = \mathfrak{Bor}_{[0,1]}$ represented by the density

$$h(x) = \text{Uniform}(x|0, 1)$$

with respect to Lebesgue measure ν restricted to $\mathfrak{Bor}_{[0,1]}$. In this case, the function h agrees with the constant function 1 on $[0, 1]$, and the

prior α also happens to agree with Lebesgue measure ν on $\mathfrak{Bor}[0, 1]$. Namely,

$$\alpha(A) = \int_A h(x) d\nu(x) \quad \text{for all } A \in \mathfrak{Bor}[0, 1].$$

According to Bayes' theorem, the Radon-Nikodym derivative of the posterior probability α_w with respect to ν has a representative, denoted here by $\Pr(x|w, N) \in \mathcal{L}_1(X, \mathfrak{A}, \nu)$, which may be called the *posterior density*:

$$\Pr(x|w, N) := \frac{f^w(x)h(x)}{\int_X f^w(x)h(x) d\nu(x)} \in \frac{d\alpha_w}{d\nu}.$$

Finally, this can also be written as:

$$\Pr(x|w, N) = \frac{\overbrace{\text{Binomial}(w|N, x)}^{f^w(x)} \overbrace{\text{Uniform}(x|0, 1)}^{h(x)}}{\int_{[0,1]} \underbrace{\text{Binomial}(w|N, x) \text{Uniform}(x|0, 1) dx}_{h(x) d\nu(x)}},$$

which is not difficult to see as a slight translation of

$$\Pr(p|w, n) = \frac{\text{Binomial}(w|n, p) \text{Uniform}(p|0, 1)}{\int \text{Binomial}(w|n, p) \text{Uniform}(p|0, 1) dp}.$$

10.3 Gaussian Model of Height

{gaussian_model}

Another example of an experiment with the joint absolutely continuous with respect to the product of the prior and predictive probability distributions, paraphrased from *Statistical Rethinking, Second Edition* [6]. The experiment is modelling a collection $\{h_i\}$ of heights. Quoting, beginning bottom page 78:

For the moment, we want a single measurement variable to model as a Gaussian distribution. There will be two parameters describing the distribution's shape, the mean μ and the standard deviation σ .

and continuing on page 82:


In most cases, priors are specified independently for each parameter, which amounts to assuming $\Pr(\mu, \sigma) = \Pr(\mu)\Pr(\sigma)$. Then we can write:

$$\begin{aligned} h_i &\sim \text{Normal}(\mu, \sigma) && [\text{likelihood}] \\ \mu &\sim \text{Normal}(178, 20) && [\mu \text{ prior}] \\ \sigma &\sim \text{Uniform}(0, 50) && [\sigma \text{ prior}] \end{aligned}$$


and in a display following, find:

$$\begin{aligned} \Pr(\mu, \sigma | h) &= \\ &= \frac{\prod_i \text{Normal}(h_i | \mu, \sigma) \text{Normal}(\mu | 178, 20) \text{Uniform}(\sigma | 0, 50)}{\int \int \prod_i \text{Normal}(h_i | \mu, \sigma) \text{Normal}(\mu | 178, 20) \text{Uniform}(\sigma | 0, 50) d\mu d\sigma} \end{aligned}$$

Let U denote the domain of the functions (μ, σ) and height h , as illustrated:




so that the underlying space U is the Cartesian product of possible means and standard deviations with n -tuples of heights. We suppose $(U, \mathcal{D}, Q)$ is a probability space such that the following diagram commutes:



Define $f : X \times Y \rightarrow \mathbb{R}$ by

$$f((r, s), t) = \prod_i \frac{1}{s\sqrt{2\pi}} \exp \left\{ -\frac{1}{2} \left(\frac{t_i - r}{s} \right)^2 \right\}$$

so the following diagram commutes:



Let λ_n denote Lebesgue measure on the sample space σ -algebra $\mathfrak{B} = \mathfrak{Bor}_{\mathbb{R}^n}$. Define a family $\{P_{(r,s)} : (r,s) \in X\}$ of probability measures on $\mathfrak{B}$ by defining $P_{(r,s)}$ on the rectangles $\prod_i [a_i, b_i]$ of $\mathfrak{B}$:

$$P_{(r,s)} \left(\prod_i [a_i, b_i] \right) = \prod_i \text{Normal}(r, s)([a_i, b_i])$$

Let us verify one of the requirements that a measure must satisfy in order to be a probability measure; namely that $P_{(r,s)}(Y) = 1$ for any $(r,s) \in X$. This verification is based on the fact that the functions

$$p : \mathbb{R} \rightarrow \mathbb{R}^+ : t \mapsto \frac{1}{s\sqrt{2\pi}} \exp \left\{ -\frac{1}{2} \left(\frac{t - r}{s} \right)^2 \right\}$$

are well-known to be a probability density functions.

Each of the sampling probabilities $P_{(r,s)}$ is absolutely continuous with respect to Lebesgue measure λ_n on $\mathfrak{Bor}_{\mathbb{R}^n}$, and the function $f_{(r,s)}$ represents the Radon-Nikodym derivative of $P_{(r,s)}$ with respect to λ_n .

Define a probability measure ρ on the rectangles $B \times C$ of the parameter space σ -algebra $\mathfrak{A} = \mathfrak{Bor}_{\mathbb{R} \times \mathbb{R}^+}$ by

$$\rho(B \times C) = \text{Normal}(178, 20)(B) \cdot \text{Uniform}(0, 50)(C).$$

See that the prior probability distribution ρ is absolutely continuous with respect to a different Lebesgue measure, call it λ_2 , on $\mathfrak{A} = \mathfrak{Bot}_{\mathbb{R} \times \mathbb{R}^+}$, and let $h \in \mathcal{L}_1(X, \mathfrak{A}, \lambda_2)$ be defined by

$$h : \mathbb{R} \times \mathbb{R}^+ \rightarrow \mathbb{R} : (r, s) \mapsto \text{Normal}(r|178, 20) \cdot \text{Uniform}(s|0, 50)$$

so that the function h represents the Radon-Nikodym derivative $d\rho/d\lambda_2$. According to Bayes' theorem, the Radon-Nikodym derivative of the posterior probability ρ_t with respect to λ_2 has a representative, denoted here by $\text{Pr}(r, s|t) \in \mathcal{L}_1(X, \mathfrak{A}, \lambda)$, where

$$\text{Pr}(r, s|t) = \frac{f^t(r, s) h(r, s)}{\int_X f^t(r, s) h(r, s) d\lambda_2(r, s)}.$$

Finally, write

$$\begin{aligned} f^t(r, s) &= \prod_i \frac{1}{s\sqrt{2\pi}} \exp \left\{ -\frac{1}{2} \left(\frac{t_i - r}{s} \right)^2 \right\} \\ &= \prod_i \text{Normal}(t_i|r, s) \end{aligned}$$

so that

$$\begin{aligned} \text{Pr}(r, s|t) &= \frac{\overbrace{\prod_i \text{Normal}(t_i|r, s)}^{f^t(r, s)} \overbrace{\text{Normal}(r|178, 20) \text{Uniform}(s|0, 50)}^{h(r, s)}}{\int_{\mathbb{R} \times \mathbb{R}^+} \prod_i \text{Normal}(t_i|r, s) \underbrace{\text{Normal}(r|178, 20) \text{Uniform}(s|0, 50) d\lambda_2(r, s)}_{h(r, s) d\lambda_2(r, s)}} \end{aligned}$$

which is our slight translation of

$$\begin{aligned} \text{Pr}(\mu, \sigma|h) &= \\ &= \frac{\prod_i \text{Normal}(h_i|\mu, \sigma) \text{Normal}(\mu|178, 20) \text{Uniform}(\sigma|0, 50)}{\int \int \prod_i \text{Normal}(h_i|\mu, \sigma) \text{Normal}(\mu|178, 20) \text{Uniform}(\sigma|0, 50) d\mu d\sigma}. \end{aligned}$$

10.4 The Linear Model

{linear_model}


The linear model we will use here can be seen as an extension of the Gaussian model in Example 10.3 by including weight, assuming a relation between height and weight. The word *relation* is used in the mathematical sense; it is a subset of a Cartesian product. The model is given in this quote from *Statistical Rethinking*, page 93:

Now how do we get weight into a Gaussian model of height? Let x be the name for the column of weight measurements, `d2$weight`. Let the average of the x values be $\bar{x}$, “ex bar”. Now we have a predictor variable x , which is a list of measures of the same length as h . To get weight into the model, we define the mean μ as a function of the values in x . This is what it looks like, with explanation to follow:

$$\begin{aligned} h_i &\sim \text{Normal}(\mu_i, \sigma) && [\text{likelihood}] \\ \mu_i &= \alpha + \beta(x_i - \bar{x}) && [\text{linear model}] \\ \alpha &\sim \text{Normal}(178, 20) && [\alpha \text{ prior}] \\ \beta &\sim \text{Normal}(0, 10) && [\beta \text{ prior}] \\ \sigma &\sim \text{Uniform}(0, 50) && [\sigma \text{ prior}] \end{aligned}$$

See that this model includes no probability distribution for the weight measurements x_i , but the measurable functions h_i , α , β , and σ do have probability distributions included in the model. We consider the x_i as known values.

Let U denote the domain of the functions (α, β, σ) and h , as illustrated:



We suppose $(U, \mathcal{D}, Q)$ is a probability space such that the following

diagram commutes:

$$\begin{array}{ccc}
 & Q \circ (\alpha, \beta, \sigma)^{\leftarrow} & \mathfrak{A} = \mathfrak{Bot}_{\mathbb{R}^2 \times \mathbb{R}^+} \\
 & \searrow (\alpha, \beta, \sigma)^{\leftarrow} & \downarrow \pi_1^{\leftarrow} \\
 [0, 1] & \xleftarrow{Q} \mathfrak{D} & \sigma(\mathfrak{A}, \mathfrak{B}) \\
 & \nwarrow h^{\leftarrow} & \uparrow \pi_2^{\leftarrow} \\
 & Q \circ h^{\leftarrow} & \mathfrak{B} = \mathfrak{Bot}_{\mathbb{R}^n}
 \end{array}$$

Define $f : X \times Y \rightarrow \mathbb{R}$ by

$$\begin{aligned}
 f((a, b, s), t) &= \prod_i \frac{1}{s\sqrt{2\pi}} \exp \left\{ -\frac{1}{2} \left(\frac{t_i - (a \cdot 1 + b(x_i - \bar{x}))}{s} \right)^2 \right\} \\
 &= \prod_i \frac{1}{s\sqrt{2\pi}} \exp \left\{ -\frac{1}{2} \left(\frac{t_i - \mu_i}{s} \right)^2 \right\} \\
 &= \prod_i \text{Normal}(t_i | \mu_i, s),
 \end{aligned}$$

where $\mu_i = a \cdot 1 + b(x_i - \bar{x})$, so the following diagram commutes:

$$\begin{array}{ccc}
 X & & \\
 \pi_1 \uparrow & \searrow f^t \text{ the likelihood} & \\
 X \times Y & \xrightarrow{f} & \mathbb{R} \\
 \pi_2 \downarrow & \nearrow f_{(a,b,s)} \text{ Radon-Nikodym derivative} & \\
 Y & &
 \end{array}$$

Define a family $\{P_{(a,b,s)} : (a,b,s) \in X\}$ of measures on the sample space σ -algebra $\mathfrak{B} = \mathfrak{Bot}_{\mathbb{R}^n}$ in terms of the indefinite integrals of the collection $\{f_{(a,b,s)} : (a,b,s) \in X\}$, integrated with respect to Lebesgue measure, call it λ_n , on $\mathfrak{Bot}_{\mathbb{R}^n}$;

$$P_{(a,b,s)}(B) = \int_B f_{(a,b,s)} d\lambda_n \quad \text{for all } B \in \mathfrak{Bot}_{\mathbb{R}^n}.$$

Let us verify a requirement that a measure must satisfy in order to be a probability measure; namely that $P_{(a,b,s)}(Y) = 1$ for any $(a, b, s) \in X$. This is based on the fact that the functions

$$p : \mathbb{R} \rightarrow \mathbb{R}^+ : t \mapsto \frac{1}{s\sqrt{2\pi}} \exp \left\{ -\frac{1}{2} \left(\frac{t-r}{s} \right)^2 \right\}$$

are well-known to be a probability density functions.

For the purpose of comparison, see that the sum $a_1^2 + a_2^2 + a_3^2$ can be written as $A^T A$ with $A^T = [a_1 \ a_2 \ a_3]$. Were you to then write $a_i = y_i - (1 \cdot \beta_0 + x_i \cdot \beta_1)$, with $\beta = (\beta_0, \beta_1)$, and with X and Y now functions or vectors or variables or whatever, you would end up with something like:

$$f(\beta, \sigma^2, Y) = (2\pi\sigma^2)^{-n/2} \exp \left\{ -\frac{1}{2\sigma^2} [Y - X\beta]^T [Y - X\beta] \right\},$$

which you could compare with equation (3.1) on page 51 of Marin & Robert, *Bayesian Core*, which says the likelihood of the *ordinary normal linear model* is

$$l(\beta, \sigma^2 | y, X) = (2\pi\sigma^2)^{-n/2} \exp \left\{ -\frac{1}{2\sigma^2} (y - X\beta)^T (y - X\beta) \right\}.$$

The translation between their point of view and ours is not exact. For one thing, our a and b would be part of their β . For another, if the particular data included the y -values at known X values, say when X is equal to X_0 , then our f would relate to their l by something like:

$$f(\beta, \sigma^2, y) = l(\beta, \sigma^2 | y, X) \Big|_{X=X_0}$$

Define a probability measure ρ on the rectangles $A \times B \times C$ of the parameter space σ -algebra $\mathfrak{A} = \mathfrak{Bor}_{\mathbb{R}^2 \times \mathbb{R}^+}$ by

$$\begin{aligned} \rho(A \times B \times C) &= \\ &= \text{Normal}(178, 20)(A) \cdot \text{Normal}(0, 10)(B) \cdot \text{Uniform}(0, 50)(C). \end{aligned}$$

See that the prior probability distribution ρ is absolutely continuous with respect to a different Lebesgue measure, call it λ , on $\mathfrak{A} = \mathfrak{Bor}_{\mathbb{R}^2 \times \mathbb{R}^+}$,

and let $h : \mathbb{R}^2 \times \mathbb{R}^+ \rightarrow \mathbb{R}$ in $\mathcal{L}_1(X, \mathfrak{A}, \lambda)$ be defined by

$$h(a, b, s) = \text{Normal}(a|178, 20) \cdot \text{Normal}(b|0, 20) \cdot \text{Uniform}(s|0, 50)$$

so that h represents the Radon-Nikodym derivative $d\rho/d\lambda$. According to Bayes' theorem, the Radon-Nikodym derivative of the posterior probability ρ_t with respect to λ has a representative, denoted here by $\text{Pr}(a, b, s | t) \in \mathcal{L}_1(X, \mathfrak{A}, \lambda)$, where

$$\text{Pr}(a, b, s | t) = \frac{f^t(a, b, s) h(a, b, s)}{\int_X f^t(a, b, s) h(a, b, s) d\lambda(a, b, s)},$$

with

$$\mu_i = a \cdot 1 + b(x_i - \bar{x})$$

and with

$$f^t(a, b, s) = \prod_i \frac{1}{s\sqrt{2\pi}} \exp \left\{ -\frac{1}{2} \left(\frac{t_i - \mu_i}{s} \right)^2 \right\}.$$

10.5 Logistic Regression

{logistic_regression}

The logit function is defined by

$$\text{logit} : (0, 1) \rightarrow \mathbb{R} : p \mapsto \ln \left(\frac{p}{1-p} \right),$$

and the logistic function is defined by

$$\text{logistic} : \mathbb{R} \rightarrow (0, 1) : a \mapsto \frac{\exp(a)}{1 + \exp(a)}.$$

The logit function is the inverse of the logistic function, that is,

$$\text{logit}(p) = a \quad \Leftrightarrow \quad p = \text{logistic}(a),$$

which says

$$\ln \left(\frac{p}{1-p} \right) = a \quad \Leftrightarrow \quad p = \frac{\exp(a)}{1 + \exp(a)}.$$

The expression $p/(1-p)$ can be seen as the odds ratio.

The logistic regression model we will look at here, quoted from the tentative model displayed in *Statistical Rethinking*, page 327, is:

$$\begin{aligned} L_i &\sim \text{Binomial}(1, p_i) \\ \text{logit}(p_i) &= \alpha_{\text{ACTOR}[i]} + \beta_{\text{TREATMENT}[i]} \\ \alpha_j &\sim \text{to be determined} \\ \beta_k &\sim \text{to be determined} \end{aligned}$$

The “ $\alpha_j \sim \text{to be determined}$ ” will later be $\alpha_j \sim \text{Normal}(0, 1.5)$, and “ $\beta_k \sim \text{to be determined}$ ” will later be $\beta_k \sim \text{Normal}(0, 0.5)$.

Let $(\alpha, \beta) : U \rightarrow X$ and $(L, \text{ACTOR}, \text{TREATMENT}) : U \rightarrow Y$ be maps such that the following diagrams commute:

$$\begin{array}{ccc} & & X \\ & \nearrow^{(\alpha, \beta)} & \\ U & \longrightarrow & X \times Y \\ & \searrow_{(L, \text{ACTOR}, \text{TREATMENT})} & \\ & & Y \end{array} \quad \begin{array}{ccc} & \mathfrak{A} & \\ & \nwarrow^{(\alpha, \beta)} & \\ \mathfrak{D} & \xrightarrow{\sigma(\mathfrak{A}, \mathfrak{B})} & \mathfrak{B} \\ & \swarrow_{(L, \text{ACTOR}, \text{TREATMENT})} & \end{array}$$

$\begin{array}{ccc} & \uparrow \pi_1 & \\ & \downarrow \pi_2 & \\ & & \end{array} \quad \begin{array}{ccc} & \downarrow \pi_1 & \\ & \uparrow \pi_2 & \end{array}$

with

$$X = \mathbb{R}^7 \times \mathbb{R}^4$$

and

$$Y = \{0, 1\}^n \times \{1, \dots, 7\}^n \times \{1, 2, 3, 4\}^n,$$

where $\mathfrak{A} = \mathfrak{Bot}_{\mathbb{R}^7 \times \mathbb{R}^4}$ and $\mathfrak{B} = \mathfrak{P}(Y)$. We are supposing $(U, \mathfrak{D}, Q)$ is a

probability space such that the following diagram commutes:

$$\begin{array}{ccc}
 & & \mathfrak{A} = \mathfrak{Bor}_{\mathbb{R}^7 \times \mathbb{R}^4} \\
 & \swarrow^{Q \circ (\alpha, \beta)^\leftarrow} & \downarrow \pi_1^\leftarrow \\
 [0, 1] & \xleftarrow{Q} \mathfrak{D} & \sigma(\mathfrak{A}, \mathfrak{B}) \\
 & \nwarrow_{(L, \text{etc.})^\leftarrow} & \uparrow \pi_2^\leftarrow \\
 & & \mathfrak{B} = \mathfrak{P}(Y) \\
 & \nwarrow_{Q \circ (L, \text{ACTOR, TREATMENT})^\leftarrow} &
 \end{array}$$

Define $f : X \times Y \rightarrow \mathbb{R}$ by

$$\begin{aligned}
 ((a, b), (q, r, s)) &\mapsto \prod_i (\text{logistic}(a_{r_i} + b_{s_i}))^{q_i} (1 - \text{logistic}(a_{r_i} + b_{s_i}))^{1-q_i} \\
 &= \prod_i \text{Binomial}(q_i | 1, p_i), \quad \text{with } p_i = \text{logistic}(a_{r_i} + b_{s_i})
 \end{aligned}$$

so the following diagram commutes

$$\begin{array}{ccc}
 X & & \\
 \pi_1 \uparrow & \searrow^{f(q, r, s)} \text{the likelihood} & \\
 X \times Y & \xrightarrow{f} & \mathbb{R} \\
 \pi_2 \downarrow & \nearrow_{f(a, b)} \text{Radon-Nikodym derivative} & \\
 Y & &
 \end{array}$$

Define a family $\{P_{(a, b)} : (a, b) \in X\}$ of measures on the sample space σ -algebra $\mathfrak{B} = \mathfrak{P}(Y)$ in terms of the indefinite integrals of the collection $\{f_{(a, b)} : (a, b) \in X\}$, integrated with respect to counting measure on

$\mathfrak{P}(Y)$, calling counting measure δ here;

$$\begin{aligned} P_{(a,b)}(B) &= \int_B f_{(a,b)} d\delta \\ &= \sum_{(q,r,s) \in B} f_{(a,b)}(q, r, s) \\ &= \sum_{(q,r,s) \in B} \prod_i \text{Binomial}(q_i | 1, p_i), \end{aligned}$$

with $p_i = \text{logistic}(a_{r_i} + b_{s_i})$. These measures are probability measures because they are defined in terms of the density

$$\prod_i \text{Binomial}(q_i | 1, p_i)$$

of a probability measure. Roughly.

Define a probability measure ρ on the rectangles $A \times B$ of the parameter space σ -algebra $\mathfrak{A} = \mathfrak{Bor}_{\mathbb{R}^7 \times \mathbb{R}^4}$ by

$$\rho(A \times B) = \text{Normal}(0, 1.5)(A) \cdot \text{Normal}(0, 0.5)(B).$$

The prior probability distribution ρ is absolutely continuous with respect to a Lebesgue measure, call it λ , on $\mathfrak{A} = \mathfrak{Bor}_{\mathbb{R}^7 \times \mathbb{R}^4}$. Let $h \in \mathcal{L}_1(X, \mathfrak{A}, \lambda)$ be defined by

$$h : \mathbb{R}^7 \times \mathbb{R}^4 \rightarrow \mathbb{R} : (a, b) \mapsto \text{Normal}(a | 0, 1.5) \cdot \text{Normal}(b | 0, 0.5).$$

so that h represents the Radon-Nikodym derivative $d\rho/d\lambda$. According to Bayes' theorem, the Radon-Nikodym derivative of the posterior probability $\rho_{(q,r,s)}$ with respect to λ has a representative, denoted here by $\text{Pr}(a, b | q, r, s) \in \mathcal{L}_1(X, \mathfrak{A}, \lambda)$, where

$$\text{Pr}(a, b | q, r, s) = \frac{f^{(q,r,s)}(a, b) h(a, b)}{\int_X f^{(q,r,s)}(a, b) h(a, b) d\lambda(a, b)}.$$

11 Conditional Independence

WE BARELY touch on a few points concerning conditional independence.

{conditionally_independent}

Definition 11.1. Let $(X, \mathfrak{A}, \mu)$ be a probability space, and let $\mathfrak{B}$ be a σ -subalgebra of $\mathfrak{A}$. Also, let $\mathfrak{C}$ and $\mathfrak{D}$ be σ -subalgebras of $\mathfrak{A}$. Then $\mathfrak{C}$ and $\mathfrak{D}$ are **conditionally independent σ -subalgebras** given $\mathfrak{B}$ if

$$\mathcal{P}_{\mathfrak{B}}(C \cap D) = \mathcal{P}_{\mathfrak{B}}(C) \cdot \mathcal{P}_{\mathfrak{B}}(D)$$

for all $C \in \mathfrak{C}$ and for all $D \in \mathfrak{D}$. The notation $\mathfrak{C} \perp\!\!\!\perp \mathfrak{D} \mid \mathfrak{B}$ may be used.

Example 11.2. Let us try to understand conditional independence in two extreme cases, namely

- case $\mathfrak{B} = \{X, \emptyset\}$, where conditional independence given $\mathfrak{B}$ is equivalent to independence with respect to the measure μ^* , and
- case $\mathfrak{B} = \mathfrak{A}$, where all σ -subalgebras of $\mathfrak{A}$ are conditionally independent given $\mathfrak{B}$.

Case $\mathfrak{B} = \{X, \emptyset\}$. Verify that $\{X, \emptyset\}$ is independent of every σ -subalgebra of $\mathfrak{A}$ with respect to μ^* , for if $\mathfrak{T}$ were another σ -subalgebra, we see that

$$\mu^*(A \cap B) = \mu^*(A) \cdot \mu^*(B) \quad \text{for all } A \in \mathfrak{T} \text{ and all } B \in \{X, \emptyset\}.$$

It follows then by Proposition 7.28 that

$$\mathcal{E}_{\mathfrak{B}} \bar{f}^{\mu} = (\mathbf{E} \bar{f}^{\mu}) \cdot \bar{1}^{\mu}$$

for every $\bar{f}^{\mu} \in L_1(X, \mathfrak{A}, \mu)$. For $C \in \mathfrak{C}$ and for $D \in \mathfrak{D}$, the condition

$$\mathcal{P}_{\mathfrak{B}}(C \cap D) = \mathcal{P}_{\mathfrak{B}}(C) \cdot \mathcal{P}_{\mathfrak{B}}(D)$$

is equivalent by definition to

$$\mathcal{E}_{\mathfrak{B}} \overline{\chi_{C \cap D}}^\mu = \mathcal{E}_{\mathfrak{B}} \overline{\chi_C}^\mu \cdot \mathcal{E}_{\mathfrak{B}} \overline{\chi_D}^\mu,$$

which in turn is equivalent to

$$\begin{aligned} (\mathbb{E} \overline{\chi_{C \cap D}}^\mu) \cdot \bar{1}^\mu &= (\mathbb{E} \overline{\chi_C}^\mu) \cdot \bar{1}^\mu \cdot (\mathbb{E} \overline{\chi_D}^\mu) \cdot \bar{1}^\mu \\ &= (\mathbb{E} \overline{\chi_C}^\mu) (\mathbb{E} \overline{\chi_D}^\mu) \cdot \bar{1}^\mu, \end{aligned}$$

implying that

$$\mu^*(C \cap D) = \mu^*(C) \cdot \mu^*(D).$$

This shows that conditional independence given $\mathfrak{B} = \{X, \emptyset\}$ is equivalent to independence with respect to μ^* .

Case $\mathfrak{B} = \mathfrak{A}$. Then for all $C \in \mathfrak{C}$ and for all $D \in \mathfrak{D}$, the functions $\chi_{C \cap D}$, χ_C , and χ_D are $(\mathfrak{B}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable. It follows by Remark 7.2 that the condition

$$\mathcal{E}_{\mathfrak{B}} \overline{\chi_{C \cap D}}^\mu = \mathcal{E}_{\mathfrak{B}} \overline{\chi_C}^\mu \cdot \mathcal{E}_{\mathfrak{B}} \overline{\chi_D}^\mu$$

is equivalent to

$$\overline{\chi_{C \cap D}}^\mu = \overline{\chi_C}^\mu \cdot \overline{\chi_D}^\mu,$$


which is equivalent to

$$\chi_{C \cap D} = \chi_C \cdot \chi_D \quad \mu^*\text{-almost everywhere on } X.$$

This is always true. This shows that all σ -subalgebras of $\mathfrak{A}$ are conditionally independent given $\mathfrak{A}$.

{conditionally_independent_functions}

Notation 11.3. Let $(U, \mathfrak{D}, \lambda)$ be a probability space, and let functions $f : (U, \mathfrak{D}) \rightarrow (X, \mathfrak{A})$, $g : (U, \mathfrak{D}) \rightarrow (Y, \mathfrak{B})$, and $h : (U, \mathfrak{D}) \rightarrow (Z, \mathfrak{C})$ be measurable:



Say that f and g are conditionally independent given h if

$$f^{\leftarrow}(\mathfrak{A}) \perp\!\!\!\perp g^{\leftarrow}(\mathfrak{B}) \mid h^{\leftarrow}(\mathfrak{C}).$$

{abs_cont_reg}

Proposition 11.4. *Let $(U, \mathfrak{D}, \lambda)$ be a probability space, and let functions $f : (U, \mathfrak{D}) \rightarrow (X, \mathfrak{A})$, $g : (U, \mathfrak{D}) \rightarrow (Y, \mathfrak{B})$, and $h : (U, \mathfrak{D}) \rightarrow (Z, \mathfrak{C})$ be measurable, with σ -finite measures μ , ν , and ρ on respective σ -algebras $\mathfrak{A}$, $\mathfrak{B}$, and $\mathfrak{C}$. Set $\mathfrak{H} = h^{\leftarrow}(\mathfrak{C})$. If $\lambda \circ (f, g, h)^{\leftarrow} \ll (\mu \times \nu \times \rho)^*$ on $\sigma(\mathfrak{A}, \mathfrak{B}, \mathfrak{C})$, then the conditional distributions $\mathcal{P}_{\mathfrak{H}} \circ f^{\leftarrow}$, $\mathcal{P}_{\mathfrak{H}} \circ g^{\leftarrow}$, and $\mathcal{P}_{\mathfrak{H}} \circ (f, g)^{\leftarrow}$ are regular.*

Proof. Set $\mathfrak{F} = f^{\leftarrow}(\mathfrak{A})$, and $\mathfrak{G} = g^{\leftarrow}(\mathfrak{B})$.

Verify that if

$$\lambda \circ (f, g, h)^{\leftarrow} \ll (\mu \times \nu \times \rho)^*$$

on $\sigma(\mathfrak{A}, \mathfrak{B}, \mathfrak{C}) = \sigma(\mathfrak{A}, \sigma(\mathfrak{B}, \mathfrak{C}))$, then by applying the proof of Proposition 5.25, you get

$$\lambda \circ (g, h)^{\leftarrow} \ll (\nu \times \rho)^*$$

on $\sigma(\mathfrak{B}, \mathfrak{C})$. In particular, by applying the proof, you show

$$\frac{d(\lambda \circ (g, h)^{\leftarrow})}{d(\nu \times \rho)^*} : Y \times Z \rightarrow \mathbb{R} : (y, z) \mapsto \int_X \frac{d(\lambda \circ (f, g, h)^{\leftarrow})}{d(\mu \times \nu \times \rho)^*}(x, y, z) d\mu(x).$$

Then by Proposition 5.26,

$$\lambda \circ (g, h)^{\leftarrow} \ll ((\lambda \circ g^{\leftarrow}) \times (\lambda \circ h^{\leftarrow}))^*,$$

and so by Proposition 8.26, the conditional distribution $\mathcal{P}_{\mathfrak{H}} \circ g^{\leftarrow}$ is regular.

Symmetrically, the conditional distribution $\mathcal{P}_{\mathfrak{H}} \circ f^{\leftarrow}$ must also be regular.

To show that $\mathcal{P}_{\mathfrak{H}} \circ (f, g)^{\leftarrow}$ is regular, again, by Proposition 5.26,

$$\lambda \circ (f, g, h)^{\leftarrow} \ll ((\lambda \circ (f, g)^{\leftarrow}) \times (\lambda \circ h^{\leftarrow}))^*$$

on $\sigma(\mathfrak{A}, \mathfrak{B}, \mathfrak{C}) = \sigma(\sigma(\mathfrak{A}, \mathfrak{B}), \mathfrak{C})$, and so by Proposition 8.26, the conditional distribution $\mathcal{P}_{\mathfrak{H}} \circ (f, g)^{\leftarrow}$ is regular. ■

Notation 11.5. Let $(U, \mathfrak{D}, \lambda)$ be a probability space, with functions $f : (U, \mathfrak{D}) \rightarrow (X, \mathfrak{A})$, $g : (U, \mathfrak{D}) \rightarrow (Y, \mathfrak{B})$ and $h : (U, \mathfrak{D}) \rightarrow (Z, \mathfrak{C})$ measurable. Let μ , ν , and ρ be σ -finite measures on $\mathfrak{A}$, $\mathfrak{B}$, and $\mathfrak{C}$, respectively, and let $\lambda \circ (f, g, h)^{\leftarrow} \ll (\mu \times \nu \times \rho)^*$ on $\sigma(\mathfrak{A}, \mathfrak{B}, \mathfrak{C})$. Set:

{densities}

$$\begin{aligned} j &= \frac{d(\lambda \circ (f, g, h)^{\leftarrow})}{d(\mu \times \nu \times \rho)^*} : X \times Y \times Z \rightarrow \mathbb{R}, \\ m_{f,h} &= \frac{d(\lambda \circ (f, h)^{\leftarrow})}{d(\mu \times \rho)^*} : X \times Z \rightarrow \mathbb{R}, \\ m_{g,h} &= \frac{d(\lambda \circ (g, h)^{\leftarrow})}{d(\nu \times \rho)^*} : Y \times Z \rightarrow \mathbb{R}, \\ m_h &= \frac{d(\lambda \circ h^{\leftarrow})}{d\rho} : Z \rightarrow \mathbb{R}. \end{aligned}$$

The letter ‘ j ’ was intended to roughly correspond with ‘joint density,’ and likewise the letter ‘ m ’ with ‘marginal density.’ I have seen notation such as ‘ $p(x, y | z)$ ’ being used to try to indicate the quotient $\frac{j(x, y, z)}{m_h(z)}$. Similarly, something like ‘ $p(x | z)$ ’, ‘ $p(y | z)$ ’, and ‘ $p(x | y, z)$ ’ might be used to try to indicate the respective quotients $\frac{m_{f,h}(x, z)}{m_h(z)}$, $\frac{m_{g,h}(y, z)}{m_h(z)}$, and $\frac{j(x, y, z)}{m_{g,h}(y, z)}$. Let us use that notation here.

{conditional_independence_equivalence}

Proposition 11.6. Let $(U, \mathfrak{D}, \lambda)$ be a probability space, and let functions $f : (U, \mathfrak{D}) \rightarrow (X, \mathfrak{A})$, $g : (U, \mathfrak{D}) \rightarrow (Y, \mathfrak{B})$, and $h : (U, \mathfrak{D}) \rightarrow (Z, \mathfrak{C})$ be measurable, with σ -finite measures μ , ν , and ρ on respective σ -algebras $\mathfrak{A}$, $\mathfrak{B}$, and $\mathfrak{C}$. Set $\mathfrak{H} = h^{\leftarrow}(\mathfrak{C})$. Let $\lambda \circ (f, g, h)^{\leftarrow} \ll (\mu \times \nu \times \rho)^*$ on $\sigma(\mathfrak{A}, \mathfrak{B}, \mathfrak{C})$, and let k , l , and m be Markov kernels corresponding to the respective regular conditional distributions $\mathcal{P}_{\mathfrak{H}} \circ f^{\leftarrow}$, $\mathcal{P}_{\mathfrak{H}} \circ g^{\leftarrow}$, and $\mathcal{P}_{\mathfrak{H}} \circ (f, g)^{\leftarrow}$.

The following are equivalent:

1)

$$f^{\leftarrow}(\mathfrak{A}) \perp\!\!\!\perp g^{\leftarrow}(\mathfrak{B}) \mid h^{\leftarrow}(\mathfrak{C}),$$

2)

$$m_u(A \times B) = k_u(A) \cdot l_u(B)$$

 for all $A \in \mathfrak{A}$, $B \in \mathfrak{B}$, and λ^* -almost all $u \in U$,

3)

$$p(x, y \mid z) = p(x \mid z) \cdot p(y \mid z)$$

 for $(\mu \times \nu)^*$ -almost all $(x, y) \in X \times Y$ and $(\lambda \circ h^\leftarrow)$ -almost all $z \in Z$,
and

4)

$$p(x \mid y, z) = p(x \mid z)$$

 for $(\mu \times \nu)^*$ -almost all $(x, y) \in X \times Y$ and $(\lambda \circ h^\leftarrow)$ -almost all $z \in Z$.

Proof. Set $\mathfrak{F} = f^\leftarrow(\mathfrak{A})$, and $\mathfrak{G} = g^\leftarrow(\mathfrak{B})$. We know the conditional distributions are regular by Proposition 11.4, so they do have corresponding Markov kernels by Proposition 8.21. The fact that (1) $\Leftrightarrow$ (2) is nothing more than wading through a definition or two.

(1) $\Rightarrow$ (2) Let $f^\leftarrow(\mathfrak{A}) \perp\!\!\!\perp g^\leftarrow(\mathfrak{B}) \mid h^\leftarrow(\mathfrak{C})$, or using $\mathfrak{H}$ in place of $h^\leftarrow(\mathfrak{C})$,

$$f^\leftarrow(\mathfrak{A}) \perp\!\!\!\perp g^\leftarrow(\mathfrak{B}) \mid \mathfrak{H}.$$

By the very Definition 11.1 of conditionally independent σ -subalgebras, this is equivalent to saying

$$\mathcal{P}_{\mathfrak{H}}(f^\leftarrow(A) \cap g^\leftarrow(B)) = \mathcal{P}_{\mathfrak{H}}(f^\leftarrow(A)) \cdot \mathcal{P}_{\mathfrak{H}}(g^\leftarrow(B))$$

for all $A \in \mathfrak{A}$ and for all $B \in \mathfrak{B}$. This is an equality of classes in $L_1(U, \mathfrak{D}, \lambda)$, and can be rewritten as

$$(11.1) \quad \mathcal{P}_{\mathfrak{H}} \circ (f, g)^\leftarrow(A \times B) = \mathcal{P}_{\mathfrak{H}} \circ f^\leftarrow(A) \cdot \mathcal{P}_{\mathfrak{H}} \circ g^\leftarrow(B).$$

Recall that two classes in $L_1(U, \mathfrak{D}, \lambda)$ are equal precisely when they have representatives that are equal for λ^* -almost all u in U . Such representatives in this case are easy to find since the corresponding Markov kernels have the property that:

$$m^{A \times B} \in \mathcal{P}_{\mathfrak{H}} \circ (f, g)^\leftarrow(A \times B),$$

$$k^A \in \mathcal{P}_{\mathfrak{H}} \circ f^{\leftarrow}(A),$$

and

$$l^B \in \mathcal{P}_{\mathfrak{H}} \circ g^{\leftarrow}(B).$$

It follows that

$$m^{A \times B}(u) = k^A(u) \cdot l^B(u) \quad \text{for } \lambda^*\text{-almost all } u \in U.$$

Since k , l , and m are Markov kernels,

$$m_u(A \times B) = k_u(A) \cdot l_u(B)$$

for all $A \in \mathfrak{A}$, $B \in \mathfrak{B}$, and for λ^* -almost all $u \in U$, as required. ■

(1) $\Leftarrow$ (2) Let

$$m_u(A \times B) = k_u(A) \cdot l_u(B)$$

for all $A \in \mathfrak{A}$, $B \in \mathfrak{B}$, and λ^* -almost all $u \in U$. Since k , l , and m are Markov kernels, so equalities like

$$m_u(A \times B) = m^{A \times B}(u)$$

hold, it follows that

$$m^{A \times B}(u) = k^A(u) \cdot l^B(u)$$

for all $A \in \mathfrak{A}$, $B \in \mathfrak{B}$, and for λ^* -almost all $u \in U$. Check that the function $m^{A \times B}$ is bounded, and is $(\mathfrak{H}, \mathfrak{B}_{\text{ot}\mathbb{R}})$ -measurable, implying it is λ^* -measurable, so by Proposition 3.38 is integrable with respect to λ . It follows that the function $k^A \cdot l^B$ is integrable with respect to λ , even by the very Definition 3.34 of integrability. Then according to the Definition 4.1 of the product of classes,

$$\begin{aligned} \overline{k^A \cdot l^B} &:= \overline{k^A \cdot l^B} \\ &= \overline{m^{A \times B}}. \end{aligned}$$

Because these Markov kernels correspond to their regular conditional distributions, and so for example $\overline{k^A} = \mathcal{P}_{\mathfrak{H}} \circ f^{\leftarrow}(A)$, it follows from the equality

$$\overline{k^A} \cdot \overline{l^B} = \overline{m^{A \times B}}$$

that

$$\mathcal{P}_{\mathfrak{H}} \circ f^{\leftarrow}(A) \cdot \mathcal{P}_{\mathfrak{H}} \circ g^{\leftarrow}(B) = \mathcal{P}_{\mathfrak{H}} \circ (f, g)^{\leftarrow}(A \times B),$$

which is equivalent to $f^{\leftarrow}(\mathfrak{A}) \perp\!\!\!\perp g^{\leftarrow}(\mathfrak{B}) \mid h^{\leftarrow}(\mathfrak{C})$, as required. $\blacksquare$

(2) $\Rightarrow$ (3) Let

$$m_u(A \times B) = k_u(A) \cdot l_u(B)$$

for all $A \in \mathfrak{A}$, $B \in \mathfrak{B}$, and λ^* -almost all $u \in U$. By Proposition 8.31,

$$\begin{aligned} m_u(A \times B) &= \int_{A \times B} p(x, y \mid z) d(\mu \times \nu)(x, y), \\ k_u(A) &= \int_A p(x \mid z) d\mu(x), \\ l_u(B) &= \int_B p(y \mid z) d\nu(y). \end{aligned}$$

Since $m_u(A \times B) = k_u(A) \cdot l_u(B)$, it follows by substitution that

$$\int_{A \times B} p(x, y \mid z) d(\mu \times \nu)(x, y) = \int_A p(x \mid z) d\mu(x) \cdot \int_B p(y \mid z) d\nu(y)$$

And by Fubini's Theorem 5.20,

$$\int_A p(x \mid z) d\mu(x) \cdot \int_B p(y \mid z) d\nu(y) = \int_{A \times B} p(x \mid z) \cdot p(y \mid z) d(\mu \times \nu)(x, y).$$

By transitivity of equality,

$$\int_{A \times B} p(x, y \mid z) d(\mu \times \nu)(x, y) = \int_{A \times B} p(x \mid z) \cdot p(y \mid z) d(\mu \times \nu)(x, y)$$

for all $A \times B$ in the semiring $\sigma(\mathfrak{A}, \mathfrak{B})$, and so by Proposition 3.41,

$$p(x, y \mid z) = p(x \mid z) \cdot p(y \mid z)$$

for $(\mu \times \nu)^*$ -almost all (x, y) in $X \times Y$. (But what is the qualifier on z ? Likely γ^* -almost all $z \in Z$.) Although it is true that $\sigma(\mathfrak{A}, \mathfrak{B})$ is a σ -algebra, it is regarded here as a semiring, for that is all the structure which is actually needed.

(2) $\Leftarrow$ (3) Let

$$p(x, y \mid z) = p(x \mid z) \cdot p(y \mid z)$$

for $(\mu \times \nu)^*$ -almost all (x, y) in $X \times Y$, and for γ^* -almost all $z \in Z$.

(3) $\Leftrightarrow$ (4) Divide both sides of (3) by $p(y \mid z)$. ■

12 Expectation Operators as Projections

AN ORTHOGONAL projection on a Hilbert space is a linear idempotent whose kernel is orthogonal to its range. There is a one-to-one correspondence between the closed subspaces of a Hilbert space and its orthogonal projections. Let $\mathfrak{H}$ be a Hilbert space, and let $\mathfrak{M}$ be a closed subspace of $\mathfrak{H}$. To each $x \in \mathfrak{H}$ there corresponds a unique element of $\mathfrak{M}$, denoted here by $\mathfrak{P}\text{roj}(x)$, such that $x - \mathfrak{P}\text{roj}(x) \perp \mathfrak{M}$. The map $\mathfrak{H} \rightarrow \mathfrak{H}$ defined by $x \mapsto \mathfrak{P}\text{roj}(x)$ is called the orthogonal projection of $\mathfrak{H}$ onto $\mathfrak{M}$ along $\mathfrak{M}^\perp$, or simply the *orthogonal projection of $\mathfrak{H}$ onto $\mathfrak{M}$* .

Orthogonal projections are contractive in the sense that any nonzero orthogonal projection on a Hilbert space has norm equal to 1. For x and y in $\mathfrak{H}$, we let $\langle x, y \rangle$ denote their inner product.

Proposition 12.1. *Let $\mathfrak{H}$ be a Hilbert space, and let $\mathfrak{M}$ be a closed subspace of $\mathfrak{H}$, where $\mathfrak{P}\text{roj} : \mathfrak{H} \rightarrow \mathfrak{H}$ is the orthogonal projection of $\mathfrak{H}$ onto $\mathfrak{M}$. Fix $x \in \mathfrak{H}$. Then $\mathfrak{P}\text{roj}(x)$ is that element of $\mathfrak{M}$ which is closest to x , meaning that*

$$\|x - \mathfrak{P}\text{roj}(x)\| < \|x - y\|$$

for all y in the subspace $\mathfrak{M}$ such that $\mathfrak{P}\text{roj}(x) \neq y$.

Proposition 12.2. *Let $\{e_1, e_2, \dots, e_n\}$ be an orthonormal set in a Hilbert space $\mathfrak{H}$, and let $\mathfrak{P}\text{roj} : \mathfrak{H} \rightarrow \mathfrak{H}$ be the orthogonal projection of $\mathfrak{H}$ onto the closed subspace $\mathfrak{Span}\{e_1, e_2, \dots, e_n\}$. Fix $x \in \mathfrak{H}$. Then $\mathfrak{P}\text{roj}(x) = \sum_{i=1}^n \langle x, e_i \rangle e_i$. That is,*

$$\left\| x - \sum_{i=1}^n a_i e_i \right\| < \|x - y\|$$

for all y in the $\mathfrak{Span}\{e_1, e_2, \dots, e_n\}$ such that $\mathfrak{P}\text{roj}(x) \neq y$ if and only if each scalar a_i equals the Fourier coefficient $\langle x, e_i \rangle$ of x .

12.1 Expectation as Projection

Remark 12.3. Some implications may be used without much fanfare; namely, if the classes $\bar{f}^\mu$ and $\bar{g}^\mu$ are in $L_2(X, \mathfrak{A}, \mu)$, then the product of classes $\bar{f}^\mu \cdot \bar{g}^\mu := \overline{fg}^\mu$ is in $L_1(X, \mathfrak{A}, \mu)$. This is a routine application of the Cauchy-Schwarz inequality, considered to be the 2 + 2 of higher mathematics.

{projection_proposition}

Proposition 12.4. *Let $(X, \mathfrak{A}, \mu)$ be a probability space, and let $\mathfrak{B}$ be a σ -subalgebra of $\mathfrak{A}$. The conditional expectation operator $\mathcal{E}_{\mathfrak{B}}$ restricted to $L_2(X, \mathfrak{A}, \mu)$ is the unique orthogonal projection of the Hilbert space $L_2(X, \mathfrak{A}, \mu)$ onto the closed subspace $\mathfrak{I}(L_2(X, \mathfrak{B}, \mu|_{\mathfrak{B}}))$.*

Proof. Let $\mathfrak{M}$ denote the subspace $\mathfrak{I}(L_2(X, \mathfrak{B}, \mu|_{\mathfrak{B}}))$. The elements of $\mathfrak{M}$ are the classes of $L_2(X, \mathfrak{A}, \mu)$ which have a $(\mathfrak{B}, \mathfrak{Bot}_{\mathbb{R}})$ -measurable representative.

We will show that to each $\bar{f}^\mu \in L_2(X, \mathfrak{A}, \mu)$ there corresponds a unique element of $\mathfrak{M}$, namely $\mathcal{E}_{\mathfrak{B}} \bar{f}^\mu$, such that $\bar{f}^\mu - \mathcal{E}_{\mathfrak{B}} \bar{f}^\mu \perp \bar{h}^\mu$ for all $\bar{h}^\mu \in \mathfrak{M}$, for then the map $\bar{f}^\mu \mapsto \mathcal{E}_{\mathfrak{B}} \bar{f}^\mu$ is the orthogonal projection of $L_2(X, \mathfrak{A}, \mu)$ onto $\mathfrak{M}$, per the first paragraph of this chapter.

Let $\bar{f}^\mu \in L_2(X, \mathfrak{A}, \mu)$, and let $\bar{h}^\mu \in \mathfrak{M}$. We must show that

$$\langle \bar{f}^\mu - \mathcal{E}_{\mathfrak{B}} \bar{f}^\mu, \bar{h}^\mu \rangle = 0,$$

which is equivalent to showing that

$$\langle \bar{f}^\mu, \bar{h}^\mu \rangle = \langle \mathcal{E}_{\mathfrak{B}} \bar{f}^\mu, \bar{h}^\mu \rangle,$$

and in turn, this means showing that

$$\mathbb{E}(\bar{f}^\mu \cdot \bar{h}^\mu) = \mathbb{E}((\mathcal{E}_{\mathfrak{B}} \bar{f}^\mu) \cdot \bar{h}^\mu).$$

Stop work here. Go back and finish Proposition 7.32 first.

We will apply Proposition 7.32, and in order to do this, check first that $\bar{f}^\mu \in L_1(X, \mathfrak{A}, \mu)$, and this is because $f^\mu \in L_2(X, \mathfrak{A}, \mu)$, and because

$L_2(X, \mathfrak{A}, \mu) \subseteq L_1(X, \mathfrak{A}, \mu)$ since the measure space is finite. Next check that h is $(\mathfrak{B}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable since $\bar{h}^\mu \in L_2(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$. Finally check that $\bar{f}^\mu \cdot \bar{h}^\mu \in L_1(X, \mathfrak{A}, \mu)$ by Cauchy-Schwarz. Therefore, since $\bar{f}^\mu \in L_1(X, \mathfrak{A}, \mu)$ and $h : X \rightarrow \mathbb{R}$ is $(\mathfrak{B}, \mathfrak{Bor}_{\mathbb{R}})$ -measurable with $\bar{f}^\mu \cdot \bar{h}^\mu \in L_1(X, \mathfrak{A}, \mu)$, it follows by Proposition 7.32 that $E(\bar{f}^\mu \cdot \bar{h}^\mu) = E(\bar{h}^\mu \cdot \mathcal{E}_{\mathfrak{B}} \bar{f}^\mu)$, as required.

We now show the uniqueness part; namely we show that if $f \in L_2(X, \mathfrak{A}, \mu)$, then $\mathcal{E}_{\mathfrak{B}} f$ is the unique element of $L_2(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$ such that $f - \mathcal{E}_{\mathfrak{B}} f \perp \bar{h}^\mu$ for all $\bar{h}^\mu \in L_2(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$. Let $f \in L_2(X, \mathfrak{A}, \mu)$, and suppose that there is a $g \in L_2(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$ having the property that $f - g \perp \bar{h}^\mu$ for all $\bar{h}^\mu \in L_2(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$. We want to see that this implies g must equal $\mathcal{E}_{\mathfrak{B}} f$. To say that

$$f - g \perp \bar{h}^\mu \quad \text{for all } \bar{h}^\mu \in L_2(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$$

means saying that

$$\langle f - g, \bar{h}^\mu \rangle = 0 \quad \text{for all } \bar{h}^\mu \in L_2(X, \mathfrak{B}, \mu|_{\mathfrak{B}}),$$

or equivalently that

$$\langle f, \bar{h}^\mu \rangle = \langle g, \bar{h}^\mu \rangle \quad \text{for all } \bar{h}^\mu \in L_2(X, \mathfrak{B}, \mu|_{\mathfrak{B}}).$$

In terms of integrals, this says

$$\int_X f \bar{h}^\mu d\mu = \int_X g \bar{h}^\mu d\mu \quad \text{for all } \bar{h}^\mu \in L_2(X, \mathfrak{B}, \mu|_{\mathfrak{B}}).$$

In particular, for every $B \in \mathfrak{B}$, the characteristic function χ_B is in $L_2(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$, and so

$$\int_X f \chi_B d\mu = \int_X g \chi_B d\mu \quad \text{for all } B \in \mathfrak{B}.$$

Since $g \chi_B \in L_1(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$ for every $B \in \mathfrak{B}$, by Cauchy-Schwarz, it follows by Proposition 4.14 that

$$\int_X g \chi_B d\mu = \int_X g \chi_B d\mu|_{\mathfrak{B}} \quad \text{for all } B \in \mathfrak{B}.$$

Consequently,

$$\int_X f \chi_B d\mu = \int_X g \chi_B d\mu|_{\mathfrak{B}} \quad \text{for all } B \in \mathfrak{B},$$

which is the same as saying that

$$\int_B f d\mu = \int_B g d\mu|_{\mathfrak{B}} \quad \text{for all } B \in \mathfrak{B}.$$

This means that $g = \mathcal{E}_{\mathfrak{B}} f$; in fact, this is the defining feature of $\mathcal{E}_{\mathfrak{B}} f$, which completes the proof. $\blacksquare$

Example 12.5. Let $(X, \mathfrak{A}, \mu)$ be a probability space, and let $\mathfrak{B}$ be the smallest σ -subalgebra of $\mathfrak{A}$, namely $\{X, \emptyset\}$. Then the conditional expectation operator $\mathcal{E}_{\mathfrak{B}} : L_1(X, \mathfrak{A}, \mu) \rightarrow L_1(X, \mathfrak{A}, \mu)$ is equal to the expected value operator E times the class containing the constant function 1_X on $L_1(X, \mathfrak{A}, \mu)$, meaning $\mathcal{E}_{\mathfrak{B}}(\bar{g}^\mu) = E \bar{g}^\mu \cdot \overline{1_X}^\mu$ for all $\bar{g}^\mu \in L_1(X, \mathfrak{A}, \mu)$. In this case, the closed subspace $L_2(X, \mathfrak{B}, \mu|_{\mathfrak{B}})$ consists of all classes that are constant $\mu|_{\mathfrak{B}}^*$ -almost everywhere on X . See Discussion 7.29.

13 Acknowledgements

- Most of the measure and integration theory here came from *Principles of Real Analysis* [2], Aliprantis and Burkinshaw.
- Conditional distribution section started with Loeve, *Probability Theory* [5].
- Bayesian statistics section began as an elaboration of sections 1.2.1 and 1.2.2 in *Elements of Bayesian Statistics* [3], Florens, Mouchart, and Rolin.
- The Bayesian statistics examples came from *Statistical Rethinking* [6], Richard McElreath.
- Abstract algebra can be found in any elementary algebra text. Hungerford, say.

14 Afterword

Initially, when trying to learn statistics, I had been trying to understand regression models, and had a background in functional analysis and operator theory, but little in the way of statistics. The hopeful outcomes at the start were:

1. Understand Kolmogorov's *conditional mathematical expectation*.
2. Explain regression models to myself in a clear way.
3. Make real mathematical sense out of the bias-variance tradeoff.

It turns out that conditional mathematical expectation is an application of the Radon-Nikodym theorem. Regression models are nothing but projections. And there is a decomposition of conditional expectation into a conditional variance and a squared conditional bias because conditional mathematical expectation is a so-called averaging operator. Not that I was disappointed.

Bibliography

- [1] Cheng-Shang Chang. Understanding Conditional Expectation via Vector Projection. 2008. You might try <https://www.ee.nthu.edu.tw/cschang/>.
- [2] Charalambos D. Aliprantis and Owen Burkinshaw. *Principle of Real Analysis*. Academic Press, Second edition, 1990.
- [3] Jean-Pierre Florens, Michel Mouchart, and Jean-Marie Rolin. *Elements of Bayesian Statistics*. Dekker, 1990.
- [4] A. N. Kolmogorov. *Foundations of the Theory of Probability*. Chelsea, 1956.
- [5] Michel Loeve. *Probability Theory*. Van Nostrand, Second edition, 1960.
- [6] Richard McElreath. *Statistical Rethinking*. CRC Press, Second edition, 2020.

Index

- absolutely continuous, 33
- algebra of sets, 4
- atom, 72

- Bayes' Theorem, 116
- Borel subsets, 9

- change of variable, 26
- codomain, 9
- complete measure space, 15
- conditional expectation given a function, 66
- conditional expectation operator, 68
- conditional expectation with respect to a σ -algebra, 64
- conditional probability, 83
- conditional probability distribution, 93
- conditionally independent σ -subalgebras, 133
- coordinate projections, 42
- corresponding Markov kernel, 98

- data, 117

- defines a Markov kernel, 85
- defines an integrable function, 19
- density, 57
- density function, 34
- distribution, 56
- domain, 9
- Doob-Dynkin lemma, 36, 37

- expected value, 58

- finite measure space, 6
- Fubini's, 44

- image, 9
- indefinite integral, 33
- independent classes, 59
- independent collections, 59
- independent functions, 59
- information, 117
- integrable function, 19
- integral, 18
- iterated integral, 44
- iterated integral exists, 44

- joint measure, 45

-
- $L_p(X, \mathfrak{S}, \mu)$, 22
 - marginal measure, 45
 - Markov kernel, 85
 - measurable function, 10
 - measurable space, 9
 - measure, 3
 - measure induced by a function, 10
 - measure space, 3
 - μ -almost all, 5
 - μ -almost everywhere, 5
 - μ -measurable set, 4
 - μ null set, 5
 - observation, 117
 - outer measure, 4
 - outer measure generated by μ , 5
 - posterior conditional distribution, 114
 - posterior probabilities, 116
 - preimage, 9
 - probability distribution, 56
 - probability measure, 55
 - probability space, 55
 - product of classes in L_1 , 23
 - product semiring, 39
 - product σ -algebra, 41
 - Radon-Nikodym derivative, 34
 - random variable, 55
 - range, 9
 - reference measure, 45
 - regression function, 67
 - regular conditional probability, 89
 - regular conditional probability distribution, 96
 - regular version, 89
 - ring of sets, 4
 - sample, 117
 - sampling conditional distribution, 114
 - sampling probabilities, 115
 - section functions, 43
 - semiring of sets, 3
 - σ -algebra, 4
 - σ -algebra generated by a collection, 4
 - σ -algebra induced by a function, 35
 - σ -finite, 6
 - σ -subalgebra, 4
 - simple function, 16
 - step function, 17
 - strong law of large numbers, 60
 - Tonelli's, 44
 - transition, 85
 - upper function, 18
 - version, 88
 - x -section, 42
 - y -section, 42